Influence of climate and environment on the efficacy of water, sanitation, and handwashing interventions on diarrheal disease in rural Bangladesh: a re-analysis of a randomized control trial

Anna T. Nguyen[1], Jessica A. Grembi[2], Marie Riviere[1], Gabriella Barratt Heitmann[1], William D. Hutson[3], Tejas S. Athni[1], Arusha Patil[1], Ayse Ercumen[4], Audrie Lin[5], Yoshika Crider[6], Andrew Mertens[5], Leanne Unicomb[7], Mahbubur Rahman[7], John M. Colford Jr.[5], Stephen P. Luby[2], Benjamin F. Arnold[8], Jade Benjamin-Chung[1,9]

[1] Department of Epidemiology and Population Health, Stanford University, Stanford, CA, United States
[2] Division of Infectious Diseases and Geographic Medicine, Stanford University, Stanford, CA, United States
[3] Brown School, Washington University in St. Louis, St. Louis, MO, USA.
[4] Department of Forestry and Environmental Resources, North Carolina State University, Raleigh, NC, United States
[5] Division of Epidemiology and Biostatistics, University of California, Berkeley, Berkeley, CA, United States
[6] King Center on Global Development, Stanford University, Stanford, CA, United States
[7] International Centre for Diarrhoeal Disease Research, Dhaka, Bangladesh
[8] Francis I. Proctor Foundation, University of California, San Francisco, San Francisco, CA, United States
[9] Chan Zuckerberg Biohub, San Francisco, CA 94158

Abstract

Background: Climate change may influence the effectiveness of environmental interventions. We investigated if climate and environment modified the effect of low-cost, point-of-use water, sanitation, and handwashing (WASH) interventions on diarrhea and predicted intervention effectiveness under climate change scenarios.

Methods: We analyzed data from a cluster-randomized trial in rural Bangladesh that measured diarrhea prevalence in children 0-2 years from 2012-2016. We matched remote sensing data on temperature, precipitation, humidity, and surface water to households by location and measurement date. We estimated prevalence ratios (PR) for WASH interventions vs. control stratified by environmental factors using generative additive models and targeted maximum likelihood estimation. We estimated intervention effects under predicted precipitation in the study region in 2050 for climate change scenarios from different Shared Socioeconomic Pathways (SSPs).

Findings: WASH interventions more effectively prevented diarrhea under higher levels of total precipitation in the previous week and when there was heavy rain in the previous week (heavy rainfall PR = 0.38, 95% CI 0.23-0.62 vs. no heavy rainfall PR = 0.77, 0.60-0.98). We did not detect substantial effect modification by other environmental variables. WASH intervention effectiveness increased under most climate change scenarios; in a fossil-fueled development scenario (SSP5), the PR was 0.46 (0.44-0.48) compared to 0.67 (0.65-0.68) in the study.

Interpretation: WASH interventions had the strongest effect on diarrhea under higher precipitation, and effectiveness may increase under climate change without sustainable development. WASH interventions may improve population resilience to climate-related health risks.

Funding: Bill & Melinda Gates Foundation, National Institute of Allergy and Infectious Diseases, National Heart, Lung, And Blood Institute

Research in Context Panel

Evidence before this study

We searched Google Scholar using the search terms "sanitation" OR "hygiene" OR "WASH" OR "water quality"; and "heterogen*" or "effect modif*"; and "climate" OR "environmental"; and "diarrhea" OR "enteric infection"; and "risk" AND/OR "factors". In general, the effect modification of sanitation and hygiene on diarrhea or enteropathogen infection by climate and environment is not well studied. One study in Ecuador investigated different relationships between rainfall, diarrhea, and unimproved sanitation and water sources. They found that poor sanitation was most strongly associated with elevated diarrhea after precipitation minimums, whereas poor water sources were most strongly associated with elevated diarrhea after precipitation maximums. One study in Rwanda also found that high levels of runoff were protective against diarrhea only in households with unimproved toilets. In Bangladesh, one study found that access to tubewells was most effective at reducing childhood diarrhea in non-flood controlled areas. No prior studies have projected changes in WASH effectiveness under different climate change scenarios.

Added value of this study

To our knowledge, this is the first study to assess differences in WASH intervention effectiveness for multiple environmental variables and the first to predict future intervention effectiveness under different climate change scenarios. We spatiotemporally matched individual-level data from a trial in rural Bangladesh to remote sensing data on temperature, precipitation, humidity, and surface water presence. We compared the effectiveness of WASH interventions to prevent diarrhea under different levels of potential environmental effect modifiers. We then used

climate models with high spatiotemporal resolution to predict changes in intervention

effectiveness under different climate change scenarios.

Implications of all the available evidence

We found that WASH interventions were more effective under increasing precipitation

and following heavy rain (over 80th percentile) events. Prior studies have found that diarrhea

tends to increase with rainfall. Several studies have documented that precipitation-associated

increases in diarrhea occurred primarily after heavy rain that followed dry periods, but we found

that WASH intervention effectiveness in preventing diarrhea was higher even at moderate

precipitation levels that were not necessarily preceded by a dry period. Under climate change

scenarios with higher precipitation, we projected that WASH interventions may become more

effective. In settings similar to rural Bangladesh, our findings suggest that regions that are

expected to experience greater rainfall should be prioritized for intervention installation,

promotion, and maintenance and that these activities might be most impactful during the rainy

season.

Background

In 2019, the Institute for Health Metrics and Evaluation estimated that over 500,000 deaths in children under the age of five were caused by diarrheal disease.[1] Children that suffer from repeated diarrheal episodes are at high risk of malnutrition, stunted growth, and impaired cognitive development.[2] The World Health Organization estimates that over half of diarrhea deaths are directly attributable to inadequate water safety, sanitation, and/or handwashing (WASH).[3]

However, randomized controlled trials of WASH interventions in rural Kenya and Bangladesh found varying effects in these quite different contexts; the trial in Bangladesh found a 1.7 percentage point reduction in diarrhea prevalence among children who received a combined water, sanitation, and hygiene intervention but the trial in Kenya found no significant reduction.[4,5] The unique physical, cultural, and environmental conditions of each study site may explain differences in treatment effects between these two trials.

Diarrheal disease is associated with meteorological variables such as temperature[6–9], precipitation[8–11], and humidity[7]. Other environmental factors, including proximity to surface water[12], have also been shown to affect enteric pathogen transmission. There are multiple plausible pathways through which climate and environmental conditions could impact the relationship between WASH interventions and diarrhea. Extreme precipitation and flooding can damage WASH infrastructure, leading to increased drinking water contamination.[13] Severe drought conditions might reduce the availability of well water for drinking and handwashing.[14]

Few studies have examined how climate and environmental factors modify the effect of WASH interventions on diarrhea. There is some evidence that the relationships between diarrhea prevalence and unimproved water sources and sanitation systems changed under different levels

of precipitation and runoff.[10,11] In Bangladesh, one study found that tubewells were most

effective in non-flood controlled areas.[15] Another study found that reductions in diarrhea from

sanitation interventions occurred exclusively during the rainy season in Bangladesh.[16] However,

no prior work has investigated whether the effectiveness of improved WASH varies under

multiple climatic and environmental conditions in a scenario where interventions were

randomized.

Our objective was to assess whether climatic and environmental factors modify the effect

of water, sanitation, and hygiene interventions on diarrhea prevalence. We analyzed individual-

level data from a randomized trial in rural Bangladesh that measured childhood diarrhea and

spatiotemporal environmental data from remote sensors.[5] We predicted potential changes in the

effectiveness of WASH interventions under different climate change scenarios.

Methods

<u>Study Data</u>

The WASH Benefits Bangladesh trial delivered low-cost, point-of-use household WASH

interventions and estimated their effects on diarrheal disease.[5,17] The trial enrolled pregnant

women in their second or third trimester between 2012-2013 in the Gazipur, Kishoreganj,

Mymensingh, and Tangail districts of rural Bangladesh. Village clusters were block-randomized

to one of the following arms: chlorinated drinking water (W); upgraded sanitation (S); promotion

of handwashing with soap (H); nutrition education with lipid-based supplement (Nutrition);

combined water, sanitation, and handwashing (WSH); combined water, sanitation, handwashing,

and nutrition (WSH + Nutrition); or control. We excluded the Nutrition arm from this analysis

since the effectiveness of nutrition alone is unlikely to depend on environmental factors.

Additional details on enrollment criteria and intervention details have been reported elsewhere.[5]

We restricted our analysis to children in the diarrhea cohort, which included children

born to pregnant women who enrolled in the trial and children living in the same compound that

were under the age of three at the time of enrollment. Diarrhea was measured at approximately 1

and 2 years after intervention delivery through caregiver report of at least three loose stools in

24-hours or a bloody stool within the past 7 days.

<u>Environmental Data</u>

We assessed pre-specified climatic and environmental variables as possible effect

modifiers (see details on data sources and variable definitions in Appendix 1). We matched

values of environmental factors to each child in the trial by finding the closest measurement to

their household coordinates at the date that outcomes were assessed. We calculated distances

between sensor measurements and study coordinates using the haversine formula.

Temperature We pulled temperature data from the MODIS/Terra Land Surface Temperature

(MOD11A1 v006) dataset from the National Aeronautics and Space Administration (NASA).

We assessed temperature using the minimum, maximum, and average temperature in the 7-day

and 30-day period preceding each outcome measurement. Additionally, we computed antecedent

7-day averages under a 1-, 2-, and 3-week lag from outcome measurements and calculated binary

variables of whether these averages were above or below the median weekly average

temperatures across the study period.

Surface Water We obtained surface water data from the Global Surface Water Explorer dataset

from the European Commission's Joint Research Centre. We quantified the amount of adjacent

surface water by calculating the proportion of area (measured by 30x30 meter pixels) covered by

seasonal surface water (consistently present during a specific time of the year), ephemeral

surface water (not consistently present), and any surface water (seasonal, ephemeral, and

permanent surface water) within a 250-, 500-, and 750-meter radius of households. We

constructed categorical variables to indicate if the proportion of area covered by each type of

surface water was greater or less than the median proportion in the study area. Additionally, we

calculated the distance (in meters) from each household to the closest detected surface water,

seasonal surface water, and ephemeral surface water. We constructed a categorical variable

indicating the tertile of distance from each type of surface water to address data sparsity in the

continuous measures of distance.

Vapor Pressure Deficit We obtained vapor pressure deficit (VPD) data from the TerraClimate

dataset and matched outcome measurements to the closest measurement taken in the same month

and year, as the data was only available at a monthly resolution. Higher values of VPD indicate

lower humidity.

Precipitation We obtained precipitation data from the Global Unified Gauge-Based Analysis of

Daily Precipitation dataset from the National Oceanic and Atmospheric Administration. We

assessed precipitation through antecedent 7-day sums under 1-, 2-, and 3-week lags of outcome

measurements. We created binary variables indicating if these weekly sums exceeded the median

weekly total precipitation over the study period. We also calculated binary variables of heavy

rain under 1-, 2-, and 3-week lags, indicating if there was at least one day in the prior week

where total precipitation was above the 80[th] percentile of all daily totals.

<u>Statistical Analysis</u>

We assessed the effect of WASH interventions on caregiver-reported diarrhea prevalence

in the past 7 days for children in an intention-to-treat-analysis. To maximize precision, we

created an indicator for any "WASH interventions" (1 = water, sanitation, hygiene, combined

WSH, or combined WSH + Nutrition arms; 0 = control arms).

To allow for potentially non-linear relationships between continuous environmental

factors and diarrhea, we fit generalized additive models (GAM) between treatment and diarrhea,

including one environmental factor modeled using a spline in each model.[18] We specified a

binomial family with logit link functions and used restricted maximum likelihood estimation to

select smoothing parameters. We estimated simultaneous confidence intervals using a parametric

bootstrap of the variance-covariance matrix under a multivariate normal distribution.[19] We also

estimated prevalence ratios at the 10^{th} and 90^{th} percentile of each observed distribution and

estimated 95% confidence intervals using a non-parametric bootstrap with resampling at the

cluster level.

For categorical environmental factors, we estimated intervention effects using targeted

maximum likelihood estimation (TMLE). We included the simple mean, generalized linear

model (GLM), Bayesian GLM, GAM, and lasso net GLM as learners for both our treatment and

outcome ensemble models. We estimated the prevalence ratios and corresponding confidence

intervals for WASH interventions vs. control for each risk factor stratum.

To assess differences in WASH effectiveness under various climate change scenarios, we

estimated the prevalence ratio under projected distributions of precipitation in the study region in

2050. We used the UK Earth System model (UKESM1-0-LL), which best matched observed

values from the trial, to project the spatiotemporal distribution of rainfall in the study region

under different climate change scenarios.[20] We include additional details on our climate model

selection process in Appendix 2.

We examined three Shared Socioeconomic Pathways (SSPs): SSP1 represents a future

under sustainable development where CO_2 emissions are reduced to net zero by 2075. SSP2

represents a "middle of the road" approach where CO_2 emissions stay constant at current levels

until 2050 before gradually falling to net zero by 2100. SSP5 represents continued fossil-fueled

development, where CO_2 emissions triple by 2075.[21]

We extracted precipitation projections for study households by geographical coordinates

under each of these scenarios for every day in 2050. From the matched, daily data, we calculated

the weekly total precipitation for each household under lag periods for which we detected

significant effect modification in our primary analysis. We estimated diarrhea prevalence for the

WASH interventions and control groups under the projected precipitation distributions using

GAMs fitted with weekly total precipitation under the relevant lags as the independent variable.

We calculated the prevalence ratio using the parametric g-formula and constructed 95%

confidence intervals using 1,000 bootstrapped resamples taken at the cluster level.

To check for possible misclassification of the diarrhea outcome, we conducted a negative

control analysis using an outcome of caregiver-reported bruising in the past 7 days.[22]

Analyses were conducted in R (version 4.1.3). The pre-analysis plan and replication

scripts are available through the Open Science Framework (https://osf.io/yt67k/).

Role of the funding source

The funders of the study had no role in study design, data analysis, interpretation, writing

of the manuscript, or the decision to submit the manuscript for publication.

Results

Our analysis included 12,440 total diarrhea measurements for 6,921 children between 0.5

and 5.5 years of age (mean 2.0, SD 1.2). We report additional participant demographics and the

distribution of environmental factors by group in Table 1. The unadjusted diarrhea prevalence

ratio associated with the pooled WASH interventions averaging across climatic conditions in the

trial was 0.66 (95% CI 0.54 - 0.80).

<u>Temperature</u>

We found no evidence of effect modification of WASH interventions by minimum or

maximum temperatures within the prior 7- or 30- day period. Diarrhea prevalence tended to

increase with average temperature in the previous 7- and 30-days in both arms, but the increase

in prevalence per degree was lower in the WASH interventions group (Figure 1A). For weekly

average temperature under 2-3-week lags, prevalence increased with temperature in the control

group but remained similar across temperature levels in the intervention group. For a change of

average weekly temperature with a 3-week lag from 19.2°C (10th percentile) to 26.6 °C (90th

percentile), the prevalence of diarrhea changed from 3.6% (95% CI 3.0-4.6) to 3.8% (95% CI

3.4-4.7) in the WASH interventions arms and 2.4% (95% CI 1.8-3.4) to 5.1% (95% CI 4.3-6.8)

in the control arm. This corresponded to a prevalence ratio of 2.03 (95% CI 1.05, 3.81) at 19.2°C

and 0.74 (95% CI 0.56, 0.98) at 26.6°C. We found no significant differences in diarrhea

prevalence between the interventions vs. control arms for weekly average temperature under a 1-

week lag. (Figure 1B). There were no differences in prevalence ratios for weekly average

temperatures above vs. below-median under any lag period. (Figure 1C, Table A1)

<u>Surface Water</u>

The effect of WASH interventions was larger among children with below-median vs.

above-median seasonal surface water within 250 meters (0.58, 95% CI 0.45-0.74 vs. 0.92, 95%

CI 0.61-1.39). We found no effect modification by the proportion of land covered by any other

type of surface water or within any other distance. (Figure 2, Table A2)

The prevalence ratio of WASH interventions was lowest in households that were in the

33rd-66th percentile in distance to seasonal surface water (0.53, 95% CI 0.36-0.76) compared to

households in the 0th-33rd percentile (0.99, 95% CI 0.55-1.78) or in the 66th-100th percentile

(0.60, 95% CI 0.38-0.94). We found no effect modification by distance to any surface water or

ephemeral surface water. (Figure 2, Table A2)

<u>Vapor Pressure Deficit (VPD)</u>

WASH interventions were less effective at higher values of VPD. The prevalence of

diarrhea was 5.0% (95% CI 4.3-6.5) in the control arm and 3.4% (95% CI 3.1-4.1) in the WASH

interventions arm at a VPD of 0.65 kPa (10th percentile), compared to 3.0% (95% CI 2.0-4.7) in

the control and 4.5% (95% CI 3.7-6) in the intervention at 1.37 kPa (90th percentile) (Figure A1).

At 0.65 kPa, we estimated a prevalence ratio of 0.70 (95% CI 0.42, 0.95). At 1.37 kPa, we

estimated a prevalence ratio of 1.93 (95% CI 1.02, 4.08).

<u>Precipitation</u>

The effectiveness of WASH interventions increased with weekly total precipitation under

1-2-week lags. (Figure 3A). For a change of weekly total precipitation under a 1-week lag from 0

mm (10th percentile) to 125 mm (90th percentile), we estimated that diarrhea prevalence changed

from 4.0% (95% CI 3.6-4.8) to 3.2% (95% CI 2.7-4.1) in the WASH interventions arms

compared to 3.0% (95% CI 2.7-3.7) to 5.3% (95% CI 4.5-7) in the control arm. The prevalence

ratio was 1.38 (95% CI 1.08, 1.8) at 0 mm and 0.72 (95% CI 0.4, 1.43) at 125 mm. We observed

similar results under 2-3-week lags, but with smaller differences in prevalence between the

WASH interventions arms and control arms at higher values of weekly total precipitation.

The prevalence ratio associated with WASH interventions was lower when there was at

least one day of heavy rain (0.38, 95% CI 0.23-0.62) in the 8- to 14-day period before outcome

measurement (1-week lag) compared to when there were no days with heavy rain (0.77, 95% CI

0.60-0.98) (Figure 3B, Table A3). We did not observe effect modification by heavy rain in other

7-day periods under 2-3-week lags. WASH interventions were more effective when total weekly

precipitation under a 3-week lag was above-median (0.57, 95% CI 0.46-0.71) compared to

below-median (1.00, 95% CI 0.67-1.49) (Figure 3C, Table A3). We found similar trends under

the 1-2-week lag periods.

Peak precipitation was higher under climate change scenarios compared to during the

trial (Figure 4A). We projected an increase in intervention effectiveness in climate scenarios in

which the study area was predicted to have heavier, more frequent rainfall. During the study

period, the prevalence ratio for total weekly precipitation under a 1-week lag was 0.67 (95% CI

0.65-0.68) (Figure 4B, Table A4). Under SSP1, where there is a predicted decrease in the

frequency and volume of rainfall, the prevalence ratio was similar (0.66; 95% CI 0.64-0.68).

Under SSP2 and SSP5, where there is a predicted increase in rainfall, we estimated lower

prevalence ratios of 0.59 (95% CI 0.57-0.60) for SSP2 and 0.46 (95% CI 0.44-0.48) for SSP5.

(Figure 4B)

Other analyses

We did not observe any significant differences in intervention effect estimates by climate

or environment on reported bruising, a negative-control outcome. (Appendix 3)

In accordance with our pre-analysis plan, we ran analyses for water flow accumulation,

enhanced vegetative index, land use, and population density but generally did not find that these

variables modified the effect of WASH interventions on diarrhea prevalence. Results for these

analyses are available at https://osf.io/yt67k/.

Discussion

WASH interventions in rural Bangladesh were more effective under conditions with

higher total precipitation and heavy rainfall. We found that WASH interventions reduced

diarrhea prevalence by 62% (95% CI 38%-77%) following heavy rainfall and up to 43% (95%

CI 28%-56%) after above-median rainfall compared to the overall 34% (95% CI 19%-45%)

reduction observed in the original trial. We observed some heterogeneity in treatment

effectiveness under varying levels of temperature, vapor pressure deficit and surface water, but

precision of these estimates was low. Under highly unregulated fossil-fueled development that

will lead to heavier, more frequent rainfall (SSP5), we predicted that WASH interventions will

reduce diarrhea prevalence by up to 54% (95% CI 52%-56%) in a climate change scenario.

WASH interventions were more effective under higher weekly average precipitation with

a 1- and 2-week lag, in weeks with a heavy rain event with a 1-week lag, and in weeks with

above-median precipitation with all lag periods. While no prior studies have rigorously assessed

whether WASH intervention effectiveness varies with precipitation, prior studies have found

both positive and negative associations between diarrhea and precipitation; heavy rain was

associated with higher diarrhea prevalence following dry periods but not following wet periods.[23]

These differences are usually attributed to the concentration-dilution hypothesis, which posits

that diarrhea risk is highest from heavy rainfall following dry periods because pathogens that

become concentrated during dry periods are flushed into the environment.[23] We were unable to

investigate this hypothesis since most heavy rain preceded by wet periods during the study.

However, we observed higher intervention effectiveness following heavy rain regardless of

whether it was preceded by a dry period. This is consistent with past studies in Bangladesh that

found that heavy rainfall was associated with higher concentrations of diarrhea-causing *E. coli* in

wells regardless of antecedental rainfall.[9] The rainy season in Bangladesh brings continuous

heavy rainfall which may flush contaminants from unimproved latrines directly into well water

or into pond water that eventually drains to tube wells through the aquifer.[9] Improved WASH

may reduce the prevalence of diarrhea under these conditions by moderating increases in fecal

contamination that are associated with heavy rain.

Diarrhea prevalence stayed relatively constant across temperature levels among WASH

intervention recipients but increased with average temperature in the control group; however,

there was no effect modification by minimum or maximum temperature. To our knowledge, no

prior studies assessed differences in WASH effectiveness by temperature, but some have

evaluated differences in diarrhea prevalence. One study found a positive association with

temperature and bacterial diarrhea, but no relationship between temperature and viral diarrhea,

suggesting that differences in environmental survival by pathogen taxa could impact the

influence of temperature on the effect of WASH interventions.[6] Our findings suggest that WASH

interventions may mitigate increases in diarrhea prevalence associated with average

temperatures, even at less than extreme temperature levels.

We found weak evidence that WASH interventions were more effective in areas with less

seasonal surface water within 250 meters. Standing surface water is highly susceptible to

contamination by enteric pathogens, and previous qualitative studies have shown that vulnerable

communities may use surface water as drinking water for convenience or during times of water

scarcity, which could lead to increased diarrhea.[24,25]

WASH interventions were slightly more effective at lower values of VPD that correspond

to high humidity. No prior studies have evaluated the impact of humidity on WASH

effectiveness, but prior studies have found very weak associations between humidity and most

enteric bacteria and viruses.[26]

Compared to during the study period, WASH effectiveness is predicted to be similar

under a sustainable development scenario (SSP1), where average rainfall was predicted to be

lower than current levels in our climate models, and to be higher under "middle-of-the-road"

(SSP2) and fossil-fueled development scenarios (SSP5), where rainy seasons are predicted to last

longer and have higher total precipitation. No studies have assessed the future effectiveness of

WASH under climate change, but there is growing evidence that increased rainfall will lead to

higher diarrhea incidence and potential degradation of water and sanitation infrastructure. [25,27–29]

Our results suggest that during future periods of extreme rainfall, WASH interventions may be

especially impactful and increase resilience of vulnerable populations. Low cost, point-of-use

WASH interventions require prolonged uptake and maintenance, which can be difficult to

sustain; efforts to promote uptake may be most fruitful prior to or during the rainy season, when

interventions are likely to have the greatest impact. Further, our findings suggest that in areas

with limited resources, WASH interventions should be targeted to populations where rainfall is

projected to be the highest.

<u>Limitations</u>

Our study is subject to several limitations. We measured caregiver-reported diarrhea,

which is susceptible to courtesy bias. However, our negative control analysis using an alternative

caregiver-reported outcome suggested that there was no evidence of misclassification.

We were unable to investigate effect modification of specific intervention arms due to

data sparsity; pooling across intervention arms was necessary to ensure sufficient statistical

power.

The generalizability and accuracy of our predicted changes in WASH effectiveness may

be limited. Because we used precipitation projections specific to our study site, our predictions

for WASH intervention effectiveness under climate change scenarios cannot necessarily be

generalized to other sites. However, our findings may still be relevant for regions with similar

climates, environmentally-mediated transmission patterns, and WASH infrastructures.

Additionally, these climate models considered how environmental variables are expected to

change under different global scenarios but do not account for improvements to WASH

technologies or local economic development. These factors will likely impact future WASH

intervention effectiveness in ways that our study could not capture.

Conclusion

In rural Bangladesh, low-cost, household level WASH interventions more effectively

reduced diarrhea prevalence at higher average temperature levels and following periods of higher

precipitation and heavy rain. We observed up to a 62% reduction in diarrhea prevalence

following extreme rainfall compared to the overall average 34% reduction found over the entire

study period. Under likely climate change scenarios, our study suggests that WASH

interventions will be more effective due to higher levels of rainfall. In settings similar to rural

Bangladesh, WASH interventions should be prioritized in areas that are prone to extreme

precipitation and during the rainy season. In addition, WASH interventions may increase

communities' resilience against extreme weather under climate change by preventing

environmentally-mediated enteric infections.

Acknowledgments

This study was supported by the Gates Foundation (grant number OPPGD759 to the University of California, Berkeley) and the Task Force for Global Health (grant number NTDSC088G to Innovations for Poverty Action, and grant number NTDSC089G to the University of California, Berkeley). Research reported in this publication was supported in part by the National Institute of Allergy And Infectious Diseases of the National Institutes of Health under Award Numbers K01AI141616 (PI: Benjamin-Chung) and R01AI166671 (PI: Arnold), the National Heart, Lung, And Blood Institute of the National Institutes of Health under award number T32HL151323 (Nguyen), and a Stanford University School of Medicine Dean's Postdoctoral Fellowship (Grembi). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. Jade Benjamin-Chung is a Chan Zuckerberg Biohub Investigator. We also acknowledge the Stanford Research Computing Center for computational resources at the Sherlock high-performance cluster.

Role of the Funding Source

Authors' Contributions

Anna T. Nguyen: data curation, formal analysis, methodology, project administration, software, visualization, writing - original draft

Jessica A. Grembi: data curation, formal analysis, investigation, methodology, writing – review & editing.

<u>Marie Riviere</u>: data curation, formal analysis, investigation, methodology, writing – review &

editing.

<u>Gabby Barratt Heitmann, William D. Hutson</u>: visualization, formal analysis, investigation,

writing – review & editing.

<u>Tejas S. Athni</u>: formal analysis, investigation, writing – review & editing.

<u>Arusha Patil:</u> visualization, writing – review & editing.

<u>Ayse Ercumen, Audrie Lin, Leanne Unicomb, Mahbubur Rahman, John M. Colford, Jr., Stephen</u>

<u>P. Luby</u>: project administration, supervision, resources, writing - review & editing

<u>Yoshika Crider, Andrew Mertens</u>: resources, methodology, writing - review & editing

<u>Benjamin F. Arnold</u>: methodology, software, project administration, writing - review & editing

<u>Jade Benjamin-Chung</u>: conceptualization, data curation, formal analysis, funding acquisition,

methodology, project administration, resources, supervision, visualization, writing - original

draft

Ethics Committee Approval

The original trial protocol was approved by the Ethical Review Committee at The International

Centre for Diarrhoeal Disease Research, Bangladesh (PR-11063), the Committee for the

Protection of Human Subjects at the University of California, Berkeley (2011-09-3652), and the

institutional review board at Stanford University (25863).

References

1 Institute for Health Metrics and Evaluation (IHME). GBD Results. Seattle WA IHME Univ.
Wash. 2020. https://vizhub.healthdata.org/gbd-results (accessed Sept 20, 2022).

2 Global Diarrhea Burden | Global Water, Sanitation and Hygiene | Healthy Water | CDC. 2018;
published online Nov 9. https://www.cdc.gov/healthywater/global/diarrhea-burden.html
(accessed Dec 18, 2021).

3 Diarrhoeal disease. https://www.who.int/news-room/fact-sheets/detail/diarrhoeal-disease
(accessed Dec 18, 2021).

4 Null C, Stewart CP, Pickering AJ, *et al.* Effects of water quality, sanitation, handwashing, and
nutritional interventions on diarrhoea and child growth in rural Kenya: a cluster-randomised
controlled trial. *Lancet Glob Health* 2018; **6**: e316–29.

5 Luby SP, Rahman M, Arnold BF, *et al.* Effects of water quality, sanitation, handwashing, and
nutritional interventions on diarrhoea and child growth in rural Bangladesh: a cluster
randomised controlled trial. *Lancet Glob Health* 2018; **6**: e302–15.

6 Carlton EJ, Woster AP, DeWitt P, Goldstein RS, Levy K. A systematic review and meta-
analysis of ambient temperature and diarrhoeal diseases. *Int J Epidemiol* 2016; **45**: 117–30.

7 Liu Z, Liu Y, Zhang Y, *et al.* Effect of ambient temperature and its effect modifiers on
bacillary dysentery in Jinan, China. *Sci Total Environ* 2019; **650**: 2980–6.

8 Levy K, Woster AP, Goldstein RS, Carlton EJ. Untangling the Impacts of Climate Change on
Waterborne Diseases: a Systematic Review of Relationships between Diarrheal Diseases and
Temperature, Rainfall, Flooding, and Drought. *Environ Sci Technol* 2016; **50**: 4905–22.

9 Nijhawan A, Howard G. Associations between climate variables and water quality in low- and
middle-income countries: A scoping review. *Water Res* 2022; **210**: 117996.

10 Bhavnani D, Goldstick JE, Cevallos W, Trueba G, Eisenberg JNS. Impact of Rainfall on
Diarrheal Disease Risk Associated with Unimproved Water and Sanitation. *Am J Trop Med
Hyg* 2014; **90**: 705–11.

11 Mukabutera A, Thomson D, Murray M, *et al.* Rainfall variation and child health: effect of
rainfall on diarrhea among under 5 children in Rwanda, 2010. *BMC Public Health* 2016; **16**:
731.

12 Hashizume M, Wagatsuma Y, Faruque ASG, *et al.* Factors determining vulnerability to
diarrhoea during and after severe floods in Bangladesh. *J Water Health* 2008; **6**: 323–32.

13 Kohlitz J, Willetts J, Gero A, Lyons S, Boisson S, Medlicott K. Climate, Sanitation and
Health. https://www.who.int/publications-detail-redirect/climate-sanitation-and-health
(accessed Dec 19, 2021).

14 Wang P, Asare E, Pitzer VE, Dubrow R, Chen K. Associations between long-term drought
and diarrhea among children under five in low- and middle-income countries. *Nat Commun*
2022; **13**: 3661.

15 Wu J, Yunus M, Streatfield PK, *et al.* Impact of tubewell access and tubewell depth on
childhood diarrhea in Matlab, Bangladesh. *Environ Health* 2011; **10**: 109.

16 Contreras JD, Islam M, Mertens A, *et al.* Evaluation of an on-site sanitation intervention
against childhood diarrhea and acute respiratory infection 1 to 3.5 years after implementation:
Extended follow-up of a cluster-randomized controlled trial in rural Bangladesh. *PLOS Med*
2022; **19**: e1004041.

17 Arnold BF, Null C, Luby SP, *et al.* Cluster-randomised controlled trials of individual and
combined water, sanitation, hygiene and nutritional interventions in rural Bangladesh and
Kenya: the WASH Benefits study design and rationale. *BMJ Open* 2013; **3**: e003476.

18 Hastie T, Tibshirani R. Generalized Additive Models. *Stat Sci* 1986; **1**: 297–310.

19 Marra G, Wood SN. Coverage Properties of Confidence Intervals for Generalized Additive
Model Components. *Scand J Stat* 2012; **39**: 53–74.

20 Sellar AA, Jones CG, Mulcahy JP, *et al.* UKESM1: Description and Evaluation of the U.K.
Earth System Model. *J Adv Model Earth Syst* 2019; **11**: 4513–58.

21 IPCC. Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to
the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge,
UK: Cambridge University Press, 2021
https://www.ipcc.ch/report/ar6/wg1/downloads/report/IPCC_AR6_WGI_Full_Report.pdf
(accessed March 28, 2022).

22 Arnold BF, Ercumen A, Benjamin-Chung J, Colford JM. Brief Report: Negative Controls to
Detect Selection Bias and Measurement Bias in Epidemiologic Studies. *Epidemiol Camb*
Mass 2016; **27**: 637–41.

23 Kraay ANM, Man O, Levy MC, Levy K, Ionides E, Eisenberg JNS. Understanding the Impact
of Rainfall on Diarrhea: Testing the Concentration-Dilution Hypothesis Using a Systematic
Review and Meta-Analysis. *Environ Health Perspect* 2020; **128**: 126001.

24 Boithias L, Choisy M, Souliyaseng N, *et al.* Hydrological Regime and Water Shortage as
Drivers of the Seasonal Incidence of Diarrheal Diseases in a Tropical Montane Environment.
PLoS Negl Trop Dis 2016; **10**: e0005195.

25 Delpla I, Jung A-V, Baures E, Clement M, Thomas O. Impacts of climate change on surface
water quality in relation to drinking water production. *Environ Int* 2009; **35**: 1225–33.

26 Colston JM, Zaitchik BF, Badr HS, *et al.* Associations Between Eight Earth Observation-
Derived Climate Variables and Enteropathogen Infection: An Independent Participant Data

Meta-Analysis of Surveillance Studies With Broad Spectrum Nucleic Acid Diagnostics.
GeoHealth 2022; **6**: e2021GH000452.

27 Levy K, Smith SM, Carlton EJ. Climate Change Impacts on Waterborne Diseases: Moving
Toward Designing Interventions. *Curr Environ Health Rep* 2018; **5**: 272–82.

28 Patz JA, Frumkin H, Holloway T, Vimont DJ, Haines A. Climate Change: Challenges and
Opportunities for Global Health. *JAMA* 2014; **312**: 1565–80.

29 Sherpa AM, Koottatep T, Zurbrügg C, Cissé G. Vulnerability and adaptability of sanitation
systems to climate change. *J Water Clim Change* 2014; **5**: 487–95.

Table 1: Study population characteristics, by intervention group

	Control	Improved WASH*
Children	1956	4965
Observations	3466	8974
Households	1203	3051
Children per household	1.6 (1.6 - 1.7)	1.6 (1.6 - 1.7)
Diarrhea during study period	186 (9.5%)	339 (6.8%)
Age at first measurement (years)	1.6 (1.6 - 1.7)	1.6 (1.6 - 1.6)
Age at any measurement (years)	2.0 (2.0 - 2.1)	2.0 (2.0 – 2.0)
Average weekly temperature, 0 week lag (C)	23.7 (23.3 - 24.2)	23.7 (23.4 - 24)
Weekly total precipitation, 1 week lag (mm)	48.3 (37.8 - 58.7)	48 (41.4 - 54.7)
Heavy rain, 0 week lag	667 (19.24%)	1584 (17.65%)
Vapor pressure deficit (kPa)	0.8 (0.8 - 0.9)	0.8 (0.8 - 0.8)
Enhanced vegetation index	0.3 (0.3 - 0.4)	0.3 (0.3 - 0.3)
Distance from any surface water (m)	266 (225.3 - 306.6)	283.4 (258.4 - 308.5)
Area covered by any surface within 500 meters (%)	8.5% (6.2% - 10.9%)	9.1% (7.6% - 10.6%)
Population density (persons per 100 m^2)	13.0 (11.7 - 14.4)	13.0 (12.1 - 13.9)
Land Use Category		
Croplands	1032 (29.77%)	2425 (27.02%)
Natural Vegetation	2065 (59.58%)	5408 (60.26%)
Savannas	367 (10.59%)	1135 (12.65%)

Sample sizes, child demographics, diarrhea prevalence, and environmental risk factor distributions in the control group vs improved WASH group (water, sanitation, hygiene, combined WSH, or combined WSH + Nutrition arms). For categorical variables, the number of occurrences and percentages are reported. For continuous variables, the mean and 95% confidence interval are reported.

Figure 1: Differences in diarrhea prevalence by temperature, stratified by treatment group

A) *Predicted diarrhea prevalence by average, minimum, and maximum temperature in the past 7 and 30 days, stratified by treatment arm. Distribution of observed temperature values are displayed below plots showing model fits, with the median value marked by a red vertical line. Prevalence ratios for diarrhea in the pooled WASH treatment vs control group are displayed for the 10th and 90th percentiles of each distribution, with 95% CIs estimated by taking 1,000 bootstrapped resamples by cluster.*
B) *Predicted diarrhea prevalence by average weekly temperature under a 1-, 2-, and 3-week lag, stratified by treatment arm*
C) *Prevalence ratios for diarrhea in the pooled treatment vs control arms by different levels of average weekly temperature under a 1-, 2-, and 3-week lag. Confidence intervals calculated to adjust for clustering of treatment assignment.*

Figure 2: Differences in WASH intervention effect by surface water presence

Prevalence ratios for diarrhea in the pooled treatment vs control arms by different levels of any, ephemeral, and season surface water presence. Confidence intervals calculated to adjust for clustering of treatment assignment. Amount of nearby surface water describes the proportion of land within a given distance from a child's household that was covered by surface water. Tertile bounds for distance from any surface water were 0-33rd: 0-164 m, 33-66th: 164-316 m, 66-100th: 316-1818 m; from ephemeral surface water were 0-33rd: 0-261 m, 33-66th: 261-508 m, 66-100th: 508-2083 m; from seasonal surface water were 0-33rd: 0-230 m, 33-66th: 230-453 m, 66-100th: 453-1902 m.

Figure 3: Differences in diarrhea prevalence and WASH intervention effect by precipitation and treatment

A) *Predicted diarrhea prevalence by total weekly precipitation, stratified by treatment group. The upper bounds of confidence intervals are truncated at a prevalence of 20% on the plot, but values reach as high as 22% under the 1-week lag and 60% under the 2-week lag. Distribution of observed precipitation values are displayed below plots showing model fits, with the median value marked by a blue vertical line. Prevalence ratios for diarrhea in the pooled WASH treatment vs control group are displayed for the 10th and 90th percentiles of each distribution, with 95% CIs calculated by taking 1,000 bootstrapped resamples by cluster.*

B) *Prevalence ratios for diarrhea in the pooled WASH treatment vs control group by different levels of heavy rain (1+ days in the week with precipitation above the 80th percentile of daily total rainfall in the study)*

C) *Prevalence ratios for diarrhea in the pooled WASH treatment vs control group by different levels of total weekly precipitation (above/below median weekly total rainfall in the study)*

Figure 4: Predicted changes in rainfall and WASH intervention effect under different climate change scenarios

A) *Predicted trends in total weekly precipitation, by climate change scenario*
B) *Prevalence ratios for diarrhea in the pooled treatment vs control arms, by climate change scenario*