

1 **Physician burnout and symptom of anxiety and depression: Burnout in**
2 **Consultant Doctors in Ireland Study (BICDIS)**
3 **Short title: Anxiety and depression in physician burnout: implications for the**
4 **post-Covid landscape**

5 Genevieve Crudden¹

6 Fabio Margiotta²

7 Anne M. Doherty^{3,4}

8

9 ¹ Department of Psychiatry, Galway University Hospitals, Newcastle Road, Galway, Ireland

10 ² Department of Psychiatry, Townsville University Hospital, Queensland, Australia.

11 ³ Department of Psychiatry, University College Dublin, 63 Eccles Street, Dublin 7, Ireland.

12 ⁴ Department of Liaison Psychiatry, Mater Misericordiae University Hospital, 63 Eccles
13 Street, Dublin 7, Ireland.

14

15

16 **Corresponding author**

17 Anne M. Doherty

18 Department of Psychiatry

19 University College Dublin

20 63 Eccles Street,

21 Dublin 7,

22 Ireland.

23 Tel +353871385336

24 Email: anne.doherty@ucd.ie

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

25 **Abstract**

26 **Background:** Burnout is a work related syndrome comprising three major dimensions:
27 emotional exhaustion, depersonalisation and reduced professional efficacy. It is recognised
28 that burnout may be associated with psychopathological sequelae, such as symptoms of
29 anxiety and depression.

30
31 **Aims:** This study aimed to examine the relationship between burnout and symptoms of
32 depression and anxiety in senior doctors in Ireland. We examined personality traits as a
33 mediating factor on burnout, in relation to work related stress among the participants.

34
35 **Methods:** This study was a national cross sectional survey of consultants. Data included
36 basic demographic data, work-related data, burnout as measured by the Maslach Burnout
37 Inventory, personality traits, and symptoms of depression and anxiety from the Depressive
38 Anxiety Stress Scale [DASS].

39
40 **Results:** Results were obtained from 477 consultants, yielding a response rate 21.9%. Nearly
41 half (42%) reported high burnout levels. The Depression and Anxiety Stress Scale revealed
42 high levels of stress symptoms, and lower levels of anxiety symptoms. After controlling for
43 gender and age, depressive symptoms were associated with high levels of emotional
44 exhaustion and a higher clinical workload

45
46 **Conclusions:** The this study reported a high level of work related burnout, associated with
47 symptoms of depression and anxiety in senior doctors. This is concerning, and strategies are
48 required that consider the relationship between personality and burnout. Further research
49 to evaluate strategies to ensure physician wellbeing and optimal delivery of patient care are
50 required to address this serious problem.

51
52
53 **Keywords:** Burnout, depression, anxiety, psychological stress, Ireland

54 Introduction

55 Burnout is an occupational phenomenon related to stress, and has been described as “a
56 *particular response to stress involving psychological withdrawal. It is a costly coping*
57 *mechanism seen as the end-point of a process which begins with the stress response*”. [1] It
58 is defined by the World Health Organisation (WHO) in ICD-11 as “*a syndrome conceptualized*
59 *as resulting from chronic workplace stress that has not been successfully managed. It is*
60 *characterized by three dimensions: feelings of energy depletion or exhaustion; increased*
61 *mental distance from one’s job, or feelings of negativism or cynicism related to one’s job;*
62 *and reduced professional efficacy*” [2]. The WHO classify it as an occupational condition and
63 not as a medical condition.

64 It can have a significant effect on the health of the individual worker and can affect how
65 they view and approach their work: “*What starts out as an important meaningful and*
66 *challenging work becomes unpleasant, unfulfilling and meaningless.*” [3]

67 In the decades preceding the Covid-19 pandemic, physician burnout has become an area of
68 increasing concern, both from an occupational medicine standpoint, and from the
69 perspective of patient safety. First described in 1974, the construct of burnout has been
70 developed into a measurable three-dimensional model comprising emotional exhaustion,
71 cynicism (depersonalisation) and ineffectiveness (low personal accomplishment). [4] It is a
72 common problem with some studies reporting rates as high as 50%, and it may have a
73 profound effect on the lives of the physicians affected. [5]

74 Doctors have been identified as a group who are at increased risk of developing work-related
75 stress. In a large study comparing US physicians with the general population, although higher
76 educational attainment was identified as a protective factor for burnout in the general
77 population, a medical degree conferred an increased risk independent of hours worked. [5]
78 The authors concluded that the occurrence of physician burnout is set apart from the larger
79 societal trend, and that there are factors specific to physicians and their work which
80 predispose this group to poorer outcomes. Riley suggested that burnout in doctors may be
81 the end consequence of the interplay of challenging work environment, personality factors,
82 subjective lack of control and insufficient reward. [6] The author suggests that the

83 “conscientious and committed personalities” working in medicine are at a loss when faced
84 with certain environmental strains. These strains include the “demand control imbalance”
85 identified by Karasek, [7] along with effort reward imbalance, and inadequate support in the
86 work environment. [8] The extent of the role of personality in relation to burnout remains
87 unclear in general, and has not been studied in the Irish context. Previous research has
88 indicated that personality traits of neuroticism or 'negative affectivity' has been associated
89 with greater stress and burnout in doctors [9]

90

91 The impact of burnout on patient care is well established. Burnout is associated with lower
92 patient satisfaction, involvement in adverse events, and failure across domains of
93 professionalism which affect patient care, including empathy, interpersonal communication
94 and adherence to guidelines. [10-12] It has been suggested that burnout may be associated
95 with issues relating to patient safety, as both burnout and patient safety relate to
96 organisational, social and individual factors which depend on environmental resources. [13]
97 There is also evidence that medical errors and their sequelae can in turn cause or worsen
98 practitioner burnout and mental health problems, an effect that may be compounded by
99 organisational factors including an organisational culture of blame, inadequate personal
100 support and suboptimal administrative support [14, 15]. Two key systematic reviews
101 examined the relationship between burnout and patient safety. Hall et al identified a
102 relationship between poor patient safety outcomes such as errors, and noted a lack of
103 prospective studies. [14] Dewa et al reported a strong relationship between burnout and
104 problems with safety, but the relationship between burnout and patient satisfaction with
105 care is less well defined. [15]

106

107 Burnout is associated with increased risk of health problems in doctors: a range of
108 difficulties including cardiovascular disease and shorter life expectancy, problematic alcohol
109 use, broken relationships, depression, and suicide. [16] While burnout is an occupational
110 phenomenon rather than a mental illness, there are many areas of overlap with depression,
111 and in particular the experience of hopelessness, helplessness and powerlessness are found
112 in individuals with burnout and with depression. [2, 17] People who are severely affected by
113 burnout are likely to also report symptoms of depressive illness: one Finnish study reported
114 that half of those with severe burnout were also experiencing a depressive episode. [18]

115

116 This study aimed to investigate the prevalence of burnout and associated factors and their
117 relationship with the development of psychiatric morbidity in a cohort of senior physicians
118 working in the Irish Healthcare system. We sought to explore the impact of burnout and
119 personality factors on the development of depressive and anxiety symptoms.

120

121 **Methods**

122 This study utilised a cross-sectional design, to study the prevalence of burnout in a national
123 population of senior hospital doctors (consultants) in Ireland. The methods were previously
124 described by Margiotta et al. [19] We will summarise here for ease of reference: an
125 anonymous survey was distributed to members of the Irish Hospital Consultants Association
126 (IHCA) via their electronic database. The IHCA is a representative organisation for hospital
127 consultants across Ireland with 2160 members. An email was sent to the membership
128 including a hyperlink to the electronic survey, the study information sheet and the consent
129 form. Participation was voluntary, and responses were completely anonymised. Ethical
130 approval was obtained from the Clinical Research Ethics Committee of the Saolta group.
131 Data were collected between 1st September and 30th November 2016.

132 Participants were asked to provide demographic details and information and details of their
133 current working climate and lifestyle. The questions encompassed work-related
134 characteristics and stressors. These included the person's medical specialty, their type of
135 contract, satisfaction with remuneration, hours worked (weekly) , time spent face-to-face
136 with patients, and any experience of medico-legal or disciplinary action.

137 Burnout itself was measured by the Maslach Burnout inventory General Survey (MBI-GS)
138 which contains 3 subscales, namely, emotional exhaustion, depersonalization or cynicism,
139 and professional efficacy. High levels of emotional exhaustion and cynicism are associated
140 with burnout, while high levels of professional efficacy are protective. This questionnaire is
141 validated in medical populations of clinicians internationally. Although it is not
142 recommended that the three internal scales of MBI-GS be combined, previous studies have
143 created a model which adds high levels of emotional exhaustion to high levels of cynicism
144 and low professional efficacy. [20]

145

146 The Depressive Anxiety Stress Scale (DASS) is a validated self-report instrument which
147 measures symptoms of depression, anxiety and stress. [21] It measures each item on a 4-
148 point Likert scale. Items related to low positive affect, decreased self-esteem, hopelessness,
149 and poor motivation are categorised as depression; while items related to autonomic
150 arousal and fearfulness were categorised as anxiety. Items relating to irritability and over-
151 reactivity are categorised as stress. [22] It has been validated in clinical and non-clinical
152 populations. [23]

153

154 The Big Five Inventory (BFI-10) is a 10-item schedule to measure personality on the traits.
155 [24] This is a 10-item scale with questions related to the five personality traits, answered on
156 a 5 point Likert scale from “disagree strongly”, through “disagree a little”, “neither agree nor
157 disagree”, “agree a little” to “agree strongly”. It is an abbreviated version of the Big Five
158 Inventory (John et al, 1991). [25]

159

160 Data were analysed using the Statistical Package for the Social Sciences (SPSS) (Version
161 27.0). To make comparison between groups we examined basic bivariate characteristics.
162 Stepwise linear regression analysis was used to investigate the association of demographic
163 variables, personality and psychiatric morbidity with burnout after controlling for the
164 population characteristics. To investigate these hypotheses we performed further
165 multivariable linear regression analysis with the DASS outcomes (depression, anxiety and
166 stress) as predictors and the three burnout dimensions as the dependent variable.

167

168 **Results**

169 The characteristics of this group were previously described. [19] Of the 2,160 consultants
170 invited to participate from the database of the Irish Hospital Consultants Association, 472
171 commenced the survey (response rate = 21.9%), and 433 completed the questionnaire:
172 these were suitable for analysis. The socio-demographic characteristics of the group are
173 outlined in Table 1. Of the respondents, 56.6% were male and 43.3% were female doctors,
174 the mean age was 49.1 years and the majority were married or in a long-term relationship
175 (82.7%). Medical physicians were the most represented specialty (19.4%), followed by
176 psychiatrists (18.7%) and surgeons (15.7%). Just under one-third of respondents reported

177 having been involved in a work-related legal action (31.9%), and 2.1% reported an
178 experience of formal disciplinary action in the workplace.

179

180 Although mean DASS scores for the whole group were below the cut off for caseness, 25.8%
181 of hospital consultants reported levels of depressive symptoms above the cut-off and 13.8%
182 reported anxiety symptoms above the cut-off. Almost 60% of participants reported stress
183 levels above the cut off for caseness (See Table 2). The mean BFI (personality) scores are
184 also shown in Table 2 and are significantly higher than in a normative general population
185 sample where means sit between 3.2 (SD 0.83) for agreeableness and 4.1 (SD 0.69) for
186 conscientiousness. [26] In this study population these means were 7.4 (SD 1.7) and 9.0 (1.3)
187 respectively, more than double the general population scores in the literature.

188

189 We performed linear regression to examine the association between the three dimensions
190 of burnout, psychiatric morbidity, and its contributory factors. After controlling for age and
191 gender, higher scores on the emotional exhaustion burnout subscale were significantly
192 associated with elevated symptoms of depression on the DASS depressive symptom
193 subscale ($p < 0.001$). These were also associated with higher rates of face-to-face patient
194 contact ($p = 0.001$). Individual doctors with reduced satisfaction with remuneration (p
195 $= 0.001$) were more likely to develop psychiatric morbidity (table 3).

196

197 Elevated burnout scores on the cynical subscale were associated with high DASS stress
198 symptoms ($p < 0.001$) after controlling for age and gender. Greater weekly hours of face-to-
199 face patient contact was associated with higher scores on the cynicism subscale ($p = 0.022$)
200 and poor satisfaction with remuneration was likewise associated with higher scores on the
201 cynicism subscale ($p < 0.001$) as shown in table 3.

202

203 Professional efficacy is the third subscale of the MBI-GS and is a protective factor for the
204 development of burnout. In this study we found that after controlling for age and gender
205 there was an inverse relationship with the stress subscale of the DASS ($p < 0.001$), i.e.,
206 among those reporting higher professional efficacy, higher levels of stress were found.
207 Professional efficacy is associated with lower levels of burnout. It was also significantly
208 associated with elevated face-to-face time with patients (0.024), and there was no

209 association in this model with satisfaction with remuneration and professional efficacy
210 (table 3).

211

212 **Discussion**

213 This study reported a high prevalence of burnout among hospital consultants in Ireland.
214 High proportions screened positive for stress (60%), symptoms of depression (26%) and
215 symptoms of anxiety (14%). Burnout is associated with higher scores of depressive
216 symptoms and stress as measured by DASS. Higher levels of patient contact were associated
217 with increased emotional exhaustion and cynicism, but also with elevated levels of the
218 protective professional efficacy. Low levels of satisfaction with remuneration were
219 associated with increased emotional exhaustion and cynicism, but there was no association
220 with professional efficacy.

221

222 Using the composite measure of burnout, higher levels of burnout were found in this
223 sample compared with those reported by O’Dea in a previous study of Irish general
224 practitioners (GPs) (10.8% vs 6.6%). [20] O’Dea’s study reported higher levels of emotional
225 exhaustion (52.7% compared with 42.7%) and lower levels of the professional efficacy which
226 is protective for burnout (16.3% compared with 1.2%). [20] O’Dea et al found similar rates
227 of burnout to UK-based GPs according to a European study, where UK GPs scored higher
228 than those based in Greece, Turkey, Italy and Bulgaria. [27]

229

230 Compared to interns, Irish consultants appear have lower rates of burnout than more junior
231 hospital doctors. Two-fifths 42% of Irish interns reported a low sense of professional
232 efficacy, and a majority (51.5%) reported high rates of emotional exhaustion [24]. It is
233 unknown whether the working conditions of younger doctors, or more general factors
234 associated with age predispose them to burn-out. For example, there may be an association
235 between age and working hours as junior doctors face greater pressure to increase
236 knowledge and skills through practice. Furthermore, their lack of experience may mean they
237 take longer to complete tasks, thereby resulting in increased working hours. [28]

238

239 The response rate in this study compares favourably with previous response rates in similar
240 studies: Al Humadi et al recruited 225 senior doctors, a response rate of 16.3%, and Garcia
241 et al reported a response rate of 17.1% in their US based studies [29, 30].

242

243 Al Humadi et al reported rates of 6.2% depression, 6.6% suicidal ideation, and 19.6%
244 burnout during the Covid-19 pandemic, all associated with history of prior depression or
245 anxiety. Suicidal ideation and burnout were also associated with younger age. Female
246 physicians reported reduced work-life balance and higher rates of burnout. In our study
247 gender was not found to be significantly associated with burnout. A more recent study of
248 Irish consultants conducted during the Covid-19 pandemic in 2021, found that three-
249 quarters screened positive for burnout in the context of the pandemic. [31]

250

251 The association of burnout with longer time spent face-to-face with patient may be a proxy
252 for longer hours of clinical work (given that the mean amount of patient contact was 5.5
253 hours per day), or it may perhaps represent the higher emotional toll of patient care,
254 especially in certain specialities. Previous studies have found a similar association between
255 high clinical workload and burnout. [32]

256

257 High rates of work-related burnout in consultants are highly concerning. Symptoms of
258 stress, anxiety and depression indicate the high psychological burden of burnout.

259 Personality factors, particularly conscientiousness and agreeableness, appear to have an
260 influence on physician burnout. There is a need for research to identify appropriate
261 strategies (both short-term and long-term) to minimise the impact of burnout and to ensure
262 physician wellbeing. There is an urgent need for evidence-based strategies perhaps
263 modulating the relationship between personality and burnout, to mitigate the impact on
264 symptoms of depression and anxiety and to optimise doctors' health for optimal health care
265 delivery.

266

267 Any interventions must include a consideration of both organisational and individual level
268 measures. West et al conducted a systematic review of the evidence for the management of
269 burnout and found strong evidence base for the use of mindfulness, stress management
270 programmes and small group (including reflective) discussions [33]. There was limited

271 evidence for organisational interventions, mainly due to a lack of studies: it is much more
272 challenging to perform a randomised controlled trial of organisational change than a
273 mindfulness programme [33, 34]. At an individual level, doctors tend to identify
274 organisational and structural factors as the priority areas for improving their working lives
275 and addressing burnout [35].

276

277 The limitations of this study include its cross-sectional design and the use of DASS, a
278 screening rather than a diagnostic measure. The small response rate, while not dissimilar to
279 other similar studies, suggests the potential for response bias: people feeling the effects of
280 burnout may be more likely to respond to a survey on this topic, but conversely doctors who
281 are feeling especially overstretched may not feel able to take on another task. It is,
282 however, a large national sample of senior clinicians which includes broad representation
283 across geographical location and speciality.

284

285 Given these high rates of burnout and stress that were reported pre-covid, it would be
286 important to consider the changes that have occurred in healthcare since 2020. There is an
287 urgent need to consider the need to consider measures to mitigate the impact of burnout
288 and to consider the causes, especially structural causes that it may be possible to mitigate.

289

290 Burnout is common in senior doctors, and it is associated with high levels of stress and
291 symptoms of depression and anxiety. Given that physician mental health and suicide is an
292 internationally growing area of concern, these findings suggest that burnout is associated
293 with poorer mental health. There is an urgent need for more research on the relationship
294 between burnout and depression and anxiety in the post-covid world and for the
295 introduction of measures to mitigate its effect on doctors and patients.

296

297

298

299 **Key points**

300 **What is already known about this subject:**

- 301 • Burnout is a common problem among healthcare professionals especially doctors
302 and includes three domains: emotional exhaustion, cynicism and professional
303 efficacy.
304 • Burnout is associated with poorer mental and physical health, and people who meet
305 the threshold for burnout frequently endorse symptoms of depression.

306
307 **What this study adds:**

- 308 • High-levels of burnout and stress were reported in this study of senior doctors
309 (consultants) in this pre-covid study.
310 • Emotional exhaustion is associated with depressive symptoms after controlling for
311 age, gender and higher workload.

312
313 **What impact this may have on practice or policy:**

- 314 • Addressing factors which contribute to burnout may have a positive impact on
315 physician mental health.
316 • Work-related factors must be considered in any strategies designed to address
317 physician burnout.

318
319
320 **Author contributions**

321 Conceptualisation: GC, FM AMD. Data curation: GC, FM AMD. Formal analysis: GC, AMD.
322 Investigation: GC, FM AMD. Methodology: GC, FM AMD. Project administration: GC, FM.
323 Supervision: AMD. Visualisation : GC, FM AMD. Writing - original draft: GC. Writing – review
324 and editing: GC, FM, AMD.

325
326 **Acknowledgements**

327 The authors acknowledge the Irish Hospital Consultants Association for allowing access to
328 the membership email, and to its members for participating in this study.

329

330

331 **Competing interests**

332 The authors have no interests to declare.

333

334 **Funding**

335 This research did not receive any specific grant from funding agencies in the public,

336 commercial, or not-for-profit sectors.

337

338

339

340 References

- 341
- 342 1. Cherniss C. Stress, burnout, and the special services provider. *Special Services in the*
- 343 *Schools*. 1985;2(1):45-61.
- 344 2. WHO. ICD-11 for mortality and morbidity
- 345 statistics. QD85 Burn-out. Geneva: WHO; 2020.
- 346 3. Maslach C, Leiter MP. Understanding the burnout experience: recent research and its
- 347 implications for psychiatry. *World psychiatry*. 2016;15(2):103-11.
- 348 4. Maslach C, Jackson SE. The measurement of experienced burnout. *Journal of*
- 349 *organizational behavior*. 1981;2(2):99-113.
- 350 5. Shanafelt TD, Boone S, Tan L, Dyrbye LN, Sotile W, Satele D, et al. Burnout and
- 351 satisfaction with work-life balance among US physicians relative to the general US
- 352 population. *Arch Intern Med*. 2012;172(18):1377-85.
- 353 6. Riley GJ. Understanding the stresses and strains of being a doctor. *Med J Aust*.
- 354 2004;181(7):350-3.
- 355 7. Karasek Jr RA. Job demands, job decision latitude, and mental strain: Implications for
- 356 job redesign. *Administrative science quarterly*. 1979:285-308.
- 357 8. Johnson JV, Hall EM. Job strain, work place social support, and cardiovascular
- 358 disease: a cross-sectional study of a random sample of the Swedish working population. *Am J*
- 359 *Public Health*. 1988;78(10):1336-42.
- 360 9. Deary IJ, Blenkin H, Agius RM, Endler NS, Zealley H, Wood R. Models of job-
- 361 related stress and personal achievement among consultant doctors. *Br J Psychol*. 1996;87 (Pt
- 362 1):3-29.
- 363 10. Shanafelt TD, Balch CM, Bechamps G, Russell T, Dyrbye L, Satele D, et al. Burnout
- 364 and medical errors among American surgeons. *Ann Surg*. 2010;251(6):995-1000.
- 365 11. Shanafelt TD, Gradishar WJ, Kosty M, Satele D, Chew H, Horn L, et al. Burnout and
- 366 career satisfaction among US oncologists. *J Clin Oncol*. 2014;32(7):678-86.
- 367 12. Anagnostopoulos F, Liolios E, Persefonis G, Slater J, Kafetsios K, Niakas D.
- 368 Physician burnout and patient satisfaction with consultation in primary health care settings:
- 369 evidence of relationships from a one-with-many design. *J Clin Psychol Med Settings*.
- 370 2012;19(4):401-10.
- 371 13. de Lima Garcia C, Bezerra IMP, Ramos JLS, do Valle J, Bezerra de Oliveira ML,
- 372 Abreu LC. Association between culture of patient safety and burnout in pediatric hospitals.
- 373 *PLoS One*. 2019;14(6):e0218756.
- 374 14. Hall LH, Johnson J, Watt I, Tsipa A, O'Connor DB. Healthcare Staff Wellbeing,
- 375 Burnout, and Patient Safety: A Systematic Review. *PLoS One*. 2016;11(7):e0159015.
- 376 15. Dewa CS, Loong D, Bonato S, Trojanowski L. The relationship between physician
- 377 burnout and quality of healthcare in terms of safety and acceptability: a systematic review.
- 378 *BMJ Open*. 2017;7(6):e015141.
- 379 16. Kakiashvili T, Leszek J, Rutkowski K. The medical perspective on burnout. *Int J*
- 380 *Occup Med Environ Health*. 2013;26(3):401-12.
- 381 17. Wurm W, Vogel K, Holl A, Ebner C, Bayer D, Mörkl S, et al. Depression-Burnout
- 382 Overlap in Physicians. *PLoS One*. 2016;11(3):e0149913.
- 383 18. Ahola K, Honkonen T, Isometsä E, Kalimo R, Nykyri E, Aromaa A, et al. The
- 384 relationship between job-related burnout and depressive disorders--results from the Finnish
- 385 Health 2000 Study. *J Affect Disord*. 2005;88(1):55-62.
- 386 19. Margiotta F, Crudden G, Byrne D, Doherty AM. Prevalence and co-variates of
- 387 burnout in consultant hospital doctors: burnout in consultants in Ireland Study (BICDIS). *Ir J*
- 388 *Med Sci*. 2019;188(2):355-64.

- 389 20. O'Dea B, O'Connor P, Lydon S, Murphy AW. Prevalence of burnout among Irish
390 general practitioners: a cross-sectional study. *Ir J Med Sci*. 2017;186(2):447-53.
- 391 21. Lovibond PF, Lovibond SH. The structure of negative emotional states: comparison
392 of the Depression Anxiety Stress Scales (DASS) with the Beck Depression and Anxiety
393 Inventories. *Behav Res Ther*. 1995;33(3):335-43.
- 394 22. Clark LA, Watson D. Tripartite model of anxiety and depression: psychometric
395 evidence and taxonomic implications. *J Abnorm Psychol*. 1991;100(3):316-36.
- 396 23. Antony MM, Bieling PJ, Cox BJ, Enns MW, Swinson RP. Psychometric properties of
397 the 42-item and 21-item versions of the Depression Anxiety Stress Scales in clinical groups
398 and a community sample. *Psychological assessment*. 1998;10(2):176.
- 399 24. Rammstedt B, John OP. Measuring personality in one minute or less: A 10-item short
400 version of the Big Five Inventory in English and German. *Journal of research in Personality*.
401 2007;41(1):203-12.
- 402 25. John OP, Donahue EM, Kentle RL. Big five inventory. *Journal of Personality and*
403 *Social Psychology*. 1991.
- 404 26. Rammstedt B. The 10-item Big Five Inventory: Norm values and investigation of
405 sociodemographic effects based on a German population representative sample. *European*
406 *Journal of Psychological Assessment*. 2007;23(3):193-201.
- 407 27. Soler JK, Yaman H, Esteva M, Dobbs F, Asenova RS, Katic M, et al. Burnout in
408 European family doctors: the EGPRN study. *Fam Pract*. 2008;25(4):245-65.
- 409 28. Amofo E, Hanbali N, Patel A, Singh P. What are the significant factors associated
410 with burnout in doctors? *Occupational medicine*. 2015;65(2):117-21.
- 411 29. Al-Humadi S, Bronson B, Muhrad S, Paulus M, Hong H, Cáceda R. Depression,
412 Suicidal Thoughts, and Burnout Among Physicians During the COVID-19 Pandemic: a
413 Survey-Based Cross-Sectional Study. *Acad Psychiatry*. 2021;45(5):557-65.
- 414 30. Garcia LC, Shanafelt TD, West CP, Sinsky CA, Trockel MT, Nedelec L, et al.
415 Burnout, Depression, Career Satisfaction, and Work-Life Integration by Physician
416 Race/Ethnicity. *JAMA Netw Open*. 2020;3(8):e2012762.
- 417 31. Doherty AM, Colleran GC, Durcan L, Irvine AD, Barrett E. A pilot study of burnout
418 and long covid in senior specialist doctors. *Ir J Med Sci*. 2021:1-5.
- 419 32. Garcia CL, Abreu LC, Ramos JLS, Castro CFD, Smiderle FRN, Santos JAD, et al.
420 Influence of Burnout on Patient Safety: Systematic Review and Meta-Analysis. *Medicina*
421 (Kaunas). 2019;55(9).
- 422 33. West CP, Dyrbye LN, Erwin PJ, Shanafelt TD. Interventions to prevent and reduce
423 physician burnout: a systematic review and meta-analysis. *Lancet*. 2016;388(10057):2272-81.
- 424 34. De Simone S, Vargas M, Servillo G. Organizational strategies to reduce physician
425 burnout: a systematic review and meta-analysis. *Aging Clin Exp Res*. 2021;33(4):883-94.
- 426 35. Dillon EC, Tai-Seale M, Meehan A, Martin V, Nordgren R, Lee T, et al. Frontline
427 Perspectives on Physician Burnout and Strategies to Improve Well-Being: Interviews with
428 Physicians and Health System Leaders. *J Gen Intern Med*. 2020;35(1):261-7.

429
430

431 **Table 1: Demographics and work related characteristics**

Characteristics		
Personal characteristics		
Age (years), mean (SD)		49.14 (6.9)
Gender, n (%)	Male Female	253 (56.6) 194 (43.4)
Region, n (%)	Dublin Munster Connacht Leinster Ulster	177 (39.2) 139 (30.6) 63 (14.0) 57 (12.6) 15 (3.3)
Relationship status, n (%)	Married Long term relationship Single Divorced/separated Widowed	374 (82.7) 16 (3.5) 33 (7.3) 25 (5.5) 4 (0.9)
Habits, n (%)	Smoking Alcohol Exercise	17 (3.7) 389 (87.8) 355 (78.9)
Work characteristics		
Specialty, n (%)	Medicine Psychiatry Surgery Anaesthesia Radiology Paediatrics Pathology Obstetrics & Gynaecology Emergency Medicine Ophthalmology	86 (19.9) 81 (18.7) 68 (15.7) 47 (10.9) 46 (10.6) 30 (6.9) 29 (6.7) 25 (5.8) 16 (3.7) 5 (1.2)
Hours spent in “face-to-face” with patients per day, mean (SD)		5.5 (4.3)
Public or private practice, n (%)	Wholly in the public system Part public/ private Part academic/ public/ private Solely private Part academic/ part public Retired Locum	206 (46.5) 173 (39.1) 28 (6.3) 18 (4.1) 12 (2.7) 4 (0.9) 2 (0.5)
Satisfaction with remuneration, n (%)		209 (47.6)
Formal disciplinary action at work, n (%)		9 (2.1)
Work-related lawsuit, n (%)		140 (31.9)
Hours worked/week, mean (SD)	Total Public setting Private Managerial Academic Other	51.8 (11.5) 30.9 (18.2) 7.25 (11.4) 3.76 (6.5) 2.1 (4.6) 0.83 (3.7)

432

433

434

435 *Table 2: Mean scores on scales measuring depression, anxiety and stress (DASS) and*
 436 *personality traits (BFI)*
 437

Scale/ Subscale	Mean (SD)	N (%) above cut-off
DASS total	10.9 (9.7)	
DASS depression subscale	3.3 (4.0)	107 (25.8%)
DASS anxiety subscale	1.6 (2.5)	58 (13.8%)
DASS stress subscale	6.0 (4.4)	245 (59.2%)
BFI - extraversion	6.3 (2.1)	
- agreeableness	7.4 (1.7)	
- conscientiousness	9.0 (1.3)	
- Neuroticism	5.0 (2.2)	
- Openness	7.4 (1.5)	
DASS = Depressive Anxiety Stress Scale BFI – Big Five Inventory		

438
 439
 440
 441
 442
 443
 444

445 *Table 3: Linear regression models with Emotional exhaustion, Cynicism and Personal Efficacy*
 446 *as the dependent variables*

447
 448

	B	p-value	Confidence Interval
<i>Emotional exhaustion</i>			
Age	-0.046	0.313	-0.136 – 0.044
Gender	0.946	0.136	-0.3 – 2.192
DASS depression	1.032	<0.001	0.882 – 1.182
Face-to-face	0.219	0.001	0.085 - 0.352
Satisfaction with remuneration	2.111	0.001	0.889 – 3.333
<i>Cynicism</i>			
Age	0.014	0.746	2.799 – 13.922
Gender	0.742	0.227	-0.073 – 0.102
DASS stress	1.033	<0.001	0.9 – 1.167
Face-to-face	0.152	0.022	0.022 – 0.281
Satisfaction with remuneration	2.681	<0.001	1.501 – 3.861
<i>Professional Efficacy</i>			
Age	-0.047	0.342	-0.143 – 0.05
Gender	-0.992	0.14	-2.312 – 0.328
DASS stress	-0.459	<0.001	-0.607 – -0.311
Face-to-face	0.196	0.024	0.026 – 0.367
Satisfaction with remuneration	-0.948	0.15	-2.241 – 0.345

449