Clinical profiling of specific diagnostic subgroups of women with chronic pelvic pain

Lysia Demetriou*[1], Michal Krassowski[1], Pedro Abreu Mendes[2], Kurtis Garbutt[1], Allison F. Vitonis[3ab], Elizabeth Wilkins[1], Lydia Coxon[1], Lars Arendt-Nielsen[4,5], Qasim Aziz[6], Judy Birch[7], Andrew W Horne[8], Anja Hoffman[9], Lone Hummelshoj[10], Claire E Lunde[1,13ab], Jane Meijlink[11], Danielle Perro[1], Nilufer Rahmioglu[1], Kathryn L. Terry[3abc], Esther Pogatzki-Zahn[12], Christine B Sieberg[13abc], Rolf-Detlef Treede[14], Christian M Becker[1], Francisco Cruz[2], Stacey A Missmer[15abcd], Krina T Zondervan[1], Jens Nagel[16], Katy Vincent[1]

1. Oxford Endometriosis Centre, Nuffield Department of Women's and Reproductive Health, University of Oxford, UK
2. IBMC/I3S and Faculty of Medicine of Porto/ Hospital S João, Porto, Portu
3. a. Boston Center for Endometriosis, Brigham and Women's Hospital and Boston Children's Hospital, Boston, Massachusetts, USA, b. Department of Obstetrics and Gynaecology, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts, USA and c. Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA
4. Center for Neuroplasticity and Pain (CNAP), SMI, Department of Health Science and Technology, School of Medicine, Aalborg University, Aalborg, Denmark
5. Department of Medical Gastroenterology, Mech-Sense, Aalborg University Hospital, Aalborg, Denmark
6. Centre for Neuroscience, Surgery and Trauma, Blizard Institute, Wingate Institute of Neurogastroenterology. Barts and The London School of Medicine and Dentistry, Queen Mary University of London UK
7. Pelvic Pain Support Network, Poole, UK
8. MRC Centre for Reproductive Health, University of Edinburgh, Edinburgh, UK
9. Research & Development, Pharmaceuticals Experimental Medicine, Bayer AG, Berlin, Germany
10. Endometriosis.org, UK
11. International Painful Bladder Foundation

12. Department of Anesthesiology, Intensive Care and Pain Medicine, University Hospital Muenster, Muenster, Germany

13. a. Biobehavioral Pain Innovations Lab, Department of Psychiatry & Behavioral Sciences, Boston Children's Hospital, Boston, MA; b. Pain and Affective Neuroscience Center, Department of Anesthesiology, Critical Care, & Pain Medicine, Boston Children's Hospital, Boston, MA; c. Department of Psychiatry, Harvard Medical School, Boston, MA

14. Department of Neurophysiology, Mannheim Center for Translational Neuroscience (MCTN), Heidelberg University, Mannheim, Germany

15. a. Department of Obstetrics, Gynecology, and Reproductive Biology; College of Human Medicine, Michigan State University, Grand Rapids, MI, USA. b. Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA, USA. c. Division of Adolescent and Young Adult Medicine, Department of Pediatrics, Boston Children's Hospital and Harvard Medical School, Boston, MA, USA. d. Boston Center for Endometriosis, Brigham and Women's Hospital and Boston Children's Hospital, Boston, MA, USA

16. Pharmaceuticals Division, Research and Early Development, Therapeutic Area Endocrinology, Metabolism and Reproductive Health, Exploratory Pathobiology, Bayer AG, Wuppertal, Germany.

Correspondence:

*Lysia Demetriou

lysimachi.demetriou@wrh.ox.ac.uk

Abstract

Chronic pelvic pain (CPP) is a common condition affecting up to 26.6% of women, with

many suffering for several years before diagnosis and/or treatment. Its clinical presentation is

varied and there are frequently comorbid conditions both within and outside the pelvis. We

aim to explore whether specific subgroups of women with CPP report different clinical

symptoms and differing impact of pain on their quality of life (QoL).

The study is part of the Translational Research in Pelvic Pain (TRiPP) project which is a

cross-sectional observational cohort study. The study includes 769 female participants of

reproductive age who completed an extensive set of questions derived from standardised

WERF EPHect questionnaires. Within this population we defined a control group (reporting

no pelvic pain, no bladder pain syndrome, and no endometriosis diagnosis, N=230) and four

pain groups: endometriosis-associated pain (EAP, N=237), interstitial cystitis/bladder pain

syndrome (BPS, N=72), comorbid endometriosis-associated pain and BPS (EABP, N=120),

and pelvic pain only (PP, N=127).

Clinical profiles of women with CPP (13-50 years old) show variability of clinical symptoms.

The EAP and EABP groups scored higher than the PP group (p<0.001) on the pain intensity

scales for non-cyclical pelvic pain and higher than both the BPS and PP groups (p<0.001) on

the dysmenorrhoea scale. The EABP group also had significantly higher scores for

dyspareunia (p<0.001), even though more than 50% of sexually active participants in each

pain group reported interrupting and/or avoiding sexual intercourse due to pain in the last 12

months.

Scores for the QoL questionnaire (SF-36) reveal that CPP patients had significantly lower

QoL across all SF-36 subscales ($p<0.001$). Significant effects were also observed between the

pain groups for pain interference with their work ($p<0.001$) and daily lives ($p<0.001$), with

the EABP suffering more compared to the EAP and PP groups ($p<0.001$).

Our results demonstrate the negative impact that chronic pain has on CPP patients' QoL and

reveal an increased negative impact of pain on the comorbid EABP group. Furthermore, it

demonstrates the importance of dyspareunia in women with CPP. Overall, our results

demonstrate the need for further exploration of interventions targeting QoL more broadly and

suggest that novel approaches to classifying women with CPP are needed.

Introduction

Chronic pelvic pain (CPP) is common, affecting up to 26.6% of women (1–4), yet it remains difficult to treat. For some women an associated pathology such as endometriosis, adenomyosis, inflammatory bowel disease or an entrapment neuropathy can be identified, but for many others diagnostic investigations will be normal and the label chronic pelvic pain syndrome (CPPS) applied. However, even in those women where, for example, endometriosis is identified the lack of correlation between measures of disease severity and pain symptoms (5,6) makes it challenging to understand the extent to which symptoms are actually caused by endometriosis and to predict the benefit of treatments targeting the ectopic tissue.

There is increasing evidence that chronic pain conditions share many features whatever the underlying cause (7,8), leading some to suggest that chronic pain should be treated as a condition in itself (9,10). Whilst there is relatively limited research on this in the context of chronic pelvic pain, studies that have looked at these features confirm similarities with other chronic pain conditions such as irritable bowel syndrome (IBS), rheumatoid arthritis and fibromyalgia (11,12). However, current clinical practice guidelines are still relatively focussed on the identification and treatment of underlying pelvic pathology (13).

The Translational Research in Pelvic Pain (TRiPP) project (https://www.imi-paincare.eu/PROJECT/TRIPP/) aims to better understand the mechanisms generating and maintaining pelvic pain with a particular focus on endometriosis and interstitial cystitis/bladder pain syndrome (IC/BPS) (14). Here, we use the baseline data from the TRiPP project to determine whether women with CPP in different diagnostic groups report different clinical symptoms and impact on their lives or whether in fact these diagnostic labels do not

help us stratify this patient group. A better understanding of the relationship between clinical

symptoms, diagnostic grouping and quality of life (QoL) would allow the development of

refined clinical pathways and the prioritisation of specific subgroups of women for further

research. Given the diversity of symptoms described by women with all forms of CPP we do

not expect to see differences in specific clinical symptoms between those with and without

endometriosis or with and without IC/BPS (with the exception of bladder symptoms

themselves). However, we hypothesise that those women with comorbid pain and bladder

symptoms will describe a poorer quality of life.

Methods

Ethical Approval

The study has received ethical approval from Yorkshire & The Humber – South Yorkshire

Research Ethics Committee (19/YH/0030) with local site approvals and was being conducted

in accordance with the principles of the Declaration of Helsinki and with relevant regulations

and Good Clinical Practice. Informed consent was obtained from all the participants, and they

were informed that they are free to withdraw their data from the study at any time.

Study population

As described in the TRiPP protocol paper (14), participants were identified from two existing

endometriosis cohort studies in Oxford (EndOX: A study to identify possible biomarkers in

women with endometriosis, Oxford REC ref 09/H0604/58) (N=276) and Boston (The

Women's Health Study from Adolescence to Adulthood (A2A), IRB-P00004267) (N=494)

plus 16 BPS participants who were recruited at Hospital São João/Instituto de Biologia

Molecular e Celular (IBMC) in Porto (Supplementary I). It is important to note that both

parent studies also included control participants with no pelvic pain symptoms.

The study comprised of five study groups and participants were assigned according to

specific inclusion and exclusion criteria for each study group (14) (Table I): endometriosis-

associated pain (EAP: N=237), comorbid endometriosis associated- and bladder pain

syndrome (EABP: N= 120), bladder pain syndrome (BPS: N=72), pelvic pain only (PP:

N=127) and controls (CON: N=230)). Information relating to these criteria was available in

the baseline questionnaires participants completed when recruited to the parent study in

UOXF and BCH. At IBMC participants were clinically diagnosed with BPS and it was

reconfirmed after completion of the baseline questionnaire that they met the criteria for

inclusion in this group.

All participants were women of reproductive age (13-50 years). Pregnant, lactating or women

planning pregnancy during the course of the study were not included as per the study

exclusion criteria.

Study design

The TRiPP study design is described in Supplementary I (and more fully elsewhere (14)), the

present report focuses on Phase I in which participants completed expanded World

Endometriosis Research Foundation (WERF) Biobanking Harmonisation Project (EPHect)

questionnaires (15). Here we explore a selected set of outcomes as described below that will

also be used in the analysis of phases II and III of the TRiPP project.

Study Data

<u>Demographics and Reproductive History</u>

Demographic information collected included participants' age, body mass index (BMI), race,

education, work and relationship status. Information about the participants' gynaecological

history consisted of: occurrence and frequency of pelvic pain, menstrual history (e.g.

menarche and dysmenorrhoea onset), use of hormones in the last 3 months and information

on fertility and dyspareunia. Data from a number of questions were analysed to give a fuller

picture of sexual activity, associated pain and functional impact. The questions on

dyspareunia were not included in the first version of the questionnaire that was completed by

67 participants in Oxford and for participants younger than 18 years old (N=79) from the

Boston cohort.

Pelvic pain

Pain was assessed using Numerical Rating Scales (NRS) for intensity ranging from 0 = "no

pain" to 10 = "worst imaginable pain". Participants were asked to complete NRS for the

experience of pelvic pain at its "worst" in the last three months (for non-cyclical pelvic pain),

during the last period (for dysmenorrhoea), worst bladder pain in the last 7 days and pain

during and 24hours post sexual intercourse (for dyspareunia).

The Pain Catastrophising Scale (PCS) (16) was used to assess pain-related worry.

Catastrophizing is the single most important risk factor that impairs the effectiveness of pain-

relieving interventions (17,18), and it statistically mediates the prospective influence of

factors such as anxiety on pain outcomes (19). The PCS is a validated/standardised 13-item

questionnaire that uses a 5-point scale ranging from 0 (not at all) to 4 (all the time), the total

PCS score was computed by summing the responses to all 13 items (16). The scores for the

three subscales: Rumination (e.g.,"I can't stop thinking about how much it hurts."),,

Magnification (e.g., "I worry that something serious may happen.") and Helplessness (e.g.,

"There is nothing I can do to reduce the intensity of my pain.") were computed by summing

responses of specific items as outlined in Sullivan (16). Scores above the "clinical cut-off

points" were considered as clinically relevant levels of catastrophising (PCS Total: 30,

Rumination: 11, Magnification: 5, Helplessness: 13).

<u>Medical comorbidities</u>

Participants were provided with a list of medical conditions and reported whether they had

received a medical diagnosis for any of those at any point in their lives. The list included

autoimmune, gynaecological, mental health, chronic pain, endocrine and cardiovascular

disorders.

Furthermore, given the prevalence of bowel symptoms in association with both endometriosis

and IC/BPS, irritable bowel syndrome (IBS) was assessed based on the Rome III criteria

using a question about the participants' bowel movements/stools when they experienced non-

cyclical pelvic pain in the last 3 months. The symptoms consisted of: a) pelvic pain getting

worse/better after bowel movement, b) more/less frequent bowel movement when pain

started and c) looser/harder stools when the pain started. Participants who reported having

two or more bowel symptoms for "most of the time" or "always" met the criteria for IBS.

<u>Quality of life</u>

The participants' health status was assessed using the Short Form Health Survey (SF-36)

which consists of 36 questions covering eight health domains: physical functioning, physical

role limitations, bodily pain, general health, emotional role limitations and mental health. The

questionnaire was scored as per instructions by Rand Corporation

(https://www.rand.org/health-care/surveys_tools/mos/36-item-short-form/scoring.html) and

Burholt et al. (20). A higher score indicates a better health status.

To assess pain interference with the participants' lives we used a question assessing the

impact of pain on six different aspects (work or school, daily activities at home, sleep, sexual

intercourse, exercise/sports and social activities). Participants had to rate the extent of pain

interference using a 4-point Likert scale ranging from "Not at all" to "Extremely" for the last

3 months. The scores were recoded into four groups: low, medium, high and no pain

interference or not applicable. Additionally, two NRSs were used to assess the severity of the

impact of pelvic pain on the participants' work and personal daily activities productivity

during the past 4 weeks.

Factors worsening and relieving pain

Two multiple-choice questions were used to assess/identify factors that participants believe

worsen or relieve their pelvic pain. Among others, the options included visceral functions

such as bowel movement or bladder emptying; behavioural including exercising,

standing/walking or sitting; wellbeing factors like stress and meditation; or other factors e.g.

time of day and having a full meal.

Statistical analysis

The data were entered into a REDCap database (21) and analysed using IBM SPSS Statistics

software, version 27. The analysis was conducted in 2 phases. First, descriptive analysis was

run to identify the characteristics of each study group. Then, statistical testing was undertaken

to investigate any effects between the study groups for the study outcomes. Statistical testing

for the NRS pelvic and bladder pain did not include the control group since they were

recruited based on their NRS scores being <3/10. One-way ANOVAs and Bonferroni

corrected post-hoc tests were used to compare between study groups for all the continuous

variables while frequencies, percentages and non-parametric tests (chi-square, Mann-

Whitney U tests) were employed for all the categorical variables. Pearson's correlations were

run to explore relationships between the variables. Bonferroni correction was used to account

for multiple comparisons.

Results

<u>Demographics and Reproductive history</u>

A total of 785 participants were included in this phase of the study with a mean age of 27.6

years (STD: 8.1) and mean BMI of 24.8 (STD: 5.2) (Table II). A one-way ANOVA showed

significant differences for age but not for BMI between groups (Age: $F_{(4,781)}=4.12$,

$p=0.003$, BMI: $F_{(4, 757)}=3.28$, $p=0.011$); post-hoc independent-samples t-tests showed that

the BPS group was significantly older than the EABP ($p=0.01$) and Controls ($p=0.04$).

The age of menarche (mean=12.2 years) ranged from 8 to 16 years old (One-way ANOVA:

$F_{(4,768)}=0.715$, $p=0.582$) and did not significantly differ among groups (Table II). Likewise,

the age at which dysmenorrhoea commenced was similar among all study groups (mean=14.3

years, $F_{(4,631)}=1.10$, $p=0.356$) (Table II). On average, patients suffered for approximately 8

years with non-cyclical pelvic pain, and there were no significant differences between the

patient groups, even though the PP group had the lowest mean number of years (mean=4.6

years), while the EABP and BPS groups had the highest mean number of years (EABP:

mean=9.7 years, BPS: mean=9.9 years) ($F_{(4,204)}=3.906$, $p=0.004$). A Pearson's correlation

showed that the number of years suffering with pelvic pain positively correlated with the age

of the participants ($r=0.411$, $p<0.001$).

More participants in the PP (90.5%) and Controls (92.6%) group reported having periods

during the last three months than participants in the EAP (76.9%), EABP (63.9%) and BPS

(71.8%) (Table III). Of the participants who reported having periods, more than 53.8% of the

controls and 35.0% of the pain groups reported having periods whilst using exogenous

hormones in the last 3 months. The most common reason for not having periods across the

groups was taking hormones continuously (91.8%). Regardless of whether they had periods,

many participants reported using exogenous hormones in the last three months (EAP: 52.7%,

EABP: 61.7%, BPS: 61.1%, PP: 39.4%, Controls: 57%) (Table III).

<u>Pelvic pain</u>

The pain rating for non-cyclical pain at its worst during the last three months was

significantly higher in the EAP and EABP compared to the PP group (Table IV). When asked

about dysmenorrhoea at its worst the EABP and EAP groups had a significantly higher mean

score on pain severity compared to both the BPS and PP groups ($p<0.001$). Results for the

NRS scale on bladder pain during the last seven days confirmed significantly higher mean

pain scores for the BPS and EABP groups compared to the EAP and PP groups ($p<0.001$) as

defined by the inclusion criteria (Figure I).

Regarding questions on dyspareunia, there were missing responses for more than 50% of our

participants, reasons for missingness include: a) 4.8% (n=38) of all the participants did not

wish to respond to the questions, b) 7.5% (n=59) reported never having sexual intercourse c)

these questions were not available to 10.1% (n=79) due to their age at time of completion as

described in the methods section above and d) 20.2% (n=150) reported not having sex in the

last 12 months and therefore were only included in the analysis focussed on ever having

experienced pain with or after intercourse.

From the available data sexual intercourse data however, most participants in each of the pain

groups (>58%) reported that they have experienced pain during intercourse or in the 24 hours

following vaginal sexual intercourse (Table V). Of those, more than 60% experienced

dyspareunia during the last month across all the pain groups. Almost half of those

participants (49.2%) in the EABP group said that they had always experienced dyspareunia

during intercourse in the last 12 months as opposed to the BPS and PP groups in which most

participants reported occasional dyspareunia in the last year. In the EAP group >77% of

participants reported experiencing dyspareunia often, usually or always (Table V).

Concerning the location of dyspareunia, most participants in each pain group reported feeling

pain deep inside the vagina, the second most common location was in the abdomen/pelvis.

When asked whether they have been avoiding or interrupting sexual intercourse during the

last 12 months due to pelvic pain, more than half of the participants (>54%) in all four pain

groups said that they have interrupted sexual intercourse while more than half (>51%) of the

participants in the EAP, EABP and BPS groups reported avoiding it.

Further analysis showed effects between the groups for the two dyspareunia NRS scales (pain

during intercourse: $F_{(4,240)}=12.77$, $p<0.001$, pain in the 24hrs post-intercourse:

$F_{(4,242)}=12.33$, $p<0.001$) (Figure I). The EABP group had significantly higher mean ratings

on the NRS scales than all other pain groups ($p<0.001$) for the pain felt during intercourse,

and significantly higher mean rating for pain severity in the 24hours after the sexual

intercourse than the EAP and PP groups (Table IV and Figure I).

<u>Medical comorbidities</u>

Across all pain groups the most common comorbidities were depression, anxiety, migraine,

asthma, and irritable bowel syndrome (IBS) (Table VI). Within the EAP, EABP and BPS

groups there were higher percentages of depression, anxiety and migraine diagnosis (>28%)

than in the PP group (<22%), while the frequency of asthma ranged between 20% - 27%

across all pain groups (Table VI). Statistical analysis using Mann-Whitney U tests revealed

that when compared to the control group only the EABP group has significantly higher

frequency of anxiety diagnosis (anxiety: $(U(N_{EABP}=114, N_{CON}=221)=10510, z=-3.39,$

$p<0.001$), and depression: $(U(N_{EABP}=120, N_{CON}=230)=10855, z=-4.36, p<0.001)$). However,

statistical analysis for migraine showed that it was significantly more common in EAP

(28.7%), EABP (34.2%) and BPS (40.2%) groups than the control group (9.6%)

$((U(N_{EAP}=237, N_{CON}=230)=22042, z=-5.23, p<0.001), (U(N_{EABP}=120, N_{CON}=230)=10405,$

$z=-5.68, p<0.001), (U(N_{BPS}=72, N_{CON}=230)=5737, z=-6.60, p<0.001))$ groups. Similarly, IBS

diagnosis was also significantly more common in the EAP (18.6%), EABP (24.2%) and BPS

(20.8%) groups when compared to the control group (4.8%) $((U(N_{EAP}=237,$

$N_{CON}=229)=23402, z=-4.60, p<0.001), (U(N_{EABP}=120, NCON=230)=11079, z=-5.39,$

$p<0.001), (U(NBPS=72, NCON=229)=6922, z=-4.22, p<0.001))$.

IBS diagnosis was also less common in the PP group (10%) compared to the EAP, EABP and

BPS groups (>18%).

Further analysis of the bowel symptoms revealed that more than 11% of the participants in

the pain groups met the criteria for IBS regardless of whether they reported having a medical

diagnosis for it in the comorbidities question. The EABP and PP groups had the higher

percentages of participants meeting the IBS criteria (>20%) (Table VI).

<u>Quality of life</u>

Analysis of the SF-36 questionnaire using a one-way ANOVA revealed significant effects

between the five study groups across all eight SF-36 domains ($p<0.001$). Specifically, the

EABP group suffered the most while the control group had the best scores across all sub-

scales ($p<0.001$) (Table VII). Significant effects were also observed between the four pain

groups for pain interference with activities at work and daily life (work: $F(3,209)=9.76$,

$p<0.001$; daily activities: $F(3,244)=10.51$, $p<0.001$) (Figure II). Post-hoc tests revealed that

the EABP group had significantly higher pain interference scores than the EAP and PP

groups at both scales ($p<0.001$) (Table IV).

<u>Pain Catastrophising</u>

Pain catastrophising was higher in the EABP group both in the mean total score and in the

mean scores of all three subscales ($p<0.001$). The EAP and BPS groups also had a higher

pain catastrophising score than both the PP and Controls groups ($p<0.001$). According to the

clinical-cut-offs more than 19% of the participants in the EAP, EABP and BPS had clinical

levels of pain catastrophising. The EABP group had the highest percentage of people above

the clinical cut-off (36.0%) (Table VII). A Pearson's correlation between all the pain

catastrophising and all the NRS pain scales across all CPP patients revealed strong positive

correlations ($p<0.001$).

<u>Factors worsening and relieving pain</u>

On average, participants from all groups chose three to four factors that either relieve or

worsen their pelvic pain. Across the four pain groups the three most common factors for

worsening pelvic pain were: stress (23.6%), full bladder/urinating (23.3%) and exercising

(20.2%). The most frequently reported factors for relieving pelvic pain were: pain medication

(31.4%), lying down (31.0%) and heating pad (29.5%) (Table IX). Analysis of bowel

movement and bladder emptying as visceral factors, shows that less people in the EAP and

the PP groups report these factors as relieving (bowel movement: EAP: 19.8% and PP 15.7%;

emptying bladder: EAP: 9.3% and PP 4.7%) or worsening (bowel movement: EAP: 6.7% and

PP 2.5%; emptying bladder: EAP: 24.1% and PP 10.2%), while in the EABP and BPS groups

more participants report visceral factors as worsening (bowel movement: EABP: 27.7% and

BPS 9.5%; emptying bladder: EABP: 65.8% and BPS 44.4%) rather than relieving (bowel

movement: EABP: 32.5% and BPS 20.8%; emptying bladder: EAP: 34.2% and BPS 23.6%).

Discussion

This study phenotypically describes the clinical profiles of women with CPP, with a

particular focus on endometriosis and IC/BPS, and contrasts them with controls. In line with

existing literature, we have demonstrated that regardless of the aetiology, CPP has a negative

impact on the lives of people who suffer with it. However, our results show that there are

important differences between different clinical subgroups of those with CPP and those with

comorbid endometriosis and bladder pain syndrome (EABP) are particularly severely

affected. The heterogeneity in factors exacerbating and relieving pelvic pain, highlights the

complexity of the condition but suggests that there may be a variety of different underlying

pain mechanisms contributing to the similar clinical presentation. Further evaluation of these

underlying mechanisms may provide more clinically relevant approaches to patient

stratification than focussing on diagnosis alone.

Importantly, we highlight how common painful sexual intercourse is in women with CPP and

that half of those who answered these questions reported interrupting or avoiding it due to

pain. This is an important issue for women and their partners and it can have a negative effect

on more than one aspect of a woman's life, including mental health, body image, relationship

with their partner and fertility (22,23). Dyspareunia is a commonly described symptom of

endometriosis, but there is less of a focus on this issue in IC/BPS. It is therefore particularly

notable that the highest dyspareunia scores were reported by those with bladder symptoms in

our cohort. This is in line with previous work in women with endometriosis, demonstrating

that the presence of tenderness of the bladder/pelvic floor or a diagnosis of IC/BPS is

associated with more severe dyspareunia in those with all stages of endometriosis (24). A

better understanding of the mechanisms underlying dyspareunia in all women with CPP not

just those with deep endometriosis is clearly an important future need and will ultimately

improve the clinical care of those suffering with this important symptom (25).

It is perhaps unsurprising that those with comorbid endometriosis and bladder pain symptoms

scored highly on NRSs for all types of pelvic pain (non-cyclical pelvic pain, bladder pain,

dysmenorrhoea and dyspareunia). However, our data demonstrate that these women also have

poorer quality of life and report greater pain interference. Whilst there are studies exploring

the prevalence and comorbidity of endometriosis and subsequent diagnosis of IC/BPS (26–

28), few studies focus on those with comorbid symptoms, either in terms of exploring clinical

features or considering specific treatment regimes. Moreover, for many clinical studies the

presence of a comorbidity is an exclusion criteria and thus very limited data exists, this is

particularly true for clinical trials. Our findings suggest this may be a priority group to

consider in future work.

Our study focuses on two specific diagnoses, endometriosis and IC/BPS, however there are

many other diagnoses associated with CPP. The EPHect Clinical Covariates questionnaire

captures some of these: IBS, fibroids, pelvic inflammatory disease, inflammatory bowel

disease and a limited number of musculoskeletal conditions (spine problems including

scoliosis, rheumatoid arthritis and fibromyalgia). However, for most of these the numbers

were relatively small within our cohort and therefore it is not possible to explore their

contribution to the clinical picture in any detail. There are also a number of other factors that

have been associated with chronic pain more broadly and with CPP specifically, including

psychological and social factors, adverse childhood events and other traumatic experiences

(13). The data collected within the EPHect questionnaire does not allow these to be taken into

consideration here. Additional information is collected on some of these measures for a

smaller cohort in Phase II of TRiPP and will be interrogated to better understand their

relationship to the pain experience and mechanisms underlying any observed relationship

(14).

Given that viscero-visceral referral is a relatively well understood phenomenon (29) and that

visceral pain conditions are often comorbid (30), we had expected to see higher rates of IBS

in those with bladder symptoms. This was the case when questioning previous diagnosis of

IBS (EABP 24% and BPS 21% vs EAP 19% and PP 10%). However, considering the

answers to the questions comprising the Rome IV criteria, a higher proportion of those in the

EABP and PP groups would be considered to meet IBS criteria than in the other groups. It is

of course difficult to know whether this reflects ongoing effective treatment in at least a

proportion of those with a past diagnosis of IBS. Nonetheless, the high rates of IBS across the

cohort as a whole highlight the importance of assessing symptoms from all abdominopelvic

organs in women with CPP.

Interestingly the impact of visceral function (bladder and bowel emptying specifically) on

pelvic pain appears to be very variable. For some participants these visceral functions

worsened their pain, whilst for others they could be relieving. In fact, there was wide

variability in the factors identified as either worsening or relieving pelvic pain even within

diagnostic groups and it would be interesting to consider whether patterns within these data

may reflect underlying pain mechanisms rather than for example the presence or absence of

endometriosis or bladder symptoms. Whilst it can feel challenging in a clinical consultation

to raise the topics of mental health and the stressors of daily life, particularly when women

may have had a long journey to get their pain taken seriously, stress was the most commonly

identified worsening factor whereas relaxation was one of the most frequent relieving factors.

Moreover, anxiety and depression were commonly reported comorbidities and those in the

EAP, EABP and BPS groups all scored significantly lower than controls on the emotional

function dimension on the quality of life questionnaire. This evidence suggests that patients

could benefit from treatments focusing on managing their stress and developing better coping

mechanisms for their pelvic pain.

Whilst not an assessment of psychological wellbeing, catastrophic thinking about pain has

been shown to be an important predictor of outcomes both in the transition to chronic pain

after an acute injury and in the response to a variety of therapeutic options (31,32). In line

with other studies (33,34) we observed that the chronic pain groups (EAP, EABP and BPS)

scored higher in pain catastrophising compared to the control group with a particularly high

percentage of EABP patients meeting clinical levels of catastrophising. This highlights the

need to include behavioural interventions that specifically target pain-related worry in the

management of CPP in the same way as they are considered essential to the management of

other chronic pain conditions (35,36).

Strengths and weakness

There are a number of strengths to the TRiPP cohort which include the size of the sample and

the detailed phenotypic information available for these participants. Moreover, our study

focuses on chronic pelvic pain and therefore includes women with a variety of underlying

diagnoses rather than studying only a single condition such as endometriosis. This strategy

allows us to compare the similarities between different clinical subgroups but also to

understand differences between them. We have a particular interest in IC/BPS and therefore

this cohort is relatively unusual in allowing the comparison between endometriosis and

IC/BPS as well as those with comorbid symptoms. Unfortunately, our BPS group is smaller

than we initially planned. Neither of the original cohorts from which the TRiPP sample was

formed had specifically recruited IC/BPS patients (although many did meet the diagnostic

criteria) and therefore we had planned to expand these with a large number of new recruits

from specialist centres. However, the COVID-19 pandemic halted our ability to recruit to the

study. Overall, we still consider our sample size sufficient to draw meaningful conclusions

(192 participants with bladder pain, 120 of whom have coexisting endometriosis).

A further strength of our study is that participants were recruited from three different centres

rather than a single site. However, the duration of recruitment to the original cohorts has

meant that the baseline questionnaires have been reviewed and updated such that not all

participants completed the same version. This has reduced the number of variables we could

analyse across the full cohort. Nonetheless the EPHect questionnaire is very comprehensive

and the data we do have available allows us to phenotypically characterise our participants in

a very detailed manner.

<u>Conclusion</u>

Overall, our results demonstrate the negative impact that chronic pain has on the lives of

those with CPP and particularly the importance of dyspareunia. Whilst we do see similarities

across the specific clinical subgroups that we explored our data highlight the difference

between these groups and importantly illustrate how severely impacted the group with

comorbid endometriosis and IC/BPS are. The heterogeneity seen in many of our measures,

especially the factors exacerbating and relieving pain suggests that multiple different

mechanisms may underlie the similar clinical presentations and emphasise the importance of

better characterising those with CPP to identify clinically meaningful methods of patient

stratification. Further studies within the TRiPP project will explore these mechanisms in

greater detail towards this aim.

Conflict of interest

LD, MK, EW, LC, DP, NR, AFV, LAN, QA, JB, AH, LH, CEL, JM, CS, PAM, KG: declare

no competing interests

AWH: : reports grant funding from the MRC, NIHR, CSO, Wellbeing of Women, Roche

Diagnostics, Astra Zeneca, Ferring, Charles Wolfson Charitable Trust and Standard Life. His

employer has received consultancy fees from Roche Diagnostics, AbbVie, Nordic Pharma

and Ferring, outside the submitted work. In addition, AWH has a patent for a serum

biomarker for endometriosis pending.

AH: Employee of Bayer AG, Germany

EPZ: received financial support from Grunenthal and Mundipharma for research activities

and advisory and lecture fees from Grünenthal, Novartis and Mundipharma. In addition, she

receives scientific support from the German Research Foundation (DFG), the Federal

Ministry of Education and Research (BMBF), the German Federal Joint Committee (G-BA)

and the Innovative Medicines Initiative (IMI) 2 Joint Undertaking under grant agreement No

777500. This Joint Undertaking receives support from the European Union's Horizon 2020

research and innovation programme and EFPIA. All money went to the institution E.M.P.-Z.

is working for.

RDT: Ad board for BAYER, IASP task force on chronic pain classification

CMB: Research Grants from Bayer Healthcare, MDNA Life Sciences, Roche Diagnostics,

European Commission, NIH. His employer has received consulancy fees from Myovant and

ObsEva for work outside of this project

FC: Consultant and/or investigator for Allergan (Abbvie), Astellas, Bayer, Ipsen and

Recordati

SAM: has been an advisory board member for AbbVie and Roche and receives research

funding from the National Institutes of Health, the US Department of Defense, the J. Willard

and Alice S. Marriott Foundation, and AbbVie; none are related to the presented work; the J.

Willard and Alice S. Marriott Foundation supported enrollment of and data collection from

the A2A cohort in Boston from which TRiPP data were sampled.

KTZ: reports grant funding from EU Horizon 2020, NIH US, Wellbeing of Women, Bayer

AG, Roche Diagnostics, Evotec-Lab282, MDNA Life Sciences, outside the submitted work.

JN: Employee and shareholder of Bayer AG, Germany

KV: declares research funding from Bayer Healthcare and UKRI and honoraria for

consultancy and talks and associated travel expenses from Bayer Healthcare, Grunenthal

GmBH, AbbVie and Eli Lilly.

Author contributions

Funding

The study was funded by Innovative Medicines Initiative 2 Joint Undertaking (JU) under

grant agreement No 777500. The JU receives support from the European Union's Horizon

2020 research and innovation programme and EFPIA. Financial support was provided by the

J. Willard and Alice S. Marriott Foundation for establishment of and baseline data collection

within the A2A cohort - from which the Boston-based TRiPP population was sampled.

References

1. Eskenazi B, Warner ML. Epidemiology of endometriosis. Obstet Gynecol Clin North Am. 1997;24(2):235–58.

2. Zondervan KT, Yudkin PL, Vessey MP, Jenkinson CP, Dawes MG, Barlow DH, et al. The community prevalence of chronic pelvic pain in women and associated illness behaviour. Br J Gen Pract. 2001;51(468):541–7.

3. Zondervan KT, Becker CM, Koga K, Missmer SA, Taylor RN, Viganò P. Endometriosis. Nat Rev Dis Prim. 2018;4(1).

4. Ahangari A. Prevalence of chronic pelvic pain among women: an updated review. Pain Physician. 2014;17(2):E141–7.

5. Vercellini P, Trespidi L, De Giorgi O, Cortesi I, Parazzini F, Crosignani PG. Endometriosis and pelvic pain: relation to disease stage and localization. Fertil Steril. 1996;65(2):299–304.

6. Porpora MG, Koninckx PR, Piazze J, Natili M, Colagrande S, Cosmi E V. Correlation between endometriosis and pelvic pain. J Am Assoc Gynecol Laparosc. 1999;6(4):429–34.

7. Ayorinde AA, Macfarlane GJ, Saraswat L, Bhattacharya S. Chronic pelvic pain in women: an epidemiological perspective. Women's Heal. 2015;11(6):851–64.

8. Steege JF, Siedhoff MT. Chronic pelvic pain. Obstet Gynecol. 2014;124(3):616–29.

9. Niv D, Devor M. Chronic pain as a disease in its own right. Vol. 4, Pain Practice. Wiley Online Library; 2004. p. 179–81.

10. Taylor AM, Phillips K, Taylor JO, Singh JA, Conaghan PG, Choy EH, et al. Is chronic pain a disease in its own right? Discussions from a pre-OMERACT 2014 workshop on chronic pain. The Journal of Rheumatology; 2015.

11. Brawn J, Morotti M, Zondervan KT, Becker CM, Vincent K. Central changes associated with chronic pelvic pain and endometriosis. Hum Reprod Update. 2014;20(5):737–47.

12. Kaya S, Hermans L, Willems T, Roussel N, Meeus M. Central sensitization in urogynecological chronic pelvic pain: a systematic literature review. Pain Physician. 2013;16(4):291–308.

13. Lamvu G, Carrillo J, Ouyang C, Rapkin A. Chronic pelvic pain in women: a review. Jama.

571  2021;325(23):2381–91.

572 14. Demetriou L, Coxon L, Krassowski M, Rahmioglu N, Arendt-Nielsen L, Aziz Q, et al. Deep phenotyping of

573  women with endometriosis-associated pain and bladder pain syndrome: the TRiPP (Translational

574  Research in Pelvic Pain) study protocol. medRxiv. 2022;

575 15. Vitonis AF, Vincent K, Rahmioglu N, Fassbender A, Buck Louis GM, Hummelshoj L, et al. World

576  Endometriosis Research Foundation Endometriosis Phenome and biobanking harmonization project: II.

577  Clinical and covariate phenotype data collection in endometriosis research. Fertil Steril.

578  2014;102(5):1223–32.

579 16. Sullivan MJL, Bishop SR, Pivik J. The pain catastrophizing scale: development and validation. Psychol

580  Assess. 1995;7(4):524.

581 17. Hill JC, Lewis M, Sim J, Hay EM, Dziedzic K. Predictors of poor outcome in patients with neck pain

582  treated by physical therapy. Clin J Pain. 2007;23(8):683–90.

583 18. Karels CH, Bierma-Zeinstra SMA, Burdorf A, Verhagen AP, Nauta AP, Koes BW. Social and psychological

584  factors influenced the course of arm, neck and shoulder complaints. J Clin Epidemiol. 2007;60(8):839–

585  48.

586 19. Pinto PR, McIntyre T, Almeida A, Araújo-Soares V. The mediating role of pain catastrophizing in the

587  relationship between presurgical anxiety and acute postsurgical pain after hysterectomy. Pain.

588  2012;153(1):218–26.

589 20. Burholt V, Nash P. Short Form 36 (SF-36) Health Survey Questionnaire: normative data for Wales. J

590  Public Health (Bangkok) [Internet]. 2011 Dec 1;33(4):587–603. Available from:

591  https://doi.org/10.1093/pubmed/fdr006

592 21. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture

593  (REDCap)—a metadata-driven methodology and workflow process for providing translational research

594  informatics support. J Biomed Inform. 2009;42(2):377–81.

595 22. Seehusen DA, Baird D, Bode D V. Dyspareunia in women. Am Fam Physician. 2014;90(7):465–70.

596 23. Ferrero S, Esposito F, Abbamonte LH, Anserini P, Remorgida V, Ragni N. Quality of sex life in women

597  with endometriosis and deep dyspareunia. Fertil Steril. 2005;83(3):573–9.

598 24. Orr NL, Noga H, Williams C, Allaire C, Bedaiwy MA, Lisonkova S, et al. Deep Dyspareunia in

599  Endometriosis: Role of the Bladder and Pelvic Floor. J Sex Med [Internet]. 2018;15(8):1158–66.

600   Available from: https://doi.org/10.1016/j.jsxm.2018.06.007

601   25.   Vercellini P, Meana M, Hummelshoj L, Somigliana E, Viganò P, Fedele L. Priorities for endometriosis

602   research: a proposed focus on deep dyspareunia. Reprod Sci. 2011;18(2):114–8.

603   26.   Wu C, Chung S, Lin H. Endometriosis increased the risk of bladder pain syndrome/interstitial cystitis: A

604   population-based study. Neurourol Urodyn. 2018;37(4):1413–8.

605   27.   Chung MK, Chung RP, Gordon D, Jennings C. The evil twins of chronic pelvic pain syndrome:

606   endometriosis and interstitial cystitis. JSLS J Soc Laparoendosc Surg. 2002;6(4):311.

607   28.   DiVasta AD, Zimmerman LA, Vitonis AF, Fadayomi AB, Missmer SA. Overlap between irritable bowel

608   syndrome diagnosis and endometriosis in adolescents. Clin Gastroenterol Hepatol. 2021;19(3):528–37.

609   29.   Aziz Q, Giamberardino MA, Barke A, Korwisi B, Baranowski AP, Wesselmann U, et al. The IASP

610   classification of chronic pain for ICD-11: chronic secondary visceral pain. Pain. 2019;160(1):69–76.

611   30.   Affaitati G, Costantini R, Tana C, Cipollone F, Giamberardino MA. Co-occurrence of pain syndromes. J

612   Neural Transm. 2020;127(4):625–46.

613   31.   Martinez-Calderon J, Jensen MP, Morales-Asencio JM, Luque-Suarez A. Pain catastrophizing and

614   function in individuals with chronic musculoskeletal pain. Clin J Pain. 2019;35(3):279–93.

615   32.   Oosterhaven J, Wittink H, Dekker J, Kruitwagen C, Devillé W. Pain catastrophizing predicts dropout of

616   patients from an interdisciplinary chronic pain management programme: a prospective cohort study. J

617   Rehabil Med. 2019;51(10):761–9.

618   33.   Sewell M, Churilov L, Mooney S, Ma T, Maher P, Grover SR. Chronic pelvic pain–pain catastrophizing,

619   pelvic pain and quality of life. Scand J pain. 2018;18(3):441–8.

620   34.   Martin CE, Johnson E, Wechter ME, Leserman J, Zolnoun DA. Catastrophizing: a predictor of persistent

621   pain among women with endometriosis at 1 year. Hum Reprod. 2011;26(11):3078–84.

622   35.   Schütze R, Rees C, Smith A, Slater H, Campbell JM, O'Sullivan P. How can we best reduce pain

623   catastrophizing in adults with chronic noncancer pain? A systematic review and meta-analysis. J Pain.

624   2018;19(3):233–56.

625   36.   de C Williams AC, Fisher E, Hearn L, Eccleston C. Psychological therapies for the management of

626   chronic pain (excluding headache) in adults. Cochrane database Syst Rev. 2020;(8).

629   **Figures**

631   ***Figure 1.*** *Mean scores and distribution of the Numerical Rating Scales (NRS) for the five study groups on dysmenorrhea*
632   *(EAP: N=204, EBAP: N=100, BPS: N=59, PP: N=115), non-cyclical pelvic pain (EAP: N=151, EBAP: N=95, BPS: N=42, PP: N=56),*
633   *dyspareunia (EAP: N=101, EABP: N=61, BPS: N=29, PP: N=41) and post-coital pain (EAP: N=101, EBAP: N=63, BPS: N=29, PP:*
634   *N=41).*

Figure 2. Mean scores and distribution of the Numerical Rating Scales (NRS) (impact of pain on work (EAP: N=114, EBAP: N=41, BPS: N=20, PP: N=38, CNP: N=18) and personal life (EAP: N=125, EBAP: N=55, BPS: N=23, PP: N=45, CNP: N=19)); and the physical functioning and emotional wellbeing sub-scales of the SF-36 questionnaire (physical functioning (EAP: N=232, EBAP: N=119, BPS: N=72, PP: N=126, CNP: N=227) and emotional wellbeing (EAP: N=230, EBAP: N=120, BPS: N=72, PP: N=125, CNP: N=226)) for the five study groups.

	Surgical confirmation of endometriosis	Urinary Symptoms	Pain perceived as arising from the bladder	Pelvic pain score on NRS (at least one)
EAP (N=237)	Yes	No	No	> 4/10
EABP (N=120)	Yes	Yes	Yes	> 4/10
BPS (N=72)	No	Yes	Yes	> 4/10
PP (N=127)	No	No	No	> 4/10
Control (N=230)	No	No	No	< 3/10

Table I. Study design and inclusion criteria for the five study groups: endometriosis-associated pain (EAP, endometriosis-associated and bladder pain (EABP), bladder pain syndrome (BPS), pelvic pain only (PP) and controls).

	EAP			EABP			BPS			PP			CNP		
	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD
Age	237	28.11	9.53	120	26.45	8.91	72	30.43	9.07	127	28.06	6.63	230	26.57	6.05
BMI	225	24.82	4.83	116	26.06	5.84	71	24.13	4.44	118	25.19	5.74	228	24.10	4.93
Menarche Age	231	12.14	1.59	117	12.09	1.41	72	12.14	1.59	124	12.26	1.34	229	12.32	1.36
Age period pain commenced	184	14.36	4.33	108	13.65	3.08	60	14.23	3.50	102	14.37	4.03	182	14.55	3.04

Table II. Key participant characteristics (age, body mass index (BMI), age of menarche and age period pain commenced). Data shown as numbers (N), mean and standard deviation (SD).

	EAP		EABP		BPS		PP		CNP	
	N	%	N	%	N	%	N	%	N	%
Period in the last 3 months										
No	54	23.1	43	36.1	20	28.2	12	9.5	17	7.4
Yes	180	76.9	76	63.9	51	71.8	114	90.5	213	92.6
Hormone use in the last 3 months	125	52.7	74	61.7	44	61.1	50	39.4	131	57
Reason for no periods in last 3 months										
Taking hormones continusously	52	94.5	43	100	12	60	11	100	16	94.1
Pregnant/breastfeeding	0	0	0	0	1	5	0	0	1	5.9
Unsure	1	1.8	0	0	2	10	0	0	0	0
Other	2	3.6	0	0	5	25	0	0	0	0
Reason for using hormones										
Birth Control	127	53.6	62	51.7	34	47.2	79	62.2	136	59.1
Pelvic or other period pain	153	64.6	94	78.3	21	29.2	23	18.1	27	11.7
Irregular periods	61	25.7	47	39.2	9	12.5	9	16.5	39	17
Heavy periods	105	44.3	76	63.3	19	26.4	28	22	31	13.5
Acne	19	9.7	16	14.4	8	11.1	18	16.4	38	17.1
Polycystic Ovary Syndrome	12	5.2	7	6	3	4.2	6	4.8	7	3
Ovarian Cyst	20	10.5	19	17.3	1	1.4	1	1	4	1.8
Periods in the last 3 months										
Natural	109	60.9	44	57.9	27	51.9	74	65.5	98	46.2
Hormone Induced	70	39.1	32	42.1	25	48.1	39	34.5	114	53.8
Ever tried to get pregnant for more than 6 months										
No	154	71.6	87	79.8	51	94.4	102	83.6	210	93.3
Yes	61	28.4	22	20.2	3	5.6	20	16.4	15	6.7
Ever seek treatment for infertility in any clinic										
No	178	84	95	90.5	56	98.2	106	90.6	205	94.5
Yes	34	16	10	9.5	1	1.8	11	9.4	12	5.5
Pelvic pain										
No	64	32.7	20	18	25	37.3	60	54.5	206	93.2
Yes	132	67.3	91	82	42	62.7	50	45.5	15	6.8
How long did you have pelvic pain in the last 3 months										
Less than one day	21	17.8	26	29.2	19	46.3	8	21.1	1	14.3
One day	4	3.4	4	4.5	4	9.8	3	7.9	2	28.6
Two to three days	26	22	13	14.6	5	12.2	14	36.8	2	28.6
One day a week	11	9.3	7	7.9	1	2.4	1	2.6	1	14.3
More than one day a week	40	33.9	23	25.8	11	26.8	12	31.6	1	14.3
Every day	16	13.6	16	18	1	2.4	0	0	0	0

Table III. Gynecological History. Data shown as percentages (%) by study group.

	EAP			EABP			BPS			PP			CNP		
	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD
Dysmenorrhea															
Worst period pain during last period	204	6.24	3.23	100	5.92	3.66	59	3.93	3.16	115	5.37	2.91	188	2.11	2.19
Pelvic Pain (non-cyclical)															
Pelvic pain worst severity last 3 months	151	7.37	2.07	95	7.88	1.74	42	7.24	2.12	56	6.82	2.50	12	2.42	1.38
Pelvic pain worst severity during last gynaecological examination	87	5.24	2.87	54	7.35	2.20	27	6.70	3.10	32	5.75	2.62	7	0.86	0.91
Bladder Pain															
Bladder pain worst severity last 7 days	95	0.95	1.61	56	6.61	1.56	29	6.52	1.53	327	0.78	1.40	9	0.11	0.33
Dyspareunia															
Worst pain severity during last intercourse	101	5.51	2.46	61	6.77	2.25	29	6.34	2.16	41	4.76	2.51	13	2.23	1.01
Worst pain severity 24hrs post last intercourse	101	3.97	2.66	63	6.05	2.46	29	4.79	2.82	41	4.29	2.70	13	1.15	1.21
Impact on work and personal life															
Work productivity during past 4 weeks	114	3.70	3.70	41	5.88	2.14	20	5.65	3.05	38	3.39	2.99	18	0.22	0.65
Daily activities productivity during past 4 weeks	125	4.06	2.81	55	6.31	2.02	23	5.57	2.78	45	4.04	4.04	19	0.26	0.73

Table IV. Mean score of the Numerical Rating Scales (NRS) on dysmenorrhea, non-cyclical pelvic pain, bladder pain, dyspareunia and impact of pain on work and personal life. Data shown as numbers (N), mean and standard deviation (SD).

	EAP		EABP		BPS		PP		CNP	
	N	%	N	%	N	%	N	%	N	%
Ever had pain during intercourse or in the 24hrs following vaginal sexual intercourse/penetration?										
No	35	25.2	7	10.0	13	21.0	39	41.9	132	76.7
Yes	104	71.8	63	90.0	49	79.0	54	58.1	40	23.3
When did you last have vaginal intercourse										
In the last month	74	70.5	41	64.1	31	62.0	44	77.2	27	69.2
1-3 months ago	13	12.4	10	15.6	6	12.0	2	3.5	6	15.4
4-12 months ago	7	6.7	10	15.6	8	16.0	4	7.0	2	5.1
More than 12 months ago	11	10.5	3	4.7	5	10.0	7	12.3	4	10.3
Pain location during last vaginal intercourse/penetration										
Entrance of the vagina	15	6.4	21	17.9	12	16.7	6	4.8	5	2.2
Deep inside the vagina	61	26.2	49	41.9	24	33.3	22	17.5	7	3
Entrance and deep inside the vagina	5	13.5	3	50.0	0	0	3	18.8	0	0
In the abdomen/pelvis	55	28.1	52	46.8	19	26.4	24	21.8	3	1.4
Other location	5	2.6	9	8.1	2	2.8	2	1.8	0	0
Frequency of pelvic pain during or 24hrs after vaginal intercourse/penetration in the last 12 months										
Never	1	0.9	-	-	3	7.0	1	1.8	5	13.5
Occasionally (less than a quarter of times)	23	21.7	8	12.3	18	41.9	28	50.9	24	64.9
Often (a quarter to half of the times)	29	27.4	10	15.4	4	9.3	9	16.4	5	13.5
Usually (more than a half of the times)	27	25.5	15	23.1	7	16.3	11	20.0	3	8.1
Always (every time)	26	24.5	32	49.2	11	25.6	6	10.9	-	-
Ever interrupted vaginal intercourse/penetration in because of pelvic pain, the last 12 months										
No	36	37.1	20	32.8	14	45.2	18	45.0	16	94.1
Yes	61	62.9	41	67.2	17	54.8	22	55.0	1	5.9
Ever avoided vaginal intercourse/penetration in because of pelvic pain, the last 12 months										
No	39	39.8	17	27.4	16	48.5	24	48.5	16	94.1
Yes	59	60.2	45	72.6	17	51.5	18	51.5	1	5.9

Currently living with a partner										
No, not in a relationship	16	11.9	9	16.4	4	13.8	6	12.2	4	20
No, but in a relationship	15	11.1	6	10.9	2	6.9	5	10.2	2	10

Table V. History of sexual life and details on dyspareunia experiences. Data shown as numbers (N), and percentages (%) by study group.

	EAP		EABP		BPS		PP		Controls	
	N	%	N	%	N	%	N	%	N	%
Depression requiring medication or therapy	67	28.27	47	39.17*	26	36.11	27	21.26	41	17.83
Anxiety requiring medication or therapy	57	28.5	39	34.21*	22	30.56	22	19.82	39	17.73
Migraine	68	28.69*	41	34.17*	29	40.28*	21	16.54*	22	9.57
Asthma	64	27	29	24.17	15	20.83	31	24.41	35	15.22
Irritable Bowel Syndrome (IBS)	44	18.57*	29	24.17*	15	20.83*	13	10.24*	11	4.82
Eczema	33	13.92	20	16.67	11	15.28	17	13.39	29	12.61
Glandular Fever	16	11.76	6	10.71	4	13.33	4	8	2	10
Polycystic Ovary Syndrome	22	9.28	11	9.17	8	11.11	15	11.81	11	4.78
Scoliosis (curvature of the spine)	17	7.17	9	7.5	4	5.56	6	4.72	17	7.46
High blood pressure	4	4.04	4	8	3	10	2	5.88	1	8.33
Thyroid disease	7	3	5	4.27	4	5.56	5	3.97	14	6.09
Spine problems (excluding scoliosis)	7	2.95	6	5	8	11.11	2	1.57	4	1.75
Fibroid uterus	12	5.06	6	5	0	0	3	2.36	3	1.3
Pelvic inflammatory disease	3	1.29	11	9.4	5	6.94	4	3.17	0	0
Fibromyalgia	3	1.27	5	4.17	2	2.78	2	1.57	0	0
Rheumatoid Arthritis	3	1.27	4	3.33	0	0	1	0.79	2	0.88
Chronic Fatigue Syndrome (CFS) / Myalgic encephalomyelitis (ME)	4	1.69	3	2.5	2	2.78	1	0.79	0	0
Cardiovascular disease	2	1	1	0.88	0	0	2	1.8	1	0.45
Graves' disease	1	0.74	0	0	1	3.33	0	0	0	0
Mitral valve prolapse	1	0.42	2	1.67	1	1.39	0	0	1	0.44
Ulcerative Colitis	3	1.27	0	0	0	0	0	0	1	0.43
Diabetes requiring insulin or tablets	2	0.86	0	0	1	1.39	0	0	0	0
Hashimoto's disease	0	0	0	0	1	3.33	0	0	0	0
Diabetes requiring diet control	1	0.43	0	0	1	1.39	0	0	0	0
Sjogren's syndrome	1	0.42	0	0	1	1.39	0	0	0	0
SLE (Lupus)	0	0	2	1.67	0	0	0	0	0	0
Multiple Sclerosis	0	0	0	0	0	0	0	0	1	0.44

Crohn's disease	0	0	0	0	1	1.39	0	0	0	0
IBS based of bowel symptoms	16	11.3	23	25.0	5	13.5	11	20.8	1	8.33

Table VI. Comorbidities occurrence in our population. Data shown as numbers (N) and percentages (%). * represents significant difference between pain groups and the control group (p<0.001) using a Mann-Whitney U test.

		EAP			EABP			BPS			PP			CNP	
	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD
SF-36															
Physical functioning	232	83.79	22.09	119	72.13	27.01	72	77.39	25.94	126	91.28	15.61	227	96.96	8.34
Limitations physical health	231	51.05	45.71	119	30.67	40.45	72	50.35	47.19	126	76.65	38.36	226	95.91	16.24
Limitations emotional problems	230	57.83	43.88	120	32.22	41.43	71	47.42	45.66	126	74.87	38.11	227	80.03	34.24
Energy fatigue	231	51.02	21.99	120	35.33	20.46	72	49.84	21.58	125	61.92	19.29	226	64.82	17.42
Emotional well being	230	64.99	19.40	120	50.92	21.97	72	59.68	23.61	125	72.56	16.74	226	71.28	17.20
Social functioning	232	54.60	28.45	120	34.39	22.19	72	39.79	20.88	126	58.90	22.69	227	50.30	16.07
Pain	231	53.73	25.84	120	40.29	24.26	72	56.74	30.46	125	73.10	25.01	227	90.50	12.08
General Health	232	56.91	22.06	120	42.51	21.86	72	52.05	23.65	126	68.89	21.10	227	74.20	18.01
PCS															
Total Score	193	21.36	14.82	111	29.32	15.08	72	20.58	15.07	110	14.07	12.31	221	9.97	8.38
Rumination	193	7.46	5.52	111	9.89	5.40	72	7.31	5.44	110	5.69	4.83	221	4.21	3.70
Magnification	193	4.02	3.40	111	5.58	3.46	72	4.04	3.40	110	2.64	2.73	221	2.27	2.16
Helplessness	193	9.89	7.20	111	13.85	7.41	72	9.24	7.36	110	5.75	5.75	221	3.50	3.75

Table VII. Quality of life SF-36 and Pain Catastrophising Questionnaire (PCS) scores by study group. Data shown as numbers (N), mean and standard deviation (SD).

	EAP		EABP		BPS		PP	
	N	%	N	%	N	%	N	%
Pain interference with:								
Work or school								
Low	19	16.1	12	13.8	7	17.5	13	31.7
Medium	16	13.6	20	23	8	20	4	9.8
High	52	44.1	46	52.9	13	32.5	8	19.5
Daily activities at home								
Low	26	22.2	12	13.6	9	22.5	10	24.4
Medium	23	19.7	22	25	9	22.5	8	19.5
High	47	40.2	50	56.8	14	35	8	19.5
Sleep								
Low	29	25.4	15	16.7	9	22.5	11	27.5
Medium	24	21.1	17	18.9	6	15	7	17.5
High	36	31.6	55	61.1	18	45	7	17.5
Sexual Intercourse								
Low	15	12.9	4	4.7	5	12.8	6	14.6
Medium	12	10.3	5	5.8	5	12.8	6	14.6
High	38	32.8	52	60.5	18	46.2	7	17.1
Exercise/sports								
Low	23	19.5	9	10.1	10	25.6	16	39
Medium	25	21.2	17	19.1	2	5.1	5	12.2
High	52	44.1	50	56.2	19	48.7	7	17.1
Social activities								
Low	31	26.7	15	17.4	10	25.6	14	34.1
Medium	20	17.2	16	18.6	9	23.1	6	14.6
High	42	36.2	49	57	13	33.3	7	17.1

Table VIII. Information for pain interference on work or school, daily activities at home, sleep, sexual life, exercise or sports and social activities. Data shown as numbers (N), mean and standard deviation (SD).

	EAP		EABP		BPS		PP	
	N	%	N	%	N	%	N	%
Worsening factors of pelvic pain								
Sitting	43	21.5	45	39.5	14	19.4	9	8.1
Full Bladder/Urinating	57	24.1	79	65.8	32	44.4	13	10.2
Bowel movement	7	6.7	18	27.7	4	9.5	2	2.5
Constipation	54	22.8	51	42.5	13	18.1	14	11
Urination	7	5.1	18	25.7	4	9.5	2	2.2
Intercourse/Orgasm	41	17.3	43	35.8	16	22.2	9	7.1
Standing/Walking	49	20.7	25	20.8	4	5.6	18	14.2
Exercise	65	27.4	51	42.5	14	19.4	24	18.9
Stress	71	30	60	50	22	30.6	25	19.7
Time of day	23	9.7	22	18.3	7	9.7	8	6.3
Full meal	36	15.2	34	28.3	11	15.3	12	9.4
Weather	2	1	11	9.9	2	2.8	2	1.8
Contact with clothing	18	9.2	12	10.8	8	11.1	4	3.6
Coughing/Sneezing	30	14.6	38	33	11	15.3	10	8.7
Nothing makes pain worse	10	4.9	1	0.9	1	1.4	9	8
Relieving factors of pelvic pain								
Pain medication	116	48.9	71	59.2	23	31.9	29	22.8
Relaxation	59	24.9	38	31.7	15	20.8	18	14.2
Lying down	112	47.3	66	55	27	37.5	31	24.4
Massage	27	11.4	19	15.8	5	6.9	9	7.1
Heating pad	105	44.3	71	59.2	20	27.8	28	22
Bowel Movement	47	19.8	39	32.5	15	20.8	20	15.7
Hot bath	64	27	56	46.7	14	19.4	22	17.3
Laxatives	6	2.5	8	6.7	3	4.2	5	3.9
TENS	8	3.4	6	5				

Emptying Bladder	22	9.3	41	34.2	17	23.6	6	4.7
Nothing helps	11	4.6	8	6.7	4	5.6	8	6.3

Table IX. Worsening and relieving factors for pelvic pain by study group. Data shown as numbers (N), mean and standard deviation (SD).

NRS for worst period pain during last period

NRS for worst period pain during last period

Mean score for SF-36 Physical Functioning

Mean score for SF-36 Emotional Wellbeing

No pain

Worst imaginable pain

No pain

Worst imaginable pain

Interquartile Range

Data point

Median

NRS for worst period pain during last period

NRS for worst pelvic pain pain during the last 3 months

Interquartile Range

Data point

Median

NRS for worst pain during last intercourse

NRS for worst pain during the 24hrs post last intercourse