Clinical profiling of subgroups of women with chronic pelvic pain

Authors:

Lysia Demetriou[1], Michal Krassowski[1], Pedro Abreu Mendes[2], Kurtis Garbutt[1], Allison F. Vitonis[3ab], Elizabeth Wilkins[1], Lydia Coxon[1], Lars Arendt-Nielsen[4,5], Qasim Aziz[6], Judy Birch[7], Andrew W Horne[8], Anja Hoffman[9], Lone Hummelshoj[10], Claire E Lunde[1,13ab], Jane Meijlink[11], Danielle Perro[1], Nilufer Rahmioglu[1], Kathryn L. Terry[3ab], Esther Pogatzki-Zahn[12], Christine B Sieberg[13abc], Rolf-Detlef Treede[14], Christian M Becker[1], Franscisco Cruz[2], Stacey A Missmer[15abcd], Krina T Zondervan[1], Jens Nagel[16], Katy Vincent[1]

Authors affiliations:

1. Oxford Endometriosis Centre, Nuffield Department of Women's and Reproductive Health, University of Oxford, UK

2. IBMC/I3S and Faculty of Medicine of Porto/ Hospital S João, Porto, Portu

3. a. Boston Center for Endometriosis, Brigham and Women's Hospital and Boston Children's Hospital, Boston, MA, USA and b. Department of Obstetrics and Gynecology, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, 02115, USA

4. Center for Neuroplasticity and Pain (CNAP), SMI, Department of Health Science and Technology, School of Medicine, Aalborg University, Aalborg, Denmark

5. Department of Medical Gastroenterology, Mech-Sense, Aalborg University Hospital, Aalborg, Denmark

6. Centre for Neuroscience, Surgery and Trauma, Blizard Institute, Wingate Institute of Neurogastroenterology. Barts and The London School of Medicine and Dentistry, Queen Mary University of London UK

7. Pelvic Pain Support Network, Poole, UK

8. MRC Centre for Reproductive Health, University of Edinburgh, Edinburgh, UK

9. Research & Development, Pharmaceuticals Experimental Medicine, Bayer AG, Berlin, Germany

10. Endometriosis.org, UK

11. International Painful Bladder Foundation

12. Department of Anesthesiology, Intensive Care and Pain Medicine, University Hospital Muenster, Muenster, Germany

13. a. Biobehavioral Pediatric Pain Lab, Department of Psychiatry & Behavioral Sciences, Boston Children's Hospital, Boston, MA; b. Pain and Affective Neuroscience Center, Department of Anesthesiology, Critical Care, & Pain Medicine, Boston Children's Hospital, Boston, MA; c. Department of Psychiatry, Harvard Medical School, Boston, MA

14. Department of Neurophysiology, Mannheim Center for Translational Neuroscience (MCTN), Heidelberg University, Mannheim, Germany

15. a. Department of Obstetrics, Gynecology, and Reproductive Biology; College of Human Medicine, Michigan State University, Grand Rapids, MI, USA. b. Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA, USA. c. Division of Adolescent and Young Adult Medicine, Department of Pediatrics, Boston Children's Hospital and Harvard Medical School, Boston, MA, USA. d. Boston Center for Endometriosis, Brigham and Women's Hospital and Boston Children's Hospital, Boston, MA, USA

16. Pharmaceuticals Division, Research and Early Development, Therapeutic Area Endocrinology, Metabolism and Reproductive Health, Exploratory Pathobiology, Bayer AG, Wuppertal, Germany.

Word Count: 4,167

Abstract

Study question

Do subgroups of women with chronic pelvic pain (CPP) report different clinical symptoms and differing impact of pain on their quality of life?

Summary answer

Clinical profiles of women with CPP show variability of clinical symptoms both within and between subgroups. However, there is an obvious negative impact of pain on the patients' lives across all subgroups with the comorbid endometriosis and bladder pain symptoms group (EABP) presenting with the higher pain intensities and the lower quality of life.

What is known already

CPP is a common condition affecting up to 26.6% of women, with many suffering for several years before diagnosis and/or treatment. The clinical presentation of CPP is varied and there are frequently comorbid conditions both within and outside the pelvis. Evidence from the literature show that there is an overlap of symptoms in chronic pain conditions whatever the underlying cause which suggests that chronic pain could be a condition itself.

Study design, size, duration

The study is part of The Translational Research in Pelvic Pain (TRiPP) project (https://www.imi-paincare.eu/PROJECT/TRIPP/) which is a cross-sectional observational cohort study. The present study includes 769 female participants sampled from two existing endometriosis-focused cohort studies in Oxford, UK and Boston, MA, USA and newly recruited from the Instituto de Biologia Molecular e Celular (IBMC)) in Porto. The participants

completed an extensive set of questions derived from standardised WERF EPHect questionnaires. Within this study population we defined a control group (reporting no pelvic pain, no bladder pain syndrome (BPS), and no endometriosis diagnosis, N=230) and four pain groups: endometriosis-associated pain (EAP, N=237), (BPS, N=72), comorbid endometriosis-associated pain and BPS (EABP, N=120), and pelvic pain only (PP, N=127).

Participants/materials, setting, methods

All participants were women of reproductive age (13-50 years) and were recruited at three different sites: Oxford (University of Oxford), Boston (Boston Center for Endometriosis (BCE)) and Porto (Instituto de Biologia Molecular e Celular (IBMC)).

The questionnaire included: demographics; reproductive history; pelvic pain intensity assessed using 10-point numerical rating scales (NRS) for dysmenorrhoea, non-cyclical pain, dyspareunia and bladder pain; medical comorbidities; factors relieving and worsening pain; quality of life assessed using the SF-36 questionnaire; and pain catastrophising.

Main results and the role of chance

The EAP (Mean:7.37) and EABP (Mean:7.88) groups scored higher on the pain intensity scales for non-cyclical pelvic pain than the PP (Mean:6.82) group ($p<0.001$) and higher on the dysmenorrhoea scale than both the BPS and PP groups ($p<0.001$). The EABP (Mean:6.61) and BPS (Mean:6.52) groups had significantly higher bladder pain scores than the EAP (Mean:0.95) and PP (Mean:0.78) ($p<0.001$). The EABP group also had significantly higher pain scores for dyspareunia ($p<0.001$), even though more than 50% of participants (who were sexually active) in each of the pain groups reported interrupting and/or avoiding sexual intercourse due to pain in the last 12 months.

Exploring the factors reported to worsen or relieve pain found that across the pain groups the three most reported factors for worsening pelvic pain were: stress (23.6%), full bladder/urinating (23.3%) and exercising (20.2%). The most common factors for relieving pelvic pain were: pain medication (31.4%), lying down (31.0%), and use of a heat pad (29.5%).

Analysis of the quality-of-life questionnaire (SF-36) subscales revealed significant differences between the study groups across all SF-36 subscales (p<0.001). In line with the pain results the EABP group reported the negative highest impact across all the health measures while the PP group's profile was closest to the control group's profile. Significant effects were also observed between the pain groups for pain interference with their work (F(3,209)=9.76, p<0.001) and daily lives (F(3,244)=10.51, p<0.001), with the EABP suffering more compared to the EAP and PP groups (p<0.001).

Limitations, reasons for caution

Data for this study were derived predominantly from existing cohorts where data have been collected over time and thus different versions of questionnaires have been used. Thus, for some questions only a subgroup may have had an opportunity to complete the measure of interest. Recruitment of participants was impacted due to the COVID-19 pandemic. As a result, sample sizes overall were smaller than originally designed, and our BPS group was predominantly identified from gynaecological rather than urological clinics making it potentially different from other published BPS cohorts.

Wider implications of the findings

Overall, our results demonstrate the negative impact that chronic pain has on CPP patients' quality of life and suggests that further exploration of interventions targeting quality of more

broadly is important. Furthermore, it demonstrates the importance of dyspareunia in women with CPP, highlighting the need for more research in this area.

Importantly, we show significant differences between the sub-groups of CPP suggesting the need for better patient stratification in future clinical studies and trials. However, the marked variability both within and between CPP sub-groups raises the question whether subgrouping on the basis of clinical diagnosis is the most appropriate strategy or whether alternative approaches could be identified allowing prioritisation of treatments better suited to the individual patient.

Study funding/competing interest(s)

This project has received funding from the Innovative Medicines Initiative 2 Joint Undertaking under grant agreement No 777500. This Joint Undertaking receives support from the European Union's Horizon 2020 research and innovation programme and EFPIA Companies. Financial support was provided by the J. Willard and Alice S. Marriott Foundation for establishment of and baseline data collection within the A2A cohort - from which the Boston-based TRiPP population was sampled.

Trial registration number

NCT04001244

Introduction

Chronic pelvic pain (CPP) is common, affecting up to 26.6% of women (1–4), yet it remains difficult to treat. For some women an associated pathology such as endometriosis can be identified, but for many others diagnostic investigations will be normal and the label chronic pelvic pain syndrome (CPPS) applied. However, even in those women where endometriosis is identified the lack of correlation between measures of disease severity and pain symptoms (5,6) makes it challenging to understand the extent to which symptoms are actually caused by endometriosis and to predict the benefit of treatments targeting the ectopic tissue.

There is increasing evidence that chronic pain conditions share many features whatever the underlying cause (7,8), leading some to suggest that chronic pain should be treated as a condition in itself (9,10). Whilst there is relatively limited research on this in the context of chronic pelvic pain, studies that have looked at these features confirm similarities with other chronic pain conditions such as irritable bowel syndrome (IBS), rheumatoid arthritis and fibromyalgia (11,12). However, current clinical practice guidelines are still relatively focussed on the identification and treatment of underlying pelvic pathology (13).

The Translational Research in Pelvic Pain (TRiPP) project (https://www.imi-paincare.eu/PROJECT/TRIPP/) aims to better understand the mechanisms generating and maintaining pelvic pain with a particular focus on endometriosis and interstitial cystitis/bladder pain syndrome (IC/BPS) (14). Here, we use the baseline data from the TRiPP project to determine whether different subgroups of women with CPP report different clinical features symptoms and differing impacts of pain on their lives. A better understanding of CPP would allow the development of refined clinical pathways and the prioritisation of specific subgroups of women for further research. Given the diversity of symptoms described by women with all

forms of CPP we do not expect to see differences in specific clinical symptoms between those with and without endometriosis or with and without IC/BPS (with the exception of bladder symptoms themselves). However, we hypothesise that those women with comorbid pain and bladder symptoms will describe a poorer quality of life.

Methods

Ethical Approval

The study has received ethical approval from Yorkshire & The Humber – South Yorkshire Research Ethics Committee (19/YH/0030) with local site approvals and was being conducted in accordance with the principles of the Declaration of Helsinki and with relevant regulations and Good Clinical Practice. Informed consent was obtained from all the participants, and they were informed that they are free to withdraw their data from the study at any time.

Study population

As described in the TRiPP protocol paper (14), participants were identified from two existing endometriosis cohort studies in Oxford (EndOX: A study to identify possible biomarkers in women with endometriosis, Oxford REC ref 09/H0604/58) (N=276) and Boston (The Women's Health Study from Adolescence to Adulthood (A2A), IRB-P00004267) (N=494) plus 16 BPS participants who were recruited at Hospital São João/Instituto de Biologia Molecular e Celular (IBMC) in Porto (Supplementary I).

The study comprised of five study groups and participants were assigned according to specific inclusion and exclusion criteria for each study group (14) (Table I): endometriosis-associated pain (EAP: N=237), comorbid endometriosis associated- and bladder pain syndrome (EABP: N= 120), bladder pain syndrome (BPS: N=72), pelvic pain only (PP: N=127) and controls

(CON: N=230)). All participants were women of reproductive age (13-50 years). Pregnant, lactating or women planning pregnancy during the course of the study were not included as per the study exclusion criteria.

Study design

The TRiPP study design is described in Supplementary I (and more fully elsewhere (14)), the present report focuses on Phase I in which participants completed expanded World Endometriosis Research Foundation (WERF) Biobanking Harmonisation Project (EPHect) questionnaires (15). Here we explore a selected set of outcomes as described below that will also be used in the analysis of phases II and III of the TRiPP project.

Study Data

Demographics and Reproductive History

Demographic information collected included participants' age, body mass index (BMI), race, education, work and relationship status. Information about the participants' gynaecological history consisted of: occurrence and frequency of pelvic pain, menstrual history (e.g. menarche and dysmenorrhoea onset), use of hormones in the last 3 months and information on fertility and dyspareunia. Data from a number of questions were analysed to give a fuller picture of sexual activity, associated pain and functional impact. The questions on dyspareunia were not included in the first version of the questionnaire that was completed by 67 participants in Oxford and for participants younger than 18 years old (N=79) from the Boston cohort.

Pelvic pain

Pain was assessed using numerical rating scales (NRSs) for intensity ranging from 0 = "no pain" to 10 = "worst imaginable pain". Participants were asked to complete NRS for the

experience of pelvic pain at its "worst" in the last three months (for non-cyclical pelvic pain), during the last period (for dysmenorrhoea), worst bladder pain in the last 7 days and pain during and 24hours post sexual intercourse (for dyspareunia).

The pain catastrophising scale (PCS) (16) was used to assess pain-related worry. Catastrophizing is the single most important risk factor that impairs the effectiveness of pain-relieving interventions (17,18), and it statistically mediates the prospective influence of factors such as anxiety on pain outcomes (19). The PCS is a validated/standardised 13-item questionnaire that uses a 5-point scale ranging from 0 (not at all) to 4 (all the time), the total PCS score was computed by summing the responses to all 13 items (16). The scores for the three subscales: Rumination (e.g.,"I can't stop thinking about how much it hurts."),, Magnification (e.g., "I worry that something serious may happen.") and Helplessness (e.g., "There is nothing I can do to reduce the intensity of my pain.") were computed by summing responses of specific items as outlined in Sullivan (16). Scores above the "clinical cut-off points" were considered as clinically relevant levels of catastrophising (PCS Total: 30, Rumination: 11, Magnification: 5, Helplessness: 13).

Medical comorbidities

Participants were provided with a list of medical conditions and reported whether they had received a medical diagnosis for any of those at any point in their lives. The list included autoimmune, gynaecological, mental health, chronic pain, endocrine and cardiovascular disorders.

Furthermore, given the prevalence of bowel symptoms in association with both endometriosis and IC/BPS, irritable bowel syndrome (IBS) was assessed based on the Rome III criteria using

a question about the participants' bowel movements/stools when they experienced non-cyclical pelvic pain in the last 3 months. The symptoms consisted of: a) pelvic pain getting worse/better after bowel movement, b) more/less frequent bowel movement when pain started and c) looser/harder stools when the pain started. Participants who reported having two or more bowel symptoms for "most of the time" or "always" met the criteria for IBS.

Quality of life

The participants' health status was assessed using the Short Form Health Survey (SF-36) which consists of 36 questions covering eight health domains: physical functioning, physical role limitations, bodily pain, general health, emotional role limitations and mental health. The questionnaire was scored as per instructions by Rand Corporation (https://www.rand.org/health-care/surveys_tools/mos/36-item-short-form/scoring.html) and Burholt et al. (20). A higher score indicates a better health status.

To assess pain interference with the participants' lives we used a question assessing the impact of pain on six different aspects (work or school, daily activities at home, sleep, sexual intercourse, exercise/sports and social activities). Participants had to rate the extent of pain interference using a 4-point Likert scale ranging from "Not at all" to "Extremely" for the last 3 months. The scores were recoded into four groups: low, medium, high and no pain interference or not applicable. Additionally, two NRSs were used to assess the severity of the impact of pelvic pain on the participants' work and personal daily activities productivity during the past 4 weeks.

Factors worsening and relieving pain

Two multiple-choice questions were used to assess/identify factors that participants believe worsen or relieve their pelvic pain. Among others, the options included visceral functions such as bowel movement or bladder emptying; behavioural including exercising, standing/walking or sitting; wellbeing factors like stress and mediation; or other factors e.g. time of day and having a full meal.

Statistical analysis

The data were entered into a REDCap database (21) and analysed using IBM SPSS Statistics software, Version 27. The analysis was conducted in 2 phases. First, descriptive analysis was run to identify the characteristics of each study group. Then, statistical testing was undertaken to investigate any effects between the study groups for the study outcomes. Statistical testing for the NRS pelvic and bladder pain did not include the control group since they were recruited based on their NRS scores being <3/10. One-way ANOVAs and Bonferroni corrected post-hoc tests were used to compare between study groups for all the continuous variables while frequencies, percentages and non-parametric tests (chi-square, mann-whitney u tests) were employed for all the categorical variables. Pearson's correlations were run to explore relationships between the variables. Bonferroni correction was used to account for multiple comparisons.

Results

Demographics and Reproductive history

A total of 785 participants were included in this phase of the study with a mean age of 27.6 years (STD: 8.1) and mean BMI of 24.8 (STD: 5.2) (Table II). A one-way ANOVA showed significant differences for age and BMI between groups (Age: $F_{(4,781)}=4.12$, $p=0.003$, BMI:

F(4, 757)=3.28, p=0.011); and post-hoc independent-samples t-tests showed that the BPS group was significantly older than the EABP (p=0.01) and Controls (p=0.04) and had a significantly higher mean BMI than the Controls (p=0.008) (Table II).

The mean age at menarche (mean=12.2 years) ranged from 8 to 16 years old (One-way ANOVA: F(4,768)=0.715, p=0.582) and did not significantly differ among groups (Table II). Likewise, the age at which dysmenorrhoea commenced was similar among all study groups (mean=14.3 years, F(4,631)=1.10, p=0.356) (Table II). On average, patients suffered for approximately 8 years with non-cyclical pelvic pain, and there were no significant differences between the patient groups, even though the PP group had the lowest mean number of years (mean=4.6 years), while the EABP and BPS groups had the highest mean number of years (EABP: mean=9.7 years, BPS: mean=9.9 years) (F(4,204)=3.906, p=0.004). A Pearson's correlation showed that the number of years suffering with pelvic pain positively correlated with the age of the participants (r=0.411, p<0.001).

More participants in the PP (90.5%) and Controls (92.6%) group reported having periods during the last three months than participants in the EAP (76.9%), EABP (63.9%) and BPS (71.8%) (Table III). Of the participants who reported having periods, more than 53.8% of the controls and 35.0% of the pain groups reported having periods whilst using exogenous hormones in the last 3 months. The most common reason for not having periods across the groups was taking hormones continuously (91.8%). Regardless of whether they had periods, many participants reported using exogenous hormones in the last three months (EAP: 52.7%, EABP: 61.7%, BPS: 61.1%, PP: 39.4%, Controls: 57%) (Table III).

Pelvic pain

The mean pain rating for non-cyclical pain at its worst during the last three months was significantly higher in the EAP and EABP compared to the PP group (Table IV). When asked about dysmenorrhoea at its worst the EABP and EAP groups had a significantly higher mean score on pain severity compared to both the BPS and PP groups (p<0.001). Results for the NRS scale on bladder pain during the last seven days revealed significantly higher mean pain scores for the BPS and EABP groups compared to the EAP and PP groups (p<0.001) (Figure I).

Regarding questions on dyspareunia, there were missing responses for more than 50% of our participants, because: a) 4.8% of all the participants did not wish to respond to the questions, b) 7.5% of all the participants reported never having sexual intercourse or c) these questions were not available to 10.1% of the participants as described in the methods section above.

From the available data sexual intercourse data however, most participants in each of the pain groups (>58%) reported that they have experienced pain during intercourse or in the 24 hours following vaginal sexual intercourse (Table V). Of those, more than 60% experienced dyspareunia during the last month across all the pain groups. Almost half of those participants (49.2%) in the EABP group said that they had always experienced dyspareunia during intercourse in the last 12 months as opposed to the BPS and PP groups in which most participants reported occasional dyspareunia in the last year. In the EAP group >77% of participants reported experiencing dyspareunia often, usually or always (Table V).

Concerning the location of dyspareunia, most participants in each pain group reported feeling pain deep inside the vagina, the second most common location was in the abdomen/pelvis except for the PP group in which 50.9% reported feeling pain at the entrance of the vagina.

When asked whether they have been avoiding or interrupting sexual intercourse during the last 12 months due to pelvic pain, more than half of the participants ($>54\%$) in all four pain groups said that they have interrupted sexual intercourse while more than half ($>51\%$) of the participants in the EAP, EABP and BPS groups reported avoiding it.

Further analysis showed effects between the groups for the two dyspareunia NRS scales (pain during intercourse: $F(4,240)=12.77$, $p<0.001$, pain in the 24hrs post-intercourse: $F(4,242)=12.33$, $p<0.001$) (Figure I). The EABP group had significantly higher mean ratings on the NRS scales than all other pain groups ($p<0.001$) for the pain felt during intercourse, and significantly higher mean rating for pain severity in the 24hours after the sexual intercourse than the EAP and PP groups (Table IV and Figure I).

Medical comorbidities

Across all pain groups the most common comorbidities were depression, anxiety, migraine, asthma, and irritable bowel syndrome (IBS) (Table VI). Within the EAP, EABP and BPS groups there were higher percentages of depression, anxiety and migraine diagnosis ($>28\%$) than in the PP group ($<22\%$), while the frequency of asthma ranged between 20% - 27% across all pain groups (Table VI). Statistical analysis using Mann-Whitney U tests revealed than when compared to the control group only the EABP group has significantly higher frequency of anxiety and depression diagnosis (anxiety: ($U(N_{EABP}=114$, $N_{CON}=221)=10510$, $z=-3.39$, $p<0.001$), depression: ($U(N_{EABP}=120$, $N_{CON}=230)=10855$, $z=-4.36$, $p<0.001$)). However, statistical analysis for migraine showed that it was significantly more common in EAP (28.7%), EABP (34.2%) and BPS (40.2%) groups than the control group (9.6%) (($U(N_{EAP}=237$, $N_{CON}=230)=22042$, $z=-5.23$, $p<0.001$), ($U(N_{EABP}=120$, $N_{CON}=230)=10405$, $z=-5.68$, $p<0.001$), ($U(N_{BPS}=72$, $N_{CON}=230)=5737$, $z=-6.60$, $p<0.001$)) groups. Similarly, IBS diagnosis was also

significantly more common in the EAP (18.6%), EABP (24.2%) and BPS (20.8%) groups when compared to the control group (4.8%) ((U(N_{EAP}=237, N_{CON}=229)=23402, z=-4.60, p<0.001), (U(N_{EABP}=120, N_{CON}=230)=11079, z=-5.39, p<0.001), (U(N_{BPS}=72, N_{CON}=229)=6922, z=-4.22,   p<0.001)).

IBS diagnosis was also less common in the PP group (10%) compared to the EAP, EABP and BPS groups (>18%). Correlation analysis between the participants' age and the different comorbidities showed only a weak but significant positive correlation between age and occurrence of spinal problems excluding scoliosis (r=0.014, p=0.001).

Further analysis of the bowel symptoms revealed that more than 11% of the participants in the pain groups met the criteria for IBS regardless of whether they reported having a medical diagnosis for it in the comorbidities question. The EABP and PP groups had the higher percentages of participants meeting the IBS criteria (>20%) (Table VI).

Quality of life

Analysis of the SF-36 questionnaire using a one-way ANOVA revealed significant effects between the five study groups across all eight SF-36 domains (p<0.001). Specifically, the EABP group suffered the most while the control group had the best scores across all sub-scales (p<0.001) (Table VII). Significant effects were also observed between the four pain groups for pain interference with activities at work and daily life (work: $F_{(3,209)}$=9.76, p<0.001; daily activities: $F_{(3,244)}$=10.51, p<0.001) (Figure II). Post-hoc tests revealed that the EABP group had significantly higher pain interference scores than the EAP and PP groups at both scales (p<0.001) (Table IV).

Pain Catastrophising

Pain catastrophising was higher in the EABP group both in the mean total score and in the mean scores of all three subscales (p<0.001). The EAP and BPS groups also had a higher pain catastrophising score than both the PP and Controls groups (p<0.001). According to the clinical-cut-offs more than 19% of the participants in the EAP, EABP and BPS had clinical levels of pain catastrophising. The EABP group had the highest percentage of people above the clinical cut-off (36.0%) (Table VII). A Pearson's correlation between all the pain catastrophising and all the NRS pain scales across all CPP patients revealed strong positive correlations (p<0.001).

Factors worsening and relieving pain

On average, participants from all groups chose three to four factors that either relieve or worsen their pelvic pain. Across the four pain groups the three most common factors for worsening pelvic pain were: stress (23.6%), full bladder/urinating (23.3%) and exercising (20.2%). The most frequently reported factors for relieving pelvic pain were: pain medication (31.4%), lying down (31.0%) and heating pad (29.5%) (Table IX). Analysis of bowel movement and bladder emptying as visceral factors, shows that less people in the EAP and the PP groups report these factors as relieving (bowel movement: EAP: 19.8% and PP 15.7%; emptying bladder: EAP: 9.3% and PP 4.7%) or worsening (bowel movement: EAP: 6.7% and PP 2.5%; emptying bladder: EAP: 24.1% and PP 10.2%), while in the EABP and BPS groups more participants report visceral factors as worsening (bowel movement: EABP: 27.7% and BPS 9.5%; emptying bladder: EABP: 65.8% and BPS 44.4%) rather than relieving (bowel movement: EABP: 32.5% and BPS 20.8%; emptying bladder: EAP: 34.2% and BPS 23.6%).

Discussion

This study phenotypically describes the clinical profiles of women with CPP and contrasts them with controls. In line with existing literature, we have demonstrated that regardless of the aetiology, CPP has a negative impact on the lives of people who suffer with it. However, our results show that there are important differences between different clinical subgroups of those with CPP and those with comorbid endometriosis and bladder pain syndrome (EABP) are particularly severely affected. The heterogeneity in factors exacerbating and relieving pelvic pain, highlights the complexity of the condition but suggests that there may be a variety of different underlying pain mechanisms contributing to the similar clinical presentation. Further evaluation of these underlying mechanisms may provide more clinically relevant approaches to patient stratification.

Importantly, we highlight how common painful sexual intercourse is in women with CPP and that half of those who answered these questions reported interrupting or avoiding it due to pain. This is an important issue for women and their partners and it can have a negative effect on more than one aspect of a woman's life, including mental health, body image, relationship with their partner and fertility (22,23). Dyspareunia is a commonly described symptom of endometriosis, but there is less of a focus on this issue in BPS. It is therefore particularly notable that the highest dyspareunia scores were reported by those with bladder symptoms in our cohort. A better understanding of the mechanisms underlying dyspareunia in all women with CPP not just those with deep endometriosis is clearly an important future need and will ultimately improve the clinical care of those suffering with this important symptom (24).

It is perhaps unsurprising that those with comorbid endometriosis and bladder pain symptoms scored highly on NRSs for all types of pelvic pain (non-cyclical pelvic pain, bladder pain,

dysmenorrhoea and dyspareunia). However, our data demonstrate that these women also have poorer quality of life and report greater pain interference. Whilst there are studies exploring the prevalence and comorbidity of endometriosis and subsequent diagnosis of IC/BPS (25–27), few studies focus on those with comorbid symptoms, either in terms of exploring clinical features or considering specific treatment regimes. Moreover, for many clinical studies the presence of a comorbidity is an exclusion criteria and thus very limited data exists, this is particularly true for clinical trials. Our findings suggest this may be a priority group to consider in future work.

Given that viscero-visceral referral is a relatively well understood phenomenon (28) and that visceral pain conditions are often comorbid (29), we had expected to see higher rates of IBS in those with bladder symptoms. This was the case when questioning previous diagnosis of IBS (EABP 24% and BPS 21% vs EAP 19% and PP 10%). However, considering the answers to the questions comprising the Rome IV criteria, a higher proportion of those in the EABP and PP groups would be considered to meet IBS criteria than in the other groups. It is of course difficult to know whether this reflects ongoing effective treatment in at least a proportion of those with a past diagnosis of IBS. Nonetheless, the high rates of IBS across the cohort as a whole highlight the importance of assessing symptoms from all abdominopelvic organs in women with CPP.

Interestingly the impact of visceral function (bladder and bowel emptying specifically) on pelvic pain appears to be very variable. For some participants these visceral functions worsened their pain, whilst for others they could be relieving. In fact, there was wide variability in the factors identified as either worsening or relieving pelvic pain even within diagnostic groups and it would be interesting to consider whether patterns within these data may reflect

underlying pain mechanisms rather than for example the presence or absence of endometriosis or bladder symptoms. Whilst it can feel challenging in a clinical consultation to raise the topics of mental health and the stressors of daily life, particularly when women may have had a long journey to get their pain taken seriously, stress was the most commonly identified worsening factor whereas relaxation was one of the most frequent relieving factors. Moreover, anxiety and depression were commonly reported comorbidities and those in the EAP, EABP and BPS groups all scored significantly lower than controls on the emotional function dimension on the quality of life questionnaire. This evidence suggest that patients could benefit from treatments focusing on managing their stress and developing better coping mechanisms for their pelvic pain.

Whilst not an assessment of psychological wellbeing, catastrophic thinking about pain has been shown to be an important predictor of outcomes both in the transition to chronic pain after an acute injury and in the response to a variety of therapeutic options (30,31). In line with other studies (32,33) we observed that the chronic pain groups (EAP, EABP and BPS) scored higher in pain catastrophising compared to the control group with a particularly high percentage of EABP patients meeting clinical levels of catastrophising. This highlights the need to include behavioural interventions that specifically target pain-related worry in the management of CPP in the same way as they are considered essential to the management of other chronic pain conditions (34,35).

Strengths and weakness

There are a number of strengths to the TRiPP cohort which include the size of the sample and the detailed phenotypic information available for these participants. Moreover, our study focuses on chronic pelvic pain and therefore includes women with a variety of underlying

diagnoses rather than studying only a single condition such as endometriosis. This strategy allows us to compare the similarities between different clinical subgroups but also to understand differences between them. We have a particular interest in IC/BPS and therefore this cohort is relatively unusual in allowing the comparison between endometriosis and IC/BPS as well as those with comorbid symptoms. Unfortunately, our BPS group is smaller than we initially planned. Neither of the original cohorts from which the TRiPP sample was formed had specifically recruited IC/BPS patients (although many did meet the diagnostic criteria) and therefore we had planned to expand these with a large number of new recruits from specialist centres. However, the COVID-19 pandemic halted our ability to recruit to the study. Overall, we still consider our sample size sufficient to draw meaningful conclusions (192 participants with bladder pain, 120 of whom have coexisting endometriosis).

A further strength of our study is that participants were recruited from three different centres rather than a single site. However, the duration of recruitment to the original cohorts has meant that the baseline questionnaires have been reviewed and updated such that not all participants completed the same version. This has reduced the number of variables we could analyse across the full cohort. Nonetheless the EPHect questionnaire is very comprehensive and the data we do have available allows us to phenotypically characterise our participants in a very detailed manner.

Conclusion

Overall, our results demonstrate the negative impact that chronic pain has on the lives of those with CPP and particularly the importance of dyspareunia. Whilst we do see similarities across the clinical subgroups that we explored our data highlight the difference between these groups and importantly illustrate how severely impacted the group with comorbid endometriosis and

IC/BPS are. The heterogeneity seen in many of our measures, especially the factors exacerbating and relieving pain suggests that multiple different mechanisms may underlie the similar clinical presentations and emphasise the importance of better characterising those with CPP to identify clinically meaningful methods of patient stratification. Further studies within the TRiPP project will explore these mechanisms in greater detail towards this aim.

Author's roles

LD and KV drafted the manuscript. All other authors contributed to the original design of the project and reviewed and edited the manuscript.

Funding

The study was funded by Innovative Medicines Initiative 2 Joint Undertaking (JU) under grant agreement No 777500. The JU receives support from the European Union's Horizon 2020 research and innovation programme and EFPIA. Financial support was provided by the J. Willard and Alice S. Marriott Foundation for establishment of and baseline data collection within the A2A cohort - from which the Boston-based TRiPP population was sampled.

Conflict of interest

LD, MK, EW, LC, DP, NR, AVF, LAN, QA, JB, AH, LH, CEL, JM, CS, PAM, KG: declare no competing interests

AWH: : reports grant funding from the MRC, NIHR, CSO, Wellbeing of Women, Roche Diagnostics, Astra Zeneca, Ferring, Charles Wolfson Charitable Trust and Standard Life. His employer has received consultancy fees from Roche Diagnostics, AbbVie, Nordic Pharma and Ferring, outside the submitted work. In addition, AWH has a patent for a serum biomarker for endometriosis pending.

AH: Employee of Bayer AG, Germany

EPZ: received financial support from Grunenthal and Mundipharma for research activities and advisory and lecture fees from Grünenthal, Novartis and Mundipharma. In addition, she receives scientific support from the German Research Foundation (DFG), the Federal Ministry of Education and Research (BMBF), the German Federal Joint Committee (G-BA) and the Innovative Medicines Initiative (IMI) 2 Joint Undertaking under grant agreement No

777500. This Joint Undertaking receives support from the European Union's Horizon 2020 research and innovation programme and EFPIA. All money went to the institution E.M.P.-Z. is working for.

RDT: Ad board for BAYER, IASP task force on chronic pain classification

CMB: Research Grants from Bayer Healthcare, MDNA Life Sciences, Roche Diagnostics, European Commission, NIH. His employer has received consulancy fees from Myovant and ObsEva for work outside of this project

FC: Consultant and/or investigator for Allergan (Abbvie), Astellas, Bayer, Ipsen and Recordati

SAM: has been an advisory board member for AbbVie and Roche and receives research funding from the National Institutes of Health, the US Department of Defense, the J. Willard and Alice S. Marriott Foundation, and AbbVie; none are related to the presented work; the J. Willard and Alice S. Marriott Foundation supported enrollment of and data collection from the A2A cohort in Boston from which TRiPP data were sampled.

KTZ: reports grant funding from EU Horizon 2020, NIH US, Wellbeing of Women, Bayer AG, Roche Diagnostics, Evotec-Lab282, MDNA Life Sciences, outside the submitted work.

JN: Employee and shareholder of Bayer AG, Germany

KV: declares research funding from Bayer Healthcare and UKRI and honoraria for consultancy and talks and associated travel expenses from Bayer Healthcare, Grunenthal GmBH, AbbVie and Eli Lilly.

References

1.	Eskenazi B, Warner ML. Epidemiology of endometriosis. Obstet Gynecol Clin North Am. 1997;24(2):235–58.

2.	Zondervan KT, Yudkin PL, Vessey MP, Jenkinson CP, Dawes MG, Barlow DH, et al. The community prevalence of chronic pelvic pain in women and associated illness behaviour. Br J Gen Pract.

2001;51(468):541–7.

3. Zondervan KT, Becker CM, Koga K, Missmer SA, Taylor RN, Viganò P. Endometriosis. Nat Rev Dis Prim. 2018;4(1).

4. Ahangari A. Prevalence of chronic pelvic pain among women: an updated review. Pain Physician. 2014;17(2):E141–7.

5. Vercellini P, Trespidi L, De Giorgi O, Cortesi I, Parazzini F, Crosignani PG. Endometriosis and pelvic pain: relation to disease stage and localization. Fertil Steril. 1996;65(2):299–304.

6. Porpora MG, Koninckx PR, Piazze J, Natili M, Colagrande S, Cosmi E V. Correlation between endometriosis and pelvic pain. J Am Assoc Gynecol Laparosc. 1999;6(4):429–34.

7. Ayorinde AA, Macfarlane GJ, Saraswat L, Bhattacharya S. Chronic pelvic pain in women: an epidemiological perspective. Women's Heal. 2015;11(6):851–64.

8. Steege JF, Siedhoff MT. Chronic pelvic pain. Obstet Gynecol. 2014;124(3):616–29.

9. Niv D, Devor M. Chronic pain as a disease in its own right. Vol. 4, Pain Practice. Wiley Online Library; 2004. p. 179–81.

10. Taylor AM, Phillips K, Taylor JO, Singh JA, Conaghan PG, Choy EH, et al. Is chronic pain a disease in its own right? Discussions from a pre-OMERACT 2014 workshop on chronic pain. The Journal of Rheumatology; 2015.

11. Brawn J, Morotti M, Zondervan KT, Becker CM, Vincent K. Central changes associated with chronic pelvic pain and endometriosis. Hum Reprod Update. 2014;20(5):737–47.

12. Kaya S, Hermans L, Willems T, Roussel N, Meeus M. Central sensitization in urogynecological chronic pelvic pain: a systematic literature review. Pain Physician. 2013;16(4):291–308.

13. Lamvu G, Carrillo J, Ouyang C, Rapkin A. Chronic pelvic pain in women: a review. Jama. 2021;325(23):2381–91.

14. Demetriou L, Coxon L, Krassowski M, Rahmioglu N, Arendt-Nielsen L, Aziz Q, et al. Deep phenotyping of women with endometriosis-associated pain and bladder pain syndrome: the TRiPP (Translational Research in Pelvic Pain) study protocol. medRxiv. 2022;

15. Vitonis AF, Vincent K, Rahmioglu N, Fassbender A, Buck Louis GM, Hummelshoj L, et al. World Endometriosis Research Foundation Endometriosis Phenome and biobanking harmonization project: II. Clinical and covariate phenotype data collection in endometriosis research. Fertil Steril.

2014;102(5):1223–32.

16. Sullivan MJL, Bishop SR, Pivik J. The pain catastrophizing scale: development and validation. Psychol Assess. 1995;7(4):524.

17. Hill JC, Lewis M, Sim J, Hay EM, Dziedzic K. Predictors of poor outcome in patients with neck pain treated by physical therapy. Clin J Pain. 2007;23(8):683–90.

18. Karels CH, Bierma-Zeinstra SMA, Burdorf A, Verhagen AP, Nauta AP, Koes BW. Social and psychological factors influenced the course of arm, neck and shoulder complaints. J Clin Epidemiol. 2007;60(8):839–48.

19. Pinto PR, McIntyre T, Almeida A, Araújo-Soares V. The mediating role of pain catastrophizing in the relationship between presurgical anxiety and acute postsurgical pain after hysterectomy. Pain. 2012;153(1):218–26.

20. Burholt V, Nash P. Short Form 36 (SF-36) Health Survey Questionnaire: normative data for Wales. J Public Health (Bangkok) [Internet]. 2011 Dec 1;33(4):587–603. Available from: https://doi.org/10.1093/pubmed/fdr006

21. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. J Biomed Inform. 2009;42(2):377–81.

22. Seehusen DA, Baird D, Bode D V. Dyspareunia in women. Am Fam Physician. 2014;90(7):465–70.

23. Ferrero S, Esposito F, Abbamonte LH, Anserini P, Remorgida V, Ragni N. Quality of sex life in women with endometriosis and deep dyspareunia. Fertil Steril. 2005;83(3):573–9.

24. Vercellini P, Meana M, Hummelshoj L, Somigliana E, Viganò P, Fedele L. Priorities for endometriosis research: a proposed focus on deep dyspareunia. Reprod Sci. 2011;18(2):114–8.

25. Wu C, Chung S, Lin H. Endometriosis increased the risk of bladder pain syndrome/interstitial cystitis: A population-based study. Neurourol Urodyn. 2018;37(4):1413–8.

26. Chung MK, Chung RP, Gordon D, Jennings C. The evil twins of chronic pelvic pain syndrome: endometriosis and interstitial cystitis. JSLS J Soc Laparoendosc Surg. 2002;6(4):311.

27. DiVasta AD, Zimmerman LA, Vitonis AF, Fadayomi AB, Missmer SA. Overlap between irritable bowel syndrome diagnosis and endometriosis in adolescents. Clin Gastroenterol Hepatol. 2021;19(3):528–37.

28. Aziz Q, Giamberardino MA, Barke A, Korwisi B, Baranowski AP, Wesselmann U, et al. The IASP

classification of chronic pain for ICD-11: chronic secondary visceral pain. Pain. 2019;160(1):69–76.

29. Affaitati G, Costantini R, Tana C, Cipollone F, Giamberardino MA. Co-occurrence of pain syndromes. J Neural Transm. 2020;127(4):625–46.

30. Martinez-Calderon J, Jensen MP, Morales-Asencio JM, Luque-Suarez A. Pain catastrophizing and function in individuals with chronic musculoskeletal pain. Clin J Pain. 2019;35(3):279–93.

31. Oosterhaven J, Wittink H, Dekker J, Kruitwagen C, Devillé W. Pain catastrophizing predicts dropout of patients from an interdisciplinary chronic pain management programme: a prospective cohort study. J Rehabil Med. 2019;51(10):761–9.

32. Sewell M, Churilov L, Mooney S, Ma T, Maher P, Grover SR. Chronic pelvic pain–pain catastrophizing, pelvic pain and quality of life. Scand J pain. 2018;18(3):441–8.

33. Martin CE, Johnson E, Wechter ME, Leserman J, Zolnoun DA. Catastrophizing: a predictor of persistent pain among women with endometriosis at 1 year. Hum Reprod. 2011;26(11):3078–84.

34. Schütze R, Rees C, Smith A, Slater H, Campbell JM, O'Sullivan P. How can we best reduce pain catastrophizing in adults with chronic noncancer pain? A systematic review and meta-analysis. J Pain. 2018;19(3):233–56.

35. de C Williams AC, Fisher E, Hearn L, Eccleston C. Psychological therapies for the management of chronic pain (excluding headache) in adults. Cochrane database Syst Rev. 2020;(8).

	Endometriosis	Urinary Symptoms	Pain perceived as arising from the bladder	Pelvic pain score on NRS (at least one)
EAP (N=237)	Surgical diagnosis	No	No	> 4/10
EABP (N=120)	Surgical diagnosis	Yes	Yes	> 4/10
BPS (N=72)	No history	Yes	Yes	> 4/10
PP (N=127)	No endometriosis (surgically confirmed)	No	No	> 4/10
Control (N=230)	No history	No	No	< 3/10

NRS for worst period pain during last period

NRS for worst pelvic pain pain during the last 3 months

NRS for worst pain during last intercourse

NRS for worst pain during the 24hrs post last intercourse

NRS for worst period pain during last period

NRS for worst period pain during last period

Mean score for SF-36 Physical Functioning

Mean score for SF-36 Emotional Wellbeing

	EAP			EABP			BPS			PP			CNP		
	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD
Age	237	28.11	9.53	120	26.45	8.91	72	30.43	9.07	127	28.06	6.63	230	26.57	6.05
BMI	225	24.82	4.83	116	26.06	5.84	71	24.13	4.44	118	25.19	5.74	228	24.10	4.93
Menarche Age	231	12.14	1.59	117	12.09	1.41	72	12.14	1.59	124	12.26	1.34	229	12.32	1.36
Age period pain commenced	184	14.36	4.33	108	13.65	3.08	60	14.23	3.50	102	14.37	4.03	182	14.55	3.04

	EAP		EABP		BPS		PP		CNP	
	N	%	N	%	N	%	N	%	N	%
Period in the last 3 months										
No	54	23.1	43	36.1	20	28.2	12	9.5	17	7.4
Yes	180	76.9	76	63.9	51	71.8	114	90.5	213	92.6
Hormone use in the last 3 months	125	52.7	74	61.7	44	61.1	50	39.4	131	57
Reason for no periods in last 3 months										
Taking hormones continusously	52	94.5	43	100	12	60	11	100	16	94.1
Pregnant/breastfeeding	0	0	0	0	1	5	0	0	1	5.9
Unsure	1	1.8	0	0	2	10	0	0	0	0
Other	2	3.6	0	0	5	25	0	0	0	0
Reason for using hormones										
Birth Control	127	53.6	62	51.7	34	47.2	79	62.2	136	59.1
Pelvic or other period pain	153	64.6	94	78.3	21	29.2	23	18.1	27	11.7
Irregular periods	61	25.7	47	39.2	9	12.5	9	16.5	39	17
Heavy periods	105	44.3	76	63.3	19	26.4	28	22	31	13.5
Acne	19	9.7	16	14.4	8	11.1	18	16.4	38	17.1
Polycystic Ovary Syndrome	12	5.2	7	6	3	4.2	6	4.8	7	3
Ovarian Cyst	20	10.5	19	17.3	1	1.4	1	1	4	1.8
Periods in the last 3 months										
Natural	109	60.9	44	57.9	27	51.9	74	65.5	98	46.2
Hormone Induced	70	39.1	32	42.1	25	48.1	39	34.5	114	53.8
Ever tried to get pregnant for more than 6 months										
No	154	71.6	87	79.8	51	94.4	102	83.6	210	93.3
Yes	61	28.4	22	20.2	3	5.6	20	16.4	15	6.7
Ever seek treatment for infertility in any clinic										
No	178	84	95	90.5	56	98.2	106	90.6	205	94.5
Yes	34	16	10	9.5	1	1.8	11	9.4	12	5.5
Pelvic pain										
No	64	32.7	20	18	25	37.3	60	54.5	206	93.2
Yes	132	67.3	91	82	42	62.7	50	45.5	15	6.8
How long did you have pelvic pain in the last 3 months										
Less than one day	21	17.8	26	29.2	19	46.3	8	21.1	1	14.3
One day	4	3.4	4	4.5	4	9.8	3	7.9	2	28.6
Two to three days	26	22	13	14.6	5	12.2	14	36.8	2	28.6
One day a week	11	9.3	7	7.9	1	2.4	1	2.6	1	14.3
More than one day a week	40	33.9	23	25.8	11	26.8	12	31.6	1	14.3
Every day	16	13.6	16	18	1	2.4	0	0	0	0

	EAP			EABP			BPS			PP			CNP		
	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD
Dysmenorrhea															
Worst period pain during last period	204	6.24	3.23	100	5.92	3.66	59	3.93	3.16	115	5.37	2.91	188	2.11	2.19
Pelvic Pain (non-cyclical)															
Pelvic pain worst severity last 3 months	151	7.37	2.07	95	7.88	1.74	42	7.24	2.12	56	6.82	2.50	12	2.42	1.38
Pelvic pain worst severity during last gynaecological examination	87	5.24	2.87	54	7.35	2.20	27	6.70	3.10	32	5.75	2.62	7	0.86	0.91
Bladder Pain															
Bladder pain worst severity last 7 days	95	0.95	1.61	56	6.61	1.56	29	6.52	1.53	327	0.78	1.40	9	0.11	0.33
Dyspareunia															
Worst pain severity during last intercourse	101	5.51	2.46	61	6.77	2.25	29	6.34	2.16	41	4.76	2.51	13	2.23	1.01
Worst pain severity 24hrs post last intercourse	101	3.97	2.66	63	6.05	2.46	29	4.79	2.82	41	4.29	2.70	13	1.15	1.21
Impact on work and personal life															
Work productivity during past 4 weeks	114	3.70	3.70	41	5.88	2.14	20	5.65	3.05	38	3.39	2.99	18	0.22	0.65
Daily activities productivity during past 4 weeks	125	4.06	2.81	55	6.31	2.02	23	5.57	2.78	45	4.04	4.04	19	0.26	0.73

	EAP		EABP		BPS		PP		CNP	
	N	%	N	%	N	%	N	%	N	%
Ever had pain during intercourse or in the 24hrs following vaginal sexual intercourse/penetration?										
No	35	25.2	7	10.0	13	21.0	39	41.9	132	76.7
Yes	104	71.8	63	90.0	49	79.0	54	58.1	40	23.3
When did you last have vaginal intercourse										
In the last month	74	70.5	41	64.1	31	62.0	44	77.2	27	69.2
1-3 months ago	13	12.4	10	15.6	6	12.0	2	3.5	6	15.4
4-12 months ago	7	6.7	10	15.6	8	16.0	4	7.0	2	5.1
More than 12 months ago	11	10.5	3	4.7	5	10.0	7	12.3	4	10.3
Pain location during last vaginal intercourse/penetration										
Entrance of the vagina	15	6.4	21	17.9	12	16.7	6	4.8	5	2.2
Deep inside the vagina	61	26.2	49	41.9	24	33.3	22	17.5	7	3
Entrance and deep inside the vagina	5	13.5	3	50.0	0	0	3	18.8	0	0
In the abdomen/pelvis	55	28.1	52	46.8	19	26.4	24	21.8	3	1.4
Other location	5	2.6	9	8.1	2	2.8	2	1.8	0	0
Frequency of pelvic pain during or 24hrs after vaginal intercourse/penetration in the last 12 months										
Never	1	0.9	-	-	3	7.0	1	1.8	5	13.5
Occasionally (less than a quarter of times)	23	21.7	8	12.3	18	41.9	28	50.9	24	64.9
Often (a quarter to half of the times)	29	27.4	10	15.4	4	9.3	9	16.4	5	13.5
Usually (more than a half of the times)	27	25.5	15	23.1	7	16.3	11	20.0	3	8.1
Always (every time)	26	24.5	32	49.2	11	25.6	6	10.9	-	-
Ever interrupted vaginal intercourse/penetration in because of pelvic pain, the last 12 months										
No	36	37.1	20	32.8	14	45.2	18	45.0	16	94.1
Yes	61	62.9	41	67.2	17	54.8	22	55.0	1	5.9
Ever avoided vaginal intercourse/penetration in because of pelvic pain, the last 12 months										
No	39	39.8	17	27.4	16	48.5	24	48.5	16	94.1
Yes	59	60.2	45	72.6	17	51.5	18	51.5	1	5.9
Currently living with a partner										
No, not in a relationship	16	11.9	9	16.4	4	13.8	6	12.2	4	20
No, but in a relationship	15	11.1	6	10.9	2	6.9	5	10.2	2	10

	EAP		EABP		BPS		PP		Controls	
	N	%	N	%	N	%	N	%	N	%
Depression requiring medication or therapy	67	28.27	47	39.17*	26	36.11	27	21.26	41	17.83
Anxiety requiring medication or therapy	57	28.5	39	34.21*	22	30.56	22	19.82	39	17.73
Migraine	68	28.69*	41	34.17*	29	40.28*	21	16.54*	22	9.57
Asthma	64	27	29	24.17	15	20.83	31	24.41	35	15.22
Irritable Bowel Syndrome (IBS)	44	18.57*	29	24.17*	15	20.83*	13	10.24*	11	4.82
Eczema	33	13.92	20	16.67	11	15.28	17	13.39	29	12.61
Glandular Fever	16	11.76	6	10.71	4	13.33	4	8	2	10
Polycystic Ovary Syndrome	22	9.28	11	9.17	8	11.11	15	11.81	11	4.78
Scoliosis (curvature of the spine)	17	7.17	9	7.5	4	5.56	6	4.72	17	7.46
High blood pressure	4	4.04	4	8	3	10	2	5.88	1	8.33
Thyroid disease	7	3	5	4.27	4	5.56	5	3.97	14	6.09
Spine problems (excluding scoliosis)	7	2.95	6	5	8	11.11	2	1.57	4	1.75
Fibroid uterus	12	5.06	6	5	0	0	3	2.36	3	1.3
Pelvic inflammatory disease	3	1.29	11	9.4	5	6.94	4	3.17	0	0
Fibromyalgia	3	1.27	5	4.17	2	2.78	2	1.57	0	0
Rheumatoid Arthritis	3	1.27	4	3.33	0	0	1	0.79	2	0.88
Chronic Fatigue Syndrome (CFS) / Myalgic encephalomyelitis (ME)	4	1.69	3	2.5	2	2.78	1	0.79	0	0
Cardiovascular disease	2	1	1	0.88	0	0	2	1.8	1	0.45
Graves' disease	1	0.74	0	0	1	3.33	0	0	0	0
Mitral valve prolapse	1	0.42	2	1.67	1	1.39	0	0	1	0.44
Ulcerative Colitis	3	1.27	0	0	0	0	0	0	1	0.43
Diabetes requiring insulin or tablets	2	0.86	0	0	1	1.39	0	0	0	0
Hashimoto's disease	0	0	0	0	1	3.33	0	0	0	0
Diabetes requiring diet control	1	0.43	0	0	1	1.39	0	0	0	0
Sjogren's syndrome	1	0.42	0	0	1	1.39	0	0	0	0
SLE (Lupus)	0	0	2	1.67	0	0	0	0	0	0
Multiple Sclerosis	0	0	0	0	0	0	0	0	1	0.44
Crohn's disease	0	0	0	0	1	1.39	0	0	0	0
IBS based of bowel symptoms	16	11.3	23	25.0	5	13.5	11	20.8	1	8.33

	EAP			EABP			BPS			PP			CNP		
	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD
SF-36															
Physical functioning	232	83.79	22.09	119	72.13	27.01	72	77.39	25.94	126	91.28	15.61	227	96.96	8.34
Limitations physical health	231	51.05	45.71	119	30.67	40.45	72	50.35	47.19	126	76.65	38.36	226	95.91	16.24
Limitations emotional problems	230	57.83	43.88	120	32.22	41.43	71	47.42	45.66	126	74.87	38.11	227	80.03	34.24
Energy fatigue	231	51.02	21.99	120	35.33	20.46	72	49.84	21.58	125	61.92	19.29	226	64.82	17.42
Emotional well being	230	64.99	19.40	120	50.92	21.97	72	59.68	23.61	125	72.56	16.74	226	71.28	17.20
Social functioning	232	54.60	28.45	120	34.39	22.19	72	39.79	20.88	126	58.90	22.69	227	50.30	16.07
Pain	231	53.73	25.84	120	40.29	24.26	72	56.74	30.46	125	73.10	25.01	227	90.50	12.08
General Health	232	56.91	22.06	120	42.51	21.86	72	52.05	23.65	126	68.89	21.10	227	74.20	18.01
PCS															
Total Score	193	21.36	14.82	111	29.32	15.08	72	20.58	15.07	110	14.07	12.31	221	9.97	8.38
Rumination	193	7.46	5.52	111	9.89	5.40	72	7.31	5.44	110	5.69	4.83	221	4.21	3.70
Magnification	193	4.02	3.40	111	5.58	3.46	72	4.04	3.40	110	2.64	2.73	221	2.27	2.16
Helplessness	193	9.89	7.20	111	13.85	7.41	72	9.24	7.36	110	5.75	5.75	221	3.50	3.75

	EAP		EABP		BPS		PP	
	N	%	N	%	N	%	N	%
Pain interference with:								
Work or school								
Low	19	16.1	12	13.8	7	17.5	13	31.7
Medium	16	13.6	20	23	8	20	4	9.8
High	52	44.1	46	52.9	13	32.5	8	19.5
Daily activities at home								
Low	26	22.2	12	13.6	9	22.5	10	24.4
Medium	23	19.7	22	25	9	22.5	8	19.5
High	47	40.2	50	56.8	14	35	8	19.5
Sleep								
Low	29	25.4	15	16.7	9	22.5	11	27.5
Medium	24	21.1	17	18.9	6	15	7	17.5
High	36	31.6	55	61.1	18	45	7	17.5
Sexual Intercourse								
Low	15	12.9	4	4.7	5	12.8	6	14.6
Medium	12	10.3	5	5.8	5	12.8	6	14.6
High	38	32.8	52	60.5	18	46.2	7	17.1
Exercise/sports								
Low	23	19.5	9	10.1	10	25.6	16	39
Medium	25	21.2	17	19.1	2	5.1	5	12.2
High	52	44.1	50	56.2	19	48.7	7	17.1
Social activities								
Low	31	26.7	15	17.4	10	25.6	14	34.1
Medium	20	17.2	16	18.6	9	23.1	6	14.6
High	42	36.2	49	57	13	33.3	7	17.1

	EAP		EABP		BPS		PP	
	N	%	N	%	N	%	N	%
Worsening factors of pelvic pain								
Sitting	43	21.5	45	39.5	14	19.4	9	8.1
Full Bladder/Urinating	57	24.1	79	65.8	32	44.4	13	10.2
Bowel movement	7	6.7	18	27.7	4	9.5	2	2.5
Constipation	54	22.8	51	42.5	13	18.1	14	11
Urination	7	5.1	18	25.7	4	9.5	2	2.2
Intercourse/Orgasm	41	17.3	43	35.8	16	22.2	9	7.1
Standing/Walking	49	20.7	25	20.8	4	5.6	18	14.2
Exercise	65	27.4	51	42.5	14	19.4	24	18.9
Stress	71	30	60	50	22	30.6	25	19.7
Time of day	23	9.7	22	18.3	7	9.7	8	6.3
Full meal	36	15.2	34	28.3	11	15.3	12	9.4
Weather	2	1	11	9.9	2	2.8	2	1.8
Contact with clothing	18	9.2	12	10.8	8	11.1	4	3.6
Coughing/Sneezing	30	14.6	38	33	11	15.3	10	8.7
Nothing makes pain worse	10	4.9	1	0.9	1	1.4	9	8
Relieving factors of pelvic pain								
Pain medication	116	48.9	71	59.2	23	31.9	29	22.8
Relaxation	59	24.9	38	31.7	15	20.8	18	14.2
Lying down	112	47.3	66	55	27	37.5	31	24.4
Massage	27	11.4	19	15.8	5	6.9	9	7.1
Heating pad	105	44.3	71	59.2	20	27.8	28	22
Bowel Movement	47	19.8	39	32.5	15	20.8	20	15.7
Hot bath	64	27	56	46.7	14	19.4	22	17.3
Laxatives	6	2.5	8	6.7	3	4.2	5	3.9
TENS	8	3.4	6	5				
Emptying Bladder	22	9.3	41	34.2	17	23.6	6	4.7
Nothing helps	11	4.6	8	6.7	4	5.6	8	6.3