Title: Estimating the potential impact and diagnostic requirements for SARS-CoV-2

test-and-treat programs

Authors: Alvin X. Han[1], Emma Hannay[2], Sergio Carmona[2], Bill Rodriguez[2], Brooke E.

Nichols[1,2,3,†], Colin A. Russell[1,3,†]

Affiliations:

[1]Department of Medical Microbiology & Infection Prevention, Amsterdam University

Medical Center, University of Amsterdam, Amsterdam, The Netherlands

[2]Foundation for Innovative New Diagnostics (FIND), Geneva, Switzerland

[3]Department of Global Health, School of Public Health, Boston University, Boston, MA,

USA

†Contributed equally

Correspondence: Alvin X. Han (x.han@amsterdamumc.nl) and Colin A. Russell

(c.a.russell@amsterdamumc.nl)

Abstract (201/200 words):

Oral antivirals have the potential to reduce the public health burden of COVID-19. However, now that we have exited the emergency phase of the COVID-19 pandemic, declining SARS-CoV-2 clinical testing rates (average testing rates = ≪10 tests/100,000 people/day in low-and-middle income countries; <100 tests/100,000 people/day in high-income countries; September 2023) make the development of effective test-and-treat programs challenging. We used an agent-based model to investigate how testing rates and strategies affect the use and effectiveness of oral antiviral test-to-treat programs in four country archetypes of different income levels and demographies. We find that in the post-emergency phase of the pandemic, in countries where low testing rates are driven by limited testing capacity, significant population-level impact of test-and-treat programs can only be achieved by both increasing testing rates and prioritizing individuals with greater risk of severe disease. However, for all countries, significant reductions in severe cases with antivirals are only possible if testing rates were substantially increased with high willingness of people to seek testing. Comparing the potential population-level reductions in severe disease outcomes of test-to-treat programs and vaccination shows that test-and-treat strategies are likely substantially more resource intensive requiring very high levels of testing (>>100 tests/100,000 people/day) and antiviral use suggesting that vaccination should be a higher priority.

Main Text

Introduction

Antiviral therapies such as anti-SARS-CoV-2 monoclonal antibodies, replication inhibitors,

protease inhibitors, and host-directed therapies can be used to treat COVID-19, reducing the

probability of severe disease to varying degrees.[1] Direct-acting antiviral drugs, such as

molnupiravir[2] and nirmatrelvir–ritonavir (Paxlovid),[3] have the potential to substantially lower

disease burden given their efficacy and convenience of oral dosing. Nirmatrelvir–ritonavir, in

particular, can reduce incidence of adverse events in high-risk individuals (i.e. ≥60 years of

age (over-60y) or an adult ≥18 years with a relevant comorbidity) by 46-89%.[3,4] Given their

ability to lower viral load,[3] these drugs could also potentially be used to control SARS-CoV-2

transmission.[5] To achieve maximum impact, these drugs must typically be administered

within a few days of symptom onset. Given limited resources and the relatively high cost of

these drugs,[6] along with the need to administer drugs quickly after symptom onset,[2,3]

diagnostic testing remains an essential first step for identifying suitable drug recipients.

Oral antivirals (the term "*antivirals*" refers only to oral direct antivirals for the rest of this

article) have the potential to reduce the disease burden of COVID-19 outbreaks. Various

studies have estimated ~10% – 40% reduction in severe disease outcomes if antivirals were

distributed to 20% – 50% of all symptomatic infected individuals.[5,7,8] However, none of these

studies have accounted for the diagnostic capacity required to identify and treat these cases

with antivirals. There have been substantial gaps in COVID-19 testing equity across country

income groups throughout the pandemic. Between January 2020 and March 2022, LMICs

were only testing at an average of 27 tests/100,000 people/day (tests/100K/day) as compared

to >800 tests/100K/day in high-income countries (HICs).[9] In the post-public health

emergency phase of the pandemic, testing rates have dwindled down to less than 10

tests/100K/day and 100 tests/100K/day on average for LMICs and HICs respectively (as of

September 2023).[9] Low testing rates severely underestimate COVID-19 cases,[10] which not

only complicate antiviral demand forecasts but also create additional barriers to the effective

distribution and use of antivirals.

Here, we used an agent-based model (PATAT)[11,12] to demonstrate how testing rates and

strategies affect the use and impact of antivirals. In the model, we focused on antigen rapid

diagnostic tests (Ag-RDTs) which can easily be performed at point-of-care or be used as self-

tests with short turnaround time needed to quickly identify high-risk infected individuals.[13]

We computed the potential impact of test-and-treat programs on infections, severe cases, and

deaths averted in three LMIC archetypes with distinct demographic structures – Brazil,

Georgia, and Zambia – and the Netherlands as an HIC example, all under varying levels of

vaccination coverage. The LMIC archetypes were selected as the age demography of their

populations were largely representative of the 132 other LMICs as classified by the World

Bank (Figure 1).[14,15] Our findings highlight the limits and expected outcomes of COVID-19

oral antiviral treatment programs under realistic testing and vaccination landscapes.

Results

Dynamic epidemic simulations with PATAT

We first provide key details of the PATAT model and assumptions to contextualize our

results. See Methods and Supplementary Information for full description of the model and

parameters. We simulated SARS-CoV-2 epidemics in each country under a range of average

effective reproduction number (i.e. R_e = 0.9, 1.2 (doubling time = 6-9 days), 1.5 (doubling

time = 3-5 days), and 2.0 (doubling time = 1-3 days)) during the first week of each

simulation. These doubling times coincide the range reported for prominent Omicron

subvariants as well, including BA.2 (~3 days)[16], BA.5 (5-6 days)[17] and XBB.1.5 (9-10

days)[18] (Figure S1). All simulations were initialized with 1% of the population infected at the

start of the epidemic. We did not model varying levels of population immunity due to the

lack of comprehensive country-specific infection data and complexities in parameterizing the

proportion and protection conferred to individuals with infections by different variant

infection histories in the past. Instead, the different R_e values should be viewed as the

collective outcome of population immunity from previous infections, intrinsic

transmissibility of the variant virus as well as effects of any existing any public health

interventions other than vaccination and oral antivirals. For each R_e value and country, we

performed two sets of simulations – one with and the other without the distribution of

antivirals. For each set of simulations, we assumed three vaccination coverage: 10%, 50%

and 90%. We randomly assigned vaccination status across the simulated population but

assumed that vaccination was age-tiered such that the older individuals were vaccinated first.

For comparability between countries and as a simplification, we assumed that protection rates

against infection and severe disease were 29% and 70% respectively, which were based on

the more conservative, lower average estimates of vaccine effectiveness against BA.1 across

different vaccines (i.e. mRNA and ChAdOx1 nCoV-19 vaccine) and doses (i.e. 1-3 doses).[19–

21]

The relative susceptibility of individuals to infection,[22,23] probability of becoming

symptomatic, [24,25] probability of developing severe disease,[24,25] and the probability of death

[26,27] depend on the age of the individual (Table S1). We assumed that only high-risk

individuals (i.e. ≥60 years of age (over-60y) or an adult ≥18 years with a relevant

comorbidity) who tested positive at clinics (e.g. a self-reported self-test would be insufficient

to access antivirals) would receive a course of antivirals. We also randomly assigned 20% of

the population to have a 40% increase in relative risk to developing severe disease because of

pre-existing comorbidities (e.g. obesity, diabetes, people living with HIV, etc.).[28,29] As a

simplification, we assumed that the prevalence of comorbidities was independent of age.

Although the phase 2/3 trial of nirmatrelvir–ritonavir reported 89% relative risk reduction

among unvaccinated high-risk patients infected by the Delta variant-of-concern,[3] we assumed

that an antiviral course conferred a 46% risk reduction for infected high-risk individuals to

severe disease outcomes based on a separate cohort study on the effectiveness of

nirmatrelvir–ritonavir among high-risk patients infected by Omicron BA.1 independent of

their vaccination status.[4]

Impact of test-and-treat

We simulated the implementation of test-and-treat programs during SARS-CoV-2 epidemic

waves in three different LMICs (Brazil, Georgia, and Zambia) with distinct population

demography (Figure 1) and the Netherlands under different levels of vaccine coverage (10%,

50% or 90%) and average test availability (10, 100 or 500 tests/100K/day). We assumed that

tests were only available at health clinics and that 65% of individuals with mild symptoms

would likely seek testing at clinics based on surveys of testing behaviour during the

pandemic.[30,31] Test-seeking individuals would, however, only be tested if tests were

available. From our simulations, we found that the likelihood of detecting an infection ranged

between 0.06% and 64.6%, depending on the country simulated, epidemic intensity,

vaccination coverage and test availability (Figure S2). Generally, detection is more likely

with a larger proportion of over-60y individuals (i.e. the more likely cases will be

symptomatic and seek testing), lower reproduction rate R_e, higher vaccination coverage and

greater test availability (i.e. any of the aforementioned factors directly or indirectly increases

the surplus of tests available for symptomatic individuals).

At 10 tests/100K/day, test-and-treat programs are unlikely to have any population-level

impact on disease transmission in all countries (Figure S3). At higher testing rates ($\geq$100

tests/100K/day) and lower R_e ($\leq$1.5), there were modest differences between simulated

countries. We found that antivirals largely only have a limited impact on total infections

averted (Figure S3), in large part because 58-67% of all transmission events were attributed

to asymptomatic and pre-symptomatic individuals (Figure S4A). However, in Georgia and

the Netherlands where >30% of the population are over-60y and high-risk individuals

transmitted almost half of all infections (Figure S4B), increasing testing rates to 100 (500)

tests/100K/day, accompanied by uncapped distribution of antivirals, could reduce total

infections by ~12% (~22-24%). On the other hand, regardless of testing rates, infections

averted were <12% and <4% in Brazil and Zambia respectively, both of which have smaller

over-60y populations (i.e. Brazil: 15%; Zambia: 6% of population; Figure S3A) and where

most infections are transmitted by low-risk individuals (Figure S4B). Across all settings and

testing rates, increasing vaccination coverage did not change the proportion of infections

averted by antivirals substantially.

If testing rates increased to 500 tests/100K/day, the proportion of severe cases averted due to

antivirals would depend on the proportion of over-60y in the population, with Zambia, Brazil,

Georgia and the Netherlands, maximally reducing up to an average of 46%, 55%, 67% and

68% of severe cases respectively through test-and-treat strategies (Figure 2). Linking

antiviral treatment to testing programs at a rate of 10 tests/100K/day did not generate any

impact under any scenario, including when 90% of the population were vaccinated. Raising

testing rates to 100 tests/100K/day – a widely publicized global target during the pandemic –

and treating all high-risk, test-positive patients with antivirals substantially increased the

proportion of severe cases averted at lower R_e (i.e. proportion of severe cases averted at $R_e =$

0.9 (1.2) with 10-90% vaccination coverage: Zambia, 17-20% (3-4%); Brazil, 24-55% (6-

14%); Georgia, 50-65% (13-30%) and the Netherlands, 48-67 (12-31%); Figure 2). The

impact was greatest in Georgia and the Netherlands given their substantial >60y population.

As R_e increases (≥1.5), the likely population demand for tests also increased, and

correspondingly >100 tests/100K/day was needed to ensure that high-risk individuals could

be identified to initiate treatment (i.e. proportion of severe cases averted at $R_e = 1.5$ (2.0)

with 10-90% vaccination coverage at 100 tests/100K/day: Zambia, 2-4% (0-3%); Brazil, 1-

4% (0-1%); Georgia, 3-9% (1-2%); Netherlands, 3-10% (0-2%). At 500 tests/100K/day:

Zambia, 9-16% (7-9%); Brazil, 11-36% (6-9%); Georgia, 24-66% (8-14%); Netherlands, 28-

65% (6-18%); Figure 2). Although we did not model the impact of antivirals in reducing the

likelihood of death, developing severe disease precedes dying from COVID-19 in our model

(see Methods), the number of deaths averted thus follow similar trends as severe cases

averted (Figure S5).

At testing rates of ≤10 tests/100,000 people/day, use of antivirals made negligible

contributions to reducing severe disease at all levels of vaccine coverage. At testing rates

≥100 tests/100,000/people/day, higher vaccination coverage was associated with a smaller

absolute number of severe cases averted by antivirals. However, at higher testing rates, the

proportion of severe cases averted by antivirals relative to no distribution of antivirals is

larger at higher vaccination coverage. This is because as infections decrease with higher

vaccination coverage, a greater percentage of severe cases could also be detected and treated

by antivirals assuming that the quantity of test availability is a constraining factor and that

demand in low vaccination scenarios would exceed supply.

Distribution of test and antivirals to high-risk household contacts of test-positive individuals

As antivirals must be administered quickly after symptom onset, one way to promptly

identify and treat infected high-risk individuals is to secondarily distribute self-tests to high-

risk household contacts who were exposed to the test-positive individuals. This would,

however, also result in a faster depletion of available test stocks under limited test

availability. We repeated our simulations with high-risk household contacts receiving Ag-

RDTs from clinics to perform self-test over the ensuing three days, initiating antiviral

treatment upon a positive diagnosis. In this scenario, however, there was little reduction in

total infections due to antivirals (Figure S6). In fact, when R_e was low (≤ 1.2) and at 100

tests/100K/day, distributing tests to high-risk household contacts for self-tests diverted away

test stocks that would otherwise be used to diagnose test-seeking symptomatic individuals

(which would, in turn, change their behavior to reduce transmission if tested positive). At 100

tests/100K/day across all R_e values, or at 500 tests/100K/day and higher R_e, the proportion of

severe cases and in turn, deaths averted diminished substantially by a factor of two- to ten-

fold (Figure S7-8) relative to no secondary distribution of Ag-RDTs to high-risk household

contacts (Figure 2 and Figure S5). Unless testing rates were increased to ≥ 500

tests/100K/day, 100 tests/100K/day remains inadequate to meet the testing demand of both

symptomatic individuals and high-risk household contacts to derive greater impact from test-

and-treat.

Restricting symptomatic testing to high-risk individuals

Given the modest impact of antivirals in reducing transmissions, testing could be targeted to

high-risk individuals only in order to distribute antivirals to as many infected high-risk

individuals as possible. This strategy can be effective in reducing severe cases and deaths by

test-and-treat when Ag-RDT availability is inadequate to test all symptomatic individuals

who seek testing, which was a common scenario in LMICs during the pandemic. Otherwise,

if most individuals only isolate themselves after a positive test, the testing restriction would

lead to excess tests available that are not effectively used to alter the behaviour of low-risk

infected individuals that curb onward transmissions.

In our model, restricting testing to high-risk groups when there are ample amount of tests to

diagnose non-high-risk symptomatic individuals as well resulted in more transmissions

(Figure S9) and severe cases (Figure 3). We estimated that there can be up to 56% more

infections at $R_e \leq 1.5$ if test availability was 500 tests/100K/day but were restricted to high-

risk individuals only. In Georgia, for example, restricting testing to high-risk groups would

reduce 52% of severe cases by antivirals at $R_e = 1.5$, 500 tests/100K/day and 90%

vaccination coverage as opposed to 66% under the same scenario but without testing

restrictions. On the other hand, when operating under limited test availability relative to R_e,

restricting symptomatic testing to high-risk individuals could be an effective strategy to

further reduce severe cases (i.e. Fold increase in proportion of severe cases averted relative to

no symptomatic testing restrictions when $R_e \geq 1.5$, across all vaccination coverages and

countries simulated: 100 tests/100K/day, median 4.9-fold (IQR = 3.3-6.4); 500

tests/100K/day, median 3.2-fold (IQR = 2.4-5.1)) and in turn, deaths as well (Figure S10). Of

the test distribution strategies simulated in this study, restricting testing to high-risk-groups-

only also substantially reduced the number of tests performed per antiviral distributed to

median 6 tests (IQR = 5-8 tests; Figure S11). In contrast, a median 20 tests (IQR = 15-33

tests) would be required per antiviral distributed if symptomatic testing was performed

without restrictions about risk status.

Oral antiviral need

Assuming that only symptomatic high-risk individuals who sought testing received an

antiviral course upon a positive test, and that there were two 90-day epidemic waves in a

year, we estimated that one antiviral course is needed for every 73-251 (14-154) persons per

year on average if testing rate was 100 (500) tests/100K/day across all simulated countries

and vaccination coverage (Figure 4). We assumed that vaccine protection against infection

was low (29%) and that antivirals were distributed regardless of vaccination status. As such,

increasing vaccination coverage did not lower antiviral need substantially (median 0.93-fold

change (IQR = 0.70-1.00) when vaccination coverage increased from 10% to 90%).

Conversely, the amount of antivirals distributed depends on R_e (median 2.60-fold change

(IQR = 0.97-4.35) when R_e increases from 0.9 to 2.0), country demographics (median 1.72-

fold change (IQR=1.02-2.04) when distributing antivirals in Georgia relative to Zambia),

testing rates (median 4.31-fold change (IQR = 1.49-5.77) when increasing from 100 to 500

tests/100K/day), and how tests were targeted (median 2.57-fold change (IQR = 1.52-4.55)

when testing only high-risk as opposed to all symptomatic individuals).

Impact of oral antivirals with over-the-counter self-tests

Unlike LMICs, over-the-counter Ag-RDTs were readily available in high-income countries

during and after the emergency phase of the pandemic. In a separate analysis for the

Netherlands, we assumed that over-the-counter Ag-RDTs for self-testing were widely

available (i.e. with no-cap on availability) such that only 10% of symptomatic individuals

seek clinic-provided testing directly. We also assumed that 80% of symptomatic individuals

who did not seek clinic-provided testing may perform a self-test using over-the-counter Ag-

RDTs instead. This effectively means that up to 82% of *all* symptomatic individuals would

perform either a clinic-provided or over-the-counter self-test. All high-risk individuals who

tested positive using self-tests would then seek reflexive testing at clinics on the same day to

be administered antivirals. Clinic-provided testing would only be performed if they were still

available under the average test availability of either 100 or 500 tests/100K/day.

Under these assumptions, we found that in combination with clinic-provided testing rate of

500 tests/100K/day, distribution of antivirals could avert 56-59% of severe cases and 67-70%

of deaths on average, regardless of the epidemic intensity (Figure 5). Reduction in infections

due to antivirals was similarly modest and did not amount to more than an average of 13%.

However, if mean clinic-provided testing rates fell to 100 tests/100K/day, the mean

proportion of severe cases and deaths averted would also drop precipitously to as low as 14%

and 19% respectively when $R_e \geq 1.5$. Across both testing rates and R_e, we found that one

antiviral course was distributed for every 4-69 individuals for two 90-day epidemic waves in

a year.

Since antivirals must be administered promptly upon a positive diagnosis, we also computed

the proportion of high-risk, symptomatic individuals that would miss the treatment window if

they had sought reflexive testing late. Regardless of clinical testing rate and R_e, for ≥90% of

high-risk symptomatic individuals who were able to avert severe disease outcomes by

antivirals to be treated with the drug, they must not seek reflexive testing at clinics (if

reflexive testing is required) later than two days after being tested positive with over-the-

counter self-tests (Figure S12).

Effectiveness of test-and-treat strategies

To further compare the effectiveness of the test-and-treat strategies we investigated, we

plotted efficiency curves of the number of severe cases averted by antivirals against the

number of antivirals administered across all R_e values and countries (Figure S13). As we

assumed that there was no cap on antiviral availability, the limited test availability thus

determines the number of antivirals distributed and in turn, the maximum number of severe

cases averted by antivirals. We found that testing and treating test-positive, high-risk

household contacts alongside the test-positive index individual ("Symptomatic + HR

household" in Figure S13) was the least efficient test-and-treat strategy in our analyses. This

is because allocating tests to screen high-risk household contacts, who may or may not be

infected, under limited test availability reduced the number of tests that would otherwise have

been used to identify symptomatic infected high-risk individuals for antiviral administration.

Restricting tests to high-risk individuals only ("HR symptomatic only") was similarly

effective to no restriction in access to tests for all symptomatic individuals ("Symptomatic")

as it is an essentially a workaround of the latter strategy to increase the number of high-risk

infected individuals who are tested and treated under limited test-availability. In short, the

greater the access high-risk individuals have to testing, the more likely they could be

identified for timely treatment by antivirals. This could also be achieved when we test all

symptomatic individuals but ensuring the wide availability of over-the-counter self-tests

alongside large clinic-based test availability ("OTC self-test").

Sensitivity analyses

We performed several sensitivity analyses in Georgia, owing to the relatively greater impact

of antivirals among the simulated LMICs, and investigated the extent to which our results

may deviate under different key assumptions. First, unvaccinated individuals could have

shared socio-demographic traits[32] and consequently vaccinated individuals would not

necessarily be randomly distributed across the population. As an approximation, we assumed

that vaccinated individuals, while still tiered by age, cluster among members from the same

household. Although reduction in infections remained similarly modest even when

vaccinated individuals tended to be clustered (Figure S14), a larger proportion of severe cases

were averted by antivirals (50% vaccinated: 8% (random) vs. 30% (clustered); 90%

vaccinated: 14% (random) vs. 44% (clustered)) at the highest epidemic intensity simulated

(R_e=2) but only if testing rates were large enough to support the distribution of antivirals

(500 tests/100,000 people/day; Figure S15). However, the greater impact of antivirals on

severe cases here is attributed to the increased number of severe cases stemming from

vaccinated individuals being clustered (Figure S15B). We found that severe cases increased

by 15-170% across all simulated scenarios if vaccinated individuals were clustered by

households as opposed to being randomly assigned. This correspondingly led to greater oral

antiviral demand as well with one antiviral course distributed for every 53-128 (5-104)

persons per year if testing rate was 100 (500) tests/100K/day. In short, while oral antivirals

could alleviate the greater disease burden associated with clustering among vaccinated

individuals, it is only facilitated by large enough testing rates and the need for greater

antiviral supply. The more critical factor towards lowering severe cases is to minimize spatial

bias among vaccinated individuals.

Second, we had assumed low average estimates of vaccine effectiveness (i.e. 29% and 70%

protection against infection and severe disease respectively). However, vaccine effectiveness

can be improved by updating the vaccine strains to match circulating viruses or through

booster shots. We repeated our simulations with vaccines conferring greater effectiveness,

including known average protection against Delta-like (i.e. 52% and 96% protection against

infection and severe disease respectively) and wild-type SARS-CoV-2 viruses (i.e. 75% and

97% protection against infection and severe disease).[19–21] Similar to our original results for

low vaccine effectiveness, use of antivirals could reduce transmissions in Georgia by up to

~20% but only if testing rates were high (500 tests/100K/day; Figure S16). In contrast, the

proportion of severe cases averted due to antivirals became increasingly uncertain (i.e. wider

error bars in Figure S17). This was because improved vaccine effectiveness, on top of wider

vaccination coverage, substantially reduced the number of severe cases. Nonetheless,

regardless of vaccine effectiveness and coverage, meaningful reductions in severe cases by

antivirals could only be achieved with higher testing rates ($\geq$ 100 tests/100K/day) to support

the administration of antivirals for infected high-risk individuals.

Third, we lowered the epidemic seeding condition from 1% to 0.1% such that antivirals were

distributed and used by the population earlier akin to the situation where Paxlovid is readily

available in certain countries. Although reduction in infections by antivirals continues to be

achieved only at higher testing rates, if antivirals were distributed earlier (i.e. starting from a

lower seeding condition), infections could be lowered by up to 30% even when test

availability was 100 tests/100K/day (e.g. At 100 tests/100K/day and 50% vaccination

coverage, only an average of 1% of infections were averted due to antivirals when $R_e = 1.5$

if the seeding condition was set to 1% but increased to 21% if seeding proportion was

lowered to 0.1%; Figure S18). The reduction in infections compounded the impact of

antivirals on severe case reduction: the proportion of severe disease averted due to antivirals

increased with improved outcomes at higher vaccination coverage (e.g. At 100

tests/100K/day and 50% vaccination coverage, only an average of 4% of severe cases were

averted due to antivirals when $R_e = 1.5$ if seeding condition was at 1% but increased to 47%

if seeding proportion was lowered to 0.1%; Figure S19). The lower seeding condition also led

to a fair proportion of severe cases averted at 10 tests/100K/day but with large uncertainty

(i.e. wider error bars in Figure S19A) and mostly only when $R_e < 1$ or at high vaccination

coverage (90%). This suggests that the benefit of antivirals can be further augmented by early

widespread adoption of test-and-treat programs.

Finally, the results above were predicated on crisis-period willingness-to-test behavior.

However, in many countries, the willingness of people to test has waned substantially in the

post-emergency phase of the pandemic. To investigate the consequences of this decline, we

repeated our simulations for all countries assuming that the likelihood a symptomatic

individual seek testing was 10% instead of the 65% assumed in the prior results (Figure S20).

Low willingness to test would substantial reduce the impact of potential test-and-treat

programs under all test availabilities, averting no more than ~10% of severe cases on average

in any country for all simulated R_e values and vaccination coverage.

Discussion

Individual-level data on the effectiveness of antivirals for reducing severe disease[3,4] and the

modelling work presented here highlight that substantial reductions in COVID-19 disease

burden could arise from population-level test-and-treat programs. However, the low testing

rates in the post-emergency phase of the pandemic represent a profound impediment for

realizing the benefits of such programs. Most of the analyses described here focused on test

availability as the functional constraint on the development of test-and-treat programs and

this remains an issue in many LMICs. However, in many countries, regardless of socio-

economic status, the willingness of people to pursue testing for respiratory virus disease is

either low or declining and this presents challenges even when tests are available.

Given that antivirals are unlikely to have substantial impact on population-level

transmission,[5] if the main objective of testing is to maximize the distribution of antivirals to

infected high-risk individuals, restricting clinic-based testing to only high-risk symptomatic

individuals at testing rates of 100 tests/100K/day could lead to 3.3-6.4-fold increase in

proportions of severe cases averted relative to the default scenario where no restrictions to

clinic-provided testing was imposed, provided that people are proactively seek testing.

It is also possible to require asymptomatic, high-risk household contacts of test-positive

symptomatic individuals to perform self-tests in order to initiate as many high-risk infected

individuals to early antiviral treatment as possible. However, setting aside tests to screen

high-risk household contacts under test availability constraints diminish the utility of tests

that would have otherwise been used to test symptomatic individuals who sought testing. In

turn, the proportion of severe cases and deaths averted due to antiviral distribution decrease

by a relative factor of two to ten-fold under this strategy. A potential workaround could be to

distribute antivirals to high-risk household contacts of test-positive individuals without the

need to confirm if the high-risk contacts were infected themselves by testing. However, this

would also increase the number of antiviral courses needed as well as result in wastage

among individuals who were not infected. A cost effectiveness analysis could be performed

to identify the most resource effective strategy but is beyond the scope of this work.

Assuming high willingness to test, ensuring the wide-availability of over-the-counter self-

tests could also lead to substantial reductions in severe cases (56-59%) and deaths (67-70%)

at $R_e \geq 1.5$ (e.g. BA.1 or BA.5 variant-like events). However, if reflexive testing is needed

for administration of antivirals, these reductions would only be possible if clinic-provided

testing is maintained at the mean HIC rate of 500 tests/100K/day. If clinical testing volumes

drop to 100 tests/100K/day, the expected reduction in severe cases and deaths attributable to

antivirals would fall to only 14% and 19% respectively in an epidemic wave initializing at

$R_e = 2.0$ (e.g. BA.1 variant-like event).

Our results suggest that regardless of the (test and antiviral) distribution strategy, an effective

test-and-treat program in any country requires large testing rates ($\gg$100 tests/100K/day) that

are far beyond testing rates reported globally since 2023.[9] In turn, increasing vaccination is

likely a more viable approach to lower severe cases than implementing large-scale test-and-

treat programs. To compare the vaccination coverage and the resource requirements needed

for test-and-treat to achieve the same reduction in disease burden, we computed the

additional vaccination coverage needed to halve the number of severe cases at different R_e

under 10% starting vaccination coverage. We also estimated the equivalent number of tests

and antivirals distributed to half the number of severe cases (Table 2). Across all countries

and R_e, we estimated that an additional 24%-67% of the population must be vaccinated to

reduce the number of severe cases by half without antivirals. Conversely, ~9,000 – 400,000

courses of antivirals per 1,000,000 people would be needed to avert the same number of

severe cases by antivirals for one epidemic wave. Furthermore, we estimated that ~200 –

7,000 tests must be performed per 100,000 people per day to support the distribution of those

antivirals. While these testing rates were achieved by some high-income countries during the

COVID-19 pandemic, no countries are testing at anywhere near these rates in the post-

emergency phase, suggesting that vaccination would likely be the more efficient option for

reducing severe disease burden.

There have been other modelling efforts that estimated substantial reductions in disease

burden by distributing antivirals to 20% - 50% of symptomatic infected individuals.

However, from our analyses, doing so would also require testing rates that are far greater than

500 tests/100,000 people/day. First, Leung et al.[7] estimated that distributing antivirals to 50%

of all symptomatic infected individuals regardless of risk status would only reduce

hospitalizations by 10-13% in a population with high vaccination coverage (70-90%). For the

Netherlands, we simulated a population with 80% vaccination coverage and large test

availability, that included both clinic-based and over-the-counter self-tests, such that at least

50% of all symptomatic individuals were diagnosed. We estimated that 56-59% of severe

cases could be averted if only high-risk symptomatic individuals were administered

antivirals. When we reconfigured our simulations to now distribute antivirals to 50% of all

symptomatic infected individuals, the proportion of severe cases averted lower to only 18%

which is more in line with Leung et al.

Second, Matrajt et al. found that initiating 20% of infected individuals that were >65 years of

age on antivirals daily could avert 32-43% of deaths in an Omicron-like wave ($R_e \geq 2$) for an

unvaccinated population in LMICs such as Kenya and Mexico.[5] We had estimated that 31-

62% of deaths could be averted at $R_e = 2$ at low (10%) vaccination coverage in LMICs but

only if test availability was 500 tests/100K/day and clinic-provided symptomatic testing were

restricted to high-risk individuals, which would mean a daily average of 19-20% of high-risk

infected individuals being initiated on treatment each day. If there are no restrictions on

access to clinic-provided tests, testing rate must be at least 750 tests/100K/day to initiate 20%

of infected >65-years on antivirals daily with >95% probability, indicating that the previous

from Martrajt et al. predicated on very high testing rates.

Finally, Brault et al. estimated that 11% of hospitalizations could be averted if antivirals with

50% effectiveness were administered to half of all high-risk cases in Wallis and Futuna,

where ~70% of individuals have at least two doses of vaccines, during an epidemic wave

with a doubling time of 2-3 days.[8] In the closest scenario we had simulated (i.e $R_e = 2$, 46%

effectiveness of antivirals, 50% vaccination coverage and 500 tests/100K/day), we estimated

that severe cases could be reduced by 7% in Brazil (Figure 2B), which has a similar

demography to Wallis and Futuna (i.e. median age = 33 and 35 years in Brazil and Wallis and

Futuna respectively; proportion of individuals ≥65 years = 10% and 13% respectively).

However, like the two preceding examples, this is only possible at testing rates that are many-

fold higher than those performed in most LMICs both during and after the emergency-phase

of the pandemic.

There are limitations to our work: First, our simulations were based on the estimated

effectiveness of nirmatrelvir–ritonavir. We did not consider the clinical benefits of other oral

antivirals as nirmatrelvir–ritonavir was the most efficacious antiviral available during the

development of this work.

Second, as a simplification, we assumed that individuals with pre-existing comorbidities that

augment the risk of severe COVID-19 disease outcomes were randomly distributed across the

population. The prevalence of certain comorbidities is known to correlate with socio-

economic and demographic factors,[33,34] resulting in the clustering of severe cases with similar

socio-economic backgrounds. However, there is limited country-specific data on the

prevalence and distribution of comorbidities across the population, especially for LMICs. We

would also need to stratify the simulation population socio-economically which is beyond the

scope of this study.

Third, we had assumed that clinical testing for disease and administration of treatment occur

on the same day in our simulations. However, any practical barriers that limit timely access to

antivirals (e.g. inadequate supply and distribution, limited access to healthcare providers,

acceptance of antiviral therapy) can substantially reduce the estimated impact of test-and-

treat programs.[35] As shown in Figure S12, even under a large test availability scenario (with

self-tests), if administration of antivirals was delayed by >2 days, <20% of high-risk treated

individuals received their antiviral courses within the 5 days post-symptom onset window

when Paxlovid was reported to be efficacious. As such, even if testing rates could sufficiently

support test-and-treat programs, delays in accessing antivirals, which had been reported in

various LMICs,[36] must be minimized for these programs to remain effective. Ideally, testing

and treatment of infected patients should occur at the same clinical interaction.

Next, others have showed that with greater vaccine effectiveness against infection (60%), a

high vaccination coverage (~70-80%) coupled with antivirals that have an effect in lowering

transmissions could synergistically reduce infections in the population.[5] However, for only

~20% of infections to be averted in an Omicron-like wave (i.e. doubling time of 2-3 days[37]),

the antiviral must block onward transmission completely after initiating treatment and 30% of

symptomatic infected adults must be administered antivirals daily.[5] Even if an antiviral that is

100% effective in truncating transmissions exist and there was high willingness to test, the

testing rate must at least be 764 tests/100K/day to initiate 30% of symptomatic infected

individuals to treatment daily with >95% probability based on our estimates.

Finally, we did not factor in changes to individual immunity levels due to previous infections

or immune waning. As a simplification, we assumed that these effects have been implicitly

captured by various initial R_e values and were able to simulate epidemics with prevalence

ranges similar to those reported during the spread of Omicron subvariants BA.5 and XBB.1.5

(Figure S1). However, it is currently unclear how changing immunity dynamics in the future

could affect severe disease outcomes.

Taken all together, while test-and-treat programs have substantial theoretical utility for

reducing population-level burden of disease, there remain fundamental challenges in terms of

the availability of diagnostics and the willingness of people to seek testing in general and

particularly within the relatively short window of effectiveness of the antivirals considered

here. The potential benefits and resource requirements of test-and-treat programs must also

be carefully considered if budget constraints make vaccination programs a competing

interest.

Methods

The PATAT simulation model

PATAT creates an age-structured population of individuals within contact networks of multi-

generational households, schools, workplaces, regular mass gatherings (e.g. religious

gatherings) and random community settings with country-specific demographic data (see

Supplementary Text). Epidemic simulations begin with 1% of the population infected with

SARS-CoV-2 and compute transmissions between individuals across different contact

networks each day. The computational flow of a PATAT simulation is summarised as

follows: First, an age-structured population of agents is created. Close contact networks are

subsequently created based on the given demographic data. The simulation is then initialised

and iterates over a given period of time where each time step corresponds to a day. The

operations during each timestep encompass updating the disease progression of infected

individuals, the status of isolated/quarantined agents, application of community testing

strategies and computation of transmission events within contact networks.

PATAT implements a SEIRD epidemic model where the simulated population is

distinguished between five compartments: susceptible, exposed (i.e. infected but is not

infectious yet; latent phase), infected (which include the presymptomatic infectious period for

symptomatic agents), recovered and dead. The infected compartments are further stratified by

their presented symptoms, including asymptomatic, presymptomatic, symptomatic mild or

severe. All symptomatic agents will also first undergo an infectious presymptomatic period

after the exposed latent period. They will either develop mild symptoms who will always

recover from the disease or experience severe infection which could either lead to death or

recovery. As a simplification, PATAT assumes that all agents presenting severe symptoms

are sufficiently isolated from the population (e.g. through hospitalization) that they are

unlikely to contribute to further transmissions.

When an infectious agent i comes into contact with a susceptible individual j, the probability

of transmission ($p_{transmission,(i,j)}$) is given by:

$p_{transmission,(i,j)} = \beta \times \Phi_i \times f_c \times f_{asymp,i} \times f_{load,i} \times f_{immunity,j} \times f_{susceptiblity,j} \times \rho_i \times \rho_j$

where β is the base transmission probability per contact, Φ_i is the overdispersion factor

modelling individual-level variation in secondary transmissions (i.e. superspreading events),

f_c is a relative weight adjusting β for the network setting c where the contact has occurred,

$f_{asymp,i}$ is the assumed relative transmissibility factor if infector i is asymptomatic,

$f_{immunity,j}$ measures the immunity level of susceptible j against the transmitted virus (i.e.

$f_{immunity,j} = 1$ if completely naïve; $f_{immunity,j} = 0$ if fully protected), $f_{susceptiblity,j}$ is the

age-dependent susceptibility of j, ρ_i and ρ_j are the contact rates of infector i and susceptible

j respectively.

Φ_i is randomly drawn from a negative binomial distribution with mean of 1.0 and shape

parameter of 0.45.[38] As evidence have been mixed as to whether asymptomatic agents are

less transmissible, we conservatively assume there is no difference relative to symptomatic

patients (i.e. $f_{asymp,i} = 1$). The age-structured relative susceptibility values $f_{susceptiblity,j}$ are

derived from odds ratios reported by Zhang et al.[22] (Table S1).

β is determined by running initial test simulations with a range of values on a naïve

population with no interventions that would satisfy the target reproduction number as

computed from the resulting exponential growth rate and distribution of generation

intervals.[39] f_c is similarly calibrated during these test runs such that the transmission

probabilities in households, workplaces, schools, and all other community contacts are

constrained by a relative weighting of 10:2:2:1.[23]

The total duration of infection since exposure depends on the symptoms presented by the

patient and is comprised of different phases (i.e. latent, asymptomatic, presymptomatic,

onset-to-recovery/death). The time period of each phase is drawn can be found in Table S1.

For each infected individual, PATAT randomly draws a within-host viral load trajectory over

the duration of infection, which impacts the sensitivity of Ag-RDTs[40], based on known

distributions for Omicron BA.1[41]. Similar viral load trajectories were drawn for both

asymptomatic and symptomatic infected individuals[42] using a stochastic model modified

from the one previously developed by Quilty et al.[43] A baseline Ct value ($Ct_{baseline}$) of 40 is

established upon exposure. The infected agent becomes infectious upon the end of the latent

period and their Ct value is assumed to be ≤ 30. A peak Ct value is then randomly drawn

from a normal distribution (Table S1). Peak Ct is assumed to occur upon symptom onset for

symptomatic agents and one day after the latent period for asymptomatic individuals.

Cessation of viral shedding (i.e. return to $Ct_{baseline}$) occurs upon recovery or death. PATAT

assumes that the transition rate towards peak Ct value should not be drastically different to

that when returning to baseline upon cessation (i.e. there should be no sharp increase to

baseline Ct value after gradual decrease to peak Ct value or vice versa). As such, the time

periods of the different phases of infection are randomly drawn from the same quintile of

their respective sample distribution. The viral load trajectory is then simulated by fitting a

cubic Hermite spline to the generated exposed ($t_{exposed}$, $Ct_{baseline}$), latent (t_{latent},

$Ct_{latent} = 30$), peak (t_{peak}, Ct_{peak}) and cessation values ($t_{recovered/death}$, $Ct_{baseline}$). The

slope of the fitted curve is assumed to be zero for all of them except during t_{latent} where its

slope is assumed to be $\frac{Ct_{peak} - Ct_{baseline}}{t_{peak} - t_{exposed}}$. PATAT then uses the fitted trajectory to linearly

interpolate the viral load transmissibility factor ($f_{load,i}$) of an infectious agent i assuming that

they are twice as transmissible at peak Ct value (i.e. $f_{load} = 2$) relative to when they first

become infectious (i.e. Ct value = 30; $f_{load} = 1$).

Unlike PCR which is highly sensitive due to prior amplification of viral genetic materials, the

sensitivity of Ag-RDT will depend on the viral load of the tested patient. While the

specificity of Ag-RDT is assumed to be 98.9%, its sensitivity depends on the Ct values of the

tested infected agent: Ct > 35 (0%); 35 – 30 (20.9%); 29 – 25 (50.7%); Ct ≤ 24 (95.8%).[40]

We assumed that agents would change their behaviour when (i) they start to present

symptoms and go into self-isolation (10% compliance assumed, 71% endpoint adherence)[27];

(ii) they test positive and are isolated for 10 days (50% compliance assumed, 86% endpoint

adherence)[27]; or (iii) they are household members (without symptoms) of positively-tested

agents and are required to be in quarantine for 14 days (50% compliance assumed, 28%

endpoint adherence)[27]. Once an agent goes into isolation/quarantine, we linearly interpolate

their probability of adherence to stay in isolation/quarantine over the respective period. Given

the lack of infrastructure and resources to set up dedicated isolation/quarantine facilities in

many LMICs, we assumed that all isolated and quarantined individuals would do so at home.

We simulated 90-day epidemic waves in a community of 1,000,000 individuals using

demographic data collected from three LMICs (i.e. Brazil, Georgia, Zambia) and the

Netherlands as a HIC counterpart. We simulated different vaccination coverage (10%, 50%

and 90%) for all countries for comparability. In the separate analysis examining how

widespread availability of over-the-counter self-tests could impact test-and-treat programs in

HICs, we assumed that 80% of the population was vaccinated in the Netherlands based on

estimates on July 2022,[44] which is largely comparable to other HICs.[45] As a simplification,

we assumed that vaccination protection rates against infection and severe disease were 29%

and 70% respectively, which were based on the more conservative, lower average estimates

of vaccine effectiveness against BA.1 across different vaccines (i.e. mRNA and ChAdOx1

nCoV-19 vaccine) and doses (i.e. 1-3 doses).[19–21] We did not assume a specific protection

rate against death since the referenced studies had reported effectiveness estimates against

severe disease outcomes which include hospitalization and/or death. Nonetheless, protection

of deaths is implicitly accounted for since individuals could only die from COVID-19 if they

had progressed to severe disease in the model.

Diagnostic testing

In the model, individuals with symptomatic COVID-19 have a probability of seeking testing

at a healthcare facility. We also estimated symptomatic testing demand from individuals

without COVID-19 who sought clinic-provided testing (e.g. individuals who present with

similar respiratory symptoms): Based on the range of test positivity rates reported by various

countries during the second-half of 2021 (when community testing was assumed to still be

prevalent in most countries),[45] we assumed that test positivity rate was 10% at the start as

well as end of an epidemic wave, and a 20% test positivity rate at the peak, linearly

interpolating the demand for periods between these time points.[11,12]

We also simulated scenarios where household contacts of clinic-provided positively-tested

individuals were given Ag-RDTs for self-testing for three consecutive days following the

positive clinical test of the latter. Adherence (likelihood) to testing by asymptomatic

household contacts was assumed to decrease linearly to 50% by the third day. We also

simulated an alternative test distribution strategy where we restricted clinic-provided

symptomatic testing to high-risk individuals only.

We performed simulations under three levels of average test availability at healthcare clinics:

10 (mean LMIC testing rate as of Q2/2022),[9] 100 and 500 (mean HIC testing rate as of

Q2/2022)[9] tests/100K/day. Regardless if symptomatic individuals choose to self-isolate, after

$\tau_{delay,symp-test}$ days from symptom onset, the symptomatic agent may decide to get tested

with a Bernoulli probability of $p_{symp-test}$. PATAT assumes that agents who have decided

against symptomatic testing (i.e. failed Bernoulli trial) or received negative test results will

not seek symptomatic testing again. We assumed that average $p_{symp-test}$ = 65% on average

based on surveys of test-seeking behaviour during the COVID-19 pandemic.[30,31] In other

words, there is an average 65% chance that an individual with mild symptom would seek

clinic-provided testing and were only tested if there were available test stocks. We lowered

$p_{symp-test}$ to 10% in a sensitivity analysis to estimate the impact of test-and-treat programs

under waning willingness to test.

In the separate analysis where over-the-counter self-tests were available in the Netherlands,

we assumed that only 10% of mild symptomatic individuals in the Netherlands would seek

clinic-provided testing upon symptom onset based on average daily testing rates reported by

all Dutch municipal health services in 2021-Q1/2022 (i.e. approximately up to the end of the

Omicron BA.1 wave; 7551 tests/100K/day) and Q2/2022 (post Omicron BA.1 wave; 641

tests/100K/day).[46] We assumed that 50% or 80% of individuals who opted not to seek clinic-

provided testing would perform a self-test using an over-the-counter Ag-RDT. We assumed

that all high-risk individuals who tested positive would then seek reflexive testing at clinics

to be disbursed an antiviral course.

Oral antivirals

Regardless of their vaccination status (per WHO guidance),[47] all high-risk individuals who

tested positive within five days after symptom onset were eligible for a course of antiviral

therapy.[3,4] We did not impose any caps on antiviral availability as we wanted to estimate the

potential number of antiviral courses needed and thus their maximum achievable impact on

epidemic outcomes under different levels of test availability and antiviral distribution

strategies. We did not factor any risk reduction in transmissions or deaths given the lack and

low certainty of evidence of the impact of oral antivirals on protection against transmission

and mortality respectively.[47] However, individuals could only die from COVID-19 if they

had progressed to severe disease in our model.

For individuals who were treated with antivirals that were simulated to result in severe

disease, we performed a Bernoulli trial with the probability of averting severe disease (i.e.

46%), provided that they were currently in the presymptomatic phase or were experiencing

mild disease. If the Bernoulli trial succeeded, we re-simulated their disease progression and

within-host viral dynamics using the procedures above but now under the assumption that

they would develop only mild disease and conditioning that the maximum viral load is lower

than before. Changes were only made to the upcoming phases of disease progression from the

then current phase of infection. The average recovery period of having mild symptoms was

assumed to be 5.4 days as opposed to 18.1 days when presenting severe disease.[41,48] In turn,

while we did not parameterize the impact on transmission reduction by antivirals, the

shortened recovery, and thus infectious period as well as lower maximum viral load of

individuals who were effectively treated could result indirect reduction in onward

transmissions.

We performed five independent simulations for each combination of parameters described

above. All key parameters are tabulated in Table S1. Further details of PATAT are described

in Han et al.[11,12] and the Supplementary Information. The PATAT model source code is

available at https://github.com/AMC-LAEB/PATAT-sim.

Data and code availability:

All data relevant to the study are included in the Article, the Supplementary Information and

the GitHub repository (https://github.com/AMC-LAEB/PATAT-sim). The PATAT model

source code can also be found in the GitHub repository

(https://github.com/AMC-LAEB/PATAT-sim).

Acknowledgements:

This work was supported by the European Research Council [NaviFlu 818353 to A.X.H. and

C.A.R.], the National Institutes of Health [5R01AI132362-04 to C.A.R.] and the Dutch

Research Council (Nederlandse Organisatie voor Wetenschappelijk Onderzoek) [Vici

09150182010027 to C.A.R.]. This work was supported by the Rockefeller Foundation, and

the Governments of Germany, Canada, UK, Australia, Norway, Saudi Arabia, Kuwait,

Netherlands and Portugal [all authors]. The authors are pleased to acknowledge that all

computational work reported in this paper was performed on the Shared Computing Cluster

which is administered by Boston University's Research Computing Services

(www.bu.edu/tech/support/research/).

Authors' contributions:

Conceptualization: A.X.H., E.H., S.C., B.R., B.E.N., C.A.R.

Methodology: A.X.H., B.E.N., C.A.R.

Software: A.X.H.

Validation: A.X.H., E.H., S.C., B.R., B.E.N., C.A.R.

Formal analysis: A.X.H.

Investigation: A.X.H., B.E.N., C.A.R.

Resources: B.E.N., C.A.R.

Data curation: A.X.H., B.E.N., C.A.R.

Writing – original draft: A.X.H., B.E.N., C.A.R.

Writing – review & editing: A.X.H., E.H., S.C., B.R., B.E.N., C.A.R.

Visualization: A.X.H., B.E.N., C.A.R.

Supervision: B.E.N., C.A.R.

Project administration: B.E.N., C.A.R.

Funding acquisition: E.H., S.C., B.R., B.E.N., C.A.R.

Declaration of interests:

The authors declare that they have no competing interests.

 Figure legends

Figure 1: Demography of simulated countries. (a) Bar plot shows the age distribution of

each simulated country archetype (Low and middle-income countries (LMICs): Brazil,

Georgia, Zambia; High-income country: Netherlands) stratified in 5-year bins. Each dashed

black line in the Brazil, Georgia and Zambia plots denotes the age distribution of one of 132

other LMICs[14] that best matches (i.e. lowest mean absolute error) the age distribution of the

simulated country archetype. Age distribution of the population in each country is

downloaded from World Population Prospects compiled by the United Nations[15]. (b)

Heatmap showing the normalized relative contact rates between individuals of different age

groups in 5-year bins averaged across all contact networks generated by the PATAT

simulation model.

Figure 2: Impact of test-and-treat on severe cases. No restrictions on access to

symptomatic testing at clinics (i.e. all symptomatic individuals who sought testing at clinics

would receive one if in stock) and high-risk household contacts of test-positive individuals

are not tested. All eligible high-risk individuals (i.e. ≥60 years of age or an adult ≥18 years

with a relevant comorbidity) who tested positive were given a course of oral antivirals. Line

plots (left y-axis) show the percentage change in severe cases relative to no distribution of

antivirals under different levels of mean test availability (different shades of color) after a 90-

day epidemic wave in a population of 1,000,000 individuals with (a) 10%, (b) 50%, and (c)

90% vaccination coverage for different epidemic intensities (measured by the initial effective

reproduction number (R_e); x-axis). Bar plots (right y-axis) show the number of severe cases

in each corresponding scenario. The dotted outline of each bar shows the number of severe

cases of each scenario when no antivirals were distributed.

Figure 3: Impact of test-and-treat on severe cases when restricting symptomatic testing

to high-risk individuals only. High-risk household contacts of test-positive individuals are

not tested. All eligible high-risk individuals (i.e. ≥60 years of age or an adult ≥18 years with

a relevant comorbidity) who tested positive were given a course of oral antivirals. Line plots

(left y-axis) show the percentage change in severe cases relative to no distribution of

antivirals under different levels of mean test availability (different shades of color) after a 90-

day epidemic wave in a population of 1,000,000 individuals with (a) 10%, (b) 50%, and (c)

90% vaccination coverage for different epidemic intensities (measured by the initial effective

reproduction number (R_e); x-axis). Bar plots (right y-axis) show the number of severe cases

in each corresponding scenario. The dotted outline of each bar shows the number of severe

cases of each scenario when no antivirals were distributed.

Figure 4: Estimated need of oral antivirals. Line plots show the ratio of estimated oral

antiviral courses needed to number of people per year (expressed as 1 oral antiviral course

per n number of individuals; assuming two epidemic waves a year) in simulated countries

(color) under different simulated scenarios (i.e. testing rate at 100 or 500 tests/100,000

people/day (shading and linestyle) and distribution modality (left plot panel: test all

symptomatic individuals who sought testing at clinics; middle plot panel: test all

symptomatic individuals who sought testing as well as distributing clinic-provided self-tests

to high-risk asymptomatic household contacts of test-positive individuals; right plot panel:

test only high-risk symptomatic individuals who sought testing at clinics). All test-positive

eligible high-risk individuals from clinic-provided testing would receive a course of oral

antivirals. (a) 10%, (b) 50% and (c) 90% vaccination coverage assumed for the simulated

population.

Figure 5: Impact of test-and-treat in a high-income country (Netherlands) with wide

availability of over-the-counter self-tests. No restrictions on access to symptomatic testing

at clinics (i.e. all symptomatic individuals who sought testing at clinics would receive one if

in stock) and high-risk household contacts of test-positive individuals are not tested. Over-

the-counter antigen rapid diagnostic tests (Ag-RDTs) are assumed to be widely available with

unlimited stocks. As such, we assumed that only 10% of symptomatic individuals would seek

clinical testing directly while 80% of those who opted not to seek clinic-provided testing

would perform self-testing using over-the-counter Ag-RDTs. All high-risk individuals who

tested positive through self-testing would seek reflexive testing at clinics on the same day.

All eligible high-risk individuals (i.e. ≥60 years of age or an adult ≥18 years with a relevant

comorbidity) who tested positive at clinics, either directly or through reflexive testing, were

given a course of oral antivirals. Line plots (left y-axis) show the percentage change in (a)

total infections, (b) severe cases and (c) deaths relative to no distribution of antivirals under

different clinical testing rates (different shades of color) after a 90-day SARS-CoV-2

epidemic wave in a population of 1,000,000 individuals with 80% vaccination coverage for

different epidemic intensities (measured by the initial effective reproduction number (R_e); x-

axis). Bar plots (right y-axis) show the number of severe cases in each corresponding

scenario. The dotted outline of each bar shows the number of severe cases of each scenario

when no antivirals were distributed.

 Tables

806 **Table 1: Number and proportion of severe cases averted due to distribution of oral**

807 **antivirals at 10% and 90% vaccination coverage**. No restrictions on access to symptomatic

808 testing at clinics (i.e. all symptomatic individuals who sought testing at clinics would receive

809 one if in stock) and high-risk household contacts of test-positive individuals are not tested.

Country	Testing rate (tests/100,000 people/day)	R_e	10% vaccination coverage		90% vaccination coverage	
			No. of severe cases averted	Proportion of severe cases averted	No. of severe cases averted	Proportion of severe cases averted
Zambia	10	0.9	29	2.7	3	1.1
		1.2	16	0.3	20	1.1
		1.5	0	0.0	0	0.0
		2.0	22	0.2	32	0.7
	100	0.9	147	16.9	48	20.1
		1.2	170	3.3	72	4.4
		1.5	172	1.9	131	3.7
		2.0	0	0.0	0	0.0
	500	0.9	242	35.0	102	45.6
		1.2	1053	26.5	441	38.0
		1.5	780	9.0	527	16.4
		2.0	824	7.2	433	8.8
Brazil	10	0.9	0	0.0	0	0.0
		1.2	0	0.0	0	0.0
		1.5	0	0.0	0	0.0
		2.0	0	0.0	0	0.0
	100	0.9	303	23.7	133	54.6
		1.2	623	6.4	195	13.6
		1.5	245	1.2	196	3.7
		2.0	0	0.0	0	0.0
	500	0.9	511	47.4	134	55.3
		1.2	2739	43.1	545	55.4
		1.5	2005	10.7	1404	35.7
		2.0	1553	6.1	872	9.4
Georgia	10	0.9	0	0.0	0	0.0
		1.2	0	0.0	0	0.0
		1.5	163	0.4	80	1.2

		2.0	0	0.0	0	0.0
	100	0.9	635	49.9	170	65.3
		1.2	1459	13.4	336	29.6
		1.5	929	2.6	483	8.7
		2.0	0	0.0	0	0.0
	500	0.9	792	63.4	167	63.5
		1.2	4344	65.3	597	66.9
		1.5	7481	24.3	2270	66.1
		2.0	4435	8.4	2115	13.6
Netherlands	10	0.9	0	0.0	0	0.0
		1.2	0	0.0	0	0.0
		1.5	0	0.0	21	0.2
		2.0	0	0.0	0	0.0
	100	0.9	598	48.2	171	66.6
		1.2	2333	20.1	362	30.7
		1.5	953	2.7	419	7.5
		2.0	0	0.0	0	0.0
	500	0.9	811	63.1	185	67.0
		1.2	4857	68.5	604	67.3
		1.5	8947	28.0	2123	65.1
		2.0	3190	5.9	2960	18.1

Table 2: Vaccination coverage and test-and-treat requirements to half the number of severe cases at 10% pre-existing vaccination coverage. For vaccination, the additional number of individuals per 1,000,000 people that must be vaccinated on top of the 10% pre-existing vaccination coverage is tabulated. For test-and-treat, there are no restrictions on access to symptomatic testing at clinics (i.e. all symptomatic individuals who sought testing at clinics would receive one if in stock) and high-risk household contacts of test-positive individuals are not tested.

Country	R_e	Vaccination	Test-and-treat	
		No. of vaccinated individuals per 1,000,000 people	No. of antivirals distributed per 1,000,000 people	Testing rate (tests/100,000 people/day)
Zambia	1.2	496,192	18,436	545
	1.5	586,432	60,370	2,185
	2.0	665,567	113,835	4,383
Brazil	1.2	312,095	15,494	316
	1.5	437,067	82,093	1,760
	2.0	566,068	197,465	4,751
Georgia	1.2	253,988	9,246	166
	1.5	317,855	52,499	721
	2.0	477,855	211,977	3,471
Netherlands	1.2	240,271	8,758	206
	1.5	319,695	49,952	701
	2.0	474,469	394,414	6,802

Supplemental information

Document S1. Supplementary Text; Figures S1-S20; Table S1.

References

1.   Singh, M. & de Wit, E. Antiviral agents for the treatment of COVID-19: Progress and

challenges. *Cell Rep Med* **3**, 100549 (2022).

2.   Jayk Bernal, A. *et al.* Molnupiravir for Oral Treatment of Covid-19 in Nonhospitalized

Patients. *New England Journal of Medicine* **386**, 509–520 (2022).

3.   Hammond, J. *et al.* Oral Nirmatrelvir for High-Risk, Nonhospitalized Adults with

Covid-19. *New England Journal of Medicine* **386**, 1397–1408 (2022).

4.   Najjar-Debbiny, R. *et al.* Effectiveness of Paxlovid in Reducing Severe Coronavirus

Disease 2019 and Mortality in High-Risk Patients. *Clinical Infectious Diseases* (2022)

doi:10.1093/CID/CIAC443.

5.   Matrajt, L., Brown, E. R., Cohen, M. S., Dimitrov, D. & Janes, H. Could widespread

use of antiviral treatment curb the COVID-19 pandemic? A modeling study. *BMC*

Infect Dis **22**, 1–16 (2022).

6.   Usher, A. D. The global COVID-19 treatment divide. *The Lancet* **399**, 779–782

(2022).

7.   Leung, K., Jit, M., Leung, G. M. & Wu, J. T. The allocation of COVID-19 vaccines

and antivirals against emerging SARS-CoV-2 variants of concern in East Asia and

Pacific region: A modelling study. *Lancet Reg Health West Pac* **21**, 100389 (2022).

8.   Brault, A. *et al.* Modelling the end of a Zero-COVID strategy using

nirmatrelvir/ritonavir, vaccination and NPIs in Wallis and Futuna. *Lancet Reg Health*

West Pac **30**, (2023).

9.   FIND. Test tracker - FIND. https://www.finddx.org/covid-19/test-tracker/ (2022).

10.   Gill, C. J. *et al.* What is the prevalence of COVID-19 detection by PCR among

852   deceased individuals in Lusaka, Zambia? A postmortem surveillance study. *BMJ Open*

853   **12**, (2022).

854   11.   Han, A. X. *et al.* Strategies for using antigen rapid diagnostic tests to reduce

855   transmission of SARS-CoV-2 in low- and middle-income countries: a mathematical

856   modelling study applied to Zambia. *Clinical Infectious Diseases* (2022)

857   doi:10.1093/CID/CIAC814.

858   12.   Han, A. X. *et al.* SARS-CoV-2 diagnostic testing rates determine the sensitivity of

859   genomic surveillance programs. *Nature Genetics 2023 55:1* **55**, 26–33 (2023).

860   13.   Wroe, E. B., Seung, K. J., Baker, B. K. & Farmer, P. E. Test and treat: a missing link

861   in the global fight against COVID-19. *Lancet Glob Health* **10**, e181–e182 (2022).

862   14.   World Bank Country and Lending Groups – World Bank Data Help Desk.

863   https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-

864   country-and-lending-groups.

865   15.   World Population Prospects - Population Division - United Nations.

866   https://population.un.org/wpp/Download/Standard/MostUsed/.

867   16.   Mefsin, Y. M. *et al.* Epidemiology of Infections with SARS-CoV-2 Omicron BA.2

868   Variant, Hong Kong, January–March 2022 - Volume 28, Number 9—September 2022

869   - Emerging Infectious Diseases journal - CDC. *Emerg Infect Dis* **28**, 1856–1858

870   (2022).

871   17.   Tegally, H. *et al.* Emergence of SARS-CoV-2 Omicron lineages BA.4 and BA.5 in

872   South Africa. *Nat Med* **28**, (2022).

873   18.   UK Health Security Agency. SARS-CoV-2 variants of concern and variants under

874   investigation in England: Technical briefing 50. (2023).

19.   Andrews, N. *et al.* Covid-19 Vaccine Effectiveness against the Omicron (B.1.1.529)

Variant. *New England Journal of Medicine* **386**, 1532–1546 (2022).

20.   Buchan, S. A. *et al.* Effectiveness of COVID-19 vaccines against Omicron or Delta

symptomatic infection and severe outcomes. *medRxiv* 2021.12.30.21268565 (2022)

doi:10.1101/2021.12.30.21268565.

21.   Tseng, H. F. *et al.* Effectiveness of mRNA-1273 against SARS-CoV-2 Omicron and

Delta variants. *Nature Medicine 2022 28:5* **28**, 1063–1071 (2022).

22.   Zhang, J. *et al.* Changes in contact patterns shape the dynamics of the COVID-19

outbreak in China. *Science (1979)* **368**, 1481–1486 (2020).

23.   Kerr, C. C. *et al.* Covasim: An agent-based model of COVID-19 dynamics and

interventions. *PLoS Comput Biol* **17**, e1009149- (2021).

24.   Verity, R. *et al.* Estimates of the severity of coronavirus disease 2019: a model-based

analysis. *Lancet Infect Dis* **20**, 669–677 (2020).

25.   Ferguson, N. M. *et al.* Report 9: Impact of non-pharmaceutical interventions (NPIs) to

reduce COVID-19 mortality and healthcare demand. (2020) doi:10.25561/77482.

26.   O'Driscoll, M. *et al.* Age-specific mortality and immunity patterns of SARS-CoV-2.

Nature 2020 590:7844 **590**, 140–145 (2020).

27.   Brazeau, N. F. *et al.* Report 34 - COVID-19 Infection Fatality Ratio Estimates from

Seroprevalence | Faculty of Medicine | Imperial College London. (2020)

doi:10.25561/83545.

28.   Rawshani, A. *et al.* Severe COVID-19 in people with type 1 and type 2 diabetes in

Sweden: A nationwide retrospective cohort study. *The Lancet Regional Health -*

Europe **4**, (2021).

29.   Bertagnolio, S. *et al.* Clinical features of, and risk factors for, severe or fatal COVID-
899   19 among people living with HIV admitted to hospital: analysis of data from the WHO
900   Global Clinical Platform of COVID-19. *Lancet HIV* **9**, e486–e495 (2022).

901   30.   Rijksinstituut voor Volksgezondheid en Milieu. Research on behavioural rules and
902   well-being: round 3. https://www.rivm.nl/en/behavioural-science-and-covid-19/results-
903   of-study-behavioural-measures-and-well-being/round-3 (2020).

904   31.   Siegler, A. J. *et al.* Willingness to seek diagnostic testing for SARS-CoV-2 with home,
905   drive-through, and clinic-based specimen collection locations. *Open Forum Infect Dis*
906   **7**, (2020).

907   32.   Alvarez-Zuzek, L. G., Zipfel, C. M. & Bansal, S. Spatial clustering in vaccination
908   hesitancy: The role of social influence and social selection. *PLoS Comput Biol* **18**,
909   (2022).

910   33.   Ante-Testard, P. A. *et al.* Temporal trends in socioeconomic inequalities in HIV
911   testing: an analysis of cross-sectional surveys from 16 sub-Saharan African countries.
912   *Lancet Glob Health* **8**, (2020).

913   34.   Mendenhall, E., Kohrt, B. A., Norris, S. A., Ndetei, D. & Prabhakaran, D. Non-
914   communicable disease syndemics: poverty, depression, and diabetes among low-
915   income populations. *The Lancet* vol. 389 Preprint at https://doi.org/10.1016/S0140-
916   6736(17)30402-6 (2017).

917   35.   Gold, J. A. W. *et al.* Dispensing of Oral Antiviral Drugs for Treatment of COVID-19
918   by Zip Code–Level Social Vulnerability — United States, December 23, 2021–May
919   21, 2022. *MMWR Morb Mortal Wkly Rep* **71**, (2022).

920   36.   World Health Organization. Update on the rollout of COVID-19 tools: a report from
921   the ACT-A Tracking & Monitoring Task Force - 15 February 2023.

https://www.who.int/publications/m/item/update-on-the-rollout-of-covid-19-tools--a-

report-from-the-act-a-tracking---monitoring-task-force---15-february-2023 (2023).

37.   Viana, R. *et al.* Rapid epidemic expansion of the SARS-CoV-2 Omicron variant in

southern Africa. *Nature 2022* 1–10 (2022) doi:10.1038/s41586-022-04411-y.

38.   Endo, A., Abbott, S., Kucharski, A. J. & Funk, S. Estimating the overdispersion in

COVID-19 transmission using outbreak sizes outside China. *Wellcome Open Research*

2020 5:67 **5**, 67 (2020).

39.   Wallinga, J. & Lipsitch, M. How generation intervals shape the relationship between

growth rates and reproductive numbers. *Proceedings of the Royal Society B:*

Biological Sciences **274**, 599–604 (2006).

40.   Brümmer, L. E. *et al.* Accuracy of novel antigen rapid diagnostics for SARS-CoV-2: A

living systematic review and meta-analysis. *PLoS Med* **18**, e1003735- (2021).

41.   Hay, J. A. *et al.* Quantifying the impact of immune history and variant on SARS-CoV-

2 viral kinetics and infection rebound: A retrospective cohort study. *Elife* **11**, (2022).

42.   Boyton, R. J. & Altmann, D. M. The immunology of asymptomatic SARS-CoV-2

infection: what are the key questions? *Nature Reviews Immunology 2021 21:12* **21**,

762–768 (2021).

43.   Quilty, B. J. *et al.* Quarantine and testing strategies in contact tracing for SARS-CoV-

2: a modelling study. *Lancet Public Health* **6**, e175–e183 (2021).

44.   National Institute For Public Health and The Environment (RIVM). Figures on the

COVID-19 vaccination programme. https://www.rivm.nl/en/covid-19-

vaccination/figures-vaccination-programme (2022).

45.   Ritchie, H. *et al.* Coronavirus Pandemic (COVID-19). *Our World in Data* (2020).

46.   National Institute for Public Health and the Environment (RIVM). COVID-19 dataset.

https://data.rivm.nl/covid-19/ (2022).

47.   World Health Organization. Therapeutics and COVID-19: living guideline.

948   https://www.who.int/publications/i/item/WHO-2019-nCoV-therapeutics-2022.3

949   (2022).

950   48.   Wölfel, R. *et al.* Virological assessment of hospitalized patients with COVID-2019.

951   *Nature 2020 581:7809* **581**, 465–469 (2020).

a) Vaccination coverage = 10%

b) Vaccination coverage = 50%

c) Vaccination coverage = 90%

a) Vaccination coverage = 10%

b) Vaccination coverage = 50%

c) Vaccination coverage = 90%

a) Vaccination coverage = 10%

b) Vaccination coverage = 50%

c) Vaccination coverage = 90%

a) Total infections **b)** Severe cases **c)** Deaths

100 tests/100,000 people/day

500 tests/100,000 people/day