

## Estimating national-level measles case fatality ratios: an updated systematic review and modelling study

Alyssa N. Sbarra<sup>1,2\*</sup>, Jonathan F. Mosser<sup>2,3</sup>, Mark Jit<sup>1</sup>, Allison Portnoy<sup>4</sup>

<sup>1</sup> Department of Infectious Disease Epidemiology, London School of Hygiene and Tropical Medicine

<sup>2</sup> Institute of Health Metrics and Evaluation, University of Washington

<sup>3</sup> Department of Health Metrics Sciences, University of Washington

<sup>4</sup> Center for Health Decision Science, Harvard T.H. Chan School of Public Health

\* Corresponding author

## Abstract

**Background:** To understand current and prevent future measles mortality burden, it is critical to have robust estimates of measles case fatality. Dynamic estimates of measles case fatality ratios (CFR) that are age-, location-, and time-specific are essential to capture variation in underlying population-level factors that contribute systematically to increases or decreases in case fatality, such as vaccination coverage and measles incidence. In this study, we updated estimates of measles CFR by expanding upon previous systematic reviews and implementing a meta-regression model.

**Methods:** We conducted a literature review of all available data on measles case fatality from 1980 through 2019 from low- and middle-income countries and ages and extracted the most granular information available on measles cases and deaths. Using this data and a suite of covariates related to measles CFR, we implemented a Bayesian meta-regression model to produce estimates of measles CFR by location and age from 1990 to 2019.

**Findings:** We identified 245 literature sources that contained information on measles cases and deaths. In 2019, we estimated a mean CFR among community-based settings of 1.32% (95% Uncertainty Interval (UI): 1.28 – 1.36%) and among hospital-based settings a mean of 5.35% (95% UI: 5.08 – 5.64%). In community-based settings, we estimated 2019 CFRs to be 3.03% (95% UI: 2.89 – 3.16%) among under-one year olds, 1.63% (95% UI: 1.58 – 1.68%) among 1 to 4 year olds, 0.84% (95% UI: 0.80 – 0.87%) among 5 to 9 year olds, and 0.67% (95% UI: 0.64 – 0.70%) among 10 to 14 year olds. Between 1990 and 2019, we estimated measles vaccination has averted approximately 49 million deaths due to measles mortality.

**Interpretation:** While CFRs have improved, there are still large heterogeneities across locations and ages. Our updated methodologic framework and estimates can be used to evaluate measles control and vaccination programs to reduce preventable measles mortality burden.

## Research in context

### *Evidence before this study*

Two previous systematic reviews have synthesized individual studies of measles CFR. The first review, by Wolfson et al., was published in 2009. It used 58 community-based studies in 29 countries to provide global estimates of measles CFR. Using descriptive methods, the Wolfson analysis published results suggesting global estimates of CFR with a mean of 3.3%, with a median of 3.9%, and range from 0 – 40.1%. For outbreak investigations, results suggested a median CFR of 5.2% (95% CI: 2.6 – 11.6%). These results were the first to produce figures of measles CFR beyond single country-year studies, reports, and investigations; however, this study only included community-based studies, did not produce estimates for other locations or years, and did not further stratify by other underlying determinants of mortality, such as development status of each country.

The later review by Portnoy and others was published in 2019 and included data from 1980 to 2016 from low- and middle-income countries; studies included reports from both community-based (n=85) and hospital-based (n=39) settings. Following the review, authors used a log-linear prediction model with a select set of covariates, understood to be related to measles CFR (previous vaccination history [first dose MCV coverage used as a proxy], estimated measles attack rate) and indirectly associated with measles CFR (under-5 mortality [U5M], total fertility rate, proportion of population living in urban areas, population density). The authors reported predicted CFR stratified by year, country-development status, under-5 mortality rate, community versus hospital, and for ages under- or over-5 years, as well as by calendar year from 1990 to 2030. Results predicted a mean CFR of 2.2% (95% CI: 0.7 – 4.5%) for years 1990–2015, with stratification for community (CFR: 1.5, 95% CI: 0.5 – 3.1%) and hospital-based studies (CFR: 2.9, 95% CI: 0.9 – 6.0%).

### *Added value of this study*

Our study updates the existing body of evidence to assess measles CFR on a global scale. To synthesize these data along with a suite of covariates demonstrated to be related to measles CFR at the population level according to the published literature, we implemented a Bayesian meta-regression modelling framework, with a flexible spline component to better capture variation in CFR by age, to produce age-, location- and year-specific estimates of measles CFR from 1990 to 2019.

### *Implications of all the available evidence*

This model, along with corresponding estimates, can contribute to a deeper understanding of measles CFR and allow for a more robust assessment of vaccination programmes and other interventions to reduce measles mortality burden.

## Introduction

In 2020, over 60,700 deaths were estimated to be attributable to measles.<sup>1</sup> However, the exact figure cannot be directly measured because there is a lack of reliable data on measles mortality from most high-burden settings. Instead, measles mortality is usually estimated by combining information or estimates of measles incidence and measles case fatality ratios (CFRs).<sup>2</sup> Hence an accurate understanding of measles CFRs over different times and geographies is vital for disease burden estimation. Additionally, a robust understanding of CFRs can help to identify opportunities for health system strengthening and inform assessments of the impact of vaccination programs. Cohort-based and cross-sectional studies and outbreak investigations provide reports of measles CFR in the literature, but are often limited to specific countries, years, and vaccine delivery strategies.

Previous work has reviewed the body of available published data on measles CFR<sup>3</sup>; more recently an additional study<sup>4</sup> has also modelled estimates of measles CFR for low- and middle-income countries (LMICs) among persons aged under and over 5-years old and in both community- and hospital-based settings. These time-varying estimates of measles CFR are critical to understanding evolving patterns of measles mortality across time and location and have been instrumental in understanding acute measles deaths and the impact of various vaccination scenarios.<sup>5</sup> While a major advancement, these previously published estimates do not include CFR data and associated covariates from the most recent years.

Additionally, CFR estimates stratified by broad age categories may obscure important variation within these age groups, particularly for younger children. Both previous systematic review studies have shown that measles CFR is lower in persons aged at least five years compared which children younger than age five. However, there are likely to be critically important age-specific variations in infants and younger children related to maternal antibody presence, immune system maturation and vaccination status, among other factors, that go uncaptured in a composite estimate of CFR among all under-five year olds.<sup>6</sup> Given that measles incidence tends to be highest among young children who are unvaccinated<sup>7</sup>, an accurate understanding of CFR among these ages is critical for understanding measles mortality burden and developing targeted interventions.

To address these previous limitations, in this study, we have expanded the literature search of previous reviews by additionally including data from non-English studies, as well as extending the scope of years to include data published through 2020 (representing cases occurring through 2019), to conduct a full literature review of measles CFR data representing both community and hospital cases in LMICs. Additionally, we developed a Bayesian meta-regression modelling framework to produce location-, year-, and age-specific estimates of measles CFR from 1990 to 2019.

## Methods

### *Literature review and data extraction*

We extended previously published systematic literature reviews on measles CFRs in LMICs to additionally include data through 2020, and non-English studies. To do this, we searched PubMed from January 1, 1980 through December 31, 2020 using the following search string:

```
((((measles[MeSH Terms] OR measles) AND (mortality[MeSH Terms] OR mortality OR "case fatality rate" OR "case fatality ratio" OR "case fatality"))))
```

In addition to the literature search, we added studies from previous systematic reviews<sup>4,8</sup> and the Global Burden of Diseases, Injuries, and Risk Factors study<sup>9</sup> prior to de-duplicating and screening. We included studies if they were included in the previous systematic reviews or, if upon screening, they contained primary data on measles cases and deaths from hospital-, community-, or surveillance-based reports,

including outbreak investigations. We excluded studies if they were not in humans, contained non-original or non-primary data (i.e., reported on outcomes of another study), reported on data from global or regional surveillance, or did not contain relevant information on measles cases and deaths. Additionally, like the previous reviews, we excluded studies: (1) only reporting on measles cases and deaths among restricted populations, such as from communities of internally displaced persons or among persons living with HIV; (2) only reporting on long-term measles mortality, such as from subacute sclerosing panencephalitis; or (3) were from a high-income country as defined by the World Bank country income classification in 2017.

We extracted the following data from each study: number of measles cases, number of measles deaths, study year, age, geographic location, whether the source was from a hospital- or community-based setting, and whether the source was reporting on an outbreak setting. As available, we also extracted location-specific geographic (state, town, or region) and socioeconomic (urban or rural) information, laboratory confirmation of measles cases, and length of time following onset of rash for a death to be considered attributable to measles. For each study, we computed the annual age-specific case-fatality ratios; we included all suspected measles cases and considered all deaths within 30 days of rash onset, unless cases or acute deaths were defined otherwise in each study.

#### *Covariate selection and analysis*

An overview of our entire covariate selection and modelling process can be found in Supplementary Figure 1. We identified measles incidence and age to be critical covariates when assessing measles CFR. We additionally selected covariates based on a systematic review and expert consultation<sup>10</sup> that identified five possible underlying mechanisms contributing to systematic increases or decreases in measles CFR (health system access and care seeking behaviours, health system quality, nutritional status, measles control and epidemiology, and risk of secondary infection) and related population-level indicators with evidence of an association with measles CFR (average household size, educational attainment, first-dose coverage of measles-containing vaccine (MCV1), human immunodeficiency virus (HIV) prevalence, level of health care available, second-dose coverage of measles-containing vaccine (MCV2), stunting prevalence, surrounding conflict, travel time to nearest health care facility, under-five mortality rate, underweight prevalence, vitamin A deficiency prevalence, vitamin A treatment prevalence, and wasting prevalence).

We used estimates of country-specific annual measles incidence that were generated using previously described methods.<sup>11</sup> For the remaining possible covariates, we searched databases of health-related indicators (World Health Organization, United Nations, World Bank, Global Burden of Disease Study) to identify possible covariate sets that could be used to represent each indicator. For the following indicators, we were able to find an appropriate covariate set available for nearly all countries and years from 1980 to 2019: HIV prevalence<sup>9</sup>, MCV1 coverage<sup>12</sup>, under-five mortality rate<sup>13</sup>, vitamin A deficiency prevalence<sup>9</sup>, and wasting prevalence<sup>9</sup>. If a covariate set was not available, we identified a proxy covariate set if an appropriate alternative existed based on expert group review. Proxy covariate sets include the following: gross domestic product (GDP) per capita<sup>13</sup> (for level of health care available), maternal education<sup>9</sup> (for educational attainment), proportion living in an urban setting<sup>13</sup> (for travel time to nearest health facility), total fertility rate<sup>13</sup> (for average household size), and mortality rate due to war and terrorism<sup>9</sup> (for surrounding conflict). For vitamin A treatment, we were unable to identify an appropriate proxy covariate set; therefore, vitamin A treatment was excluded as a covariate in our model.

If country-years were missing in covariate sets, we either computed an interpolated or projected value if there was less than 20% missingness per country for the covariate or used the regional average of covariate values for a country if there was more than 20% missingness. For covariates with less than 20%

missingness per country, we linearly interpolated missing years using values from adjacent available years of covariate values. If missing covariate values were at the beginning or end of the covariate time series, we used an annualized rate of change with exponential weights to compute the projected covariate values either forwards or backwards in time to complete the full time series, such that weights are more representative of years in the time-series that are closer to either the most recent year for forwards projections and most historic year for backwards projections. For wasting prevalence specifically, which was only available from 1990 onwards, we held the 1990 value constant from 1980 to 1990. We assumed that all covariates took their values in 1980 for all years before 1980.

We conducted a two-step data analysis of the covariate sets to determine the strength of relationship and predictive capability of each covariate in describing underlying trends in CFR. To examine correlation of covariates, we tested linear correlation coefficients of each covariate per underlying possible mechanism related to CFR described in the previous review. Covariates were removed if there was a correlation above 0.8 with any other covariate in the mechanistic group; covariates were removed sequentially based on the average highest collinearity between all other covariates. Next, as a second step, we performed a simple linear regression of the remaining covariates per mechanistic group with the measles CFR dataset. Covariates were removed if they had a p-value of  $> 0.3$  in these regression models. The final list of covariates selected for inclusion were: age, a categorical indicator for community versus hospital studies, measles incidence, mortality rate due to war and terrorism, maternal education, GDP per capita, HIV prevalence, MCV1 coverage, total fertility rate, under-5 mortality rate, proportion living in urban settings, vitamin A deficiency, and wasting prevalence; specific details of this process can be found in Supplementary Information Section 1.

Finally, we selected a transformation (log, logit, or untransformed) for each covariate by fitting separate linear regressions with each version of the transformed covariate as a predictor and an outcome of logit CFR. Transformation was selected based on the corresponding model with the lowest Akaike information criterion (AIC) score. Then, to improve model stability, we standardized each transformed covariate by subtracting the mean of the transformed covariate and dividing by the standard deviation.

### *Model fitting*

To reduce bias resulting from inclusion of data without age-specificity, we first fit a model to only data for which there was age granularity representing groupings five-years wide or smaller; data included in this model ranged from persons aged 0 to 34 years old. This model used the age-granular data and transformed and standardized covariate values for each study midpoint year and fit a Bayesian fixed-effects meta-regression model<sup>14</sup> with the outcome variable as the logit CFR; for details on model selection see Supplementary Information Section 2. We computed standard error in logit space per study using the delta method transformation<sup>15</sup> and used these values as weights in the meta-regression. To represent the relationship between logit CFR and age, we used a quadratic spline with 5 knots, with 3 internal knots, placed uniformly based on data density, such that equal proportions of our input data were represented between each knot. Internal knots were placed at ages 0.68, 1.31, and 3.83 years. Next, we split cases and deaths from each input data source reporting age bins wider than 1 year differentially based on estimates of country- and age-specific incidence<sup>11</sup> and the overall relative age pattern of CFR estimated in the first stage model. We then recalculated logit of CFR and standard error per the newly adjusted number of deaths and cases per new granular age group.

Then, in a second stage model, we used the same general model formula previously described, maintaining the spline knot locations identified in the first stage model, and fit our outcome of logit CFR to all data following age-splitting (Supplementary Information Section 3). To ensure the correct direction

of association between each covariate and CFR, we placed priors on each regression coefficient to ensure the correct direction of association was described. We obtained 1000 samples of the regression coefficients from their fitted joint posterior distribution, predicted country- and age-specific measles CFR in LMICs from 1990 to 2019. We assumed CFR varied up to age 34 (the age up until which we had age-specific data) and held CFR constant for older ages.

To understand the effect of modelling changes, the addition of new covariates, and updating our dataset on subsequent estimates of CFR relative to those produced by Portnoy and colleagues<sup>4</sup>, we conducted a decomposition analysis comparing this new modelling framework to that described in this previous work (Supplementary Information Section 4). Additionally, we computed in-sample and five-fold out-of-sample cross validation metrics to assess model performance. We produced mean estimates of CFR at the age-, region-, or year-level by using the case-weighted average of age-, country-, year-specific CFR estimates. We performed all analyses and produced all figures within the R computing environment (version 5.4).

#### *Role of the funding source*

The funders of this study had no role in study design, data collection, data analysis, data interpretation, or the writing of the report.

## Results

From our literature search, we identified 2581 articles for review. Following our inclusion and exclusion criteria, we extracted information on measles cases and deaths from 245 studies (Figure 1). A full list of source-specific extractions can be found in Supplementary Information Table 5. One hundred seventy-five of the sources were from community-based settings, 66 were from hospital-based settings, and four contained observations from both community- and hospital-based settings. One hundred twenty-six studies contained granular information on age (i.e., at least one age group less than or equal to five years wide) and 67 studies only presented information across all ages. Among all sources, 57 overall unique age groups were represented. Eighty-eight sources provided information on laboratory confirmation of cases and 84 sources provided information on a definition for a measles-related death.

There were 44 studies with information on measles cases and deaths before 1980, 119 from 1980–1989, 84 from 1990–1999, 67 from 2000–2009, and 71 from 2010–2019. Seventy-five countries were represented among sources. Among 3,477,901 measles cases represented across all observations, the crude mean CFR was 6.3% and median CFR was 2.2%. Among hospital-based studies, the crude mean CFR was 8.9%, and among community-based studies the crude mean CFR were 5.1%.

In 1990, our model estimated the mean community-based CFR across all locations to be 2.60% (95% Uncertainty Interval (UI): 2.52 – 2.69%), and in 2019, the mean modeled estimate was 1.32% (95% UI: 1.28 – 1.36%). Among hospital-based settings, the mean estimated CFR across all locations in 1990 was 10.13% (95% UI: 9.67 – 10.60%), and in 2019 was 5.35% (95% UI: 5.08 – 5.64%). In all regions, estimated measles CFRs decreased from 1990 to 2019 in both community- and hospital-based settings (Table 1). In the sub-Saharan Africa region, CFRs were estimated to be the highest across all regions in 2019 in both community- and hospital-based settings.

The median country-specific case-weighted CFR estimates as well as the range in CFR estimates by country decreased across the study period (Figure 2). CFR decreases in all estimated LMICs from 1990 to 2019. As mean estimates of CFR have been case-weighted, country- and year-specific means were influenced by the underlying distribution of the ages of cases within that specific country and year; a

relative distribution of the ages of cases is shown in Supplementary Figure 6. Age-standardized CFR estimates, which show that the declining CFR trends of persist after age-standardization, can be found in Supplementary Information Section 5. Country-specific CFR results can be found in Supplementary Information Appendix 2, and validation metrics can be found in Supplementary Tables 8–9.

Additionally, we estimated CFR to be highest among infants and declining as age increased (Figure 3a). This general trend was consistent across regions (Figure 3b) and over time. Among LMICs in 2019 in community-based settings, CFR among under-one year olds was 5.33% (95% UI: 5.06 – 5.59%), among 1 to 4 year olds was 2.80% (95% UI: 2.70 – 2.90%), among 5 to 9 year olds was 1.50% (95% UI: 1.44 – 1.57%), and among 10 to 14 year olds was 0.87% (95% UI: 0.83 – 0.91%). In 2019 across LMICs in community-based settings, we estimated that CFR among under-one year olds was 3.03% (95% UI: 2.89 – 3.16%), 1.63% (95% UI: 1.58 – 1.68%) among 1 to 4 year olds, 0.84% (95% UI: 0.80 – 0.87%) among 5 to 9 year olds, and 0.67% (95% UI: 0.64 – 0.70%) among 10 to 14 year olds.

In addition to our baseline scenario, we predicted results for a scenario in which there had been no vaccine introduction (i.e., MCV1 coverage was 0% in all countries and years, and incidence values also reflect a lack of vaccination). Methods for estimating incidence in a no-vaccination scenario have been described at length elsewhere.<sup>16</sup> CFR estimates by region, year, and community- and hospital-based settings are larger relative to the baseline vaccination scenario (Table 2). As a result of these differences in CFR and measles incidence in both scenarios, we estimated that from years 2000 to 2019 there have been approximately 49 million deaths averted that are attributable to vaccination in these LMICs.

## Discussion

Understanding of measles CFR was previously limited to a former, and now outdated, systematic review among community-based settings which suggested a single metric of CFR across time. Following the initial review, an updated systematic review expanded literature to additionally include CFRs among hospital-based settings, and used a time-varying model to estimate CFR by location and year. While more comprehensive, this updated review did not include age-specificity beyond variation among under and over 5-year-olds, and covariates included were not selected via a transparent, systematic process. To address these shortcomings, our new study: (1) updated the former literature searches; (2) based covariate selection on widespread expert consultation, a literature review, and selection through a statistical process; and (3) accounted for an age distribution in the modelling process.

Our study included 40 new sources on measles CFR from 21 new countries up to year 2019. We statistically tested new covariates representing known relations to measles CFR for inclusion such as vitamin A deficiency prevalence, wasting prevalence, HIV prevalence and mortality rate due to war and terrorism. We estimated CFR in community-based settings 1990 to be 2.60% and have declined by 2019 to 1.32%. We estimated higher CFRs in hospital-based settings relative to community-based settings, an observation noted in previous work.<sup>4</sup>

Additionally, we included information on age variation among literature sources on CFR in our model which allowed us to produce age-specific estimates of measles case fatality. The relationship between age and CFR has been established previously and is a phenomenon influenced by a variety of factors including but not limited to vaccination coverage, maturation of immune response, and other age-related dependencies of other CFR risk factors. We estimated infants to have the highest CFR values. In this age group, the risk for infection with measles is influenced by the persistence of maternal antibodies, which in turn depend on gestational age and underlying maternal immunity rates.<sup>6</sup> On an individual level, the presence of maternal antibodies may also mitigate the severity of measles infection, potentially leading to



lower CFRs. In our analysis, after controlling for study-level covariates, our model suggests that population-level CFR decreases monotonically with age, consistent with previous studies.<sup>17</sup> This may be because increased measles in infants outweighs the protective effects of maternal antibodies, or because the infants who acquire measles are less likely to have substantial titers of measles antibodies, which would result in increased risk of infection. More detailed and robust data collection in these youngest ages will be critical for assessing this relationship further.

This study has several limitations. First, we did not include studies representing special populations, including some that might be particularly vulnerable to both measles infection and also increased case fatality such as refugees and internally displaced persons. These populations should not be neglected in the global measles control agenda; however, at this time there is a lack of data on these subpopulations to accurately assess their current situation. Next, we assumed that the case fatality estimates presented in each study were nationally representative. Additionally, we assumed that the age distribution of measles cases represented in studies without reported age-specificity followed the same relative age distribution of cases estimated nationally in that country and year during our age-splitting process.

Also, we were constrained by the limited number of studies reporting information on both laboratory confirmation of measles cases and a definition of what was considered a death attributable to measles. Therefore, we chose to include all available studies in the analysis regardless of their reporting on case confirmation or death definition to avoid introducing a compositional bias, stemming from differences in study level demographics, in our estimation framework.

Next, we were not able to incorporate uncertainty from our covariates, including measles incidence, and model specification in our modeling framework, and as such, our uncertainty estimates only reflect uncertainty in the CFR modeling process itself without any additional factors. Additionally, for all covariates that passed our statistical analysis checks, we included each in our modeling framework with priors to govern the direction of association estimated by the model. In this process, four covariates (mortality rate due to war and terrorism, wasting prevalence, HIV prevalence, and GDP per capita) no longer contributed significantly to our model, so they were removed. It is important to emphasize that these covariates may still be significantly related to measles CFR; their exclusion was a result of the underlying collinearity of our covariates that suggested little added predictive benefit to their inclusion in the final model. Lastly, we did not examine individual-level relationships between measles CFR and the covariates included in our modeling framework but rather assessed population-level trends for use in population-level modeling. Therefore, the presented associations between covariates and CFR should not be considered causal relationships.

Our study improves upon previous estimates of measles CFR by incorporating new data sources, systematically identifying covariates, and including improved age-specific variation. These estimates and methodology will be essential in future assessment of measles mortality and vaccination programmes.

## Acknowledgements

We thank Natasha Crowcroft, Felicity Cutts, Emily Dansereau, Deepa Gamage, Katy Gaythorpe, Katrina Kretsinger, Kevin McCarthy, Mark Papania, and Niket Thakkar for their participation in a working group of experts and for the Immunization and Vaccines related Implementation Research advisory committee (IVIR-AC) for their valuable comments on multiple versions of this work. A.N.S, J.F.M., and A.P. received funding for this work from the Bill & Melinda Gates Foundation (INV-019383). A.N.S. received additional

support from the National Institutes of Health (F31AI167535). The content of this work is the sole responsibility of the authors and does not represent official views of the National Institutes of Health.

## References

1. Dixon MG, Ferrari M, Antoni S, et al. Progress Toward Regional Measles Elimination - Worldwide, 2000-2020. *MMWR Morb Mortal Wkly Rep* 2021; **70**(45): 1563-9.
2. World Health Organization. Reducing global measles mortality. 2003.
3. Wolfson LJ, Grais RF, Luquero FJ, Birmingham ME, Strebel PM. Estimates of measles case fatality ratios: a comprehensive review of community-based studies. *Int J Epidemiol* 2009; **38**(1): 192-205.
4. Portnoy A, Jit M, Ferrari M, Hanson M, Brenzel L, Verguet S. Estimates of case-fatality ratios of measles in low-income and middle-income countries: a systematic review and modelling analysis. *Lancet Glob Health* 2019; **7**(4): E472-E81.
5. Portnoy A, Hsieh YL, Abbas K, et al. Differential health impact of intervention programs for time-varying disease risk: a measles vaccination modeling study. *BMC Medicine* 2022; **20**.
6. Guerra FM, Crowcroft NS, Friedman L, et al. Waning of measles maternal antibody in infants in measles elimination settings - A systematic literature review. *Vaccine* 2018; **36**(10): 1248-55.
7. Goodson JL, Masresha BG, Wannemuehler K, Uzicanin A, Cochi S. Changing Epidemiology of Measles in Africa. *Journal of Infectious Diseases* 2011; **204**: S205-S14.
8. Wolfson LJ, Grais RF, Luquero FJ, Birmingham ME, Strebel PM. Estimates of measles case fatality ratios: a comprehensive review of community-based studies. *International Journal of Epidemiology* 2009; **38**(1): 192-205.
9. Vos T, Lim SS, Abbafati C, et al. Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *The Lancet* 2020; **396**(10258): 1204-22.
10. Sbarra AN, Jit M, Mosser JF, et al. Population-level risk factors related to measles case fatality: a conceptual framework and systematic review of evidence. *medRxiv* 2022. <https://doi.org/10.1101/2022.09.30.22280585>.
11. Simons E, Ferrari M, Fricks J, et al. Assessment of the 2010 global measles mortality reduction goal: results from a model of surveillance data. *The Lancet* 2012; **379**(9832): 2173-8.
12. WHO. WHO/UNICEF coverage estimates for 1980-2021. 2022. <https://www.who.int/data/gho/data/themes/immunization> (accessed July 15 2022).
13. World Bank. World Development Indicators. 2022. <https://databank.worldbank.org/source/world-development-indicators> (accessed July 15 2022).
14. Zheng P, Barber R, Sorensen RJ, Murray CJ. Trimmed constrained mixed effects models: formulations and algorithms. *Journal of Computational and Graphical Statistics* 2021; **30**(3): 1-13.
15. Levy Y. The delta method and its implementation in R. 2019. <https://www.r-bloggers.com/2019/03/the-delta-method-and-its-implementation-in-r/> (accessed August 2022).
16. Toor J, Echeverria-Londono S, Li X, et al. Lives saved with vaccination for 10 pathogens across 112 countries in a pre-COVID-19 world. *eLife* 2021; **10**: e67635.
17. Aaby P, Martins CL, Garly M, Rodrigues A, Benn CS, Whittle H. The optimal age of measles immunisation in low-income countries: a secondary analysis of the assumptions underlying the current policy. *BMJ Open* 2012; **2**.


## Tables and Figures

	1990		2000		2010		2019	
	Community-based	Hospital-based	Community-based	Hospital-based	Community-based	Hospital-based	Community-based	Hospital-based
North Africa and Middle East	3.33% (3.14 – 3.53%)	12.53% (11.75 – 13.33%)	3.28% (3.08 – 3.49%)	12.32% (11.56 – 13.13%)	2.23% (2.08 – 2.38%)	8.78% (8.18 – 9.42%)	0.83% (0.79 – 0.86%)	3.42% (3.19 – 3.65%)
Sub-Saharan Africa	3.63% (3.50 – 3.75%)	13.63% (13.09 – 14.19%)	3.05% (2.97 – 3.15%)	11.75% (11.30 – 12.23%)	2.10% (2.04 – 2.17%)	8.34% (7.92 – 8.75%)	1.92% (1.86 – 1.97%)	7.67% (7.31 – 8.07%)
Central Europe, Eastern Europe, and Central Asia	0.66% (0.60 – 0.71%)	2.79% (2.52 – 3.09%)	0.28% (0.26 – 0.31%)	1.22% (1.10 – 1.35%)	0.15% (0.14 – 0.16%)	0.64% (0.58 – 0.71%)	0.18% (0.16 – 0.20%)	0.79% (0.70 – 0.89%)
South Asia	3.16% (3.03 – 3.29%)	12.34% (11.72 – 12.99%)	2.23% (2.08 – 2.27%)	8.79% (8.32 – 9.27%)	1.51% (1.44 – 1.57%)	6.18% (5.82 – 6.57%)	0.82% (0.78 – 0.86%)	3.45% (3.24 – 3.69%)
Latin America and Caribbean	1.81% (1.69 – 1.92%)	7.34% (6.76 – 7.94%)	0.87% (0.82 – 0.93%)	3.66% (3.34 – 3.98%)	0.52% (0.48 – 0.56%)	2.21% (1.99 – 2.43%)	0.35% (0.32 – 0.39%)	1.50% (1.34 – 1.67%)
Southeast Asia, East Asia, and Oceania	0.92% (0.88 – 0.97%)	3.83% (3.58 – 4.09%)	0.65% (0.61 – 0.67%)	2.73% (2.54 – 2.93%)	0.40% (0.38 – 0.42%)	1.71% (1.59 – 1.85%)	0.37% (0.34 – 0.39%)	1.56% (1.45 – 1.68%)

**Table 1.** Case-weighted CFR by region, year, and setting (community- versus hospital-based), with 95% confidence interval.


	1990		2000		2010		2019	
	Community-based	Hospital-based	Community-based	Hospital-based	Community-based	Hospital-based	Community-based	Hospital-based
North Africa and Middle East	5.04% (4.66 – 5.48%)	18.26% (17.03 – 19.65%)	4.51% (4.19 – 4.90%)	16.45% (15.39 – 17.67%)	3.52% (3.25 – 3.82%)	13.25% (12.33 – 14.29%)	1.83% (1.68 – 1.99%)	7.19% (6.65 – 7.79%)
Sub-Saharan Africa	5.32% (5.04 – 5.62%)	19.03% (18.30 – 19.87%)	4.15% (3.93 – 4.41%)	15.41% (14.73 – 16.17%)	3.67% (3.46 – 3.92%)	13.79% (13.07 – 14.65%)	2.77% (2.59 – 2.96%)	10.77% (10.16 – 11.51%)
Central Europe, Eastern Europe, and Central Asia	1.97% (1.78 – 2.14%)	7.97% (7.31 – 8.61%)	0.56% (0.50 – 0.63%)	2.39% (2.14 – 2.67%)	0.37% (0.33 – 0.42%)	1.57% (1.39 – 1.77%)	0.33% (0.29 – 0.38%)	1.42% (1.25 – 1.62%)
South Asia	4.50% (4.25 – 4.76%)	16.88% (16.03 – 17.79%)	3.42% (3.20 – 3.65%)	13.21% (12.46 – 14.05%)	2.49% (2.31 – 2.67%)	9.89% (9.25 – 10.63%)	1.61% (1.49 – 1.74%)	6.57% (6.11 – 7.11%)
Latin America and Caribbean	2.75% (2.48 – 3.01%)	10.80% (9.84 – 11.84%)	1.88% (1.68 – 2.07%)	7.58% (6.81 – 8.41%)	1.24% (1.08 – 1.40%)	5.12% (4.50 – 5.75%)	0.88% (0.77 – 1.01%)	3.66% (3.18 – 4.16%)
Southeast Asia, East Asia, and Oceania	1.64% (1.52 – 1.78%)	6.68% (6.21 – 7.21%)	1.16% (1.06 – 1.26%)	4.79% (4.40 – 5.21%)	0.83% (0.75 – 0.91%)	3.45% (3.13 – 3.79%)	0.75% (0.68 – 0.83%)	3.15% (2.86 – 3.47%)

**Table 2.** Case-weighted CFR by region, year, and setting (community- versus hospital-based), with 95% confidence interval, for no-vaccination scenario.




**Figure 1.** PRISMA diagram

The number of studies at each stage of the review is shown for the systematic review.




**Figure 2.** Box plots of country-specific, community-based case fatality ratios (CFRs) by year, with case-weighted LMIC mean CFR value by year (red line).



**Figure 3a.** Age-specific, community-based, case-weighted case fatality ratio (CFR) estimates among 0–14-year-olds, by years 1990, 2000, 2010, and 2019 for low- and middle-income countries.





**Figure 3b.** Age-specific, community-based, case-weighted case fatality ratio (CFR) estimates among 0–14-year-olds, by years 1990, 2000, 2010, and 2019 by region.