Angular reproduction numbers improve estimates of transmissibility

when disease generation times are misspecified or time-varying

Kris V Parag[1,*], Benjamin J Cowling[2] and Ben C Lambert[3]

[1]MRC Centre for Global Infectious Disease Analysis, Imperial College London, London, UK.

[2]WHO Collaborating Centre for Infectious Disease Epidemiology and Control, School of Public Health,

The University of Hong Kong, Hong Kong

[3]Department of Mathematics, College of Engineering, Mathematics and Physical Sciences, University

of Exeter, Exeter, UK.

*For correspondence: k.parag@imperial.ac.uk.

Abstract

We introduce the *angular reproduction number* Ω, which measures time-varying changes in

epidemic transmissibility resulting from variations in both the effective reproduction number R,

and generation time distribution w. Predominant approaches for tracking pathogen spread

either infer R or the epidemic growth rate r. However, R is biased by mismatches between the

assumed and true w, while r is difficult to interpret in terms of the individual-level branching

process underpinning transmission. R and r may also disagree on the relative transmissibility

of epidemics or variants (i.e., $r_A > r_B$ does not imply $R_A > R_B$ for variants A and B). We find that

Ω responds meaningfully to mismatches and time-variations in w while mostly maintaining the

interpretability of R. We prove that $\Omega > 1$ implies $R > 1$ and that Ω agrees with r on the relative

transmissibility of pathogens. Estimating Ω is no more difficult than inferring R, uses existing

software, and requires no generation time measurements. These advantages come at the

expense of selecting one free parameter. We propose Ω as complementary statistic to R and

r that improves transmissibility estimates when w is misspecified or time-varying and better

reflects the impact of interventions, when those interventions concurrently change R and w or

alter the relative risk of co-circulating pathogens.

Keywords: infectious diseases; epidemic models; reproduction numbers; generation times;

growth rates; transmission dynamics.

Introduction

Estimating the rate of spread or transmissibility of an infectious disease is a fundamental and

ongoing challenge in epidemiology [1]. Identifying salient changes in pathogen transmissibility

can contribute important information to policymaking, providing useful warnings of resurgent

epidemics, assessments of the efficacy of interventions and signals about the emergence of

new variants of concern [1–3]. The effective or instantaneous reproduction number, R, and

time-varying growth rate, *r*, are commonly used to characterise pathogen transmissibility. The

former statistic is an estimate of the average number of new infections per active (circulating)

past infection, while the latter describes the exponential rate of new infection accumulation [4].

Although *R* and *r* are important and popular means of tracking the dynamics of epidemics,

they suffer from key limitations that diminish their fidelity and interpretability. Specifically, the

meaningfulness of *R* depends on our ability to measure the generation time distribution of the

infection under study, *w*. This distribution captures the inter-event times among primary and

secondary infections [5] and is convolved with the past infections to define Λ, the time-varying

total infectiousness of the disease. The total infectiousness serves as the denominator when

inferring *R*, which is the ratio of new infections to Λ. We illustrate all key notation in **Figure 1**.

However, infection times and hence *w* are difficult to measure, requiring detailed transmission

chain data from contact tracing or transmission studies [6]. Even if these data are available,

the estimated *w* (and hence Λ) depends on how inter-event times are sampled or interpreted

(e.g., there are forward, backward, intrinsic and realised generation intervals) [7,8].

Figure 1: Definitions of transmissibility metrics. Panel A plots generation time distributions

that define how past infections cause later ones via the probabilities or weights w with support

m. This involves convolving these weights with past infection incidence I. We show in panel B

that if we represent w and I as vectors then the convolution is equivalent to a projection of w

onto the vector of I. Panels C-D illustrate that standard reproduction numbers R implicitly apply

this projection to compute the denominator Λ. This projection and hence Λ is sensitive to the

w, meaning that if the distribution switches between the two from panel A, our estimates of R

become biased (often changes in generation time are difficult to measure). Our new metric Ω

maximises the projection from panel B to reduce sensitivity (practically this involves a window

based on m) leading to a new denominator M in panel C. This maintains the branching process

interpretation of the epidemic in panel D, while improving transmissibility estimate robustness.

Workarounds, such as approximating w by the serial interval distribution [9], which describes

inter-event times between the onset of symptoms, or inferring w from this distribution [10], do

exist but suffer from related problems [6]. Consequently, w and Λ can often be misspecified,

biasing R and likely misrepresenting the true branching process dynamics of epidemics. While

r is more robust to w misspecification (it only depends on the log gradient of the smoothed

infection time series) [4], it lacks the individual-level informativeness and interpretation of R.

Given estimates of r, it is unclear how to derive the proportion of new infections that need to

be suppressed (roughly R^{-1}), herd immunity thresholds (related to $1-R^{-1}$) or the probability of

epidemic elimination and establishment (both linked to R^{-N} for N infections) [11–13]. The only

known means of attaining such information converts r into R using estimates of w [14].

Difficulties in accurately inferring generation times therefore cause practical bottlenecks that

constrain our ability to measure pathogen transmissibility. These problems are worsened as

recent studies have empirically found that generation times also vary substantially with time

(i.e., w is non-stationary) [15]. These variations may correspond to different epidemic phases

[16], emerging variants of concern [17] and coincide with the implementation of interventions

[18]. These are precisely the situations in which we also want to infer R. However, concurrent

changes in R and w are rarely identifiable, and r inextricably groups the effects of w and R on

transmissibility. While high quality, longitudinal contact tracing data [19] can potentially resolve

these identifiability issues, this is an expensive and logistically hard solution. Here we propose

another means of alleviating the above problems and complementing the insights provided by

R and r – the *angular reproduction number*, Ω.

The angular reproduction number defines transmissibility as a ratio of new infections to M, the

root mean square number of past infections over a user-defined window δ. Because it replaces

Λ with *M*, a quantity that does not require knowledge of generation times, Ω is more robust to

the problems of inferring *w*. We demonstrate that Ω is able to measure the overall changes in

transmissibility caused by fluctuations in both *R* and *w*. Moreover, we prove that Ω has similar

threshold properties to *R*, maintains much of its individual-level interpretation and is a useful

metric for communicating transmissibility. This last point follows as we only need to quote Ω

and the known window δ to generalise our estimates of transmissibility to different settings. In

contrast, the meaningfulness of *R* is contingent on the unknown or uncertain *w*. Downstream

studies sometimes use *R* outside of its generation time context [20], while dashboards aiming

at situational awareness commonly quote *R* without *w*, introducing biases and interpretability

problems into how disease spread is communicated [21].

Additionally, we demonstrate how *r* and *R* can easily disagree on relative transmissibility, both

across time and for co-circulating variants. Unmeasured changes in *w* over time can cause *R*

and *r* to vary in opposite directions (one signals an increase in transmissibility and the other a

decrease). Similarly, co-circulating pathogens with different but stationary and known *w*, may

possess contradictory *R* and *r* value rankings i.e., for variants A and B, $r_A > r_B$ does not imply

$R_A > R_B$. These issues are amplified when interventions (which can change *w*, *R* or both [18])

occur, obscuring notions of the relative risk of spread. However, we find $r_A > r_B$ guarantees Ω_A

$> \Omega_B$ and that Ω is consistent with *r* across time even when *w* changes.

Last, while we may also convert *r* into threshold statistics about 1 by using a free parameter

together with a transformation from [14], we show that Ω is more robust to choices of its free

parameter than those statistics, which implicitly make stronger assumptions (**Supplementary**

Information). These robustness and consistency properties of Ω reinforce its usefulness for

tracking and comparing outbreak spread and emerge from its maximum entropy approach to

managing uncertain generation time distributions. We propose Ω as a complementary statistic

that can be integrated with *R* and *r* to present a more comprehensive perspective on epidemic

transmissibility, especially when *w* is poorly specified or varying with time.

Results

Angular reproduction numbers

The epidemic *renewal model* [22] provides a general and flexible representation of disease

transmission. It defines how the incidence of new infections at time *t*, denoted I_t, depends on

the effective or *instantaneous reproduction number*, R_t, and the past incident time series of

infections, $I_1^{t-1} \stackrel{\text{def}}{=} \{I_1, I_2, \dots I_{t-1}\}$. This results in the conditional moment relationship in **Eq. (1)**

[9]. Generally, we use X_a^b to denote the time series $\{X_a, X_{a+1}, \dots, X_{b-1}, X_b\}$ and $\mathbf{E}[X|Y]$ for the

expectation of X over possible epidemic trajectories given known variables Y. Where obvious,

and for convenience, we sometimes drop Y in $\mathbf{E}[X|Y]$, writing $\mathbf{E}[X]$.

118
$$\mathbf{E}[I_t \mid I_1^{t-1}, w_1^m] = R_t \Lambda_t, \qquad \Lambda_t = \sum_{u=1}^{m} w_u I_{t-u}. \quad (1)$$

In this model Λ_t is known as the *total infectiousness* and summarises the weighted influence

of past infections. The set of weights w_u for all u defines the *generation time distribution* of the

infectious disease with $\sum_{u=1}^{m} w_u = 1$, and m as the support of this distribution, which we

assume to be practically finite [14]. When the time series is shorter than m we truncate and

renormalise the w_u. Commonly, the stochasticity around the expectation $R_t \Lambda_t$ is modelled

using either Poisson or negative binomial count distributions [1,12].

Although **Eq. (1)** has successfully been applied to model many diseases including COVID-19,

Ebola virus disease, pandemic influenza and measles, among others, it has one major flaw –

it assumes that the generation time distribution is fixed or stationary and known [9]. If this

assumption holds (we ignore surveillance biases [9,23] until the Discussion), **Eq. (1)** allows

epidemic transmissibility to be summarised by fluctuations of the time-varying R_t parameters.

This follows because the sign of $R_t - 1$ determines if I_t will increase or decline relative to the

total infectiousness Λ_t. This reproduction number can be linked to the *instantaneous epidemic*

growth rate, r_t, using the moment generating function of the generation time distribution [14].

Consequently, from R_t, we obtain temporal information about the rate of pathogen spread and

its mechanism i.e., we learn how many new infections we can expect per circulating infection

because $R_t = \mathbf{E}[I_t]\Lambda_t^{-1}$. As R_t is a threshold parameter, we know that we must block at least

a fraction $1 - R_t^{-1}$ of new infections to suppress epidemic growth ($R_t = 1$ signifies that $r_t =$

0). The time scale over which this suppression is achievable [14] and our ability to detect these

changes in R_t [24] in the first place, however, are determined by the generation times.

Recent works emphasise that the assumption of a known or fixed generation time distribution

is often untenable, with appreciable fluctuations caused by interventions [15,18] and emerging

pathogenic variants [17] or occurring as the epidemic progresses through various stages of

its lifetime [5]. Substantial biases in R_t can result (because its denominator Λ_t is incorrectly

specified [4]), which even impede optimal Bayesian inference algorithms [25]. As R_t is the

predominant metric of transmissibility, contributing key evidence towards infectious disease

policymaking [1], this may potentially obscure situational awareness or misinform intervention

planning. While improved and intensive contact tracing can provide updated generation time
information, this is usually difficult and expensive. We propose a robust alternative.

We redefine Λ_t by recognising it as a dot product between the vectors of generation time
probabilities $\vec{w} \stackrel{\text{def}}{=} w_1^m$ and the past incidence $\vec{I} \stackrel{\text{def}}{=} I_{t-m}^{t-1}$ over the support of the generation time
distribution, m. This gives the left equality of **Eq. (2)** with the Euclidian norm of $\vec{X}$ as $\|\vec{X}\| \stackrel{\text{def}}{=}$
$(\sum_{u=1}^m X_u^2)^{\frac{1}{2}}$ and θ_t as the time-varying angle between $\vec{w}$ and $\vec{I}$. This equality holds for non-
stationary generation times i.e., both $\vec{w}$ and $\vec{I}$ can have elements that change over time. We
illustrate this notation and elements of the subsequent derivation in **Figure 1**. **Eq. (2)** implies
that the count of new infections (for any R_t) is maximised when θ_t is minimised i.e., when the
temporal profile of past infections matches the shape of the generation time distribution.

156
$$\Lambda_t = \|\vec{w}\|\|\vec{I}\| \cos \theta_t, \qquad \mathbf{E}[I_t] = \left(\frac{\|\vec{w}\|}{\|\vec{w}_{\max}\|} R_t \cos \theta_t\right) M_t. \quad (2)$$

We can compute the root mean square of the incidence across the support of the generation
time distribution as $M_t \stackrel{\text{def}}{=} \frac{1}{\sqrt{m}} \|\vec{I}\|$. Under the constraint that $\sum_{u=1}^m w_u = 1$ (if $t - 1 < m$ we
truncate this distribution to sum to 1 – this is an edge effect of the epidemic) then the maximum
possible value of the generation time norm is $\|\vec{w}_{\max}\| = \frac{1}{\sqrt{m}}$. This is achieved by the maximum
entropy generation time distribution of $\vec{w}$, which is uniform (has m entries of $\frac{1}{m}$).

Combining these definitions with **Eq. (1)**, we derive the second expression in **Eq. (2)** for the
expected number of new infections at time t. This may seem an unnecessarily complicated
manipulation of the standard renewal model, but it admits a novel and important insight – we
can separate the influences of the reproduction numbers and the generation time distribution
(together with its changes) on epidemic transmissibility. These multiply M_t, which defines a
new denominator – the root mean square number of past infections (this is also the average
signal power of the past infection time series) – that replaces the total infectiousness Λ_t.

Consequently, we define a new metric in **Eq. (3)**, the *angular reproduction number* Ω_t, which
multiplies R_t by the scaled projection of the generation time distribution, $\frac{\|\vec{w}\|}{\|\vec{w}_{\max}\|} \cos \theta_t$, onto $\vec{I}$,
the past incidence vector (see **Figure 1**). This means that Ω_t is a time-varying ratio between
the expected infection incidence and the past root mean square incidence M_t. We use the

term reproduction number for Ω_t due to its relation to R_t, the similarity of **Eq. (2)** and **Eq. (3)**

and because of its threshold properties, which we explore in the next section.

$$\Omega_t \stackrel{\text{def}}{=} \frac{\|\vec{w}\|}{\|\vec{w}_{\max}\|} R_t \cos \theta_t \implies \Omega_t = \mathbf{E}[I_t] M_t^{-1}. \quad (3)$$

This metric captures all possible variations that impact the ability of the epidemic to transmit.

It responds to both changes in R_t and the generation time distribution. The latter would scale

$\|\vec{w}\|$ and rotate θ_t, which is why we term this angular. The benefit of compactly describing

both types of transmissibility changes does come with a trade-off in interpretability as it may

be harder to intuit the meaning behind $\mathbf{E}[I_t] = \Omega_t M_t$ than the more usual $\mathbf{E}[I_t] = R_t \Lambda_t$.

We argue that this is not the case practically because Λ_t is frequently misspecified [15,26],

obscuring the meaning of R_t. In contrast, M_t does not depend on generation time assumptions

(beyond characterising its support m). We remove structural uncertainty induced by the often

unknown w_u because M_t is a maximum entropy version of Λ_t i.e., $M_t = \max\limits_{\|\vec{w}\| \cos \theta_t} \Lambda_t =$

$\|\vec{w}_{\max}\| \|\vec{I}\|$ subject to $\sum_{u=1}^m w_u = 1$. We also find that $M_t = \Lambda_t$ and hence $\Omega_t = R_t$, when

the past incidence is flat (as then $\Lambda_t = M_t$ and w_u has no effect). This defines the important

and universal equilibrium condition $\Omega_t = R_t = 1$. There is further convergence for branching

process models [27] with timesteps at its fixed generation time, as then trivially $w_1 = 1$.

Relationship to popular transmissibility metrics

Having defined the angular reproduction number above, we explore its properties and show

why it is an interesting and viable measure of transmissibility. We examine an exponentially

growing epidemic with incidence $I_t = I_0 e^{rt}$ and constant growth rate r. This model matches

the dynamics of fundamental *compartmental models* such as the SIR and SEIR (in the limit of

an excess of susceptible individuals) and admits the known relation $gr = (R - 1)$ [28], with

g as the mean generation time. We assume growth occurs over some period of δ and compute

Ω_t as the ratio $\mathbf{E}[I_t] M_t^{-1}$ from **Eq. (3)**. Since this model is deterministic $\mathbf{E}[I_t] = I_t = I_0 e^{rt}$.

We evaluate M_t from its definition above as $\frac{1}{\sqrt{m}} \|\vec{I}\|$ with $\delta = m$ and using the continuous-time

expression for $\|\vec{I}\| = \left(\int_{t-\delta}^t I_s^2 \, ds \right)^{\frac{1}{2}}$. This yields $M_t = \left(\delta^{-1} \int_{t-\delta}^t I_s^2 \, ds \right)^{\frac{1}{2}}$ with $I_s^2 = I_0^2 e^{2rs}$ from

the exponential incidence equation and evaluates to $\sqrt{2\delta r I_0 (e^{2rt} - e^{2r(t-\delta)})}$. Substituting

this into $\Omega_t = (I_0 e^{rt}) M_t^{-1}$ results in the left relation in **Eq. (4)**.

201
$$\Omega_t^2 = \frac{2\delta r}{1 - e^{-2\delta r}} \geq 1, \qquad \Omega_t^2 = \frac{2\delta g^{-1}(R - 1)}{1 - e^{-2\delta g^{-1}(R-1)}}. \quad (4)$$

Several important points follow. First, as $x \geq 1 - e^{-x}$ for every $x \geq 0$, then $\Omega_t - 1$ and r are

positive too (an analogous argument proves the negative case). Second, we substitute for r

using the compartmental R-r relationship $gr = (R - 1)$ to get the right-side relation of **Eq. (4)**.

Applying L' Hopital's rule we find $\lim_{R \to 1} \Omega_t = 1$. We hence confirm the threshold behaviour of Ω_t

i.e., the sign of $\Omega_t - 1$ and $R_t - 1$ are always consistent (for all values of $\delta > 0$).

Third, we see that constant growth rates imply constant angular reproduction numbers. The

converse is also true, and we may input time-varying growth rates, r_t, into **Eq. (4)** to estimate

Ω_t. These properties hold for any δ, which is now a piecewise-constant time window. We plot

the ramifications of **Eq. (4)** in **Figure 2**. Further, in **Table 1** we summarise how Ω_t relates to

predominant R_t and r_t metrics. We explore some properties in this table in later sections (in

addition to reinforcing our analyses with stochastic models) and demonstrate that relationships

among r_t, R_t and Ω_t have important consequences when comparing outbreaks subject to

interventions and variations in generation times.

Metric property	Growth r	Effective R	Angular Ω
Definition of transmissibility	$r_t \stackrel{\text{def}}{=} \frac{d \log \mathbf{E}[I_t]}{dt}$	$R_t \stackrel{\text{def}}{=} \frac{\mathbf{E}[I_t]}{\Lambda_t}$	$\Omega_t \stackrel{\text{def}}{=} \frac{\mathbf{E}[I_t]}{M_t}$
Pathogen spread threshold	$r_t > 0$	$R_t > 1$	$\Omega_t > 1$
Biased by generation time $\vec{w}$ assumed, given curve I_1^t	Insensitive to the assumed $\vec{w}$	Biased when $\vec{w}$ is misspecified	Signals changes in $\vec{w}$ and R_t
Ranking risk of outbreaks or variants by spreading rate	$r_A > r_B \Rightarrow$ variant A spreads faster	$r_A > r_B \nRightarrow R_A > R_B$ (inconsistent)	$r_A > r_B \Rightarrow \Omega_A > \Omega_B$ (consistent)
Short-term predictive power	Negligible differences among metrics in prediction quality		
Non-dimensional metric	No, inverse time	Yes, both have no units, scalable	
Individual-level interpretability	Not obvious	New infections per circulating ones	

Computability if $\vec{w}$ unknown	Yes (smooth I_1^t)	Not possible	Yes, for any δ

Table 1: Summary of transmissibility metrics. We list important relationships among the instantaneous growth rate (r), the instantaneous or effective reproduction number (R) and the angular reproduction number (Ω) and assess their value as measures of transmissibility.

Note that we may also invert the relationship in **Eq. (4)** to estimate r_t from Ω_t (see Methods for details). This involves solving **Eq. (5)**, where $W_k(x)$ is the Lambert W function with index $k \in [0, -1]$ (this range results from the indicator $\mathbf{1}(y)$) [29].

$$\frac{d \log I_t}{dt} = r_t = 2\delta^{-1}\left(\Omega_t^2 + W_{-\mathbf{1}(\Omega<1)}\left(-\Omega_t^2 e^{-\Omega_t^2}\right)\right). \quad (5)$$

A central implication of **Eq. (4)** and **Eq. (5)** is that we can infer angular reproduction numbers directly from growth rates or vice versa, without requiring knowledge of the generation times.

We further comment on connections between angular and effective reproduction numbers using a deterministic *branching process* model, which is also foundational in epidemiology. We again focus on growth, which is geometric as this is a discrete-time process with time steps scaled in multiples of the mean generation time g. Here incidence is $I_t = R^t$ and $\Omega_t = \mathbf{E}[I_t]M_t^{-1} = R^t\left(\delta^{-1}\sum_{s=t-\delta}^{t-1} R^{2s}\right)^{-\frac{1}{2}}$, with window δ in units of g. If $\delta = 1$ we recover $\Omega_t = R$. If $R = 1$, then $\Omega_t = R$ for all δ. For growing epidemics, as δ increases, $\Omega_t > R$ because we reference present incidence to smaller past infections (or denominators). The opposite occurs if the epidemic declines. This may seem undesirable, but we argue that Ω_t improves overall practical transmissibility measurement because g will likely be misspecified or vary with time.

Any g mismatches bias R, limiting its interpretation, meaningfulness and making comparisons among outbreaks or pathogenic variants difficult, because we cannot be certain that our denominators correspond. This is particularly problematic when estimates of R obtained from a modelling study are incorporated as parameters into downstream studies without accounting for the generation time context on which those estimates depend. However, by additionally communicating Ω and δ, we are sure that denominators match and that we properly include the influences of any g mismatches. Choosing δ is also no worse (and more explicit) than equivalent window assumptions made when inferring R and r [4,30] In the **Supplementary Information** we perform analyses of window choices for Ω and other threshold metrics.

Figure 2: Relationships among transmissibility metrics. Panel A and B show how growth rates (r) and reproduction numbers (R) have diverse functional relationships (see [14,30]) for SEIR models with an excess of susceptible individuals and branching processes. Coloured lines indicate R at different mean generation times (g). Black lines highlight a single functional relationship between angular reproduction numbers Ω and r at all g, using a window δ of 20d. Panel C shows that while Ω varies with choice of δ (increasing from blue to red and computed from **Eq. (4)**), we have a bijective relationship with r. Panel D indicates that R and r can signify inverted changes e.g., an NPI reducing R and g may increase r, raising questions about impact (see [15,18]). Here Ω converts r into a consistent transmissibility metric (also from **Eq. (4)**).

Last, we illustrate how Ω_t relates to other key indicators of epidemic dynamics such as herd

immunity and elimination probabilities. As our derivation replaces **Eq. (1)** with $\mathbf{E}[I_t|\,I_1^{t-1},\delta]=$

$\Omega_t M_t$ for the same observed incidence, these indicators are also readily obtained. Assuming

Poisson noise, the elimination probability $\prod_{s=t}^{\infty}\mathbf{P}[I_t=0|\,I_1^{t-1},R_1^{t-1}]=e^{-\sum_{s=t}^{\infty}\Lambda_t R_t}$ is replaced

by $e^{-\sum_{s=t}^{\infty}M_t\Omega_t}$, and has analogous properties [31]. Herd immunity, which traditionally occurs

when a fraction $1-R^{-1}$ of the population is immune is approximated by $1-\Omega^{-1}$ (since both

metrics possess the same threshold behaviour) [11]. In a subsequent section we demonstrate

that one-step-ahead incidence predictions from both approaches are also comparable.

Responding to variations in generation time distributions

We demonstrate the practical benefits of Ω_t using simulated epidemics with non-stationary or

time-varying generation time distributions. Such changes lead to misspecification of Λ_t in **Eq.**

(1), making estimates of the effective reproduction number R_t, denoted $\hat{R}_t$, a poor reflection

of the true underlying R_t. In contrast variations in the estimated $\widehat{\Omega}_t$ are a feature (see **Eq. (3)**)

and not a bug (for some chosen δ we control M_t, which is not misspecified). We simulate

epidemics with Ebola virus or COVID-19 generation times from [32,33] using renewal models

with Poisson noise [9]. We estimate both the time-varying R_t and Ω_t using *EpiFilter* [25], which

applies Bayesian algorithms that minimise mean square estimation error.

Inferring Ω_t from incident infections, I_1^t, requires only that we replace the input Λ_t with M_t in

the estimation function and that we choose a window δ for computing M_t. We provide software

for general estimation of Ω_t and code for reproducing this and all other analyses in this paper

at https://github.com/kpzoo/Omega. We heuristically set $\delta\approx 2g_0$ as our window with g_0 as

the original mean generation time of each disease from [32,33]. We find (numerically) that this

δ ensures $\sum_{u=0}^{\delta}w_u\geq 0.86$ over many possible gamma distributed generation times i.e., it is

long enough to cover most of the likely probability mass of unknown changes to the generation

time distributions, which cause time-varying means g_t. In general, we find that an overly small

δ tends to neglect important dynamics, while too large a δ induces edge effects. The Methods

and **Supplementary Information** provide for more information on choosing δ.

Our results are plotted in **Figure 3**. We show that $\widehat{\Omega}_t$ responds as expected to both changes

in the true R_t and w_1^m, subject to the limits on what can be inferred [24]. In **Figure 3** we achieve

changes in w_1^m by altering the mean generation time g_t by ratios that are similar in size to

those reported from empirical data [15]. In contrast, we observe that $\hat{R}_t$ provides incorrect and

overconfident transmissibility estimates, which emerge because its temporal fluctuations also

have to encode structural differences due to the misspecification of w_1^m. These can strongly

mislead our interpretation and understanding of the risk posed by a pathogen.

Figure 3: Estimating transmissibility under temporal variations in generation times. We simulate epidemic incidence curves (black) using generation time distributions of Ebola virus disease (EVD) [32] and COVID-19 [33] in panels A and B. The means of these distributions (g) vary over time (grey piecewise, starting from original mean g_0), but we fix their variance at their original values. We find substantial bias in R estimated from the initial EVD and COVID-19 generation times (red with 95% credible intervals, true value in black). These estimates try to compensate for generation time mismatches and changes in an uncontrolled manner that obscures interpretation. However, Ω responds as we expect (blue with 95% credible intervals, window δ, true value in black) and we infer change-points due to both R and g fluctuations (subject to bounds induced by noise i.e., at low incidence inference is more difficult [24]). Our estimates derive from EpiFilter [25] with default settings and we truncate time series to start

We can derive alternative threshold statistics that relate to r_t and do not explicitly depend on

the generation time by applying monotonic transformations from [14]. In theory these should

have comparable behaviour around the critical point of 1 to both R_t and Ω_t. We investigate

these statistics in the **Supplementary Information**, computing them across the simulations

of **Figure 3**. We find that they require stronger assumptions than Ω_t (i.e., they fix distributional

formulae for generation times), possess at least as many free parameters as Ω_t and are less

robust to changes in those parameters (often strongly over-estimating transmissibility), than

Ω_t is to fluctuations in δ. This confirms that angular reproduction numbers can complement

standard metrics, improving transmissibility estimates when generation times are changing,

or unknown and forming part of a more comprehensive suite of outbreak diagnostics.

Ranking epidemics or variants by transmissibility

Misspecification of generation time distributions, and corresponding misestimation of R as in

Figure 3, also plays a crucial role when assessing the relative transmissibility of pathogens,

variants of concern or even outbreaks (where we may want to contrast the spread of contagion

among key demographic or spatial groups). As shown in **Figure 2**, these variations can mean

that increases in the growth rate r_t actually signify decreases in the effective reproduction

number R_t or that a pathogen with a larger r_t can have a smaller R_t. Here we illustrate that

these issues can persist even if the generation time distributions of pathogens are correctly

specified and remain static, obscuring our understanding of relative transmission risk.

In **Figure 4** we simulate epidemics under two hypothetical variants of two pathogens. We use

EVD and COVID-19 generation time distributions from [32,33] to define our respective base

variants. For both pathogens we specify the other variant by reducing the mean generation of

each base but fixing the variance of the generation times. Reductions of this type are plausible

and have been measured for COVID-19 variants [17]. All w_1^m distributions are stationary and

known in this analysis. We discover that changes in R_t alone can initiate inversions in the

relative growth rate of different variants or epidemics. As far as we can tell, this phenomenon

has not been explicitly investigated. Given that interventions can change R_t in isolation or in

combination with w_1^m [15,18], this effect has the potential to be widespread. We determine the

mathematical conditions for this inversion in the **Supplementary Information**.

Interestingly, the angular reproduction numbers of **Figure 4** do preserve an ordering that is
consistent with growth rates, while maintaining the interpretability (e.g., threshold properties)
of reproduction numbers. Hence, we argue that Ω_t blends advantages from both R_t and r_t [4]
and serves as a useful outbreak analytic for understanding and conveying the relative risk of
spread of differing pathogens or pathogen strains, or of spread among different spatial and
demographic groups. Recent studies have only begun to disentangle component drivers of
transmission, including the differing effects that interventions can introduce (e.g., by defining
the strength and speed of control measures [34]) and the diverse properties of antigenic
variants [17]. We believe that Ω can play a distinctive role in accelerating these investigations.

340
Figure 4: Comparing transmissibility across outbreaks, variants or even diseases. We
simulate epidemics of variant 1 in blue (with estimates of metrics also in blue) under standard
generation time distributions of Ebola virus disease (EVD) [32] and COVID-19 [33] in panels
A and B. In red (with estimates also in red) we overlay simulations in which the generation
time of these diseases is 40% and 50% shorter (than the blue epidemics), which may indicate
a new co-circulating variant 2 or another epidemic with different properties (e.g., in a higher

risk group). We demonstrate (for the first time to our knowledge) that changes in R due to an
intervention (or release of one) may invert the relative growth rates (r) of the epidemics (see
Supplementary Information for mathematical intuition for this inversion). The mismatches in
the R-r rankings alter perceptions of relative risk, making transmissibility comparisons difficult.
However, Ω classifies the risk of these epidemics in line with their realised growth rates, while
still offering the individual-level interpretability of a reproduction number. True values are in
black and all estimates (with 95% credible intervals) are outputs from EpiFilter [25] with default
settings. We truncate the time series to start from δ to remove any edge effects.

Reproduction numbers for explanation or prediction?

We highlight an important but underappreciated subtlety when inferring the transmissibility of
epidemics – that the value of accurately estimating R, r and Ω largely depends on if our aim
is to explain or predict [35] the dynamics of epidemics. The above analyses have focussed on
characterising transmissibility to explain mechanisms of spread and design interventions. For
these problems, misestimation of parameters, such as R, can bias our assessment of outbreak
risk and hence misinform the implementation of control measures. An important concurrent
problem aims to predict the likely incidence of new infections from these estimates. This
involves projecting the epidemic dynamics forward in time to infer upcoming infection patterns.

Here we present evidence that the solution of this problem, at least over short projection time
horizons, is robust to misspecification of generation times provided both the incorrect estimate
and the misspecified denominator are used in conjunction. We repeat the analyses of **Figure
3** for 200 replicate epidemics and apply EpiFilter [25] to obtain the one-step-ahead predictive
distributions $\mathbf{P}(I_t | I_1^{t-1})$ for every t. We compute the predicted mean square error (PMSE)
and the accumulated predictive error (APE). These scores, which we denote as $D(I_t | I_1^{t-1})$,
average square errors between mean predictions and true incidence and sum log probabilities
of observing the true incidence from the predicted distribution respectively [36,37]. We plot the
distributions of scores over replicates and illustrate individual predictions in **Figure 5**.

We find only negligible differences among the one-step-ahead predictive accuracies of the R
estimated given knowledge of the changing generation times ($R|w$), the R estimated assuming
an unchanged (and hence wrongly specified) w and our inferred Ω. As APE and PMSE also
measure model suitability, their similarity across the three estimates demonstrate that, if the
problem of prediction is of interest, then incorrect generation time choices are not important
as long as the erroneous denominator (Λ_t) and estimate (R_t) are used together. If this estimate
is however used outside of the context of its denominator (e.g., if it is simply input into other
studies), then inaccurate projections will occur (in addition to poor estimates). As multi-step-

ahead predictions can be composed from iterated one-step-ahead ones [38], we conjecture

that subtleties between prediction and explanation are likely to also apply on longer horizons.

Figure 5: One-step-ahead prediction accuracy and model mismatch. We simulate 200

replicates of the epidemics from **Figure 3**, which involve non-stationary changes to EVD and

COVID-19 generation times. We use estimates of effective, R, and angular, Ω, reproduction

numbers to produce successive one-step-ahead predictions and assess their accuracy to the

simulated (true) incidence. Panels A-D provide a representative example of a single simulated

epidemic (true incidence shown as black dots) and the R and Ω one-step ahead predictions

(red and blue respectively with 95% credible intervals). In panels E-F we formally compute

accuracy using distance metrics, D, based on accumulated prediction errors (APE, dashed)

and prediction mean square errors (PMSE, solid) for all 200 replicates from R, Ω and R given

knowledge of the generation time changes i.e., $R|w$. We obtain distributions of D by applying

kernel smoothing. We find negligible differences in predictive power from all approaches.

Empirical example: COVID-19 in mainland China

We complete our analysis by illustrating the practical usability of Ω on an empirical case study
where generation time changes are known to have occurred. In [15], the dynamics of COVID-
19 in mainland China are tracked across January and February 2020. Transmission pair data
indicated that the serial interval of COVID-19 shortened across this period leading to biases
in the inferred R if updated serial intervals are not used. Here serial intervals, which measure
the lag between the symptom onset times of an infector and infectee are used as a proxy for
the generation time. **Figure 6** presents our main results. We find Ω (blue), which requires no
serial interval information, behaves similarly to the R (red) inferred from the time-changing w.
Both metrics appear less biased than estimates of R (green) that assume a fixed serial interval.
This is largely consistent with the original investigation in [15].

406
Figure 6: COVID-19 transmissibility in China under non-stationary generation times. We
analyse COVID-19 data from [15], which spans 9th January 2020 to 13 February 2020 and is
known to feature a serial interval distribution that shortened in mean substantially from 7.8d
to 2.6d (change times are shown as grey vertical lines). We assume that the serial interval

approximates the generation time well and replicate the analysis from Figure 2 of [15]. In panel
A, we compare estimates (green) of effective reproduction numbers, R, using fixed generation
time distributions inferred in [15] (specified by their means g) against those of our angular
reproduction number Ω (blue). We use EpiFilter [25] to obtain all estimates (means shown
with 95% credible intervals) and find relative trends similar to those in Figure 2 of [15]. In panel
B we plot the incidence (black) and the denominators we use to compute an R that does
account for the generation time changes (Λ, red) and for Ω (M, blue). This R uses the different
distributions inferred at the grey vertical change times (their means are in panel B and are
also the fixed distributions of panel A in sequence). We plot these R and Ω estimates in panel
C. In panel D we show the growth rates that are inferred from the R and Ω estimates of C (red
and blue respectively) against that obtained from taking the smoothed log derivative (black).

We see that Ω provides a lower assessment of the initial transmissibility as compared to the
R that is best informed by the changing w but that both agree in general and in particular at
the important threshold between super- and subcritical spread. Interestingly, Ω indicates no
sharp changes at the w change-times. This follows because the incidence is too small for
those changes to shape overall transmissibility and matches the gradual w changes originally
inferred in [15]. The distributions used in **Figure 6** provide a piecewise approximation to these
variations. We also compare r estimates derived from R (red, from [14]), Ω (blue, from **Eq. (5)**)
and the empirical log gradient of smoothed incidence (black, $\frac{d \log S[I_t]}{dt}$ [4]). We find that the r
from Ω agrees more closely with the empirical growth rate than the r from R, which somewhat
by design shows jumps at the w change-points. While this analysis is not meant as a detailed
study of COVID-19 in China, it does demonstrate the practical usefulness of Ω.

Discussion

Quantifying the time-varying transmissibility of a pathogen remains an enduring challenge in
infectious disease epidemiology. Changes in transmissibility may signify shifts in the dynamics
of an epidemic of relevance to both preparedness and policymaking. While this challenge has
been longstanding, the statistics that we use to summarise transmissibility have evolved from
dispersibility [39] and incidence to prevalence ratios [40] to cohort [41] and instantaneous [22]
reproduction numbers. While the last, which we have denoted R, has become the predominant
metric of transmissibility, all of these proposed statistics ultimately involve a ratio between new
infections and a measure of active infections (i.e., the denominator). Deciding on appropriate
denominators necessitates some notion (implicit or explicit) of a generation time [42].

Difficulties in characterising these generation times and their changes substantially bias [6]

estimates of transmissibility and have motivated recent works to propose the instantaneous

growth rate, r, as a more reliable approach for inferring pathogen spread [20]. However, on its

own, r is insufficient to resolve many of the transmission questions that R can answer and its

computation may employ smoothing assumptions that are in some instances equivalent to the

generation time ones behind R [4]. We formulated the novel angular reproduction number Ω,

to merge some advantages from both R and r and to contribute to a more comprehensive view

of transmissibility. By applying basic vector algebra (**Eqs. 1-3**), we encoded both changes to

R and the generation time distribution, w, into a single time-varying metric, deriving Ω.

We found that Ω maintains the threshold properties and individual-level interpretability of R

but responds to variations in w, in a manner consistent with r (**Figure 2**). Moreover, Ω indicates

variations in transmissibility caused by R and w without requiring measurement of generation

times (**Figure 3**). This is a consequence of its denominator, which is the root mean square of

infections over a user-specified window δ that is relatively simple to tune (see Methods). We

can interpret $\Omega = a > 1$ as indicating that infections across δ need to be reduced by a^{-1}. This

reduces mean and root mean square infections by a^{-1} and causes Ω to equal 1. Further, Ω

circumvents identifiability issues surrounding the joint inference or R and w [43] by refocussing

on estimating the net changes produced by both. This improves our ability to *explain* the shifts

in transmissibility underpinning observed epidemic dynamics and means Ω is essentially a

reproduction number that provides individual-level interpretation of growth rates (**Eqs. 4-5**).

The benefits of this r-Ω correspondence are twofold. First, as interventions may alter R, w or

R and w concurrently [15,18] situations can arise where r and R disagree across time on both

the drivers and magnitude of transmissibility. While it may seem possible to minimise this issue

by constructing alternative threshold statistics by directly combining r with assumed generation

time structures, we find these statistics often exhibit worse performance and larger bias than

Ω (**Supplementary Information**). Second, this disagreement can also occur when comparing

pathogenic variants or epidemics (e.g., from diverse spatial or sociodemographic groups) with

different but known and unchanging w. This study appears to be among the earliest to highlight

these discrepancies, which can occur in multiple settings (see **Supplementary Information**).

Realistic transmission landscapes possess all of the above complexities, meaning that relying

solely on conventional measures of relative transmissibility can lead to contradictions.

We found that Ω consistently orders epidemics by growth rate while capturing notions of the

average new infections per past infection (**Figure 4**). This suggests Ω blends advantages from

R and r, with clearer assumptions (choice of window δ). However, Ω offers no advantage if we

want to *predict* epidemic dynamics (see [35] for more on prediction-explanation distinctions).

For this problem even an *R* inferred using a misspecified denominator performs equally well

(**Figure 5**). This follows as only the product of any reproduction number and its denominator

matter when determining the next incidence value. Iterations of this product underpin multi-

step ahead predictions [38]. This may explain why autoregressive models, which ignore some

characteristics of *w*, can serve as useful predictive models [44]. Other instances where Ω will

not improve analysis are at times earlier than δ (due to edge effects [9]) and in periods of near

zero incidence (there is no information to infer *R* either [24]). We summarised and compared

key properties of *R*, *r* and Ω in **Table 1**.

There are several limitations to our study. First, we only examined biases inherent to *R* due to

the difficulty of measuring the generation time accurately and across time. While this is a major

limitation of existing transmissibility metrics [15], practical surveillance data are also subject

to under-reporting and delays, which can severely diminish the quality of any transmissibility

estimates [23,43,45]. While Ω ameliorates issues due to generation time mismatch, it is as

susceptible as *R* and *r* to surveillance biases and corrective algorithms (e.g., deconvolution

methods [46]) should be applied before inferring Ω. Second, our analysis depends on renewal

and compartmental epidemic models [22]. These assume random mixing and cannot account

for realistic contact patterns. Despite this key structural uncertainty, there is evidence that well-

mixed and network models are comparable when estimating transmissibility [47].

Although the above limitations can, in some instances, reduce the added value of improving

the statistics summarising transmissibility, we believe that Ω will be of practical and theoretical

benefit, offering complementary insights to *R* and *r* and forming part of a more comprehensive

epidemic analytic toolkit. Its similarity in formulation to *R* means it is as easy to compute using

existing software and therefore can be deployed on dashboards and updated in real time to

improve situational awareness. Further, Ω improves comparison and communication of the

relative risks of circulating variants or epidemics among diverse groups, avoiding *R-r*

contradictions provided the known parameter, δ, is fixed. This supplements *R*, which is hard

to contextualise [20] when *w* is misspecified or varying and hence compare across groups, as

each group may have distinct and correspondingly poorly specified denominators. Last, Ω can

help probe analytical questions about how changes in *R* and *w* interact because it presents a

common framework for testing how variations in either influence overall transmissibility.

Methods

Inferring angular reproduction numbers across time

We outline how to estimate Ω_t given a time series of incident infections I_1^T, with T defining the

present or last available data timepoint i.e., $1 \le t \le T$. Because Ω_t simply replaces the total

infectiousness Λ_t, used for computing R_t, with the root mean square of the new infection time

series (see **Figure 1**), M_t, we can obtain Ω_t from standard R_t estimation packages with minor

changes. This requires evaluating M_t over some user-defined, backward sliding window of

size δ. Under a Poisson (Pois) renewal model this follows as in **Eq. (6)** for timepoint t.

516
$$\mathbf{P}(I_t \mid I_1^{t-1}, \delta) \equiv \mathrm{Pois}(\Omega_t M_t), \qquad M_t = \left(\frac{1}{\delta} \sum_{u=t-\delta}^{t-1} I_u^2 \right)^{\frac{1}{2}}. \quad (6)$$

The choice of δ is mostly arbitrary but should be sufficiently long to capture most of the likely

probability mass of the unknown generation time but not overly long since it induces an edge

effect (similar to the windows in [9,37]). We found a suitable heuristic to be twice or thrice the

initial expected mean generation time (g_0). We can then input M_t and I_t into packages such

as EpiEstim [9] or EpiFilter [25] to estimate Ω_t with 95% credible intervals.

Due to the similarity between computing R_t and Ω_t we only specify the latter but highlight that

replacing M_t with Λ_t yields the expressions for evaluating any equivalent quantities from R_t.

The only difference relates to how the growth rates r_t are computed. We estimate r_t from R_t

by applying the generation time, $\vec{w}$, based transformation from [14]. For a correctly specified

$\vec{w}$ this gives the same result as the smoothed derivative of the incidence curve [4]. We derive

r_t from Ω_t using **Eq. (5)**, which follows from rearranging **Eq. (4)** into $\left(2\delta r_t - \Omega_t^2 \right) e^{2\delta r_t - \Omega_t^2} =$

$-\Omega_t^2 e^{-\Omega_t^2}$. This expression then admits Lambert W function solutions. In all estimates of r_t we

propagate uncertainty from the posterior distributions (see below) over R_t or Ω_t.

We applied EpiFilter in this study due to its improved extraction of information from I_1^T. This

method assumes a random walk state model for our transmissibility metric as in **Eq. (7)** with

ϵ_{t-1} as a normally distributed (Norm) noise term and η as a free parameter (default 0.1).

533
$$\Omega_t = \Omega_{t-1} + \left(\eta \sqrt{\Omega_{t-1}} \right) \epsilon_{t-1}, \qquad \mathbf{P}(\epsilon_{t-1}) \equiv \mathrm{Norm}(0, 1). \quad (7)$$

The EpiFilter approach utilises Bayesian smoothing algorithms incorporating the models of

Eq. (6)-(7) and outputs the complete posterior distribution $\mathbf{P}(\Omega_t \mid I_1^T, \delta)$ with T as the complete

length of all available data (i.e., $1 \le t \le T$). We compute our mean estimates $\widehat{\Omega}_t$ and 95%

credible intervals from this posterior distribution and these underlie our plots in **Figures 3-4**.

EpiFilter also outputs the one-step-ahead predictive distributions $\mathbf{P}(I_t \mid I_1^{t-1}, \delta)$, which we use

in **Figure 5**. There we quantify predictive accuracy using the predicted mean square error

PMSE and the accumulated prediction error APE, defined as in **Eq. (8)** [36,37] with $\hat{I}_t$ as the

posterior mean estimate from $\mathbf{P}(I_t \mid I_1^{t-1}, \delta)$ and I_t^* as the true simulated incidence. These are

computed with $\mathbf{P}(\Omega_{t-1} \mid I_1^{t-1}, \delta)$ and not $\mathbf{P}(\Omega_t \mid I_1^T, \delta)$, ensuring no future information is used.

543
$$\text{PMSE} = \frac{1}{T-\delta} \sum_{t=\delta+1}^{T} \left(I_t^* - \hat{I}_t \right)^2, \quad \text{APE} = \sum_{t=\delta+1}^{T} -\log \mathbf{P}(I_t = I_t^* \mid I_1^{t-1}, \delta). \quad (8)$$

We collectively refer to these as distance metrics $D(I_t \mid I_1^{t-1})$ and construct their distributions,

$\mathbf{P}(D)$, over many replicates of simulated epidemics. Last, we use $\mathbf{P}(\Omega_t \mid I_1^T, \delta)$ to compute the

posterior distribution of the growth rate $\mathbf{P}(r_t \mid I_1^T, \delta)$ and hence its estimates as in **Eq. (5)**. More

details on the EpiFilter algorithms are available at [25,31,48]. We supply open-source code to

reproduce all analyses at https://github.com/kpzoo/Omega as well as functions in MATLAB

and R to allow users to estimate Ω_t from their own data.

Bibliography

1. Anderson R, Donnelly C, Hollingsworth D, Keeling M, Vegvari C, Baggaley R.
Reproduction number (R) and growth rate (r) of the COVID-19 epidemic in the UK:
methods of. The Royal Society. 2020;

2. Li Y, Campbell H, Kulkarni D, Harpur A, Nundy M, Wang X, et al. The temporal
association of introducing and lifting non-pharmaceutical interventions with the time-
varying reproduction number (R) of SARS-CoV-2: a modelling study across 131
countries. Lancet Infect Dis. 2021;21: 193–202. doi:10.1016/S1473-3099(20)30785-4

3. Volz E, Mishra S, Chand M, Barrett JC, The COVID-19 Genomics UK (COG-UK)
consortium, Johnson R, et al. Assessing transmissibility of SARS-CoV-2 lineage
B.1.1.7 in England. Nature. 2021; doi:10.1038/s41586-021-03470-x

4. Parag KV, Thompson RN, Donnelly CA. Are epidemic growth rates more informative
than reproduction numbers? J Royal Statistical Soc A. 2022; doi:10.1111/rssa.12867

5. Svensson A. A note on generation times in epidemic models. Math Biosci. 2007;208:
300–311. doi:10.1016/j.mbs.2006.10.010

6. Britton T, Scalia Tomba G. Estimation in emerging epidemics: biases and remedies. J
R Soc Interface. 2019;16: 20180670. doi:10.1098/rsif.2018.0670

7. Champredon D, Dushoff J. Intrinsic and realized generation intervals in infectious-
disease transmission. Proc Biol Sci. 2015;282: 20152026. doi:10.1098/rspb.2015.2026

8. Nishiura H. Time variations in the generation time of an infectious disease: implications
for sampling to appropriately quantify transmission potential. Math Biosci Eng. 2010;7:
851–869. doi:10.3934/mbe.2010.7.851

9. Cori A, Ferguson NM, Fraser C, Cauchemez S. A new framework and software to
estimate time-varying reproduction numbers during epidemics. Am J Epidemiol.
2013;178: 1505–1512. doi:10.1093/aje/kwt133

10. Ganyani T, Kremer C, Chen D, Torneri A, Faes C, Wallinga J, et al. Estimating the
generation interval for coronavirus disease (COVID-19) based on symptom onset data,

March 2020. Euro Surveill. 2020;25. doi:10.2807/1560-7917.ES.2020.25.17.2000257

11.   Hethcote HW. The Mathematics of Infectious Diseases. SIAM Rev. 2000;42: 599–653.
doi:10.1137/S0036144500371907

12.   Parag KV. Sub-spreading events limit the reliable elimination of heterogeneous
epidemics. J R Soc Interface. 2021;18: 20210444. doi:10.1098/rsif.2021.0444

13.   Anderson R, May R. Infectious diseases of humans: dynamics and control. Oxford
University Press; 1991.

14.   Wallinga J, Lipsitch M. How generation intervals shape the relationship between
growth rates and reproductive numbers. Proc R Soc B. 2007;274: 599–604.

15.   Ali ST, Wang L, Lau EHY, Xu X-K, Du Z, Wu Y, et al. Serial interval of SARS-CoV-2
was shortened over time by nonpharmaceutical interventions. Science. 2020;369:
1106–1109. doi:10.1126/science.abc9004

16.   Kenah E, Lipsitch M, Robins JM. Generation interval contraction and epidemic data
analysis. Math Biosci. 2008;213: 71–79. doi:10.1016/j.mbs.2008.02.007

17.   Hart WS, Miller E, Andrews NJ, Waight P, Maini PK, Funk S, et al. Generation time of
the alpha and delta SARS-CoV-2 variants: an epidemiological analysis. Lancet Infect
Dis. 2022;22: 603–610. doi:10.1016/S1473-3099(22)00001-9

18.   Favero M, Scalia Tomba G, Britton T. Modelling preventive measures and their effect
on generation times in emerging epidemics. J R Soc Interface. 2022;19: 20220128.
doi:10.1098/rsif.2022.0128

19.   Kraemer MUG, Pybus OG, Fraser C, Cauchemez S, Rambaut A, Cowling BJ.
Monitoring key epidemiological parameters of SARS-CoV-2 transmission. Nat Med.
2021;27: 1854–1855. doi:10.1038/s41591-021-01545-w

20.   Pellis L, Scarabel F, Stage HB, Overton CE, Chappell LHK, Fearon E, et al.
Challenges in control of COVID-19: short doubling time and long delay to effect of
interventions. Philos Trans R Soc Lond B, Biol Sci. 2021;376: 20200264.
doi:10.1098/rstb.2020.0264

21.   The R value and growth rate - GOV.UK [Internet]. [cited 1 Jul 2021]. Available:
https://www.gov.uk/guidance/the-r-value-and-growth-rate

22.   Fraser C. Estimating individual and household reproduction numbers in an emerging
epidemic. PLoS One. 2007;2: e758. doi:10.1371/journal.pone.0000758

23.   Parag KV, Donnelly CA, Zarebski AE. Quantifying the information in noisy epidemic
curves. Nat Comput Sci. 2022;2: 584-594 doi.org/10.1038/s43588-022-00313-1

24.   Parag KV, Donnelly CA. Fundamental limits on inferring epidemic resurgence in real
time using effective reproduction numbers. PLoS Comput Biol. 2022;18: e1010004.
doi:10.1371/journal.pcbi.1010004

25.   Parag KV. Improved estimation of time-varying reproduction numbers at low case
incidence and between epidemic waves. PLoS Comput Biol. 2021;17: e1009347.
doi:10.1371/journal.pcbi.1009347

26.   Torneri A, Libin P, Scalia Tomba G, Faes C, Wood JG, Hens N. On realized serial and
generation intervals given control measures: The COVID-19 pandemic case. PLoS
Comput Biol. 2021;17: e1008892. doi:10.1371/journal.pcbi.1008892

27.   Lloyd-Smith JO, Schreiber SJ, Kopp PE, Getz WM. Superspreading and the effect of
individual variation on disease emergence. Nature. 2005;438: 355–359.
doi:10.1038/nature04153

28.   Bettencourt LMA, Ribeiro RM. Real time bayesian estimation of the epidemic potential
of emerging infectious diseases. PLoS One. 2008;3: e2185.
doi:10.1371/journal.pone.0002185

29.   Lehtonen J. The Lambert W function in ecological and evolutionary models. Methods
Ecol Evol. 2016;7: 1110–1118. doi:10.1111/2041-210X.12568

30.   Gostic KM, McGough L, Baskerville EB, Abbott S, Joshi K, Tedijanto C, et al. Practical
considerations for measuring the effective reproductive number, Rt. PLoS Comput
Biol. 2020;16: e1008409. doi:10.1371/journal.pcbi.1008409

31.   Parag KV, Cowling BJ, Donnelly CA. Deciphering early-warning signals of SARS-CoV-
2 elimination and resurgence from limited data at multiple scales. J R Soc Interface.
2021;18: 20210569. doi:10.1098/rsif.2021.0569

32.   Van Kerkhove MD, Bento AI, Mills HL, Ferguson NM, Donnelly CA. A review of
epidemiological parameters from Ebola outbreaks to inform early public health
decision-making. Sci Data. 2015;2: 150019. doi:10.1038/sdata.2015.19

33.   Ferguson N, Laydon D, Nedjati-Gilani G, Others. Impact of non-pharmaceutical
interventions (NPIs) to reduce COVID- 19 mortality and healthcare demand. Imperial
College London; 2020.

34.   Dushoff J, Park SW. Speed and strength of an epidemic intervention. Proc Biol Sci.
2021;288: 20201556. doi:10.1098/rspb.2020.1556

35.   Shmueli G. To Explain or to Predict? Stat Sci. 2010;25: 289–310. doi:10.1214/10-
STS330

36.   Wagenmakers E-J, Grünwald P, Steyvers M. Accumulative prediction error and the
selection of time series models. J Math Psychol. 2006;50: 149–166.
doi:10.1016/j.jmp.2006.01.004

37.   Parag KV, Donnelly CA. Using information theory to optimise epidemic models for real-
time prediction and estimation. PLoS Comput Biol. 2020;16: e1007990.
doi:10.1371/journal.pcbi.1007990

38.   Marcellino M, Stock JH, Watson MW. A comparison of direct and iterated multistep AR
methods for forecasting macroeconomic time series. J Econom. 2006;135: 499–526.
doi:10.1016/j.jeconom.2005.07.020

39.   Nishiura H, Chowell G, Heesterbeek H, Wallinga J. The ideal reporting interval for an
epidemic to objectively interpret the epidemiological time course. J R Soc Interface.
2010;7: 297–307. doi:10.1098/rsif.2009.0153

40.   White PJ, Ward H, Garnett GP. Is HIV out of control in the UK? An example of
analysing patterns of HIV spreading using incidence-to-prevalence ratios. AIDS.
2006;20: 1898–1901. doi:10.1097/01.aids.0000244213.23574.fa

41.   Wallinga J, Teunis P. Different epidemic curves for severe acute respiratory syndrome
reveal similar impacts of control measures. Am J Epidemiol. 2004;160: 509–516.
doi:10.1093/aje/kwh255

42.   Yan P. Separate roles of the latent and infectious periods in shaping the relation
between the basic reproduction number and the intrinsic growth rate of infectious
disease outbreaks. J Theor Biol. 2008;251: 238–252. doi:10.1016/j.jtbi.2007.11.027

43.   Azmon A, Faes C, Hens N. On the estimation of the reproduction number based on
misreported epidemic data. Stat Med. 2014;33: 1176–1192. doi:10.1002/sim.6015

44.   Bracher J, Held L. Endemic-epidemic models with discrete-time serial interval
distributions for infectious disease prediction. Int J Forecast. 2020;
doi:10.1016/j.ijforecast.2020.07.002

45.   Dalziel BD, Lau MSY, Tiffany A, McClelland A, Zelner J, Bliss JR, et al. Unreported
cases in the 2014-2016 Ebola epidemic: Spatiotemporal variation, and implications for
estimating transmission. PLoS Negl Trop Dis. 2018;12: e0006161.
doi:10.1371/journal.pntd.0006161

46.   Goldstein E, Dushoff J, Ma J, Plotkin JB, Earn DJD, Lipsitch M. Reconstructing
influenza incidence by deconvolution of daily mortality time series. Proc Natl Acad Sci
USA. 2009;106: 21825–21829. doi:10.1073/pnas.0902958106
47.   Liu Q-H, Ajelli M, Aleta A, Merler S, Moreno Y, Vespignani A. Measurability of the
epidemic reproduction number in data-driven contact networks. Proc Natl Acad Sci
USA. 2018;115: 12680–12685. doi:10.1073/pnas.1811115115
48.   Sarrka S. Bayesian Filtering and Smoothing. Cambridge, UK: Cambridge University
Press; 2013.

681
682

Funding

KVP acknowledges funding from the MRC Centre for Global Infectious Disease Analysis (reference MR/R015600/1), jointly funded by the UK Medical Research Council (MRC) and the UK Foreign, Commonwealth & Development Office (FCDO), under the MRC/FCDO Concordat agreement and is also part of the EDCTP2 programme supported by the European Union. The funders had no role in study design, data collection and analysis, decision to publish, or manuscript preparation.

Data availability statement

All data and code underlying the analyses and figures of this work are freely available (in R and MATLAB) at: https://github.com/kpzoo/Omega

693
694