Safety and effectiveness of COVID-19 mRNA vaccination and risk factors for

hospitalisation caused by the omicron variant in 0.8 million adolescents: A nationwide

cohort study in Sweden

Prof. Peter Nordström, PhD[1], Marcel Ballin, MSc[1], Anna Nordström, PhD[1,2,3]

[1]Department of Community Medicine and Rehabilitation, Unit of Geriatric Medicine, Umeå

University, Umeå, Sweden

[2]Department of Public Health and Clinical Medicine, Section of Sustainable Health, Umeå

University, Umeå, Sweden

[3]School of Sport Sciences, UiT the Arctic University of Norway, Tromsø, Norway

Corresponding author:

Peter Nordström, Professor

Unit of Geriatric Medicine

Department of Community Medicine and Rehabilitation

Umeå University

90187 Umeå

Sweden

Phone: +4670 899 65 99

E-mail: peter.nordstrom@umu.se

Abstract

Background

Real-world evidence on the safety and effectiveness of COVID-19 vaccination against severe disease caused by the omicron variant among adolescents is sparse. In addition, evidence on risk factors for severe COVID-19 disease, and whether vaccination is similarly effective in such risk groups, is unclear.

Methods and findings

Nationwide registers were used to examine the safety and effectiveness of COVID-19 mRNA vaccination against COVID-19 hospitalisation, and risk factors for COVID-19 hospitalisation in adolescents. The safety analysis included all individuals in Sweden born between 2003-2009 (aged 11.3-19.2 years) given at least one dose of mRNA vaccine (N=645,355), and never vaccinated controls (N=199,022). Outcomes evaluated included all hospitalisations until 5 June 2022. The vaccine effectiveness (VE) against COVID-19 hospitalisation and associated risk factors was evaluated in adolescents given two doses of mRNA vaccine (N=501,945), as compared to never vaccinated controls (N=170,083), during an omicron predominant period (1 January 2022 to 5 June 2022). The safety analysis showed that COVID-19 mRNA vaccination was not associated with an increased risk of any serious adverse events resulting in hospitalisation. During follow-up, 1.69% of the vaccinated individuals were hospitalised compared to 1.71% of the controls (P=0.29). In the VE analysis, there were 21 cases (0.004%) of COVID-19 hospitalisation among 2-dose recipients and 26 cases (0.015%) among controls, resulting in an estimated VE of 75% (95% CI, 54-86, P<0.001). Strong risk factors for COVID-19 hospitalisation included previous infections (odds ratio [OR], 14.3, 95% CI, 7.7-26.6, P<0.001), and cerebral palsy/development disorders (OR, 12.0, 95% CI, 6.4-22.6, P<0.001), with similar estimates of VE in these subgroups as in the total cohort. The number needed to vaccinate with two doses to prevent one case of

COVID-19 hospitalisation was 9,007 in the total cohort and 1,031 in those with previous

infections or developmental disorders. None of the individuals hospitalised due to COVID-19

died within 30 days.

Conclusions

In this nationwide study, COVID-19 mRNA vaccination was not associated with an increased

risk of any serious adverse event in adolescents. Two doses were associated with a lower risk

of COVID-19 hospitalisation during the omicron predominant period, especially among those

with certain predisposing conditions who should be prioritized for vaccination. However,

COVID-19 hospitalisation among general adolescents was extremely rare, and additional

doses in this population may not be warranted at this stage.

Why was this study done

Evidence before this study

> There is limited evidence on the effectiveness of COVID-19 vaccination against severe outcomes during the omicron era among adolescents. In addition, there is lack of data on whether certain groups of adolescents are at greater risk of severe COVID-19 and should be prioritized in vaccination programs, and whether vaccination is equally effective in such risk groups.

> Regarding safety, some studies have indicated a link between COVID-19 mRNA vaccination and increased risk of myocarditis and pericarditis in young men, although the data appear inconsistent.

What did the researchers do and find?

> Using Swedish nationwide health registers, a cohort of 844,377 adolescents were followed until 5 June 2022 to evaluate the safety and effectiveness of COVID-19 mRNA vaccination against COVID-19 hospitalisation, and risk factors for COVID-19 hospitalisation.

> COVID-19 mRNA vaccination in adolescents was not associated with an increased risk of any diagnoses resulting in hospitalisation. In contrast, two doses of vaccine had an associated 75% effectiveness against COVID-19 hospitalisation. However, only about 6 individuals in 100,000 were hospitalised due to COVID-19 during follow-up .

> There were certain strong risk factors for COVID-19 hospitalisation, such as previous infections and different development disorders, which increased the risk of COVID-19 hospitalisation more than tenfold. Vaccine effectiveness among these individuals was similar as in the rest of the cohort.

What do these findings mean?

➤ Although COVID-19 mRNA vaccination appears safe and associated with reduced

risk of COVID-19 hospitalisation, the risk of severe COVID-19 seems to be extremely

low in general adolescents. Therefore, administration of additional doses to the

general population of adolescents may not be warranted at this stage of the pandemic.

In contrast, individuals with a high risk for severe COVID-19 should be prioritised for

vaccination.

Introduction

The omicron variant of SARS-CoV-2 led to a surge in cases among young people during 2022 when several countries lifted or alleviated their restrictions.[1] Although clinical trials showed that the BNT162b2 and mRNA-1273 vaccines had acceptable safety profiles in adolescents and reduced the risk of infection in the short-term,[2, 3] these trials were conducted before the emergence of omicron. Limited sample sizes also hindered the evaluation of the level of protection against severe COVID-19, such as hospitalisation, which could be estimated using large-scale observational studies. These studies also offer the possibility of investigating any potential links between vaccination and rare serious adverse events, such as myocarditis, for which the data appear inconsistent.[4, 5]

Currently, there is limited data on vaccine effectiveness (VE) against severe COVID-19 caused by the omicron variant among adolescents. Two case-control studies found that two doses of the BNT162b2 mRNA vaccine had about 80% VE against COVID-19 hospitalisation or death in adolescents.[6] [7] However, given their study design, it is difficult to determine how common severe disease is during the omicron era. In addition, little is known about whether certain groups of adolescents should be prioritized for vaccination because of a higher risk of severe COVID-19, and whether vaccination has a similarly protective effect in such risk groups. Moreover, although a third dose, also known as a booster dose, may increase protection against symptomatic omicron infection in adolescents,[8, 9] the risk of severe COVID-19 after a third dose relative to after the second dose is unclear.

Therefore, the present study leveraged Swedish nationwide registers to evaluate, among adolescents, 1) the risk of serious adverse events following COVID-19 mRNA vaccination, and 2) the effectiveness of COVID-19 mRNA vaccination against hospitalisation due to COVID-19 and risk factors for COVID-19 hospitalisation during an omicron predominant period.

Methods

Study design and cohorts

This nationwide, retrospective cohort study was approved by the Swedish Ethical
Review Authority (number 00094/2021), who waived the requirement of obtaining informed
consent given the retrospective study design.

The cohort considered for inclusion was compiled by Statistics Sweden (www.SCB.se);
the government agency in charge of nationwide statics in different areas and covering the total
population of Sweden. All individuals born 2003-2009 who were given at least one dose of
COVID-19 vaccine or had a documented SARS-CoV-2 infection until March 2022 were
considered for inclusion (N=692,333). To these individuals, Statistics Sweden matched one
control individual on birth year, sex, and municipality. Controls could not have a documented
SARS-CoV-2 infection at the date of documented infection in the corresponding infected
individual, and they could not have received a first dose of vaccine at the date when the
corresponding vaccinated individual had received two doses of vaccine. The cohort was
updated with vaccination data and SARS-CoV-2 infections until 2 June 2022. Because one
control could be matched to several vaccinated individuals, the total eligible cohort consisted
of 844,511 individuals, of whom 645,489 had been vaccinated with at least one dose, 600,846
had been vaccinated with two doses, and 60,475 had been vaccinated with at least three doses
(Figure 1). After excluding a few individuals vaccinated with a vector-based vaccine
(N=134), a total of 844,377 individuals remained and were included in the present study for
two sets of analyses. In a first set of analyses, named the safety analysis, all diagnoses set
during hospitalisation were evaluated in all individuals given at least one dose of vaccine
during follow-up (N=645,355) as compared to individuals never vaccinated during follow-up
(N=199,022). In a second set of analyses, VE against COVID-19 hospitalisation was
evaluated by comparing individuals given at least two doses of vaccine (N=501,945) to

individuals never vaccinated during follow-up (N=170,083), excluding all individuals with a

previous documented SARS-CoV-2 infection (Figure 1). Data on individuals vaccinated

against COVID-19 and data on documented SARS-CoV-2 infections were collected from the

Swedish Vaccination Register and the SmiNet register, respectively. Both these registers are

managed by the Public Health Agency of Sweden and all healthcare providers in Sweden

were obliged to report to these registers according to law.[10, 11]

Outcomes

In the safety analysis, based on all diagnoses set in the cohort during the hospital stay

until 5 June 2022, results are presented for 30 different diagnoses. These diagnoses were

selected based on their incidence and general interest given previous reports of links between

certain diagnoses and COVID-19 mRNA vaccination.[4, 12, 13] Only the first diagnosis was

evaluated for each individual, and individuals with this diagnosis at baseline were therefore

excluded from the prospective analyses.

In the VE analysis, the primary outcome was a main diagnosis of COVID-19 (thus,

"due to" COVID-19 rather than "with" COVID-19), set during inpatient hospital stay from 1

January 2022 until 5 June 2022. The secondary outcome was documented SARS-CoV-2

infection of any severity from the SmiNet register, from 1 January 2022 until 28 February

2022, as per when the guidelines for testing in Sweden had changed. For these two outcomes,

only cases that occurred 7 days after the second dose of vaccine and onwards were counted to

ensure the full effect of vaccination. The start of 1 January was selected on the basis that the

omicron variant was first documented in Sweden on 29 November 2021[14] and by early

January 2022 it represented >90% of sequenced cases (Supplementary Table 1). Data on all

diagnoses used in this study were obtained from the National Patient Register[15] using 10th

revision International Classification of Disease codes, starting from 1 January 2016 earliest

until 5 June 2022 latest. The definition and diagnostic codes for all diagnoses considered in

the safety analysis and for the analysis of risk factors for COVID-19 hospitalisation are shown

in Table 1. The positive predictive value for diagnoses set within the National Patient Register

differs, but has generally been found to be between 85-95%, although sensitivity is often

lower.[15] Finally, data on death due to COVID-19 or influenza, defined as death within 30

days after the diagnosis, were obtained from the National Cause of Death Register.[16] The

present study was approved by the Swedish Ethical Review Authority.

Statistical analysis

In the safety analysis, diagnoses were evaluated before and after the first dose of

vaccine in 645,355 individuals given at least one dose of vaccine and in 199,022 controls who

were unvaccinated during the follow-up time. The baseline date in vaccinated individuals was

the date of vaccination with the first dose. In the controls, a baseline date was randomly

assigned based on the mean baseline date and standard deviation among the vaccinated

individuals (10 October 2021 $\pm$ 54 days). Student's t-tests and chi-square tests were used to

compare the prevalence of different variables at baseline. To estimate hazard ratios (HRs) for

the 30 different diagnoses during follow-up, Cox regression models were used. Individuals

were censored on the date of the diagnosis of interest, death, or end of follow-up (5 June

2022), whichever came first.

In the VE analysis, logistic regression was used to estimate odds ratios (OR) for the

primary outcome of COVID-19 hospitalisation (from 1 January 2022 until 5 June 2022), and

for the secondary outcome of a SARS-CoV-2 infection (from 1 January 2022 until 28

February 2022). The ORs obtained were used to estimate VE as 1 minus the OR x 100. In all

analyses, the first model was unadjusted and the second model was adjusted for baseline date,

age, sex, and whether the individual was born in Sweden or not. Data underlying these

covariates were retrieved from Statistics Sweden.[17] To investigate whether VE differed by

the covariates, interaction analyses were performed using product terms created by

multiplying the variable coding for vaccination status at baseline (vaccinated/unvaccinated)

by each respective covariate, which was added to the logistic regression model. Given that the

interaction term was statistically significant (P<0·05) for the baseline date, VE was also

estimated in subgroups according to this covariate. The number needed to vaccinate (NNV)

with two doses to prevent one case of COVID-19 hospitalisation during follow-up was

estimated as the inverse of the absolute risk difference between the groups

(vaccinated/unvaccinated). The NNV was estimated both in the total cohort and by subgroups.

Finally, a sensitivity analysis using a negative control outcome was conducted to

explore the potential risk of bias due to unmeasured confounding.[18] Here, a logistic

regression model was performed, comparing individuals given at least two doses of COVID-

19 mRNA vaccine compared to never vaccinated individuals concerning the outcome of

hospitalisation due to influenza from 1 January 2022 until 5 June 2022. All analyses were

performed in SPSS v29·0 for Mac (IBM Corp, Armonk, NY, USA), and Stata v16·1 for Mac

(Statcorp, College Station, Texas, USA). A two-sided P-value <0·05 or Ors/HRs with 95%

CIs not crossing one were considered statistically significant.

Role of the funding source

The present study was not funded.

Results

The total cohort comprised 844,377 adolescents born 2003-2009 (age 11.3-19.2 years),

of whom 645,355 received at least one dose of COVID-19 mRNA vaccine and 199,022 never

vaccinated individuals (controls). Almost 90% of the vaccinated individuals received

BNT162b2 as a first dose, while the remaining received mRNA-1273. Baseline characteristics

are shown in Table 1. Individuals that were never vaccinated were slightly younger, more

often of male sex and born outside of Sweden, and were less often diagnosed with a previous

SARS-CoV-2 infection (P<0.001 for all). Concerning other diagnoses at baseline, differences

between the groups were marginal (Table 1).

Serious adverse events after a first dose of vaccine

In the safety analysis (N=844,377), there were a total of 19,660 hospitalisations among

14,320 individuals during follow-up. In vaccinated individuals, 1.69% (N=10,918) were

hospitalised at least once, compared to 1.71% (N=3,402) in those never vaccinated (P=0.29

for difference). There were marginal differences between the two groups in the 30 diagnoses

presented (Table 2), although statistically significant associations in favour of vaccination

were observed with respect to the risk of sepsis, thrombocytopenia, alcohol dependency, and

Crohn's disease (P<0.05 for all). None of the other associations were statistically significant.

Vaccine effectiveness against COVID-19 hospitalisation

In the analysis of VE against COVID-19 hospitalisation, 501,945 individuals vaccinated

with two doses and 170,083 never vaccinated controls were included. Between 1 January

2022 and 5 June 2022, a total of 47 individuals (6 per 100,000) were hospitalised due to

COVID-19. Of these, 21 cases were among those vaccinated with two doses (0.004%), and 26

among unvaccinated (0.015%), resulting in a VE of 75% (95% CI, 54-86, P<0.001) (Table 3).

The NNV with two doses to prevent one case of COVID-19 hospitalisation was 9,007. For

those with a second dose of vaccine earlier than 15 November, the VE was 68% (95% CI, 25-

86, P=0.009), compared to 86% (95% CI, 63-95, P<0.001) for those with a second dose of

vaccine 15 November 2021 and later (P=0.03 for interaction). When comparing individuals

vaccinated with three doses (N=41,225) compared to those vaccinated with two doses

(N=413,544), only 12 individuals were hospitalised due to COVID-19 during follow-up (VE,

13%, 95% CI, -354-84, P=0.86). There were no deaths within 30 days of hospitalisation

among the 238 individuals hospitalised due to COVID-19 in the total cohort (N=844,377)

since the beginning of the pandemic in January 2020.

Risk factors for COVID-19 hospitalisation and vaccine effectiveness by subgroups

Of the 47 individuals hospitalised due to COVID-19 in the VE analysis, 28 (60%) had

previously been hospitalised for another condition, compared to 66,756 (10%) of the

individuals in the rest of the cohort. Among the individuals hospitalised due to COVID-19, 15

(32%) had previously been diagnosed with an infection, compared to 22,110 (3.3%) in the

rest of the cohort, equal to an adjusted OR for COVID-19 hospitalisation of 14.3 (95% CI,

7.7-26.6, P<0.001), after adjustment for age, sex, baseline date, vaccination status, and

whether the individual was born in Sweden. The VE in this risk group was similar (VE, 88%,

95% CI, 57-96, P<0.001), as in the total cohort, but with a slightly lower vaccination

coverage (72.7% vs. 74.7%). In addition, 15 (32%) of the individuals hospitalised due to

COVID-19 had previously been diagnosed with cerebral palsy and/or different development

disorders, compared to 25,947 individuals (3.9%) in the rest of the cohort, resulting in an

adjusted OR for COVID-19 hospitalisation of 12.0 (95% CI, 6.4-22.6, P<0.001). Again, VE

in this subgroup was similar as in the total cohort (VE, 72%, 95% CI, 18-90, P=0.02), but

with a slightly lower vaccination coverage (71.5% vs 74.8%). The NNV with two doses to

prevent one case of COVID-19 hospitalisation in the subgroup of individuals previously

diagnosed with infections or development disorders (N=46,783) was 1,031.

Vaccine effectiveness against SARS-CoV-2 infection

The VE against SARS-CoV-2 infection of any severity was estimated among 488,441

individuals vaccinated with two doses compared to 165,626 never vaccinated controls.

Between 1 January 2022 and 28 February 2022, there were 72,631 cases of confirmed SARS-

CoV-2 infection. The VE varied by time since the last dose (P<0.001), with no VE in the total

cohort (VE, -2%, 95% CI, -4-0, P=0.02) (Table 4). There was a marginal VE among

individuals with a second dose no earlier than 1 September 2021 (VE, 4%, 95% CI, 2-6,

P<0.001), a low VE for individuals with a second dose no earlier than 1 November 2021 (VE,

21%, 95% CI, 19-23, P<0.001), and a moderate VE for individuals with a second dose no

earlier than 1 January 2022 (VE, 47%, 95% CI, 43-51, P<0.001).

Sensitivity analysis

The risk of hospitalisation due to influenza was evaluated in 600,021 individuals given at

least two doses of vaccine compared to in 199,003 never vaccinated individuals. Between 1

January 2022 and 5 June 2022, a total of 48 individuals (6 per 100,000) were hospitalised due

to influenza, with no detectable VE (VE, -6%, 95% CI, -51-119, P=0.88). Of the 130

individuals in the total cohort hospitalised for influenza since the start of the COVID-19

pandemic in January 2020, one individual died within 30 days of hospitalisation.

Discussion

In this nationwide study of more than 0.8 million adolescents, COVID-19 mRNA

vaccination was not associated with an increased risk of hospitalisation for any diagnosis,

with an overall risk of hospitalisation which was similar in vaccinated and unvaccinated

individuals. The results also showed that although vaccination was associated with lower risk

of COVID-19 hospitalisation during an omicron predominant period, the absolute risk of

being hospitalised was extremely low, except in subgroups of adolescents that had previously been diagnosed with infections, cerebral palsy or other development disorders.

Evidence on the safety and effectiveness of COVID-19 mRNA vaccination in adolescents during the omicron predominant period is limited. In this study, no association between vaccination and increased risk of hospitalisation in general, or for any specific diagnosis, was detected. This suggests that COVID-19 vaccination in adolescents is safe. These findings add to, and extend upon those from a nationwide study in Scotland, reporting no association between vaccination and increased risk of hospital stay for 17 different diagnoses among adolescents.[5] The present study also estimated that two doses of mRNA vaccine had about 75% effectiveness against hospitalisation due to COVID-19. Similar estimates were reported in two case-control studies of adolescents conducted in the US and Brazil during an omicron predominant period, where VE against COVID-19 hospitalisation or death was about 80%.[6] [7] However, given their design, these studies were unable to determine how common severe COVID-19 is. Therefore, the very low absolute risk of severe disease observed in the present cohort study based on a nationwide sample of adolescents is an important finding for decision making concerning the need for COVID-19 vaccination in adolescents. Overall, only 47 individuals, or 6 in 100,000, were hospitalised due to COVID-19 during a period of 5 months, and none of all adolescents hospitalised due to COVID-19 since the start of the pandemic died within 30 days of hospitalisation. Based on this very low risk of severe COVID-19 in the total cohort, the NNV with two doses to prevent one case during follow-up was about 9,000. Interestingly, the absolute risk of being hospitalised due to COVID-19 and influenza was similar, despite that the infection pressure from SARS-CoV-2 during follow-up was more than 100 times higher than that of influenza.[19]

Given the above, it is important to evaluate whether certain groups of adolescents are at higher risk of severe COVID-19, and if so, whether vaccination is associated with similar

protection among these individuals. This study identified two different clusters of diagnoses which increased the risk of COVID-19 hospitalisation more than tenfold. The first included previous infections and the second included cerebral palsy and other developmental disorders. An encouraging finding was that the VE in these subgroups was similar to that in the total cohort, and consequently, the NNV to prevent one case of COVID-19 hospitalisation was about 1,000 individuals. It is therefore of concern that vaccination coverage in these risk groups was somewhat lower compared to in the total cohort. Although we are unable to determine the underlying causes for this observation, factors such as recurrent infections interfering with the administration of the vaccine, and fear of adverse events, may have contributed. Taken together, these findings suggest that vaccination of adolescents during the omicron era should primarily be targeting risk groups, such as those with previous infections and development disorders.

Concerning the outcome of SARS-CoV-2 infection of any severity, the results indicated that VE from primary vaccination wanes within a few months, similar to observations made in a few other countries.[7-9] However, there is a lack of data exploring whether booster doses reduce the risk of severe COVID-19 as compared to primary vaccination.[20] The results for this comparison in the present study showed that COVID-19 hospitalisations during follow-up were once again extremely rare, implying that administration of booster doses to the general population of adolescents may not be warranted at the current stage of the pandemic. Based on these findings, it is noteworthy that the US and several countries in Europe are recommending booster doses to adolescents.[21, 22]

The present study has limitations that should be considered. Because of the observational design, conclusions based on the associations found should be made with caution. For example, despite that VE changed marginally before and after adjustment for covariates, there may be other factors that could have influenced the estimates. However, for

the analysis of serious adverse events, we evaluated all diagnoses set during hospital stay both

before and after the first dose of vaccine, thereby increasing the chance of detecting selection

bias that could interfere with the results. In addition, the sensitivity analysis wherein influenza

was used as a negative control outcome supported the lack of important confounding.

Moreover, because COVID-19 hospitalisation in this population was very rare, it was not

possible to estimate the VE of a first booster dose in this cohort with any form of precision.

Strengths of this study include that all the registers used to obtain the data used in the present

study have nationwide coverage with virtually zero loss to follow-up. Finally, the study

cohort was based on the total population of Swedish adolescents, including 0.8 million

individuals aged 11-19 years of age, which increases the possibility to generalize the findings

to other countries with similar population structures.

Conclusion

In summary, COVID-19 mRNA vaccination was not associated with an increased risk

for hospitalisation of any cause in adolescents, suggesting that they are safe to use. While

vaccination was associated with a reduced risk of COVID-19 hospitalisation during the

omicron era, the absolute risk among general adolescents was extremely low and similar to

that of influenza. In contrast, in a large proportion of adolescents hospitalised due to COVID-

19, certain risk factors were present, and the effectiveness of vaccination in these individuals

was similar to in the total cohort. These results indicate that certain vulnerable subgroups of

adolescents, rather than adolescents in general, should be prioritized for vaccination.

Contributors

PN conceived and designed the study. PN acquired the data. PN performed the

statistical analyses. PN and MB accessed and verified the underlying data. All authors

interpreted the data. PN and MB drafted the manuscript. All authors critically revised the

manuscript for intellectual content. PN and AN supervised the work. All authors gave final

approval of the version to be published. All authors had full access to all the data and had

final responsibility for the decision to submit for publication

Declaration of interests

We declare no competing interests.

Role of the funding source

There was no funding source for this study.

Data sharing statement

The data files used for the present study are publicly unavailable according to

regulations under Swedish law. However, all data used for the present study can be applied

for from the National Board of Health and Welfare, Statistics Sweden, and the Public Health

Agency of Sweden.

References

1. World Health Organization. Interim statement on COVID-19 vaccination for
children. Available from: https://www.who.int/news/item/11-08-2022-interim-statement-
on-covid-19-vaccination-for-children. Accessed 20 Sep, 2022.
2. Frenck RW, Jr., Klein NP, Kitchin N, Gurtman A, Absalon J, Lockhart S, et al.
Safety, Immunogenicity, and Efficacy of the BNT162b2 Covid-19 Vaccine in Adolescents. N
Engl J Med. 2021;385(3):239-50. Epub 2021/05/28. doi: 10.1056/NEJMoa2107456. PubMed
PMID: 34043894; PubMed Central PMCID: PMCPMC8174030.
3. Ali K, Berman G, Zhou H, Deng W, Faughnan V, Coronado-Voges M, et al.
Evaluation of mRNA-1273 SARS-CoV-2 Vaccine in Adolescents. N Engl J Med.
2021;385(24):2241-51. Epub 2021/08/12. doi: 10.1056/NEJMoa2109522. PubMed PMID:
34379915; PubMed Central PMCID: PMCPMC8385554.

4.   Mevorach D, Anis E, Cedar N, Hasin T, Bromberg M, Goldberg L, et al.
Myocarditis after BNT162b2 Vaccination in Israeli Adolescents. N Engl J Med.
2022;386(10):998-9. Epub 2022/01/27. doi: 10.1056/NEJMc2116999. PubMed PMID:
35081295; PubMed Central PMCID: PMCPMC8823652.
5.   Rudan I, Millington T, Antal K, Grange Z, Fenton L, Sullivan C, et al. BNT162b2
COVID-19 vaccination uptake, safety, effectiveness and waning in children and young people
aged 12-17 years in Scotland. The Lancet Regional Health – Europe. 2022;23. doi:
10.1016/j.lanepe.2022.100513.
6.   Price AM, Olson SM, Newhams MM, Halasa NB, Boom JA, Sahni LC, et al.
BNT162b2 Protection against the Omicron Variant in Children and Adolescents. N Engl J
Med. 2022;386(20):1899-909. Epub 2022/03/31. doi: 10.1056/NEJMoa2202826. PubMed
PMID: 35353976; PubMed Central PMCID: PMCPMC9006785.
7.   Florentino PTV, Millington T, Cerqueira-Silva T, Robertson C, de Araujo Oliveira
V, Junior JBS, et al. Vaccine effectiveness of two-dose BNT162b2 against symptomatic and
severe COVID-19 among adolescents in Brazil and Scotland over time: a test-negative case-
control study. Lancet Infect Dis. 2022. Epub 2022/08/12. doi: 10.1016/S1473-
3099(22)00451-0. PubMed PMID: 35952702; PubMed Central PMCID: PMCPMC9359673
programme from JBS SA. VdAO, VSB, MLB, and MB-N are employees of Fiocruz, a federal
public institution that manufactures Vaxzevria in Brazil through a full technology transfer
agreement with AstraZeneca. Fiocruz allocates all its manufactured products to the Ministry
of Health for public health use. SVK was Co-Chair of the Scottish Government's Expert
Reference Group on Ethnicity and COVID-19 and a member of the UK Government's
Scientific Advisory Group on Emergencies subgroup on ethnicity. IR is a member of the
Scientific Advisory Committee of the Government of Croatia and co-Editor-in-Chief of the
Journal of Global Health. CRS declares funding from the Medical Research Council, the
National Institute for Health Research, the Chief Scientist Office, and the New Zealand
Ministry for Business, Innovation and Employment and Health Research Council during the
conduct of this study. CR declares he is a Member of SPI-M, Scottish Government Scientific
Advisory Committee, MHRA COVID-19 vaccine benefit and risk expert working group. AS
declares that he is a member of the UK and Scottish Governments COVID-19 Advisory Boards
and Astra-Zeneca's Thrombotic Thrombocytopenic Taskforce. IR is Co-Editor-in-Chief of the
Journal of Global Health and President of the International Society of Global Health. All other
authors declare no competing interests.
8.   Fleming-Dutra KE, Britton A, Shang N, Derado G, Link-Gelles R, Accorsi EK, et al.
Association of Prior BNT162b2 COVID-19 Vaccination With Symptomatic SARS-CoV-2
Infection in Children and Adolescents During Omicron Predominance. JAMA.
2022;327(22):2210-9. Epub 2022/05/14. doi: 10.1001/jama.2022.7493. PubMed PMID:
35560036; PubMed Central PMCID: PMCPMC9107063.
9.   Tartof SY, Frankland TB, Slezak JM, Puzniak L, Hong V, Xie F, et al. Effectiveness
Associated With BNT162b2 Vaccine Against Emergency Department and Urgent Care
Encounters for Delta and Omicron SARS-CoV-2 Infection Among Adolescents Aged 12 to 17
Years. JAMA Netw Open. 2022;5(8):e2225162. Epub 2022/08/04. doi:
10.1001/jamanetworkopen.2022.25162. PubMed PMID: 35921109; PubMed Central PMCID:
PMCPMC9350713.
10.   Public Health Agency of Sweden. The National Vaccination Register. Available
from: https://www.folkhalsomyndigheten.se/smittskydd-
beredskap/vaccinationer/nationella-vaccinationsregistret/. Accessed 20 Sep, 2022.

11.   Public Health Agency of Sweden. SmiNet. Available from:
https://www.folkhalsomyndigheten.se/smittskydd-beredskap/overvakning-och-
rapportering/sminet/. Accessed 20 Sep, 2022.
12.   Wong HL, Hu M, Zhou CK, Lloyd PC, Amend KL, Beachler DC, et al. Risk of
myocarditis and pericarditis after the COVID-19 mRNA vaccination in the USA: a cohort study
in claims databases. Lancet. 2022;399(10342):2191-9. Epub 2022/06/13. doi:
10.1016/S0140-6736(22)00791-7. PubMed PMID: 35691322; PubMed Central PMCID:
PMCPMC9183215 KLA and JDS own stock in UnitedHealth Group. All other authors declare
no competing interests.
13.   Barda N, Dagan N, Ben-Shlomo Y, Kepten E, Waxman J, Ohana R, et al. Safety of
the BNT162b2 mRNA Covid-19 Vaccine in a Nationwide Setting. N Engl J Med.
2021;385(12):1078-90. Epub 2021/08/26. doi: 10.1056/NEJMoa2110475. PubMed PMID:
34432976; PubMed Central PMCID: PMCPMC8427535.
14.   Public Health Agency of Sweden. New virus variant of SARS-CoV-2 also in
Sweden.
https://www.folkhalsomyndigheten.se/nyheter-och-press/nyhetsarkiv/2021/november/ny-
virusvariant-av-sars-cov-2-aven-i-sverige/. Accessed 29 March 2022.
15.   Ludvigsson JF, Andersson E, Ekbom A, Feychting M, Kim JL, Reuterwall C, et al.
External review and validation of the Swedish national inpatient register. BMC Public Health.
2011;11:450. Epub 2011/06/09. doi: 10.1186/1471-2458-11-450. PubMed PMID: 21658213;
PubMed Central PMCID: PMCPMC3142234.
16.   Brooke HL, Talbäck M, Hörnblad J, Johansson LA, Ludvigsson JF, Druid H, et al.
The Swedish cause of death register. European journal of epidemiology. 2017;32(9):765-73.
17.   Ludvigsson JF, Almqvist C, Bonamy A-KE, Ljung R, Michaëlsson K, Neovius M, et
al. Registers of the Swedish total population and their use in medical research. European
Journal of Epidemiology. 2016;31(2):125-36. doi: 10.1007/s10654-016-0117-y.
18.   World Health Organization. Evaluation of COVID-19 vaccine effectiveness.
Interim guidance. March 17, 2021. Available from:
https://apps.who.int/iris/handle/10665/340301. Accessed 20 Sep, 2022.
19.   Public Health Agency of Sweden. Influensasäsongen 2021-2022. Available from:
https://www.folkhalsomyndigheten.se/contentassets/a9433fcecdc24c788e11ea0d47fdf21b
/influensasasongen-2021-2022-sasongssammanfattning-final.pdf. Accessed 29 Sep, 2022.
20.   Chiew CJ, Premikha M, Chong CY, Wei WE, Ong B, Lye DC, et al. Effectiveness of
primary series and booster vaccination against SARS-CoV-2 infection and hospitalisation
among adolescents aged 12-17 years in Singapore: a national cohort study. The Lancet
Infectious Diseases. 2022. doi: 10.1016/S1473-3099(22)00573-4.
21.   Centers for Disease Control and Prevention. Stay Up to Date with COVID-19
Vaccines Including Boosters. Available from: https://www.cdc.gov/coronavirus/2019-
ncov/vaccines/stay-up-to-date.html. Accessed 30 Sep, 2022.
22.   European Centre for Disease Prevention and Control. COVID-19 vaccine
effectiveness in adolescents aged 12–17 years and interim public health considerations for
administration of a booster dose. 8 February 2022. Available from:
https://www.ecdc.europa.eu/sites/default/files/documents/COVID-19-considerations-for-
booster-doses-in-adolescents-Feb%202022.pdf. Accessed 30 Sep, 2022.
520
521

Table 1. Baseline characteristics of the individuals at date of the first dose of vaccine and in never vaccinated individuals.

	Vaccinated with at least one dose (N=645,355)	Never vaccinated (N=199,022)	P value
Age, years (standard deviation)	15.4 (1.9)	14.8 (2.0)	<0.0001
Female sex	318,983 (49.4%)	90,957 (45.7%)	<0.0001
Born in Sweden	567,298 (87.9%)	147,056 (73.9%)	<0.0001
Mean baseline date,	10 October 2021	9 October 2021	<0.0001
Previous SARS-CoV-2 infection	91,934 (14.2%)	20,896 (10.5%)	<0.0001
Previous diagnoses (ICD-10 code)			
Gastroenteritis (A09)	610 (0.094%)	198 (0.099%)	0.27
Sepsis (A41)	130 (0.020%)	43 (0.022%)	0.35
Erysipelas (A46)	129 (0.020%)	38 (0.019%)	0.40
Bacterial infection unspecified (A49)	77 (0.012%)	16 (0.008%)	0.05
Mononucleosis (B27)	437 (0.068%)	81 (0.041%)	<0.001
Virus infection, unspecified (B34)	499 (0.077%)	153 (0.077%)	0.48
Chronic lymphocytic leukemia (C91)	78 (0.012%)	34 (0.017%)	0.06
Iron deficiency anemia (D50)	186 (0.029%)	66 (0.033%)	0.16
Thrombocytopenia (D69)	189 (0.029%)	66 (0.033%)	0.19
Agranulocytosis (D70)	81 (0.013%)	24 (0.012%)	0.43
Alcohol dependency (F10)	1520 (0.24%)	460 (0.23%)	0.36
Depressive episode (F32)	1427 (0.22%)	334 (0.17%)	<0.001
Anxiety state, unspecified (F41)	1008 (0.16%)	312 (0.16%)	0.48
Allergy or anaphylactic shock (T78)	539 (0.084%)	146 (0.073%)	0.08
Anorexia nervosa (F50)	1,034 (0.16%)	174 (0.087%)	<0.001
Epilepsy (G40)	871 (0.14%)	309 (0.16%)	0.02
Otitis media (H66)	144 (0.022%)	55 (0.028%)	0.10
Myocarditis (I40)	97 (0.015%)	30 (0.015%)	0.50
Pericarditis (I30)	51 (0.008%)	17 (0.008%)	0.39
Sinuitis (J01)	208 (0.032%)	76 (0.038%)	0.11
Tonsillitis (J03)	454 (0.070%)	129 (0.065%)	0.21
Chronic tonsillitis (J35)	1,070 (0.17%)	307 (0.15%)	0.13
Upper respiratory infection (J06)	233 (0.036%)	67 (0.034%)	0.31
Pneumonia (J15 and J18)	839 (0.13%)	236 (0.12%)	0.10
Peritonsillar abscess (J36)	337 (0.052%)	97 (0.049%)	0.28
Appendicitis (K35)	5,769 (0.89%)	1,508 (0.76%)	<0.001
Crohn's disease (K50)	207 (0.032%)	73 (0.037%)	0.17
Cutaneous abscess (L02)	215 (0.033%)	78 (0.039%)	0.12
Nephritis (N10)	588 (0.091%)	148 (0.74%)	0.01
Traumatic brain injury (S06)	4,222 (0.65%)	1,072 (0.54%)	<0.001
Combined diagnoses			
Cerebral palsy/development disorders (G80, F73, F82, F84, Q02)	23,093 (3.6%)	8,437 (4.2%)	<0.001
Infections (A49, J03, J15, J18, J35)	21,208 (3.3%)	7,302 (3.7%)	<0.001

All data are shown as N (%) unless stated otherwise.
All previous diagnoses were main diagnoses set during inpatient hospital stay, with the exception of the combined diagnoses where also open specialized health care was used.

Table 2. Hazard ratios for 30 different diagnoses during follow-up in individuals vaccinated with at least one dose compared to never vaccinated individuals.

Diagnosis (ICD-10 code)	Vaccinated with at least one dose (N=645,355) Number of cases (incidence rate)	Never vaccinated (N=199,022) Number of cases (incidence rate)	HR (95% CI)	P value
Gastroenteritis (A09)	58 (0.38)	20 (0.43)	1.02 (0.59-1.77)	0.94
Sepsis (A41)	5 (0.03)	7 (0.15)	0.18 (0.06-0.59)	0.004
Erysipelas (A46)	8 (0.05)	2 (0.04)	1.04 (0.19-5.67)	0.97
Bacterial infection unspecified (A49)	11 (0.07)	3 (0.06)	0.87 (0.24-3.21)	0.84
Mononucleosis (B27)	115 (0.76)	26 (0.55)	0.82 (0.53-1.26)	0.36
Virus infection, unspecified (B34)	42 (0.28)	7 (0.15)	1.65 (0.73-3.73)	0.23
Chronic lymphocytic leukemia (C91)	4 (0.03)	4 (0.09)	0.36 (0.08-1.50)	0.16
Iron deficiency anemia (D50)	38 (0.25)	14 (0.30)	0.94 (0.50-1.76)	0.84
Thrombocytopenia (D69)	8 (0.05)	10 (0.21)	0.22 (0.08-0.60)	0.003
Agranulocytosis (D70)	1 (0.007)	1 (0.02)	-	-
Alcohol dependency (F10)	409 (2.69)	124 (2.65)	0.79 (0.64-0.97)	0.02
Depressive episode (F32)	363 (2.39)	103 (2.19)	0.92 (0.73-1.15)	0.46
Anxiety state, unspecified (F41)	341 (2.25)	75 (1.60)	0.96 (0.74-1.24)	0.76
Allergy or anaphylactic shock (T78)	64 (0.42)	23 (0.49)	0.74 (0.45-1.21)	0.23
Anorexia nervosa (F50)	197 (1.30)	48 (1.02)	1.12 (0.81-1.55)	0.50
Epilepsy (G40)	82 (0.54)	34 (0.73)	0.89 (0.57-1.40)	0.62
Otitis media (H66)	7 (0.05)	4 (0.09)	0.61 (0.16-2.33)	0.47
Myocarditis (I40)	86 (0.57)	19 (0.41)	1.05 (0.63-1.74)	0.86
Pericarditis (I30)	21 (0.14)	3 (0.06)	0.92 (0.27-3.17)	0.89
Sinuitis (J01)	15 (0.10)	10 (0.21)	0.68 (0.27-1.69)	0.41
Tonsillitis (J03)	90 (0.59)	11 (0.23)	1.58 (0.83-3.00)	0.16
Chronic tonsillitis (J35)	94 (0.59)	19 (0.41)	1.17 (0.70-0.94)	0.56
Upper respiratory infection (J06)	24 (0.16)	5 (0.11)	1.64 (0.57-4.73)	0.36
Pneumonia (J15 and J18)	68 (0.45)	17 (0.36)	1.47 (0.83-2.60)	0.19
Peritonsillar abscess (J36)	52 (0.34)	17 (0.36)	0.58 (0.33-1.03)	0.06
Appendicitis (K35)	665 (4.38)	176 (3.76)	1.09 (0.92-1.29)	0.34
Crohn's disease (K50)	24 (0.16)	14 (0.30)	0.48 (0.24-0.97)	0.04
Cutaneous abscess (L02)	10 (0.07)	5 (0.11)	0.63 (0.21-1.90)	0.41
Nephritis (N10)	105 (0.69)	19 (0.41)	1.21 (0.73-1.99)	0.47
Traumatic brain injury (S06)	307 (2.02)	86 (1.84)	1.13 (0.88-1.45)	0.35

CI = confidence interval, HR = hazard ratio.

Incidence rates are presented as cases per one million person-years. HRs were adjusted for age, sex, baseline date, and whether the individual was born in Sweden.

Table 3. Vaccine effectiveness against COVID-19 hospitalisation from 7 days onwards after a second dose of vaccine as compared to never vaccinated individuals from 1 January 2022 until 5 June 2022, and by time since vaccination and subgroups.

	Vaccinated with two doses Number of cases (%)	Never vaccinated Number of cases (%)	Unadjusted vaccine effectiveness (95% CI)	Adjusted vaccine effectiveness (95% CI)
Total cohort (N=672,028)[a]	21 (0.004%)	26 (0.015%)	73 (51-85)	75 (54-86)
Subgroups				
Baseline date < 15 Nov 2021 (N=293,643)[b]	15 (0.007%)	14 (0.017%)	58 (13-80)	68 (25-86)
Baseline date > 14 Nov 2021 (N=378,385)[c]	6 (0.002%)	12 (0.014%)	85 (60-94)	86 (63-95)
Previously diagnosed with infections (N=22,125)[d]	4 (0.025%)	11 (0.182%)	86 (57-96)	88 (57-96)
Previously diagnosed with development disorders(N=25,962)[e]	6 (0.032%)	9 (0.121%)	73 (15-90)	72 (18-90)

CI = confidence interval.
Adjusted models were adjusted for age, baseline date, sex and whether the individual was born in Sweden.
[a]Of which 501,945 were vaccinated and 170,083 never vaccinated.
[b]Of which 210,747 were vaccinated and 82,896 never vaccinated.
[c]Of which 291,198 were vaccinated and 87,187 never vaccinated.
[d]Of which 16,083 were vaccinated and 6,042 never vaccinated.
[e]Of which 18,553 were vaccinated and 7,409 never vaccinated.

Table 4. Vaccine effectiveness against SARS-CoV-2 infection of any severity from 7 days onwards after a second dose of vaccine as compared to never vaccinated individuals from 1 January 2022 until 28 February 2022, and by time since vaccination.

	Vaccinated with two doses	Never vaccinated	Unadjusted vaccine effectiveness (95% CI)	Adjusted vaccine effectiveness (95% CI)
	Number of cases (%)	Number of cases (%)		
Total cohort (N=654,067)[a]	55,101 (11.3%)	17,530 (10.6%)	-7 (-9, 6)	-2 (-4, 0)
Subgroups				
≥ 1 September 2021 (N=622,296)[b]	51,497 (11.0%)	16,037 (10.5%)	-5 (-7, -3)	4 (2, 6)
≥ 1 October 2021 (N=487,348)[c]	32,613 (9.2%)	13,655 (10.3%)	11 (9, 13)	10 (8, 12)
≥ 1 November 2021 (N=409,833)[d]	27,213 (8.8%)	9,743 (9.7%)	11 (9, 13)	21 (19, 23)
≥ 1 December 2021 (N=245,914)[e]	12,547 (6.8%)	5,288 (8.5%)	21 (18, 23)	36 (34, 23)
≥ 1 January 2022 (N=107,773)[f]	2,087 (2.6%)	1,279 (4.5%)	42 (38, 46)	47 (43, 51)

CI = confidence interval.
Adjusted models were adjusted for age, baseline date, sex and whether the individual was born in Sweden.
[a]Of which 488,441 were vaccinated and 165,626 never vaccinated.
[b]Of which 469,627 were vaccinated and 152,669 never vaccinated.
[c]Of which 354,304 were vaccinated and 133,044 never vaccinated.
[d]Of which 309,804 were vaccinated and 100,029 never vaccinated.
[e]Of which 183,462 were vaccinated and 62,452 never vaccinated.
[f]Of which 79,161 were vaccinated and 28,612 never vaccinated.

597 **Figure legends**

598 **Figure 1.** Selection of study participants.

```
┌─────────────────────────────────────────────────┐
│      Data from the Public Health Agency of Sweden: │
│  All individuals vaccinated against COVID-19 or with a │
│          documented SARS-CoV-2 infection,          │
│               born 2003-2009                       │
│                  (N=692,546)                       │
└─────────────────────────────────────────────────┘
                        │
                        ▼
┌─────────────────────────────────────────────────┐          ┌──────────────────────────┐
│  To these individuals, Statistics Sweden matched one │          │  N=134 individuals not    │
│  control individual from the total population of     │ ──────▶  │  vaccinated with an mRNA  │
│  individuals born 2003-2009. Matched controls were not │          │  vaccine were excluded    │
│  vaccinated at the date of vaccination in the          │          └──────────────────────────┘
│  corresponding case, and did not have a documented    │
│  infection on the date of the corresponding case. One │
│  control could be matched to several cases.           │
│                  (N=844,511)                        │
└─────────────────────────────────────────────────┘
```

Safety analysis:
All individuals given at least one dose of mRNA vaccine **(N=645,355)** in the cohort was compared to all individuals not given any vaccine **(N=199,022)**. Outcomes were all diagnoses set during hospital stay after vaccination, and after a corresponding date in the controls.

Vaccine effectiveness analysis:
After excluding individuals with a documented previous SARS-CoV-2 infection at baseline, all individuals given at least two doses of mRNA vaccine (**N=501,945**) was compared to all individuals not given any vaccine (**N=170,083**). The outcome was COVID-19 hospitalisation from 7 days after the second dose of vaccine onwards, and a corresponding date in the controls.