Title: Global Menstrual Cycle Symptomatology as Reported by Users of a Menstrual Tracking

Mobile Application

Alessandra J Ainsworth, MD [1]

Kimberly Peven, PhD[2]

Ryan Bamford, MSc[2]

Liudmila Zhaunova, PhD[2]

Rodion Salimgaraev, MD[2]

Carley Prentice, BA[2]

Aidan Wickham, PhD[2]

Adam Cunningham, PhD[2]

Frederick Goddard, PhD[2]

Sonia Ponzo, PhD[2, 3]

Samir N Babayev, MD [1]

14
15 [1] Department of Obstetrics and Gynecology, Division of Reproductive Endocrinology and

16 Infertility, Mayo Clinic, Rochester, Minnesota

17 [2] Flo Health UK Limited, London, United Kingdom

18 [3] University College London, Institute of Health Informatics

19
Address Correspondence to: Samir N. Babayev, MD

Mayo Clinic

200 First Street SW

Rochester, MN 55905

(507) 284-9792

Babayev.samir@mayo.edu

Abstract

Background: Although a shared experience throughout the world, menstrual symptomatology is

underreported and often misunderstood, by both individuals who menstruate and their providers.

This study aimed to investigate symptomatology throughout the menstrual cycle, including the

frequency of symptoms reported for each phase of the cycle and the relationship between

symptoms.

Methods: This study included self-reported cycle information and symptoms during ovulatory

menstrual cycles in mobile application users. Symptom frequency was described overall, by

phase, and by day. Relationships between symptoms were examined through correlation

coefficients and term frequency-inverse document frequency analysis.

Results: 437,577 unique users and 896,051 unique cycles were included in the analysis. Somatic

symptoms were most frequently reported, logged in 88.3% of cycles. Of the total symptoms

logged, somatic, gastrointestinal, and negative mood symptoms were most prevalent in the late

luteal and bleeding days. In contrast, positive mood and discharge were most reported in the

fertile window. Mood symptoms were highly correlated with each other (energetic mood and

happy mood $r=0.62$, anxious mood and obsessive thoughts $r=0.60$). Fatigue was highly

correlated with somatic and gastrointestinal symptoms such as bloating ($r=0.62$), headache

($r=0.60$), backache ($r=0.58$), and nausea ($r=0.55$).

Conclusions: Menstrual cycle-related symptoms are very common and vary throughout the

menstrual cycle. While negative symptoms, including pain, bloating, and negative mood, are

most commonly reported, women also report positive symptoms, such as energetic and happy

mood. Self-reported data from cycle tracking and health apps can improve our understanding of

the prevalence and variability of menstrual-related symptoms.

Trial Registration: Not applicable.

52
53   **Key Words**: Menstrual symptoms, Menstruation, Period Tracking, Mobile Application

54
55   **Plain English Summary**: Our study includes data collected from users of a menstrual tracking

56   application. We found that menstrual symtpoms are common, throughout the menstrual cycle,

57   and include both positive and negative mood and physical symptoms. This large data set may

58   help to normalize menstrual symptoms and provide a basis for future research and understanding.

59   **Background:**
60   Although a shared experience by women and people who menstruate throughout the

world, menstruation represents an area of health often ignored and misunderstood by both

patients and healthcare providers. While many individuals experience menstruation without

significant symptoms, others endure pain and many other somatic, gastrointestinal, and mood

symptoms which vary throughout the cycle. Dysmenorrhea affects 45-95% of menstruating

individuals, with wide variation due to differing definitions of the condition (1). Compared with

pain-free phases of their cycles, those suffering from dysmenorrhea experience reduced quality

of life, poor mood, and decreased sleep during menstruation (1). Gastrointestinal symptoms also

vary throughout the menstrual cycle, with one-third of women experiencing symptoms during

menstruation (2), most frequently on the first day of bleeding (3).

Additionally, irritability and mood swings can affect 22-40% of women, more often

before menstruation (4). As it relates to symptom co-occurrence, individuals with mood

symptoms like depression and anxiety are more likely to experience multiple gastrointestinal

symptoms both before and during menstruation (5). Despite improved standardization in

terminology and increasing prioritization of women's health issues, these symptoms are under

reported (6–8). Importantly, menstrual symptoms may present a significant impairment to

productivity and quality of life (9).

A recent review highlighted challenges in the field of menstrual health research (10).

These challenges include how and from whom data is collected, normalization of symptoms,

stigma regarding menstrual disorders, lack of standardized assessment tools, and differing

cultural perceptions of this physiologic process (10–13). Previous studies of symptom prevalence

and variation throughout the menstrual cycle are small prospective or survey studies (4,5). Given

these limitations, there remain gaps in our knowledge of patient symptoms and opportunities for

further study.

The emerging availability of mobile applications (apps) and user-reported data provides

a novel approach to population health, in this instance, menstrual symptomatology, through

'digital epidemiology' (14–16). The impact of digital epidemiology to inform our understanding

of population norms in menstrual health recently demonstrated a luteal phase length of 12.4 ± 2

days (17). These results differ from the classical teaching of a conserved 14-day luteal phase

(17).

In this study, through in-app user-reported data, we aim to report a population-based

assessment of symptomatology throughout the menstrual cycle. Specific objectives include

describing the frequency of symptoms reported in each phase of the menstrual cycle and analysis

of the relationship between symptoms.

Methods:

This cohort study includes active users of the Flo app recording symptoms during an ovulatory

cycle starting between January 1, 2021, and January 1, 2023. Users aged 18-45, not pregnant,

and actively involved in the app (reporting at least three menstrual cycles) were included.

Menstrual cycles between 21-35 days, menstrual bleeding between 1-10 days, follicular and

luteal phase lengths of at least five days each, positive ovulation predictor kit use, and at least

one self-reported menstrual symptom at any point in the cycle were included as 'normal'

menstrual cycles.

Cycle dates and phases were calculated based on user-provided data. Users report the

start and end dates of their period and the date of positive ovulation test. Each cycle start date is

defined as the first day of bleeding. The mid-follicular phase was defined as the day after the last

day of menstrual bleeding until the start of the fertile window (Figure 1). An individual with less

than five days between the end of menstrual bleeding and the day of ovulation had no defined

mid-follicular phase. The fertile window was calculated as the five days preceding ovulation and

the day of ovulation was estimated as the day following a user-reported positive ovulation

predictor kit. The luteal phase was split evenly in half and described as the early luteal phase and

the late luteal phase.

Symptoms were self-reported and associated with the phase of the menstrual cycle as

defined above. We classified symptoms into five groups: somatic (cramps, tender breasts,

headache, fatigue, backache, acne), gastrointestinal (GI) (bloating, nausea, diarrhea, cravings,

constipation), positive mood (happy, high sex drive, energetic), negative mood (mood swings,

anxious, depressed, insomnia, stress, obsessive thoughts), and discharge (creamy, eggwhite,

sticky, watery). Users could record as many symptoms in each category as desired each day.

Individual characteristics were matched to each cycle. Age at the time of the cycle was

based on the user's birthdate and cycle dates. A user's country was assigned based on phone

settings at app installation. Where the country at app installation was unavailable, country phone

settings at the last login was used. Country prevalence of symptoms was evaluated only when

100 users or more were included.

Statistical Methods

Using a cycle-level unit of analysis, we employed descriptive statistics to describe cycle

length and symptoms experienced overall and by phase. For cycles with a length of 25-30 days,

we describe symptoms experienced by day of the cycle, where day 1 is the first day of bleeding.

We examined the correlation between all possible pairs of symptoms using a cycle-level unit of

analysis considering symptoms experienced at any point in the cycle. We reported Pearson

correlation coefficients for each possible pair. We also conducted a term frequency-inverse

document frequency (tf-idf) analysis (18) for pairs of symptoms separately by cycle phase to

describe the frequency of pairs of any symptoms, taking into account the frequency of the

individual symptoms that make up the pair. This is calculated as term frequency times inverse

document frequency, where term frequency is defined as the number of instances of a pair of

symptoms in a cycle phase, and inverse document frequency is defined as the logarithm of the

number of users logging one symptom from the pair during the relevant cycle phase divided by

the total number of symptoms logged in the phase. Data were analyzed in Python Jupyter

Notebook 6.0.1 and R version 4.2.0.

Results:

A total of 437,577 unique users and 896,051 unique, normal menstrual cycles met

inclusion criteria with an average of 2.05 ± 1.8 cycles per user (Figure 2). The mean age of users

was 30.0 ± 5.4 years; users under age 18 or over 45 were excluded from the analysis. Users were

included from 222 countries, with 113 countries having at least 100 users. The largest group of

users was in the United States (n=126,352), followed by Germany (n=45,018) and the United

Kingdom (n=35,007).

Of the 437,577 unique users included in this analysis (last cycle), the mean cycle length

was 28.4 ± 2.6 days (median 28.0 days), and the mean length of bleeding was 4.0 ± 1.1 days

(median 4.0 days). Most included cycles had a length of 25-31 days (75.7%, n =331,360),

followed by a length of $\geq$32 days (15.8%, n=69,479) and <25 days (8.4 %, n =36,738).

Calculated from the predicted day of ovulation, the mean follicular phase length amongst users

was 14.4 ± 3.0 days (median 14.0 days), and the mean luteal phase length was 14.0 ± 2.7 days

(median 14.0 days).

Of the 896,051 unique cycles, somatic symptoms were reported most often; 88.3%

(n=790,923) of cycles reported at least one somatic symptom. Comparatively, gastrointestinal

symptoms and discharge symptoms were reported in 60.8% of cycles each (n=545,107 and

544,460, respectively), negative mood symptoms were reported in 43.3% (n=388,249) of cycles,

and positive mood symptoms were reported in 35.4% (n=317,558). The type and frequency of

symptoms are shown in Figure 3. Menstrual cramps, breast tenderness, headache, and fatigue

were the most common symptoms reported.

Symptoms were next assessed by the menstrual cycle phase. Somatic, gastrointestinal,

and negative mood symptoms were most frequent in the late luteal phase and least frequent in the

mid-follicular phase (Figure 4). 57.3% of all logged instances of tender breasts occurred during

the late luteal phase compared to 1.8% in the mid-follicular phase. Discharge and positive mood

symptoms were most common during the fertile window. 63.1% of all logged instances of egg

white discharge and 45.3% of all logged instances of high sex drive were during the fertile

window.

For cycles between 25-30 days in length (n=570,123 cycles), we examined daily

symptom logs for the three symptoms with the most variability per category (Figure 5). Overall,

most symptoms are reported on day 1 of the cycle (436,812 symptoms logged from 211,021

cycles out of 570,123 total cycles); however, the fertile window is where most cycles have at

least one symptom logged. Out of 570,123 total cycles, 382,115 cycles had at least one symptom

logged during the fertile window, and 283,652 had at least one symptom logged during the

bleeding days. The prevalence of somatic, gastrointestinal, and negative mood symptoms are

highest on day 1 but decline quickly, reaching their lowest point around day 7 before slowly

increasing throughout the rest of the cycle. For positive mood symptoms, the prevalence of

energetic mood remains relatively stable throughout the cycle. In contrast, happy and high-sex-

drive moods increase slowly in the early part of the cycle, peaking around day 14 of the cycle

and reaching baseline around day 20. Discharge symptoms are rarely logged during the early part

of the cycle where bleeding occurs. Egg white discharge increases rapidly between days 6 and 13

before decreasing quickly to a low level by day 20.

For countries with at least 100 users, the number per country ranged from 103 in The

Bahamas to 126,352 in the United States (Supplemental Table 1). The prevalence of cramps

during the last cycle ranged from 30.5% in Vietnam to 74.6% in Iceland (Figure 6). Bloating

ranged from 17.6 in Thailand to 49.7% in Chile. Mood swings ranged from 13.3% in Hong Kong

to 43.1% in Iraq.

Mood symptoms were most highly correlated with each other. The anxious mood was

highly correlated with obsessive thoughts (r=0.60) and depressed mood (r=0.58). Depressed

mood was correlated with obsessive thoughts (r=0.55), and happy mood was highly correlated

with energetic mood (r=0.62, Figure 7). Similarly, fatigue was highly correlated with somatic

and gastrointestinal symptoms such as bloating (r=0.62), headache (r=0.60), backache (r=0.58),

and nausea (r=0.55).

A term frequency-inverse document frequency (tf-idf) analysis by cycle phase showed

how pairs of symptoms correlate within cycle phases, adjusting for the frequency of individual

symptoms (Figure 8 and Supplementary Tables 2 and 3). During the bleeding days and late luteal

phase when cramps are common, the top five tf-idf values all included cramps, and cramps were

most closely associated with nausea (tf-idf = 54,861 bleeding days, tf-idf = 68,467 late luteal).

Happy and energetic were the top pair in the mid-follicular phase (tf-idf=23,333) and second in

the fertile window (tf-idf=55,636).

Discussion:

In this study, we report the frequency of somatic, gastrointestinal, mood, and discharge

symptoms experienced throughout the menstrual cycle by a cohort of global, ovulating period-

tracking app users. Notably, most users were aged 30-34 and located in Europe and the United

States. Our results build on prior work evaluating the menstrual cycle characteristics of women

using period-tracking apps. The median cycle length, 28.8 days in our study, is consistent with

previous studies and period tracking app data (17,19–22).

Somatic symptoms were reported most frequently, with cramps being the most reported

symptom. Somatic, gastrointestinal, and negative mood symptoms were more common in the

late luteal phase than in the rest of the cycle. They reached a nadir in the mid-follicular phase

between bleeding and the fertile window/ovulation. The hormonal changes observed in the luteal

phase may explain these findings. There is a direct correlation between the rapid rise of

progesterone, the dominant sex hormone in the luteal phase, and the abrupt increase in somatic

symptoms. Progesterone has been shown to influence breast changes and associated breast

tenderness in the luteal phase (23). By thermoregulatory actions, progesterone has also been

shown to impact circadian rhythms leading to decreased rapid eye movement (REM) sleep in the

luteal phase and, thus a reported increase in fatigue (24). However, progesterone levels fall in the

week prior to menstruation while somatic symptoms continue to increase. Symptoms in the late

luteal phase reflect the effect of decreasing systemic sex steroid hormones. Pain disorders, such

as migraines and non-migraine headaches, have been shown to increase in the late luteal phase

when estradiol levels are lowest (25–27), in line with our findings of headaches being most

reported in this phase. These effects are modulated by estrogen effects on central pain processing

networks and cerebral vasculature (26). Conversely, decreasing progesterone levels and the

resultant increase in available androgen receptors may explain the increase in acne reported in

the late luteal phase (28).

Gastrointestinal symptoms that increase throughout the luteal phase likely reflect multiple

mechanisms (3). First, the effect of progesterone on serotonergic activity in both the central

nervous system and the GI tract slows gastrointestinal motility leading to constipation in the

early luteal phase (29). Second, increasing prostaglandin levels in the late luteal phase lead to

stimulation of smooth muscle contractions – leading to both uterine cramping and increased

gastrointestinal motility and associated diarrhea (30).

The effect of the menstrual cycle on mental health has been well studied, and

premenstrual mood disorders are well classified, with over 300 subtypes reported. Mood changes

in the luteal phase affect 50-80% of reproductive-aged women, although 3-8% of individuals

have such severe symptoms with associated functional impairments that they meet diagnostic

criteria in the DSM-V (31). Our study confirmed an increase in negative mood symptoms

throughout the luteal phase, possibly reflecting known alterations in sex steroids and the

following neurotransmitters: serotonin, noradrenaline, and gamma-aminobutyric acid (GABA)

(31,32). Negative mood symptoms decreased throughout the follicular phase, with a

corresponding increase in positive mood symptoms that reached peak levels in the late follicular

phase. Positive mood symptom frequency in the follicular phase is consistent with prior studies

reporting individual well-being to be highest at this time in the cycle (33,34). Recent work

utilizing functional magnetic resonance imaging (fMRI) has added to our understanding. Sex

hormones appear to impact communication between brain regions, in addition to their known

impact on neurotransmitter secretion (35). This impacts cognition and emotional processing,

such that estrogen seems to decrease reaction to negative stimuli while progesterone seems to

increase emotional reactions to negative stimuli (36).

The relationship between mood and somatic symptoms during the menstrual cycle has

been evaluated with mixed results. One study found an increase in mood symptoms in women

with gastrointestinal symptoms, skin changes, breast changes, and abdominal bloating. At the

same time, there was no effect of concurrent mood symptoms seen in women who reported back

and joint pain (37). We found negative mood symptoms (depressed mood, mood swings, anxious

mood) co-occurred with cramps in the bleeding days and late luteal phase while positive mood

symptoms (energetic and happy) occurred together in the mid-follicular through early luteal

phases, without strong associations with somatic symptoms.

We found small geographic differences in user-reported symptoms during the menstrual

cycle. This may reflect the impact of knowledge or sociocultural constructs, which impact

individual perceptions of symptoms and their menstrual experience (38). Historically,

menstruation has been described negatively. Previous work has shown an association between

early menstrual/menarchal experiences and current menstrual experience, body perception, and

overall health (39–42). However, other studies examining symptom experience across broad

geographic regions have not found regional differences in symptoms (43).

The strength of this study lies in the number of unique users and in the large amount of

user-reported data from a global population. However, this study is not without limitations.

Specifically, this study is subject to selection bias as not all menstruating individuals choose to

use period-tracking apps. Demographic and other individual characteristic data were not

collected, including user-reported race and ethnicity, which may be important mediators of

menstrual symptom characteristics and experiences that may impact our results. We included

only ovulatory cycles where ovulation was confirmed by self-ovulation test to more accurately

divide the cycle into phases. Users logging ovulation tests likely have a goal of conception and

may thus have a different experience of menstrual symptoms and a difference in the tendency of

logging symptoms. Additionally, irregular and non-ovulatory cycles are not represented in this

analysis, and results are not generalizable to these populations. Although this study represents a

global menstruating population, many countries remain absent or underrepresented, presenting

an opportunity for future research. Other limitations include the use of self-reported symptom-

tracking data. Users log symptoms through the app but may not log symptoms consistently

throughout the cycle (e.g., potentially more likely to log symptoms when logging other events

such as ovulation or period) and may not log all symptoms experienced. Users also may be more

likely to log symptoms that appear at the top and front of the app over symptoms that appear

lower down or require scrolling to see. This decreases the representativeness of symptoms

recorded and limits our ability to compare results with other methods of symptom data

collection.

Conclusions:

Our study presents an expanded understanding of symptoms experienced during the

menstrual cycle. These findings reflect the value of population-level menstrual cycle data, which

can help inform individual patients and clinicians while strengthening research efforts and calls

for advocacy for Women's Health.

284
Abbreviations:

GI: gastrointestinal

Tf-idf: term frequency-inverse document frequency

REM: rapid eye movement

DSM-V: Diagnostic and Statistical Manual of Mental Disorders, 5th edition

GABA: gamma-aminobutyric acid

fMRI: functional magnetic resonance imaging

Declarations:

1. IRB approval waived. Documentation attached as supporting document

2. Consent for publication: Not applicable

3. Availability of data and materials: The datasets used and/or analyzed during the current

study are available from the corresponding author on reasonable request.

4. Competing interests:

KP: Flo Health UK Limited (Employment, ownership interest, stock/stock options)

RB: Flo Health UK Limited (Employment)

LZ: Flo Health UK Limited (Employment, ownership interest, stock/stock options)

RS: Flo Health UK Limited (Employment, ownership interest, stock/stock options)

CP: Flo Health UK Limited (Employment)

AW: Flo Health UK Limited (Employment, ownership interest, stock/stock options)

AC: Flo Health UK Limited (Employment)

FG: Flo Health UK Limited (Employment)

SP: Flo Health UK Limited (Employment, ownership interest, stock/stock options).

Other authors did not report any potential conflicts of interest.

5. Author contributions:

AJA: Design, interpretation of data, drafted and substantially revised work

KP: Design; acquisition, analysis, and interpretation of data; drafted and substantially

revised work

RB: Acquisition, analysis, and interpretation of data

LZ: Design, interpretation of data, and substantively revised work

RS: Design; acquisition, analysis, and interpretation of data

CP: Analysis and interpretation of data

AW: Acquisition, analysis, and interpretation of data

AC: Acquisition, analysis, and interpretation of data

FG: Analysis and interpretation of data

SP: Interpretation of data and substantively revised work

SB: Design, interpretation of data, substantially revised work

6. Acknowledgements: None

References

1. Lacovides S, Avidon I, Baker FC. What we know about primary dysmenorrhea today: a critical
review. Hum Reprod Update. 2015 Nov 1;21(6):762–78.
2. Moore J, Barlow D, Jewell D, Kennedy S. Do gastrointestinal symptoms vary with the menstrual
cycle? Br J Obstet Gynaecol. 1998 Dec;105(12):1322–5.
3. Judkins TC, Dennis-Wall JC, Sims SM, Colee J, Langkamp-Henken B. Stool frequency and
form and gastrointestinal symptoms differ by day of the menstrual cycle in healthy adult women
taking oral contraceptives: a prospective observational study. BMC Womens Health. 2020 Jun
29;20(1):136.
4. Lete I, Häusler G, Pintiaux A, Jamin C, Nappi RE, Fiala C, et al. The inconvenience due to
women's monthly bleeding (ISY) survey: a study of premenstrual symptoms among 5728
women in Europe. Eur J Contracept Reprod Health Care Off J Eur Soc Contracept. 2017
Oct;22(5):354–9.
5. Bernstein MT, Graff LA, Avery L, Palatnick C, Parnerowski K, Targownik LE. Gastrointestinal
symptoms before and during menses in healthy women. BMC Womens Health. 2014 Jan
22;14:14.
6. Chen CX, Draucker CB, Carpenter JS. What women say about their dysmenorrhea: a qualitative
thematic analysis. BMC Womens Health. 2018 Mar 2;18(1):47.
7. Chen CX, Kwekkeboom KL, Ward SE. Beliefs About Dysmenorrhea and Their Relationship to
Self-Management. Res Nurs Health. 2016 Aug;39(4):263–76.
8. Ramos-Pichardo JD, Ortega-Galán ÁM, Iglesias-López MT, Abreu-Sánchez A, Fernández-
Martínez E. Why Do Some Spanish Nursing Students with Menstrual Pain Fail to Consult
Healthcare Professionals? Int J Environ Res Public Health. 2020 Nov 5;17(21):E8173.

9. Schoep ME, Adang EMM, Maas JWM, De Bie B, Aarts JWM, Nieboer TE. Productivity loss
due to menstruation-related symptoms: a nationwide cross-sectional survey among 32 748
women. BMJ Open. 2019 Jun 27;9(6):e026186.

10. Critchley HOD, Babayev E, Bulun SE, Clark S, Garcia-Grau I, Gregersen PK, et al.
Menstruation: science and society. Am J Obstet Gynecol. 2020 Nov;223(5):624–64.

11. Matteson KA, Zaluski KM. Menstrual Health as a Part of Preventive Health Care. Obstet
Gynecol Clin North Am. 2019 Sep;46(3):441–53.

12. Chen CX, Shieh C, Draucker CB, Carpenter JS. Reasons women do not seek health care for
dysmenorrhea. J Clin Nurs. 2018 Jan;27(1–2):e301–8.

13. Tan DA, Haththotuwa R, Fraser IS. Cultural aspects and mythologies surrounding menstruation
and abnormal uterine bleeding. Best Pract Res Clin Obstet Gynaecol. 2017 Apr;40:121–33.

14. Symul L, Wac K, Hillard P, Salathé M. Assessment of menstrual health status and evolution
through mobile apps for fertility awareness. NPJ Digit Med. 2019;2:64.

15. Mao L, Xi S, Bai W, Yao C, Zhou Y, Chen X, et al. Menstrual patterns and disorders among
Chinese women of reproductive age: A cross-sectional study based on mobile application data.
Medicine (Baltimore). 2021 Apr 23;100(16):e25329.

16. Karasneh RA, Al-Azzam SI, Alzoubi KH, Muflih SM, Hawamdeh SS. Smartphone Applications
for Period Tracking: Rating and Behavioral Change among Women Users. Obstet Gynecol Int.
2020;2020:2192387.

17. Bull JR, Rowland SP, Scherwitzl EB, Scherwitzl R, Danielsson KG, Harper J. Real-world
menstrual cycle characteristics of more than 600,000 menstrual cycles. NPJ Digit Med.
2019;2:83.

18. Leskovec J, Rajaraman A, Ullman JD. Mining of Massive Data Sets. Cambridge University
Press; 2020. 566 p.

19. Li K, Urteaga I, Wiggins CH, Druet A, Shea A, Vitzthum VJ, et al. Characterizing physiological
and symptomatic variation in menstrual cycles using self-tracked mobile-health data. NPJ Digit
Med. 2020;3:79.

20. Najmabadi S, Schliep KC, Simonsen SE, Porucznik CA, Egger MJ, Stanford JB. Menstrual
bleeding, cycle length, and follicular and luteal phase lengths in women without known
subfertility: A pooled analysis of three cohorts. Paediatr Perinat Epidemiol. 2020
May;34(3):318–27.

21. Li K, Urteaga I, Shea A, Vitzthum VJ, Wiggins CH, Elhadad N. A predictive model for next
cycle start date that accounts for adherence in menstrual self-tracking. J Am Med Inform Assoc
JAMIA. 2021 Dec 28;29(1):3–11.

22. Broad A, Biswakarma R, Harper JC. A survey of women's experiences of using period tracker
applications: Attitudes, ovulation prediction and how the accuracy of the app in predicting period
start dates affects their feelings and behaviours. Womens Health Lond Engl. 2022
Dec;18:17455057221095246.

23. Graham JD, Clarke CL. Physiological action of progesterone in target tissues. Endocr Rev. 1997
Aug;18(4):502–19.

24. Shechter A, Varin F, Boivin DB. Circadian variation of sleep during the follicular and luteal
phases of the menstrual cycle. Sleep. 2010 May;33(5):647–56.

25. LeResche L, Mancl L, Sherman JJ, Gandara B, Dworkin SF. Changes in temporomandibular
pain and other symptoms across the menstrual cycle. Pain. 2003 Dec;106(3):253–61.

26. Sacco S, Ricci S, Degan D, Carolei A. Migraine in women: the role of hormones and their
impact on vascular diseases. J Headache Pain. 2012 Apr;13(3):177–89.

27. Stening K, Eriksson O, Wahren L, Berg G, Hammar M, Blomqvist A. Pain sensations to the cold pressor test in normally menstruating women: comparison with men and relation to menstrual phase and serum sex steroid levels. Am J Physiol Regul Integr Comp Physiol. 2007 Oct;293(4):R1711-1716.

28. Raghunath RS, Venables ZC, Millington GWM. The menstrual cycle and the skin. Clin Exp Dermatol. 2015 Mar;40(2):111–5.

29. Xiao ZL, Pricolo V, Biancani P, Behar J. Role of progesterone signaling in the regulation of G-protein levels in female chronic constipation. Gastroenterology. 2005 Mar;128(3):667–75.

30. Evans J, Salamonsen LA. Inflammation, leukocytes and menstruation. Rev Endocr Metab Disord. 2012 Dec;13(4):277–88.

31. Ryu A, Kim TH. Premenstrual syndrome: A mini review. Maturitas. 2015 Dec;82(4):436–40.

32. Toffol E, Kalleinen N, Urrila AS, Himanen SL, Porkka-Heiskanen T, Partonen T, et al. The relationship between mood and sleep in different female reproductive states. BMC Psychiatry. 2014 Jun 16;14:177.

33. Sanders D, Warner P, Bäckström T, Bancroft J. Mood, sexuality, hormones and the menstrual cycle. I. Changes in mood and physical state: description of subjects and method. Psychosom Med. 1983 Dec;45(6):487–501.

34. Meaden PM, Hartlage SA, Cook-Karr J. Timing and severity of symptoms associated with the menstrual cycle in a community-based sample in the Midwestern United States. Psychiatry Res. 2005 Mar 30;134(1):27–36.

35. Toffoletto S, Lanzenberger R, Gingnell M, Sundström-Poromaa I, Comasco E. Emotional and cognitive functional imaging of estrogen and progesterone effects in the female human brain: a systematic review. Psychoneuroendocrinology. 2014 Dec;50:28–52.

36. Sakaki M, Mather M. How reward and emotional stimuli induce different reactions across the menstrual cycle. Soc Personal Psychol Compass. 2012 Jan 1;6(1):1–17.

37. Kiesner J, Pastore M. Day-to-day co-variations of psychological and physical symptoms of the menstrual cycle: insights to individual differences in steroid reactivity. Psychoneuroendocrinology. 2010 Apr;35(3):350–63.

38. Marván ML, Escobedo C. Premenstrual symptomatology: role of prior knowledge about premenstrual syndrome. Psychosom Med. 1999 Apr;61(2):163–7.

39. McPherson ME, Korfine L. Menstruation across time: menarche, menstrual attitudes, experiences, and behaviors. Womens Health Issues Off Publ Jacobs Inst Womens Health. 2004 Dec;14(6):193–200.

40. Stubbs ML. Cultural perceptions and practices around menarche and adolescent menstruation in the United States. Ann N Y Acad Sci. 2008;1135:58–66.

41. Padmanabhanunni A, Jaffer L, Steenkamp J. Menstruation experiences of South African women belonging to the ama-Xhosa ethnic group. Cult Health Sex. 2018 Jun;20(6):704–14.

42. Barrington DJ, Robinson HJ, Wilson E, Hennegan J. Experiences of menstruation in high income countries: A systematic review, qualitative evidence synthesis and comparison to low- and middle-income countries. PloS One. 2021;16(7):e0255001.

43. Dennerstein L, Lehert P, Heinemann K. Global epidemiological study of variation of premenstrual symptoms with age and sociodemographic factors. Menopause Int. 2011 Sep;17(3):96–101.

Bleeding days **Mid folicular** **Fertile window** **Early luteal** **Late luteal**

	Acne	Anxious	Backache	Bloating	Constipation	Cramps	Cravings	Creamy Discharge	Depressed	Diarrhea	Eggwhite Discharge	Energetic	Fatigue	Happy	Headache	High Sex Drive	Insomnia	Mood Swings	Nausea	Obsessive Thoughts	Sticky Discharge	Stress	Tender Breasts	Watery Discharge
Acne																								
Anxious	0.35																							
Backache	0.41	0.35																						
Bloating	0.45	0.39	0.51																					
Constipation	0.34	0.29	0.38	0.48																				
Cramps	0.35	0.29	0.5	0.49	0.33																			
Cravings	0.38	0.34	0.39	0.48	0.35	0.33																		
Creamy Discharge	0.33	0.26	0.34	0.38	0.27	0.38	0.27																	
Depressed	0.31	0.58	0.31	0.32	0.26	0.25	0.3	0.21																
Diarrhea	0.27	0.24	0.32	0.35	0.24	0.32	0.26	0.23	0.2															
Eggwhite Discharge	0.25	0.2	0.27	0.29	0.22	0.31	0.21	0.4	0.16	0.18														
Energetic	0.26	0.27	0.23	0.28	0.2	0.21	0.27	0.3	0.19	0.18	0.24													
Fatigue	0.48	0.45	0.58	0.62	0.43	0.49	0.48	0.4	0.37	0.37	0.31	0.29												
Happy	0.3	0.29	0.28	0.32	0.22	0.27	0.27	0.39	0.18	0.21	0.28	0.62	0.37											
Headache	0.44	0.37	0.54	0.49	0.36	0.49	0.38	0.35	0.33	0.32	0.28	0.25	0.6	0.31										
High Sex Drive	0.24	0.23	0.27	0.27	0.21	0.25	0.29	0.27	0.2	0.18	0.23	0.3	0.3	0.32	0.27									
Insomnia	0.32	0.34	0.36	0.4	0.36	0.3	0.38	0.24	0.31	0.27	0.19	0.2	0.46	0.2	0.37	0.21								
Mood Swings	0.4	0.46	0.42	0.44	0.32	0.36	0.42	0.34	0.44	0.26	0.26	0.3	0.5	0.31	0.44	0.31	0.35							
Nausea	0.33	0.3	0.47	0.46	0.34	0.47	0.35	0.29	0.27	0.33	0.23	0.18	0.55	0.22	0.5	0.25	0.33	0.36						
Obsessive Thoughts	0.29	0.6	0.3	0.32	0.26	0.22	0.33	0.2	0.55	0.19	0.16	0.22	0.36	0.21	0.3	0.2	0.3	0.42	0.25					
Sticky Discharge	0.29	0.24	0.28	0.32	0.23	0.3	0.24	0.43	0.19	0.19	0.33	0.26	0.34	0.33	0.3	0.23	0.21	0.29	0.23	0.19				
Stress	0.29	0.47	0.28	0.28	0.22	0.2	0.27	0.2	0.4	0.19	0.15	0.19	0.34	0.2	0.3	0.19	0.27	0.38	0.2	0.43	0.19			
Tender Breasts	0.32	0.22	0.39	0.4	0.28	0.46	0.3	0.33	0.19	0.23	0.27	0.2	0.42	0.23	0.39	0.22	0.25	0.31	0.38	0.17	0.26	0.15		
Watery Discharge	0.25	0.22	0.29	0.31	0.23	0.31	0.23	0.37	0.17	0.21	0.33	0.23	0.32	0.28	0.29	0.22	0.2	0.27	0.26	0.17	0.27	0.16	0.26	

Percentage of symptom occurrence during each phase