Investigating inequalities in HIV testing in sub-Saharan Africa: insights from a spatial analysis of 25 countries

Pearl Anne Ante-Testard[1,2][§], Gabriel Carrasco-Escobar[3,4], Tarik Benmarhnia[4], Laura Temime[1,2], Kévin Jean[1,2,5]

1 Laboratoire MESuRS, Conservatoire national des Arts et Métiers, Paris, France
2 Unité PACRI, Institut Pasteur, Conservatoire national des Arts et Métiers, Paris, France
3 Health Innovation Laboratory, Institute of Tropical Medicine "Alexander Von Humboldt", Universidad Peruana Cayetano Heredia, Lima, Peru
4 Scripps Institution of Oceanography, University of California, San Diego, California, USA
5 MRC Centre for Global Infectious Disease Analysis, Department of Infectious Disease Epidemiology, Imperial College London, United Kingdom

[§] Corresponding author: Pearl Anne Ante-Testard

Laboratoire MESuRS, Conservatoire national des Arts et Métiers
292 rue Saint Martin, 75003 Paris, France
Email: pearlannemante@gmail.com or pearl.ante@ucsf.edu

E-mail addresses of authors:

P.A.A.-T.: pearlannemante@gmail.com or pearl.ante@ucsf.edu

G.C.-E.: gabriel.carrasco@upch.pe

T.B.: tbenmarhnia@ucsd.edu

L.T.: laura.temime@lecnam.net

K.J.: kevin.jean@lecnam.net

Keywords: socioeconomic inequality, HIV, HIV testing, spatial analysis, sub-Saharan Africa, social epidemiology

Word count:

Abstract: 349/350

Main text: 3471/3500

Abstract

Introduction:

We aim to explore spatial variations in socioeconomic inequalities in self-reported recent HIV testing uptake in sub-Saharan Africa (SSA) at different geographical scales, in order to identify potential geographical hotspots of inequalities. Additionally, to evaluate the potential benefits of HIV testing programs, we assess whether local levels of HIV testing match the local levels of HIV prevalence.

Methods:

We analyzed data from 25 countries in SSA with Demographic and Health Surveys between 2011 and 2019. We quantified socioeconomic inequalities in self-reported HIV testing in the last 12 months with both the Slope Index of Inequality (SII) and Relative Index of Inequality (RII) in different geographical scales to capture sex-specific within-country spatial variations. We also conducted sampling cluster-level analyses based on the Local Indicator of Spatial Association to consider the autocorrelation in SII and RII across SSA countries. To assess the spatial efficiency of HIV testing programs, we measured the correlation between recent HIV testing uptake and HIV prevalence through Pearson correlation across geographical scales.

Results:

We observed pro-rich inequalities on both absolute and relative scales in recent HIV testing in the majority of SSA countries in female and male participants at the national level. We also identified inequalities at subnational levels. Within- and between-country heterogeneities in sex-specific inequalities on both inequality scales and their respective spatial distributions varied depending on the scale used. Clustering of high absolute and relative inequalities were mostly observed in Western and Central Africa with a few regions in Eastern and Southern Africa. Despite significant sex-specific correlations between HIV testing and HIV prevalence in all countries when assessed at the national level, we report an absence of such a correlation at fine scale in 39 of 50 sex-country combinations.

Conclusions:

These findings highlight the importance of investigating the spatial variability of various HIV

indicators and related inequalities. Results may help local, national and international

policymakers to prioritize areas and socioeconomic subgroups in need of HIV testing services.

Our results also show the need to monitor efficiency of HIV testing programs in relation to HIV

risk at subnational levels as a complementary to national estimates.

Introduction

The role of HIV testing in the fight against HIV/AIDS is crucial since it is the gateway to HIV prevention and care, especially in sub-Saharan Africa (SSA), the global epicenter of the disease. Over the years, testing has scaled up due to the increasing availability of antiretroviral therapy (ART) and in response to the UNAIDS 90-90-90 and 95-95-95 targets by 2020 and 2030, respectively [1,2].

Socioeconomic inequalities have been reported in HIV testing uptake in SSA. Numerous studies have found that people in higher socioeconomic position (SEP) were more likely to seek HIV testing or know their HIV status [3–8]. However, most of these studies assessed testing inequalities at the national level. Very few studies have analyzed the spatial distribution of these inequalities. To the best of our knowledge, such a local analysis has only been performed in a single country, Malawi [9].

Spatial analysis at local scales has proved useful for the control and prevention of infectious diseases, by uncovering spatial variations in the transmission or the access to prevention [10–12]. Regarding HIV, it has helped identify high transmission areas [13] and understand access difficulties to healthcare facilities in underserved areas in Africa [14]. However, it has not been frequently utilized in monitoring inequalities in the HIV response, particularly in HIV testing. Notably, uptake of HIV testing has been reported to be higher in countries with the greatest HIV burden at the national level. Nevertheless, whether this association also exists at a finer scale has not been empirically assessed. Observing such spatial variations is important not only for ensuring equity in epidemic control but also for prioritizing areas with the greatest burden in terms of infection and/or inequalities [15]. Mapping HIV testing uptake and their inequalities across different geographical scales and identifying their local hotspots has become especially relevant in a context of decreasing international funding for the HIV response.

115 Here, we aim: i) to explore spatial variations in socioeconomic inequalities in HIV testing uptake
116 across geographical scales; ii) to identify geographical hotspots for such inequalities in several
117 SSA countries; and iii) to assess spatial correlations between HIV testing and HIV prevalence at
118 the national and subnational levels as a way to approximate the potential benefits of existing
119 programs.

Methods

Study design and data sources

We carried out a multi-country analysis of cross-sectional surveys in SSA countries, namely the
Demographic and Health Surveys (DHS). The DHS are nationally representative surveys
regularly conducted in the Global South collecting information on a broad range of indicators
such as sociodemographic indicators, maternal and child health, malaria and HIV/AIDS. They
are based on a two-stage sampling design, with Primary Sampling Units (PSU) selected in the
first stage and households in the second. Women aged 15-49 years and men aged 15-54/59 years
in participating households are eligible. Depending on the survey, data for men for the HIV
indicators, biomarkers or both might may be collected only from a sub-sample of the selected
households. Some DHS include HIV serological surveys in which participants are asked for
consent to be tested for HIV, which is done anonymously in most of the surveys. Individuals
who consented are interviewed face-to-face by trained interviewers who use a standard
questionnaire. The agencies and research institutes that conducted the surveys were responsible
for acquiring ethical clearance in each country.

DHS Global Positioning System coordinates were obtained from the DHS database. These
coordinates were intentionally and randomly displaced to ensure confidentiality of the
respondents. Urban clusters were displaced between 0 and 2 kilometers (km), while rural clusters
were displaced between 0 and 5 km. The spatial boundaries were also obtained from the
DHS spatial database repository (https://spatialdata.dhsprogram.com/home/). The DHS data
were linked to these spatial data.

For our analysis, we selected SSA countries with available DHS surveys between 2011 and
2019 that contained the variables of interest and spatial data. We selected the more recent survey
in countries with more than one eligible survey.

Data/ Variables

SEP was defined as the relative rank of the participants in the cumulative distribution of the DHS
wealth index. The wealth index is a composite measure of living standards based on household
assets and living characteristics. The DHS divided the wealth index homogenously into quintiles
from poorest to richest.

The outcome of interest was the self-report of recent (< 12 months) HIV testing uptake.
Participants were asked if they recently had an HIV test and the time since last test. Being HIV
positive was defined as testing positive in the serological survey.

Statistical analysis

First, for each country and sex, we calculated the HIV prevalence and the proportion of recent
HIV testing uptake while accounting for survey design and sampling weights at different
geographical scales: i) national, ii) first administrative subnational level (hereafter, province),
and iii) PSU level (hereafter referred to as "fine scale").

Second, we measured national-, province-, and PSU-level socioeconomic inequalities both on the
absolute and relative scales. We estimated the Slope Index of Inequality (SII) and the Relative
Index of Inequality (RII) to assess the absolute and relative inequalities, respectively [16]. It is
highly recommended to report inequalities on both scales as conclusions may diverge depending
on the scale used [17].

At the national and province levels, both indicators were obtained by fitting a modified Poisson

regression (with robust variance) with a log link function [18] to estimate the association

between recent HIV testing at each wealth level and the hierarchical ranking of wealth.

Generalized estimating equations were used to account for the clustering of observations [19].

The SII represents the absolute difference in the predicted proportions between the richest and

the poorest participants, whereas the RII expresses the ratio of the predicted outcomes between

these two extremes.

Due to the smaller sample sizes at the PSU-level (at least 10 individuals), we fitted a linear

regression to estimate fine-scale inequality indicators. When performed on small samples,

regressions may lead to negative RII values, which we truncated to zero when estimating the

fine-scale RII.

Third, spatial autocorrelations of the fine-scale SII and RII across SSA and between sexes were

assessed using the local *Getis-Ord Gi** statistic for PSUs with a sample size of at least 10. The

Getis-Ord Gi* statistic identifies local patterns and clusters of high- or low-inequality across

countries that may not be evident when using global statistics [20].   More specifically,

a distance-based neighborhood structure was used for its computation. Neighboring PSUs were

defined based on the distance *d* that assigns a number of nearest neighbors (*k*) to each PSU. We

selected the number of nearest neighbors that gave high spatial autocorrelation based on a global

Moran's I statistic for each sex and inequality indicator. We categorized the Gi* statistic based

on the sign (cold- or hotspot for negative and positive signs, respectively) and percentile (90%,

95%, 99%) to avoid bias due to multiple and dependent tests [21].

We assessed the spatial correlation of the proportion of recent HIV testing and level of HIV

prevalence across various geographical scales through Pearson correlation. Indeed, HIV

prevalence drives, at least partly, the local risk of incident HIV infection as it reflects the

probability for one's sexual partner to be infected by HIV [22].

Sensitivity analysis

SII and RII were calculated for PSUs with a sample size of at least 20 and 30 and their local
spatial autocorrelations were assessed as sensitivity analysis. We also conducted similar analyses
for countries with surveys between 2011 and 2014 and for countries with surveys between 2015
and 2019 to assess possible temporal trends in the spatial distribution of inequalities.

Results

Study population characteristics

Twenty-five countries were eligible based on data availability between 2011 and 2019: Angola,
Burundi, Cameroon, Chad, Côte d'Ivoire, Democratic Republic of Congo (Congo DR), Ethiopia,
Gabon, Ghana, Guinea, Lesotho, Liberia, Malawi, Mali, Mozambique, Namibia, Rwanda,
Senegal, Sierra Leone, Tanzania, Togo, Uganda, South Africa, Zambia, and Zimbabwe. Table 1
shows the summary statistics by country and sex. There was a total of 472,763 participants
(311,652 women and 161,111 men) with 351,921 individuals (74.4%, 252,508 women and
99,413 men) from PSUs with a sample size of at least 10 and complete data (Table S1). The
sample size in the provinces ranged between 275 and 11,342 among women and between 135
and 3, 236 among men. At a finer scale, sample size in PSUs ranged between 10 and 96 women
and between 10 and 54 men. The distributions of PSU sample size and proportion of recent HIV
testing   are   shown   in   Figure   S1.

216   **Table 1. Summary estimates of national level HIV prevalence, HIV testing, absolute and relative inequalities in HIV testing uptake in the**

217   **previous 12 months in 25 sub-Saharan African countries between 2011 and 2019 by country and sex.**

Country	Year	N	HIV prevalence	HIV testing[†]	SII (x 100)[‡] (95% CI)	RII (95% CI)	N	HIV prevalence	HIV testing[†]	SII (x 100)[‡] (95% CI)	RII (95% CI)
				Female					Male		
AO Angola	2015-16	14379	3%	29%	40 (37; 44)	4.07 (3.54; 4.69)	5684	1%	20%	26 (21; 31)	3.23 (2.63; 3.97)
BI Burundi	2016-17	17269	1%	30%	7 (4; 10)	1.26 (1.15; 1.39)	7552	0.7%	20%	7 (4; 11)	1.42 (1.19; 1.69)
CD Congo DR	2013-14	18827	2%	9%	19 (16; 22)	18.29 (12.83; 26.08)	8656	0.5%	7%	16 (13; 19)	12.83 (8.47; 19.43)
CI Côte d'Ivoire	2011-12	5183	6%	4%	6 (3; 9)	6.04 (3.10; 11.75)	4503	3%	4%	4 (2; 6)	4.34 (2.11; 8.95)
CM Cameroon	2018	14677	4%	41%	40 (36; 44)	2.68 (2.43; 2.96)	6978	2%	35%	42 (36; 47)	3.05 (2.65; 3.52)
ET Ethiopia	2019	15 683	1%	21%	44 (39; 48)	4.73 (4.13; 5.42)	12688	0.5%	19%	35 (31; 39)	4.11 (3.52; 4.79)
GA Gabon	2012	8 422	6%	33%	16 (12; 20)	1.68 (1.45; 1.94)	5654	3%	24%	23 (20; 27)	3.52 (2.80; 4.42)
GH Ghana	2014	9 396	3%	14%	11 (8; 13)	2.11 (1.74; 2.56)	4388	1%	6%	11 (8; 15)	5.75 (3.25; 10.15)
GN Guinea	2018	10 874	2%	8%	23 (19; 26)	11.09 (8.20; 14.98)	4117	1%	6%	15 (11; 20)	12.25 (7.17; 20.95)
LB Liberia	2019	9 239	2%	20%	5 (1; 8)	1.26 (1.06; 1.49)	4118	2%	13%	14 (10; 17)	3.28 (2.30; 4.67)
LS Lesotho	2014	6 621	30%	59%	-10 (-14; -6)	0.85 (0.78; 0.91)	2931	19%	38%	23 (17; 30)	1.86 (1.57; 2.20)
ML Mali	2018	10 424	1%	6%	26 (21; 30)	22.66 (16.15; 31.78)	4399	0.9%	6%	20 (15; 25)	14.74 (8.89; 24; 44)

Country	Year	N	%	%	SII	RII	N	%	%	SII	RII
MW *Malawi*	**2015-16**	**2 4562**	11%	44%	1 (-2; 3)	1.02 (0.96; 1.08)	7478	7%	43%	-0.2 (-4; 4)	0.996 (0.91; 1.09)
MZ *Mozambique*	**2015**	**7 749**	15%	30%	27 (22; 32)	2.08 (1.82; 2.38)	5283	10%	19%	32 (26; 37)	3.35 (2.77; 4.04)
NM *Namibia*	**2013**	**10 018**	17%	51%	-4 (-8; -1)	0.91 (0.85; 0.99)	4481	11%	38%	17 (12; 22)	1.57 (1.36; 1.81)
RW *Rwanda*	**2014-15**	**13 497**	4%	39%	3 (-0.2; 7)	1.08 (0.99; 1.18)	6217	2%	37%	-3 (-7; 1)	0.92 (0.82; 1.04)
SL *Sierra Leone*	**2019**	**15 574**	2%	24%	12 (9; 16)	1.74 (1.50; 2.02)	7197	1%	13%	20 (16; 24)	4.93 (3.71; 6.56)
SN *Senegal*	**2017**	**16 787**	0.5%	14%	5 (3; 7)	1.47 (1.25; 1.74)	6977	0.4%	6%	8 (6; 11)	4.48 (3.02; 6.65)
TD *Chad*	**2014-15**	**17 719**	2%	9%	5 (3; 6)	6.82 (4.34; 10.72)	5248	1%	7%	16 (11; 20)	9.09 (5.58; 14.80)
TG *Togo*	**2013-14**	**9 480**	3%	17%	19 (15; 22)	3.15 (2.58; 3.85)	4476	2%	12%	21 (17; 25)	6.90 (4.84; 9.84)
TZ *Tanzania*	**2011-12**	**10 967**	6%	33%	12 (8; 15)	1.44 (1.29; 1.61)	8352	4%	28%	13 (9; 17)	1.61 (1.41; 1.85)
UG *Uganda*	**2011**	**12 153**	8%	12%	5 (2; 7)	1.48 (1.18; 1.85)	9588	6%	12%	5 (2; 7)	1.46 (1.17; 1.82)
ZA *South Africa*	**2016**	**8 514**	28%	61%	-8 (-12; -4)	0.87 (0.82; 0.93)	3618	14%	46%	9 (3; 14)	1.21 (1.06; 1.38)
ZM *Zambia*	**2018**	**13 683**	14%	66%	9 (5; 13)	1.15 (1.07; 1.22)	12132	8%	53%	14 (10; 18)	1.31 (1.22; 1.41)
ZW *Zimbabwe*	**2015**	**9 955**	17%	51%	-4 (-7; -1)	0.92 (0.86; 0.99)	8396	11%	37%	7 (3; 11)	1.19 (1.08; 1.32)
Average			8%	29%				5%	22%		

219 N: Total number of participants; SII: Slope index of inequality; RII: Relative index of inequality; 95% CI: 95% Confidence Interval.

220 † Self-reported recent (< 12 months) uptake of HIV testing.

221 ‡ SII values and their 95% CIs were multiplied by 100 so that they may be easily interpreted as percentage point differences (instead of proportion
222 differences).

National-level estimates

Table 1 also presents the national estimates of HIV prevalence, testing, and inequalities in recent HIV testing. Overall, at the national level, HIV prevalence ranged from 0.5% (Senegal) to 30% (Lesotho) among women and from 0.4% (Senegal) to 19% (Lesotho) among men. Self-reported uptake of recent testing ranged from 4% (Côte d'Ivoire) to 66% (Zambia) among women and from 4% (Côte d'Ivoire) to 53% (Zambia) among men. Women tended to have higher HIV prevalence and proportion of recent testing than men.

On the absolute scale, we observed pro-rich absolute inequalities in recent testing in 19 of 25 countries for women and 23 of 25 countries for men (SII > 0) (Table 1). Absolute inequalities ranged between -8 (95% Confidence Interval [95% CI] -12; -4) percentage points (% points) in South Africa and 44 (39; 48) % points in Ethiopia among women. This means that the absolute difference between the richest and poorest quintiles was -8 (-12; -4) % points (i.e., pro-poor) among women in South Africa and 44 (39; 48) % points in Ethiopia (i.e., pro-rich). Meanwhile, among men, SII ranged between -3 (-7; 1) % points in Rwanda and 42 (0.36; 0.47) % points in Cameroon.

We observed pro-rich relative inequalities in 18 of 25 countries for women and 23 of 25 countries for men (RII > 1) (Table 1). Relative inequalities ranged between 0.85 (0.78; 0.91) in Lesotho and 22.66 (16.15; 31.78) in Mali among women. This translates to the richest women being 0.85 (0.78; 0.91) times as likely to report recent HIV testing than the poorest participants in Lesotho (i.e., pro-poor), while 22.66 times (16.15; 31.78) more likely in Mali (i.e., pro-rich). Among men, it ranged between 0.92 (0.82; 1.04) in Rwanda and 14.74 (8.89; 24.44) in Mali.

Province-level estimates

The distribution of province-level HIV prevalence and proportion of recent HIV testing are mapped in Figure 1. We observed within- and between-country variations in their respective spatial distributions.

Figure 1. Province-level distribution of weighted HIV prevalence among A) female and B) male participants and weighted percentage of self-reported recent (<12 months) uptake of HIV testing among C) female and D) male participants. Dark grey colours indicate unavailability of the HIV biomarker. Missing polygons within the country indicate no data is available for this province from the Demographic and Health Surveys.

Figure 2 maps the provincial absolute and relative inequalities in recent HIV testing. We also observed spatial heterogeneities of these inequalities for all sexes and inequality scales. On the absolute scale, we observed pro-rich spatial distribution of SIIs in most of the provinces in SSA except for a few areas in Eastern and Southern Africa (ESA) such as South Africa, Namibia, and Malawi. On the relative scale, higher pro-rich relative inequalities were observed more frequently in Western and Central Africa (WCA), while lower inequalities tended to be observed in ESA.

Figure 2. Province-level distribution of wealth-related inequalities in self-reported recent (<12 months) uptake of HIV testing at the A) and C) absolute scale and B) and D) relative scale among female and male participants, respectively, across 25 sub-Saharan African countries. SII, Slope Index of Inequality; RII, Relative Index of Inequality. Capped RII values between 0.1 and 300. Missing polygons in Chad represent provinces with extreme RII values (> 300 for females) and (< 0.1 for males).

Spatial clustering analysis at fine scale

Global Moran's I showed that using 1 or 2 nearest neighbors gave the highest spatial autocorrelation for both sexes and inequality indicators (Figure S2). For uniformity, we used *k=2* to calculate for the Getis-Ord Gi* statistics. Hotspots and coldspots of inequalities across SSA depended on the inequality scale used and sex (Figure 3). Overall, hotspots on both scales were more marked in WCA and a few ESA countries such as Ethiopia (for both sexes), Mozambique and Tanzania (only on relative scales for both sexes).

Figure 3. Local spatial autocorrelation of socioeconomic inequalities in self-reported recent (<12 months)
uptake of HIV testing as Local Getis-Ord Gi* at Primary Sampling Unit (PSU) level (two nearest neighbors)
across 25 sub-Saharan African countries between 2011 and 2019. Spatial clustering at the absolute scales among
A) and C) absolute scales and B) and D) relative scales among female and male participants, respectively. Only
PSUs with a sample size of at least 10 and more than one wealth quintile were included. LISA, Local Indicator of
Spatial Autocorrelation; SII, Slope Index of Inequality; RII, Relative Index of Inequality.

On the absolute scale, we observed numerous pockets of high SII values in western Africa with a
few areas in eastern Africa. Hotspots of absolute inequalities in recent HIV testing were
observed in Cameroon, Ghana, Togo, and Ethiopia for both sexes. Coldspots or pockets of low
SIIs were mostly observed in ESA including Zambia, Zimbabwe, Burundi, Rwanda, and small
areas in Mozambique for both sexes and in South Africa for women.

On the relative scale, we observed hotspots of RII values mostly in WCA with a few areas in
eastern Africa. Coldspots were noted in ESA including Burundi and Rwanda both sexes and in
South Africa for women.

300   Moreover, we observed diverging patterns of hotspots and coldspots for each sex in the same
301   country. There were hotspots of RIIs in Uganda among women, while coldspots in a few areas
302   were noted among men. Contrastingly, areas in Namibia displayed coldspots among women,
303   while hotspots were noted among men.

Spatial correlation between HIV testing and HIV prevalence

We assessed the spatial correlation between HIV testing and HIV prevalence across geographical
scales – whether testing services are reaching those with high HIV risk in the population. At the
national level, HIV prevalence and proportion of recent HIV testing were found to be positively
correlated among women and men (Figure S3).

However, this observation was not sustained at subnational levels. Within-country correlation of
province-level HIV prevalence and proportion of testing showed that in many countries, both
variables were uncorrelated (i.e., the level of HIV testing did not always match the magnitude of
HIV prevalence). Out of 50 settings (2 sex-specific results for each of the 25 countries), we
observed only 11 that had statistically significant positive correlations (p-value < 0.05) or what
we termed as having "efficient HIV testing services" (Figure S4): both sexes (Côte d'Ivoire,
Ethiopia, and Tanzania), females (Rwanda and Zambia) and males (Liberia, Lesotho, and Sierra
Leone).

Similarly, Figure 4 shows that at the PSU level, HIV prevalence did not correlate with the level
of recent HIV testing in many of the countries for both sexes. Only in Burundi (female),
Mozambique (female), Namibia (male), Tanzania (both sexes), and Zambia (both sexes) did we
observe a significant positive correlation. Meanwhile, in Lesotho, HIV prevalence and recent
testing had a significant negative correlation (i.e., PSUs with higher HIV prevalence had lesser
uptake of recent HIV testing).

Figure 4. Correlation (Pearson correlation coefficient R and p-value) between the proportion of HIV prevalence and proportion of self-reported recent (< 12 months) uptake of HIV testing at the Primary Sampling Unit (PSU) level in 25 sub-Saharan African countries stratified by sex. Only included PSUs with a sample size equal or higher than 10 with both the HIV biomarker and HIV testing variables. There were not sufficient clusters to calculate the Pearson correlation R and p-values in Chad (TD). Each point represents a PSU. Refer to Table 1 for full country names.

Sensitivity analysis

Results of the local spatial clustering analysis were consistent when sub-setting by cluster size of at least 20 or 30 (Figures S5 and S6). Areas with pockets of high and low inequalities were also consistent when conducting spatial clustering analysis across countries with surveys between 2011 and 2014 and between 2015 and 2019 separately (Figures S7 and S8).

Discussion

In this study, we quantified and mapped absolute and relative socioeconomic inequalities of recent HIV testing at different geographical scales. We also conducted spatial clustering analysis of such inequalities and explored the spatial correlation between HIV testing and HIV prevalence at various geographical scales across SSA. Our results show existing inequalities at the national, province, and PSU levels. Heterogeneities in the spatial distribution of these inequalities at subnational levels and hotspot areas varied depending on the inequality scales and sex groups. Most hotspots of inequalities were observed in WCA with a few areas in ESA such as Ethiopia, Mozambique, and Tanzania. Meanwhile, coldspots were rather observed in ESA. We also revealed that, while HIV testing programs seemed efficient in reaching those with high risk of acquiring HIV at the national level, they seemed to be less efficient at the subnational levels in most of the countries. Indeed, the provinces and PSUs with higher recent testing uptake did not match the level of HIV prevalence in many of the countries for both sexes.

We tended to observe higher HIV prevalence and proportion of recent HIV testing among women and countries located in ESA. As expected, we report higher pro-rich inequalities among men and in WCA countries. These findings were consistent with previous studies [5,7]. Higher HIV burden among women may be explained by higher vulnerability than men in SSA due to several biological, societal, and economic factors [23].

We also highlighted that recent HIV testing was not shared equally across wealth levels in SSA (within and between countries) and that such inequalities were not randomly distributed across space. Indeed, our results also showed varying spatial patterns of recent HIV testing inequalities between absolute and relative scales and between women and men. This highlights the necessity for HIV testing programs to be tailored depending on the level of inequality to be addressed and the needs of each sex. We observed that in a few countries like Namibia, national-level inequality estimates showed low relative inequality for women and even pro-poor inequality on the absolute scale for both sexes suggesting that pro-rich inequalities are not a fatality and might be overcome. Nevertheless, hotspots of inequalities were still observed in some areas within

these countries that achieved equitable HIV testing levels. This suggests that national-level
inequality estimates may hide remaining pro-rich inequalities at lower scales.

However, one could argue that pro-rich inequalities are not necessarily unfair, especially if those
with higher SEP also tend to be the ones who are more at-risk of acquiring HIV. Several early
studies reported higher levels of HIV prevalence among higher SEP [24,25]. However, this
initial inverse social gradient, an unusual feature as compared to most of the diseases, is likely in
a process of reversion. Last-generation HIV prevalence surveys now report higher levels of HIV
prevalence among the poorest in some countries such as Côte d'Ivoire, Ethiopia, Lesotho, and
Zimbabwe. Moreover, a recent cohort study relying on a two-decade follow-up in rural Uganda
documented a changing socioeconomic gradient over time, with a higher risk of incident HIV
infection among the poorest [26]. Thus, suggesting that these pro-rich inequalities might indeed
be unfair and contributing to the reversing social gradient that is probably ongoing in some
settings.

There are various criteria for health programs that can be used to evaluate their public benefit
which include efficiency. Efficiency is concerned with the optimal production and distribution of
scarce health resources and is critical for sustainability and maximizing health gains [27]. In this
study, we did not conduct a formal impact evaluation of HIV testing programs, but rather to
investigate the spatial correlations between HIV testing and HIV prevalence across geographical
scales to capture their potential benefits. Community-level HIV prevalence has been found to be
a strong predictor of HIV incidence in a recent meta-analysis [22]. The seemingly sub-optimal
efficiency of HIV testing programs at subnational levels may suggest the failure of HIV
programs in some settings to reach those who are at higher risk of HIV. However, risk may have
been affected when people with HIV (PWH) undergo ART which ensures viral suppression, thus
preventing transmission. Another potential reason why the level of HIV testing did not match the
level of HIV risk is that in settings with high prevalence, people with known HIV infection
(possibly under treatment) do not necessarily need to seek HIV testing. A further similar survey

might be useful to assess recent HIV testing among the at-risk population only (i.e., excluding

PWH under ART), however our data do not allow us to do so.

This study carries several limitations. First, the self-reported nature of HIV testing uptake may be

subject to recall and social desirability biases resulting to underreporting. Second, differential

accuracy in self-reporting between socioeconomic groups might have biased our results.

Evidence in cancer screening suggest that overreporting of self-reported screening is common

among marginalized groups such as racial minorities [28]. If this also applies to HIV testing, this

may have led to an under-estimation of the pro-rich inequalities and over-estimation of the pro-

poor inequalities. While self-reported lifetime HIV testing was found to be highly sensitive (96-

99%) [29], testing in the past 12 months may be prone to telescoping bias that may have led to

over-reporting [30]. Third, the wealth index can only measure relative wealth within a country.

However, it can measure long-term SEP and has also been found to be stable especially in the

Global South. Lastly, some available DHS surveys were conducted before 2014 and may not

have captured more recent patterns of inequalities.

Despite these limitations, to our knowledge, this is the first study to provide a comprehensive

context of the socioeconomic inequalities in HIV testing in SSA by quantifying and mapping

them at different geographical levels on both absolute and relative scales and by assessing the

spatial efficiency of HIV testing at different levels. This study revealed the importance of

monitoring inequalities at different geographical scales. First, national estimates are often use for

funding allocations by donors, prioritization of programs and comparison of inequality metrics.

Second, province-level estimates are essential for program implementation, and for within-

country funding allocations. Lastly, fine-level estimates allow us to visualize small-scale

heterogeneities to precisely target communities in need.

Conclusions

Our results show the need to monitor inequalities and assess the efficiency of HIV testing
services in reaching those who are at-risk of HIV at smaller geographical scales, beyond national
estimates that may mask disparities. By providing estimates of such inequalities at national-,
province-, and PSU- level, and by localizing their hotspots, these findings may help
policymakers, local and international organizations to prioritize areas and groups that need HIV
testing efforts, while increasing efficiency.

Competing interests

The authors declare no competing interests.

Authors' contributions

P.A.A.-T., L.T. and K.J. conceived and discussed the study with inputs from G.C.-E. and T.B.
P.A.A.-T. collated and processed the DHS. P.A.A.-T. conducted the analysis with inputs from
G.C.-E., T.B., L.T. and K.J. P.A.A.-T. produced output figures and tables with inputs from G.C.-
E, T.B., L.T. and K.J. All authors contributed to the interpretation of the results. P.A.A.-T. wrote
the initial draft with inputs from L.T. and K.J. All authors contributed to subsequent revisions.
All authors read and approved the final version of the manuscript.

Acknowledgements

Funding: INSERM-ANRS (France Recherche Nord and Sud Sida-HIV Hépatites), grant number
ANRS-12377 B104.

Disclaimer: Funding agency had no role in the study design, data collection and analysis.

Additional files

Additional file 1: **Supplementary material**

Description:

Table S1. Total number of participants in Primary Sampling Units that have at least 10 samples.

Figure S1 (A). PSU distribution among women A) sample size and B) proportion of self-

reported uptake of recent (< 12 months) HIV testing.

Figure S1 (B). PSU distribution among men A) sample size and B) proportion of self-reported

uptake of recent (< 12 months) HIV testing.

Figure S2 (A). Global Moran's I statistic test by number of nearest neighbor (1-10) by inequality

scales among women.

Figure S2 (B). Global Moran's I statistic test by number of nearest neighbor (1-10) by gender

and inequality scales.

Figure S3. Correlation (Pearson's correlation coefficient R and p-value) between weighted HIV

prevalence and weighted self-reported recent (< 12 months) uptake of HIV testing at the national

level in 25 sub-Saharan African countries by gender.

Figure S4. Correlation (Pearson's correlation coefficient R and p-value) between weighted HIV

prevalence and weighted self-reported recent (< 12 months) uptake of HIV testing at the regional

level in 25 sub-Saharan African countries by gender.

Figure S5. Sensitivity analysis of local spatial autocorrelation of socioeconomic inequalities in

self-reported (<12 months) uptake of HIV testing as Local Getis-Ord Gi* at Primary Sampling

Unit (PSU) level (2 nearest neighbors) with at least 20 participants across sub-Saharan African

countries.

Figure S6. Sensitivity analysis of local spatial autocorrelation of socioeconomic inequalities in

self-reported (<12 months) uptake of HIV testing as Local Getis-Ord Gi* at Primary Sampling

Unit (PSU) level (2 nearest neighbors) with at least 20 participants across sub-Saharan African

countries.

Figure S7. Sensitivity analysis of local spatial autocorrelation of HIV testing socioeconomic

inequalities as Local Getis-Ord Gi* at PSU level (two nearest neighbors) across sub-Saharan

African countries with surveys between 2011 and 2014.

Figure S8. Sensitivity analysis of local spatial autocorrelation of HIV testing socioeconomic inequalities as Local Getis-Ord Gi* at PSU level (two nearest neighbors) across sub-Saharan African countries with surveys between 2015 and 2019.

References

1. UNAIDS. Fast-Track: Ending the AIDS Epidemic by 2030 [Internet]. 2014. Available from: https://www.unaids.org/sites/default/files/media_asset/JC2686_WAD2014report_en.pdf

2. UNAIDS. 90-90-90 An ambitious treatment target to help end the AIDS epidemic [Internet]. 2014. Available from: Available from: https://www.unaids.org/sites/default/files/media_asset/90-90-90_en.pdf

3. Ante-Testard PA, Benmarhnia T, Bekelynck A, Baggaley R, Ouattara E, Temime L, et al. Temporal trends in socioeconomic inequalities in HIV testing: an analysis of cross-sectional surveys from 16 sub-Saharan African countries. Lancet Glob Health. 2020 Jun;8(6):e808–18.

4. Awopegba OE, Ologunowa TO, Ajayi AI. HIV testing and self□testing coverage among men and women in South Africa: an exploration of related factors. Trop Med Int Health. 2021 Feb;26(2):214–27.

5. Hamidouche M, Ante-Testard PA, Baggaley R, Temime L, Jean K. Monitoring socioeconomic inequalities across HIV knowledge, attitudes, behaviours and prevention in 18 sub-Saharan African countries. AIDS. 2022 May 1;36(6):871–9.

6. Jean K, Anglaret X, Moh R, Lert F, Dray-Spira R. Barriers to HIV Testing in Côte d'Ivoire: The Role of Individual Characteristics and Testing Modalities. Braitstein P, editor. PLoS ONE. 2012 Jul 18;7(7):e41353.

7. Kim SW, Skordis-Worrall J, Haghparast-Bidgoli H, Pulkki-Brännström AM. Socio-economic inequity in HIV testing in Malawi. Glob Health Action. 2016 Dec 1;9(1):31730.

8. Kirakoya-Samadoulougou F, Jean K, Maheu-Giroux M. Uptake of HIV testing in Burkina Faso: an assessment of individual and community-level determinants. BMC Public Health. 2017 Dec;17(1):486.

9. Kim SW, Haghparast-Bidgoli H, Skordis-Worrall J, Batura N, Petrou S. A method for measuring spatial effects on socioeconomic inequalities using the concentration index. Int J Equity Health. 2020 Dec;19(1):9.

10. Rice BL, Golden CD, Randriamady HJ, Rakotomalala AANA, Vonona MA, Anjaranirina EJG, et al. Fine-scale variation in malaria prevalence across ecological regions in Madagascar: a cross-sectional study. BMC Public Health. 2021 Dec;21(1):1018.

11. Carrasco-Escobar G, Fornace K, Benmarhnia T. Mapping socioeconomic inequalities in malaria in Sub-Sahara African countries. Sci Rep. 2021 Dec;11(1):15121.

12. Mosser JF, Gagne-Maynard W, Rao PC, Osgood-Zimmerman A, Fullman N, Graetz N, et al. Mapping diphtheria-pertussis-tetanus vaccine coverage in Africa, 2000–2016: a spatial and temporal modelling study. The Lancet. 2019 May;393(10183):1843–55.

13. Bulstra CA, Hontelez JAC, Giardina F, Steen R, Nagelkerke NJD, Bärnighausen T, et al. Mapping and characterising areas with high levels of HIV transmission in sub-Saharan Africa: A geospatial analysis of national survey data. Barnabas RV, editor. PLOS Med. 2020 Mar 6;17(3):e1003042.

14. Kim H, Musuka GN, Mukandavire Z, Branscum A, Cuadros DF. When distance matters: Mapping HIV health care underserved communities in sub-Saharan Africa. Eke R, editor. PLOS Glob Public Health. 2021 Nov 24;1(11):e0000013.

15. Stresman GH, Mwesigwa J, Achan J, Giorgi E, Worwui A, Jawara M, et al. Do hotspots fuel malaria transmission: a village-scale spatio-temporal analysis of a 2-year cohort study in The Gambia. BMC Med. 2018 Dec;16(1):160.

16. Moreno-Betancur M, Latouche A, Menvielle G, Kunst AE, Rey G. Relative Index of Inequality and Slope Index of Inequality: A Structured Regression Framework for Estimation. Epidemiology. 2015 Jul;26(4):518–27.

17. King NB, Harper S, Young ME. Use of relative and absolute effect measures in reporting health inequalities: structured review. BMJ. 2012 Sep 3;345(sep03 1):e5774–e5774.

18. Zou G. A Modified Poisson Regression Approach to Prospective Studies with Binary Data. Am J Epidemiol. 2004 Apr 1;159(7):702–6.

19. Yelland LN, Salter AB, Ryan P. Performance of the Modified Poisson Regression Approach for Estimating Relative Risks From Clustered Prospective Data. Am J Epidemiol. 2011 Oct 15;174(8):984–92.

20. Getis A, Ord JK. The Analysis of Spatial Association by Use of Distance Statistics. Geogr Anal. 2010 Sep 3;24(3):189–206.

21. Caldas de Castro M, Singer BH. Controlling the False Discovery Rate: A New Application to Account for Multiple and Dependent Tests in Local Statistics of Spatial Association. Geogr Anal. 2006 Apr;38(2):180–208.

22. Jia KM, Eilerts H, Edun O, Lam K, Howes A, Thomas ML, et al. Risk scores for predicting HIV incidence among adult heterosexual populations in sub□Saharan Africa: a systematic

review and meta☐analysis. J Int AIDS Soc [Internet]. 2022 Jan [cited 2022 Jul 27];25(1).
Available from: https://onlinelibrary.wiley.com/doi/10.1002/jia2.25861

23. Ramjee G, Daniels B. Women and HIV in Sub-Saharan Africa. AIDS Res Ther.
2013;10(1):30.

24. Hargreaves JR, Glynn JR. Educational attainment and HIV-1 infection in developing
countries: a systematic review. Trop Med Int Health. 2002 Jun;7(6):489–98.

25. Fox AM. The Social Determinants of HIV Serostatus in Sub-Saharan Africa: An Inverse
Relationship between Poverty and HIV? Public Health Rep. 2010 Jul;125(4_suppl):16–24.

26. Santelli JS, Chen I, Makumbi F, Wei Y, Nalugoda F, Lutalo T, et al. Household wealth and
HIV incidence over time, rural Uganda, 1994–2018. AIDS. 2021 Sep 1;35(11):1835–43.

27. Ding J, Hu X, Zhang X, Shang L, Yu M, Chen H. Equity and efficiency of medical service
systems at the provincial level of China's mainland: a comparative study from 2009 to
2014. BMC Public Health. 2018 Dec;18(1):214.

28. Burgess DJ, Powell AA, Griffin JM, Partin MR. Race and the validity of self-reported
cancer screening behaviors: Development of a conceptual model. Prev Med. 2009
Feb;48(2):99–107.

29. Xia Y, Milwid RM, Godin A, Boily MC, Johnson LF, Marsh K, et al. Accuracy of self-
reported HIV-testing history and awareness of HIV-positive status in four sub-Saharan
African countries. AIDS. 2021 Mar 1;35(3):503–10.

30. Rubin DC, Baddeley AD. Telescoping is not time compression: A model. Mem Cognit.
1989 Nov;17(6):653–61.