Exploring Language-independent Digital Speech Biomarkers of Hypokinetic Dysarthria

Daniel Kovac[1], Jiri Mekyska[1], Vered Aharonson[2,3], Pavol Harar[1,4], Zoltan Galaz[1], Steven Rapcsak[5], Juan Rafael Orozco-Arroyave[6,7], Lubos Brabenec[8], and Irena Rektorova[8,9]

[1] Department of Telecommunications, Brno University of Technology, Brno, Czech Republic
[2] Afeka Center for Language Processing, Afeka Tel Aviv Academic College of Engineering, Tel Aviv, Israel
[3] School of Electrical and Information Engineering, University of the Witwatersrand, Johannesburg, South Africa
[4] Data Science Research Network, University of Vienna, Vienna, Austria
[5] Department of Neurology, College of Medicine, University of Arizona, Tucson, USA
[6] Faculty of Engineering, Universidad de Antioquia – UdeA, Medellín, Colombia
[7] Pattern Recognition Lab, Friedrich-Alexander-Universität, Erlangen-Nürnberg, Erlangen, Germany
[8] Applied Neuroscience Research Group, Central European Institute of Technology – CEITEC, Masaryk University, Brno, Czech Republic
[9] First Department of Neurology, Faculty of Medicine and St. Anne's University Hospital, Masaryk University, Brno, Czech Republic

Abstract:
Motor speech disorders in patients with Parkinson's disease (PD), collectively referred to as hypokinetic dysarthria, are the early markers of the disease. Acoustic speech features are, therefore, suitable digital biomarkers for the diagnosis and monitoring of this pathological phenomenon. At the same time, it is clear that language plays an essential role in using these features to classify speech successfully into healthy and dysarthric one. This paper focuses on the multilingual analysis of the currently used and newly proposed acoustic speech features and their use in the supportive diagnosis of PD. The goal is to explore digital speech biomarkers of PD, focusing on their language independence and high discrimination power using statistical analysis and machine learning techniques. Thirty-three acoustic features of Czech, American, Israeli, Columbian and Italian PD patients and healthy controls were analyzed using correlation and statistical tests, descriptive statistics, and the XGBoost classifier with posterior explanation by features importances and Shapley values. The features quantifying the prominence of the second formant, monopitch, and the lower number of pauses detected during text reading show the best results, both by statistical analysis and machine learning. Classification accuracies range from 67% to 85%, depending on the language. The paper introduces the concept of language robustness as a quality in which a feature behaves the same, independently of the language. By using this new concept along with other metrics, it proposes several digital speech biomarkers that have the potential to be language-independent with high discrimination power.

Keywords:
hypokinetic dysarthria; Parkinson's disease; multilingual study; acoustic speech features; statistical analysis; machine learning

1 Introduction

Hypokinetic dysarthria (HD) refers to motor disorders manifested in respiration, phonation, articulation, resonances and prosody of the speech and has a major impact on the patient's communication ability [24, 30]. Characteristic features of a person's voice with HD include tremor [35], hoarseness [4] and a large proportion of audible breath [33]. Speech is further characterised by hypernasality [15], the repetitiveness of

syllables with their shortening or lengthening [11, 17] and inappropriate silences [12]. Overall, it is relatively unintelligible [34] and quiet [1], with poor intonation and only slight variations in volume [6]. These disorders most commonly occur in patients with neurodegenerative diseases such as Parkinson's disease (PD) or various types of dementia. In other cases, stroke, hypoxia, trauma or other causes may be involved in the development of HD [9]. In patients with PD, HD is caused by progressive degeneration of dopaminergic neurons in the substantia nigra [16] and is present in up to 90% of these people [14]. The risk of PD increases with age; the second factor is mainly genetic predisposition. It is also clear that the number of PD patients grows with increasing population and life expectancy [7]. Although the science has made significant advances since the disease was first described [28], we still do not know the actual cause of PD and are not able to cure it; we can only alleviate its development. Therefore, early detection and initiation of treatment are crucial to the future course of the disease [20]. Since HD occurs in the early phases of PD [29], acoustic speech analysis can be a suitable supportive tool for automatic diagnosis or objective monitoring of the disease. Although many teams have worked on this topic, there is still no comprehensive and robust set of acoustic features quantifying the speech disorders that can be applied in practice and capture and describe HD in all domains. One factor that significantly affects these features is the speaker's language.

In 2010, Whitehill TL [37] described that Chinese PD patients show many of the same patterns of speech abnormalities as English-speaking people with HD and suggested that there may be universal acoustic features that could distinguish between healthy and HD-affected speech.

Hazan et al. [13] then published the results of a multilingual study focused on the automatic diagnosis of PD patients speaking in English and German. To distinguish the affected speech from the healthy one, they chose the articulation features based on the formants. Using the support vector machine as the machine learning model, they were able to classify the speech with an accuracy of 85%, which dropped to 75% when they trained the model by the features of one language and tested it on the other one. In the summary of the article, they mention that features that can differentiate dysarthric speech from the healthy one probably vary with the language.

In 2016, Orozco-Arroyave et al. [25] performed multilingual experiments with Spanish, German and Czech recordings of the dysarthric speech. The prediction accuracy ranged from 60% to 99%, depending on the language combination and the ratio of training to test data. When classifying PD in one language only, the accuracy ranged from 85% to 99%. The most discriminatory features were Mel frequency cepstral coefficients (MFCC) and energy in the critical Bark bands (BBE), both extracted from unvoiced segments of the reading text.

Kim and Choi [18] published the results of their descriptive study in which they describe the differences in acoustic vowel space (AVS) of Korean- and English-speaking PD patients. No differences in articulation rate were observed.

Next, in 2019 Moro-Velazquez et al. [23] reached an accuracy between 85% and 94% when classifying PD patients in a multilingual cohort including Castillian Spanish, Colombian Spanish and Czech. It dropped to the range between 72% and 82% when cross-corpora validating. For this purpose, they used an approach based on phonemic grouping. The most significant phonemes for detecting the HD were plosives and fricatives. Extraction of the features from reading text led to better results than the quantitative analysis of a diadochokinetic task.

Vásquez-Correa et al. [36] used convolution neural nets and transfer learning strategy to classify PD in Spanish, German and Czech with MFCC and BBE as input independent variables. Accuracy ranged between 70% and 77%. More accurate and more balanced in the frame of sensitivity and specificity were models trained by features of more than one language.

Rusz et al. [31] performed a speech analysis of Czech, English, German, French and Italian speakers in the early phase of PD. From the sustained phonation of vowel [a], diadochokinetic task and monologue, they extracted seven features in total. Significant role in differentiating the speech into healthy and dysarthric one had the features quantifying monopitch, length of pauses and voice onset time (VOT). According to statistical analysis, there were no differences between language groups in monopitch and length of pauses.

In 2022, Ozbolt et al. [27] analysed machine learning models trained by phonation features of Spanish, English and Italian PD patients and healthy controls. These features mainly quantify energies in different parts of spectra of the speech signal recorded during the sustained phonation. The same features were differently important in English, Spanish and Italian models. During the cross-corpora validating, the accuracy was higher when the model was trained by Spanish data and tested on Italian rather than vice

versa. Different vowels of sustained phonation were significant for each scenario.

It is evident that the language has a non-negligible effect on the accuracy of classifying speakers into those who are healthy and those affected with HD, but no study has yet looked at multilingual analysis in depth. This work aims to explore language-independent digital speech biomarkers of Parkinson's disease. It seeks to determine which acoustic speech features are important for different languages in terms of classification accuracy and which are sufficiently robust and independent of the speaker's language.

2 Materials and Methods

The diagram in Figure 1 describes the workflow used to determine the robustness of acoustic speech features to language differences and their subsequent discrimination power. The methods are further specified in the following subsections.

Figure 1: Workflow.

2.1 Speech corpus

The corpus contains speech recordings of 506 people (265 healthy controls – HC, 241 PD) and is created by grouping several datasets described in Table 1:

- Czechs (CZ) including HIDI [2], PARCZ [10] and CoBeN (only HC) [19] data,

- Americans (US) speaking American English from CoBeN project [19],

- Israelis (IL) speaking Hebrew,

- Colombians (CO) speaking Spanish [26],

- Italians (IT) – freely available dataset [8].

Every participant signed an informed consent, and the relevant ethics committee approved the study. Their age distribution can bee seen in Figure 2 and clinical data of PD patients are summarized in Table 2. It describes the time since the first symptoms occurred, medication and severity of PD assessed on the Unified Parkinson's Disease Rating Scale, part III (UPDRS III) and Hoehn and Yahr speech rating scale (H&Y). All patients in the cohort are on dopaminergic medication (ON state). Since deep brain stimulation may affect a person's speech [3], recordings of patients with this stimulation were discarded from the Israeli dataset, resulting in a slight imbalance of classes that can also be observed in the US data. Two Italian recordings containing a buzz were also removed from the corpus. We analyzed the speech and voice of subjects recorded during a reading of a short text (Read), prolonged phonation of vowel [a], and a diadochokinetic task (DDK) consisting of repeating the syllables [pa]-[ta]-[ka]. The reading task in each dataset contains a different text written in the nation's corresponding language, and there are two different texts to read in the Czech dataset. In the Italian dataset, patients and healthy controls repeat only the syllable [pa] or the syllable [ta] as a second variant of the task. In the Czech and American acquisition, participants performed all tasks only once. In the Israeli dataset, participants had two trials of vowel [a], and in the Colombian one, three trials. The Italians had each task recorded twice. We resampled all recordings to a uniform 16 kHz sampling frequency.

Table 1: Numbers of PD patients, healthy controls, men (M) and women (F) in the corpus.

| | CZ | | | US | | | IL | | | CO | | | IT | | | |
	F	M	total	F	M	total	F	M	total	F	M	total	F	M	total	altogether
HC	46	35	81	13	5	18	53	43	96	25	25	50	10	10	20	265
PD	49	84	133	3	8	11	7	12	19	25	25	50	9	19	28	241
total	95	119	214	16	13	29	60	55	115	50	50	100	19	29	48	506

Figure 2: Probability distribution of age.

Table 2: Clinical data: mean and standard deviation of LED – L-dopa equivalent daily dose; UPDRS III – Unified Parkinson's Disease Rating Scale, part III (motor examination) and H&Y – Hoehn and Yahr speech rating scale.

	CZ	US	IL	CO	IT
PD duration [year]	12.2 ± 5.1	5.0 ± 2.1	14.1 ± 6.9	11.2 ± 9.8	U
LED [mg]	1001 ± 524	286 ± 222	U	U	U
UPDRS III	22.5 ± 11.5	U	U	37.6 ± 18.1	U
UPDRS III speech	U	U	U	1.3 ± 0.8	1.0 ± 1.2
H&Y	U	U	2.8 ± 0.9	2.2 ± 0.7	U

U – unknown value

2.2 Parametrization

We extracted 33 acoustic speech features from the recordings quantifying disorders associated with HD. Due to the different signal lengths recorded during vowel [a], we extracted phonatory features from a signal duration of 1.5 s without a vowel onset. In the case of more than one trial for the speech task, the resulting feature is the average of two or three values obtained from the recordings. These biomarkers divided into phonation involving respiration, articulation involving resonance, and prosody involving pausing are listed and described in Table 3. This table also describes the expected change in the feature of people with PD. The expectation is based on experience or general knowledge of HD [22, 21, 5]. It is related to the desire to ensure sufficient clinical interpretability, and it also allows us to observe whether a feature behaves as expected in a particular dataset. After parametrization, we regressed out the effect of confounding factors such as age and gender.

2.3 Statistical analysis

According to the Shapiro-Wilk test, not all features were normally distributed; therefore, we used the Mann-Whitney U test to compare the distributions of HC and PD features in each dataset. Spearman's rank-correlation coefficients then inform us about the negative or positive correlation of the feature with the duration of PD and clinical scores. This test was performed for the data of all languages together. P-values of all tests were corrected via the False Discovery Rate (FDR) approach. We also looked at the number of patients deviating from the norm given by the distribution of the HC feature. We set the lower T_{low} and upper T_{up} thresholds to:

$$T_{\mathrm{low}} = Q_1 - 1.5 \cdot IQR, \tag{1}$$

$$T_{\mathrm{up}} = Q_3 + 1.5 \cdot IQR, \tag{2}$$

where Q_1 and Q_3 are lower and upper quartiles and IQR is the interquartile range. These outliers give us other information about the strength of the feature to detect the HD and show how the feature changes with the disease. Finally, we compared the medians of PD and HC features.

2.3.1 Feature robustness

From the results of the statistical analyses, we determined the language robustness of the features. By language robustness, the exact behaviour of the feature in all datasets is meant. A feature is determined to be robust if it satisfies the following conditions:

1. Significantly discriminate PD from HC in at least one dataset.

2. Follows the assumption in datasets, where it discriminates significantly.

3. The significant correlation of the feature with PD duration and clinical scores goes with the assumption.

Table 3: Extracted acoustic features.

Speech task	Acoustic feature	Expected change	Specific disorder	Feature definition
			PHONATION	
[a]	HNR	↓	Increased noise	Harmonics-to-noise ratio, the amount of noise in the speech signal due to incomplete vocal fold closure and/or the turbulences in the vocal tract. HNR is defined as the amplitude of noise relative to tonal components in speech.
[a]	CPP	↓	Increased breathness	Cepstral peak prominence representing the disphonia (hoarseness). CPP is defined as the difference between the cepstral peak representing the fundamental frequency and the linear regression line calculated from the magnitude-quefrency cepstra.
[a]	HRF	↓	Increased breathness	Harmonic richness factor, the amount of noise in the speech signal, mainly due to incomplete vocal fold closure. HRF is defined as the ratio between the sum of magnitudes of higher order harmonics and magnitude of the fundamental frequency.
[a]	NAQ	↑	Increased voice harshness	Mean normalised amplitude quotient, defined as $A/(D*T0)$, where A is the amplitude of the glotal flow pulse, D is the peak amplitude of the glotal flow derivative and T0 is one period of glotal flow. A higher value means a slower transition from the open to closed phase.
[a]	relNAQSD	↑	Irregularity of vocal folds activity	The standard deviation of nomralised amplitude quotient relative to its mean.
[a]	QOQ	↑	Increased voice harshness	Mean quasi-open quotient, difined as the ration between the time of opened phase and fundamental period (once cycle of the vocal fold). Higher the quotient, lower the energy of harmonics, but higher the overall intensity of the phonation.
[a]	relQOQSD	↑	Irregularity of vocal folds activity	The standard deviation of quasi-open quotient relative to its mean.
[a]	relF0SD	↑	rregular pitch fluctuations	Standard deviation of fundamental frequency relative to its mean, variation in frequency of vocal fold vibration.
[a]	Jitter (PPQ)	↑	Microperturbations in frequency	Frequency perturbation, extent of variation of the voice range. Jitter is defined as the variability of the F0 of speech from one cycle to the next.
[a]	Shimmer (APQ)	↑	Microperturbations in amplitude	Amplitude perturbation, representing rough speech. Shimmer is defined as the sequence of maximum extent of the signal amplitude within each vocal cycle.
[a]	DUV	↑	Aperiodicity	Degree of unvoiced segments, the fraction of pitch frames marked as unvoiced.
[a]	relF1SD	↑	Tremor of jaw	Standard deviation of first formant relative to its mean. Formants are related to resonances of the oro-naso-pharyngeal tract and are modified by position of tongue and jaw
[a]	relF2SD	↑	Tremor of jaw	Standard deviation of second formant relative to its mean. Formants are related to resonances of the oro-naso-pharyngeal tract and are modified by position of tongue and jaw.
			ARTICULATION	
Read	RFA1	↓	Articulatory decay	Resonant frequency attenuation defined as the distance (dB) in linear predictive coding (LPC) spectrum between resonance of second formant and the local minima before this formant.
Read	RFA2	↓	Articulatory decay	Resonant frequency attenuation defined as the distance (dB) in linear predictive coding (LPC) spectrum between resonance of second formant and the local minima after this formant.
Read	#loc_max	↓	Articulatory decay	The number of local maxima in frequency response of the the vocal tract representing the resonances.
Read	relF1SD	↓	Rigidity of tongue and jaw	Standard deviation of first formant relative to its mean.
Read	relF2SD	↓	Rigidity of tongue and jaw	Standard deviation of second formant relative to its mean.
Read	#lndmrk	↓	Imprecise articulation	The number of speech landmarks relative to total speech time representing the moments of different abrupt acoustic changes, that inform a listener what type of consonant was produced.
DDK	PR	↓	Slow alternating motion rate	Pace rate, representing the number of syllable vocalizations per second. Considering first 30 syllables.
DDK	relSDSD	↑	Erratic articulation	The sum of the standard deviations of the duration of each of the three syllable types relative to their average duration. Considering first 30 syllables.
DDK	COV	↑	Instability of diadochokinetic pace	Coefficient of variation, defined as the ratio of the standard deviation of the duration of the fourth to tenth DDK cycles to the average duration of the first three cycles
DDK	RI	↑	Instability of diadochokinetic pace	Rhythm instability, defined as sum of absolute deviations from a regression line modelling each DDK cycle duration, weighted to the total DDK performance time.
DDK	PA	↑	Acceleration of diadochokinetic pace	Pace acceleration, defined as PA = 100 x (avCycDur4_6 – avCycDur7_9) / avCycDur1_3. where avCycDurX_Y is average duration of cycles X_Y.
DDK	RA	↑	Acceleration of diadochokinetic pace	Rhythm acceleration, defined as gradient of regression line modelling DDK cycle durations (positive values mean acceleration).
			PROSODY	
Read	relF0SD	↓	Monopitch	Pitch variation, defined as a standard deviation of F0 contour relative to its mean.
Read	relSE0SD	↓	Monoloudness	Speech loudness variation, defined as a standard deviation of intensity contour relative to its mean after removing silences exceeding 50ms.
Read	EEVOL	↓	Unstable loudness	Energy evolution, defined as the slope of intensity.
Read	SPIR	↓	Irregular rhythm of speech	Number of pauses (longer than 50ms) relative to total speech time.
Read	PPR	↑	Higher proportion of silence time	Percentual pause ratio, defined as total duration of silences (longer than 50ms)/total duration of speech.
Read	DurMED	↑	Longer duration of silences	Median duration of silences longer than 50 ms.
Read	DurMAD	↑	Higher variability of silence duration	Median absolute deviation of silence duration (longer than 50ms).
Read	NST	↑	Higher proportion of silence time	Net speech time relative to total speech time.

4. No more than 10% of PD patients in any dataset deviate from the HC norm against the assumption.

The assumption is the expected change in the feature value of people with PD compared to HC (see Table 3 – Expected change).

2.4   Machine learning

We chose the Extreme Gradient Boosting (XGBoost) algorithm with random search hyperparameter tuning strategy to classify speakers into PD and HC and find the features importances. Before any mathematical modelling, we performed a data transformation first to avoid differences between the features' values across the language datasets caused by different recording conditions:

$$f_{\mathrm{T}} = \frac{f - \tilde{f}_{\mathrm{HC}}}{IQR_{\mathrm{HC}}}, \tag{3}$$

where f_{T} is the transformed feature, f is an original feature (after the adjustment for age and gender), $\tilde{f}_{\mathrm{HC}}$ is a median calculated from HC (in the given language), and IQR_{HC} is an interquartile range calculated from HC.

2.4.1   All features

To analyze the effect of the language on classification, we worked with all extracted acoustic features and used three different model validation approaches:

- Stratified 10-fold cross-validation with 20 repetitions
  - 6 scenarios (CZ, US, IL, CO, IT, all)
  - stratification ensures balanced train/test data split in the frame of HC/PD and the frame of languages in the scenario with all datasets.
- Cross-language validation technique
  - model trained on data of one dataset and tested on every other.
- Leave-one-language-out validation technique
  - model trained on all but one dataset used for testing.

Models performances were evaluated by Mathews correlation coefficient (MCC), accuracy (ACC), sensitivity (SEN) and specificity (SPE). Importances of features are obtained from models trained on all subjects in each scenario (CZ, US, IL, CO, IT, all), that is, without any split.

2.4.2   Robust features

In the next stage, we took only robust features according to statistical analysis (see section 2.3.1) and trained the models again (CZ, US, IL, CO, IT) to get features importances. Multiplying the importance coefficients of each model with subsequent normalisation gives the global importance coefficients that we ranked in order to find the best robust features. Finally, we trained the model on all datasets and analysed it with the SHAP approach based on game theory to observe the feature behaviour when classifying subjects speaking in different languages.

3 Results

3.1 Statistical analysis

Table 4 describes the results of the Mann-Whitney U test and the change in statistical parameters of PD patients compared to HC. Highlighted are the p-values of the features that significantly differentiate these classes. Table 5 shows the results of the correlation of each feature with the duration of PD, medication and clinical scores. Highlighted p-values represent significant correlations. The significance level for rejecting the null hypothesis was 0.05 for both tests. Figure 3 summarizes the features significant according to the Mann-Whitney U test and the determination of their robustness to a language.

3.2 Machine learning

The performance of each machine learning model trained by all features can be seen in Table 6: 10-fold cross-validation with 20 repetitions, Table 7: cross-language validation and Table 8: leave-one-language-out. In Figure 4, we ranked the acoustic features according to their importance in different language scenarios, and Figure 5 shows the global importance of the robust features only and their behaviour in the model trained by the robust features of all subjects in the corpus.

Table 4: Results of Mann-Whitney U test (p-values after the FDR correction) and changes in statistical parameters (mean $\bar{x}$, median $\tilde{x}$, standard deviation σ) with PD.

	CZ				US				IL				CO				IT			
	p-value	$\bar{x}$	$\tilde{x}$	σ	p-value	$\bar{x}$	$\tilde{x}$	σ	p-value	$\bar{x}$	$\tilde{x}$	σ	p-value	$\bar{x}$	$\tilde{x}$	σ	p-value	$\bar{x}$	$\tilde{x}$	σ
[a]-HNR	0.198	↑	↑	↑	0.250	↑	↑	↓	0.664	↓	↓	↑	**0.008**	↓	↓	↑	**0.008**	↑	↑	↑
[a]-CPP	0.479	↑	↑	↑	0.113	↓	↓	↓	1.000	↓	↑	↑	0.074	↑	↑	↑	0.315	↓	↓	↑
[a]-HRF	0.888	↓	↓	↑	0.773	↓	↓	↑	0.309	↓	↓	↓	0.975	↓	↑	↑	**0.045**	↓	↓	↓
[a]-NAQ	0.888	↓	↓	↑	0.307	↑	↑	↓	0.605	↓	↓	↓	0.631	↑	↑	↑	**<0.001**	↑	↑	↑
[a]-relNAQSD	0.316	↓	↓	↑	0.104	↓	↓	↓	0.749	↓	↑	↓	0.162	↑	↑	↓	**<0.001**	↓	↓	↓
[a]-QOQ	0.791	↓	↑	↑	0.277	↑	↑	↑	0.288	↓	↓	↓	0.603	↓	↓	↓	**0.007**	↑	↑	↓
[a]-relQOQSD	0.180	↓	↓	↑	0.113	↓	↓	↓	0.728	↑	↑	↑	0.201	↑	↑	↑	**<0.001**	↓	↓	↓
[a]-relF0SD	0.862	↓	↑	↑	0.535	↓	↓	↓	**0.004**	↑	↑	↑	**0.008**	↑	↑	↑	0.503	↓	↓	↑
[a]-Jitter (PPQ)	0.316	↓	↓	↑	0.251	↓	↓	↓	0.947	↑	↑	↑	**0.018**	↑	↑	↑	0.934	↑	↓	↑
[a]-Shimmer (APQ)	0.470	↓	↓	↓	0.762	↓	↑	↓	0.836	↓	↓	↓	**0.008**	↑	↑	↑	**0.001**	↓	↓	↓
[a]-DUV	**0.034**	↑	↓	↑	0.104	↓	↓	↓	0.799	↓	↓	↓	**0.031**	↑	↑	↑	**0.008**	↓	↓	↑
[a]-relF1SD	0.653	↑	↓	↑	0.946	↓	↓	↓	0.749	↑	↑	↑	**0.008**	↑	↑	↑	**0.039**	↓	↓	↑
[a]-relF2SD	0.670	↓	↓	↓	0.727	↓	↓	↓	0.347	↑	↑	↓	0.631	↑	↑	↑	0.558	↓	↓	↑
Read-RFA1	0.096	↓	↓	↓	0.104	↓	↓	↓	0.886	↓	↑	↓	**0.033**	↓	↓	↑	**0.001**	↓	↓	↓
Read-RFA2	**0.035**	↓	↓	↑	0.538	↓	↓	↓	**0.033**	↓	↓	↑	0.147	↓	↓	↑	**0.005**	↓	↓	↓
Read-#loc_max	0.658	↓	↓	↑	0.504	↑	↑	↓	0.409	↓	↓	↓	0.252	↓	↓	↑	**<0.001**	↓	↓	↑
Read-relF1SD	0.385	↓	↓	↑	0.855	↑	↑	↑	0.605	↑	↑	↑	0.056	↑	↑	↑	0.952	↓	↑	↓
Read-relF2SD	**0.034**	↓	↓	↑	0.153	↓	↓	↑	0.605	↑	↑	↑	0.858	↑	↑	↑	0.408	↓	↓	↑
Read-#lndmrk	0.862	↓	↓	↓	0.525	↓	↓	↑	**0.001**	↓	↓	↑	0.975	↓	↑	↑	0.751	↓	↓	↑
DDK-PR	0.243	↑	↑	↓	0.525	↓	↓	↓	0.728	↑	↑	↑	0.252	↓	↓	↑	0.437	↓	↓	↑
DDK-relSDSD	0.096	↑	↑	↑	0.440	↓	↓	↓	**0.004**	↑	↑	↓	0.053	↑	↑	↑	-	-	-	-
DDK-COV	0.056	↑	↑	↑	0.727	↓	↓	↓	**0.007**	↑	↑	↑	0.146	↑	↑	↑	0.170	↑	↑	↑
DDK-RI	0.245	↑	↑	↑	0.605	↓	↓	↓	**0.005**	↑	↑	↑	0.389	↑	↑	↑	0.992	↑	↓	↑
DDK-PA	0.784	↑	↑	↑	0.727	↓	↓	↑	0.464	↓	↓	↑	0.975	↑	↓	↑	0.065	↑	↑	↑
DDK-RA	0.548	↓	↓	↑	0.525	↓	↓	↓	0.065	↓	↓	↑	0.528	↑	↑	↑	0.427	↑	↑	↑
Read-relF0SD	**0.001**	↓	↓	↑	0.171	↓	↓	↓	0.249	↓	↓	↑	0.061	↓	↓	↑	**<0.001**	↓	↓	↓
Read-relSE0SD	0.888	↓	↓	↓	0.843	↓	↓	↓	0.691	↓	↑	↓	0.786	↑	↑	↓	0.095	↓	↓	↑
Read-EEVOL	0.552	↑	↑	↑	0.122	↑	↑	↓	0.664	↑	↑	↓	0.975	↑	↑	↓	0.777	↑	↑	↓
Read-SPIR	0.056	↓	↓	↑	0.104	↓	↓	↓	**0.002**	↓	↓	↑	0.075	↓	↓	↑	**0.008**	↓	↓	↑
Read-PPR	0.888	↑	↓	↑	0.605	↓	↓	↑	1.000	↓	↑	↓	0.711	↓	↓	↑	0.109	↑	↑	↑
Read-DurMED	0.056	↓	↑	↓	0.457	↑	↑	↑	**0.045**	↑	↑	↑	0.711	↑	↑	↑	0.173	↑	↑	↑
Read-DurMAD	0.050	↓	↑	↓	0.339	↑	↑	↑	**0.033**	↑	↑	↑	0.407	↑	↑	↑	0.170	↑	↑	↑
Read-NST	0.944	↓	↑	↓	0.525	↑	↑	↑	0.947	↑	↓	↓	0.772	↑	↑	↑	0.170	↓	↓	↑

Table 5: Results of Spearman's rank correlation (p-values after the FDR correction).

	PD duration		LED		UPDRS III		UPDRS III speech		H&Y	
	p-value	coeff	p-value	coeff	p-value	coeff	p-value	coeff	p-value	coeff
[a]-HNR	**0.025**	-0.209	0.489	0.117	0.645	-0.038	0.262	-0.183	0.853	0.052
[a]-CPP	**0.040**	0.193	0.820	-0.036	0.084	-0.149	0.217	-0.197	0.661	-0.139
[a]-HRF	0.412	-0.084	0.759	-0.045	**0.019**	-0.210	0.644	-0.085	0.785	-0.087
[a]-NAQ	0.412	0.087	0.629	0.073	0.084	0.152	0.210	0.205	0.706	0.108
[a]-relNAQSD	0.153	0.144	0.629	-0.071	**0.046**	0.177	0.492	0.121	0.853	-0.044
[a]-QOQ	0.443	-0.076	0.489	-0.129	0.461	0.068	0.851	-0.030	0.853	0.051
[a]-relQOQSD	0.147	0.156	0.560	-0.092	**0.046**	0.177	0.527	0.108	0.914	-0.031
[a]-relF0SD	0.153	-0.146	0.560	0.091	0.078	0.158	0.325	0.163	0.984	-0.002
[a]-Jitter (PPQ)	0.222	0.124	0.999	0.000	0.338	0.085	0.102	0.268	0.918	0.026
[a]-Shimmer (APQ)	**0.004**	0.249	0.489	-0.121	0.633	0.046	0.210	0.203	0.853	-0.067
[a]-DUV	0.882	0.016	0.999	-0.004	0.084	0.148	0.693	0.070	0.661	0.133
[a]-relF1SD	0.222	0.125	0.489	-0.180	0.097	0.139	**0.005**	0.400	0.661	0.152
[a]-relF2SD	0.882	0.018	0.489	-0.115	0.735	-0.027	0.955	0.007	0.965	0.014
Read-RFA1	0.610	0.054	0.489	-0.111	**0.004**	-0.257	0.695	-0.066	0.661	-0.142
Read-RFA2	**0.003**	-0.274	0.944	-0.020	**0.043**	-0.183	0.305	-0.170	0.853	0.060
Read-#loc_max	**0.004**	-0.252	0.999	-0.002	0.084	0.145	0.851	0.036	0.100	-0.313
Read-relF1SD	0.625	-0.049	0.560	-0.087	0.066	0.165	0.644	-0.084	0.678	0.125
Read-relF2SD	0.412	-0.083	0.560	-0.090	0.473	0.065	0.492	-0.120	0.186	0.274
Read-#lndmrk	0.166	-0.139	0.674	-0.064	**0.012**	-0.222	**0.017**	-0.354	0.661	-0.132
DDK-PR	0.286	0.107	0.489	0.126	**0.006**	0.245	0.151	0.232	0.706	0.107
DDK-relSDSD	0.286	0.106	0.489	0.134	**<0.001**	0.334	0.104	0.323	0.100	0.314
DDK-COV	0.286	0.124	0.489	0.128	**0.025**	0.234	**0.031**	0.387	0.100	0.377
DDK-RI	0.770	-0.032	0.684	0.060	**0.010**	0.231	0.151	0.228	0.661	0.179
DDK-PA	0.732	-0.055	0.489	0.244	0.850	0.023	0.851	0.056	0.984	0.007
DDK-RA	0.882	0.013	0.604	-0.080	0.645	0.042	0.731	0.057	0.661	-0.143
Read-relF0SD	0.882	0.015	0.944	-0.018	**0.004**	-0.259	0.693	-0.071	0.853	0.043
Read-relSE0SD	0.147	0.153	0.560	0.098	0.248	0.100	0.527	0.110	0.661	-0.151
Read-EEVOL	0.286	0.106	0.059	0.261	0.645	-0.039	0.892	-0.020	0.785	-0.083
Read-SPIR	0.884	0.010	0.748	0.052	**0.004**	-0.260	**<0.001**	-0.561	0.692	-0.118
Read-PPR	0.610	0.052	0.489	0.160	0.084	0.146	0.492	0.127	0.785	0.084
Read-DurMED	0.412	0.082	0.756	0.048	0.248	0.100	0.113	0.251	0.447	0.221
Read-DurMAD	0.454	0.073	0.965	-0.013	0.084	0.145	0.102	0.275	0.471	0.208
Read-NST	0.610	-0.055	0.489	-0.130	0.191	-0.114	0.492	-0.130	0.853	-0.062

4 Discussion

We processed the voice and speech recordings of 506 people speaking five different languages to explore the ability of specific biomarkers to distinguish between healthy and pathological speech.

4.1 Statistical analysis

According to the Mann-Whitney U test, most features discriminate significantly in the Italian dataset (14/33), following the Israeli (9/33) and Spanish (7/33) ones. There are four significant features in the Czech dataset and none in the American one. Of the 33 extracted acoustic speech features, 23 discriminated significantly in at least one language dataset (Table 4), after which seven did not meet the language robustness conditions (Figure 3).

In the Italian dataset, compared to healthy controls, patients had a significantly less breathy voice (based on the feature [a]-HNR) with fewer irregularities ([a]-relNAQSD, [a]-relQOQSD) and fewer perturbations in amplitude ([a]-Shimmer (APQ)). At the same time, their jaw trembled less ([a]-relF1SD), and their speech was more voiced ([a]-DUV). A significantly higher proportion of voiced segments in patients also occurred in the CZ dataset. Moreover, in the US dataset, many patients deviated from the norms of healthy controls against the assumption in the feature [a]-relNAQSD (64%), [a]-relQOQSD (27%) and [a]-DUV (91%). For these reasons, the mentioned phonation features did not meet the conditions for sufficient

Figure 3: Percentage ratio of PD patients outside the norms given by HC. The left side from zero shows the percentage of PD patients below the lower limit and the right side above the upper limit. The significant difference between HC and PD represents an asterisk, and its position indicates the direction in which the median value of the PD feature has changed compared to HC. Spearman correlation coefficient values are highlighted if they are statistically significant (p-value < 0.05). Features not following the condition for being robust are crossed out.

language robustness. We also enclose prosodic features quantifying the duration of pauses (Read-DurMED) and the variation of their duration (Read-DurMAD) due to the high number of patients deviating from the HC norms against the assumption (64% and 79%) in the CZ dataset. However, two different texts to read is probably the cause here; and thus, these features do not exhibit the task robustness rather than the language.

Most features did not match the robustness conditions due to the phonation of Italians, where patients showed better results in half of the phonation features than healthy ones. The reason for this may be that Italians manifest HD differently in their voice and breathing or just that the patients with these HD symptoms were not present in the IT dataset but come from a different subtype, as described in the study by Rusz et al. [32]. In this study they also present monopitch as a feature that is common in each subtype and because our results show language robustness in this feature, it gains a high potential in the field of objective HD assessment. Moreover, in another of their studies [31], this feature was also independent of the speaker's language based on the general least-squares linear models and had significant discrimination power.

The successful significant and language-independent biomarkers on the basis of statistical analyses hence

Table 6: Classification results: stratified cross-validation.

	MCC	ACC [%]	SEN [%]	SPE [%]
CZ	0.28 ± 0.21	67 ± 9	82 ± 11	43 ± 18
US	0.42 ± 0.54	70 ± 26	74 ± 42	68 ± 37
IL	0.37 ± 0.35	79 ± 11	58 ± 36	83 ± 11
CO	0.43 ± 0.27	70 ± 13	62 ± 21	78 ± 20
IT	0.71 ± 0.31	85 ± 5	86 ± 19	83 ± 27
all	0.49 ± 0.11	75 ± 6	73 ± 8	76 ± 7

Table 7: Classification results: cross-language validation – MCC.

Training	Testing				
	CZ	US	IL	CO	IT
CZ	-	0.55	0.24	0.25	0.43
US	0.21	-	0.07	-0.14	0.34
IL	0.13	-0.11	-	0.09	0.32
CO	0.06	-0.24	0.08	-	-0.03
IT	0.19	0.22	0.23	0.18	-

Table 8: Classification results: leave-one-language-out.

Testing	MCC	ACC [%]	SEN [%]	SPE [%]
CZ	0.19	58	53	67
US	0.48	76	45	94
IL	0.30	53	95	45
CO	0.00	50	78	22
IT	0.61	79	71	90

remain the features: increased breathiness due to incomplete vocal fold closure ([a]-HRF), increased voice harshness due to a slower transition from an open to a closed phase ([a]-NAQ) and a longer duration of an open phase ([a]-QOQ) of the vocal fold cycle, irregular pitch fluctuations ([a]-relF0SD), microperturbations in frequency ([a]-Jitter (PPQ)), articulatory decay as lower prominence of the second formant (Read-RFA1, Read-RFA2) and less local maxima (Read-#loc_max) in the LPC spectrum, the rigidity of tongue and jaw as the lower standard deviation of the second formant during the reading (Read-relF2SD), imprecise articulation due to a lower number of speech landmarks (Read-#lndmrk), erratic articulation as a higher variance of syllable duration during the diadochokinetic task (DDK-relSDSD), instability of diadochokinetic pace as an increased variance of the cycle duration at the end of the task compared to the beginning (DDK-COV) and overall variance of cycle duration (DDK-RI), monopitch (Read-relF0SD) and a lower number of pauses during the speech (Read-SPIR). These are the features that behaved the same in all language datasets.

The best results shows the feature Read-RFA2, which is the only one that discriminates significantly in three datasets (CZ, IL and IT) and strongly correlates with the UPDRS III score and the duration of the disease. There was no significant correlation of any feature with LED or H&Y score (Table 5).

Looking at the results, it is evident that the ability of acoustic speech features to detect HD is indeed language-dependent [13], but at the same time, some features manifest with the development of HD in different languages in the same way as already pointed out by Whitehill [37].

4.2 Machine learning

Based on all extracted features, we were able to detect PD with accuracy ranging from 67% to 85%, depending on the language dataset. For all languages combined, the model achieved an accuracy of 75% with a sensitivity of 73% and a specificity of 76% (Table 6). The accuracy here is approximately the average of the values of

Figure 4: Coefficients of importance of ten most important features in each language scenario (machine learning models trained by all extracted features).

each language model. With lower standard deviations of cross-validation results and higher balance in terms of sensitivity and specificity, a classifier based on gradient trees can be considered suitable when dealing with multilingual data. Vasquez-Correa et al. [36] also observed a lower difference between sensitivity and specificity and lower standard deviation of cross-validation results when fine-tuning the deep machine learning model. This implies that the model trained on more data has a more balanced classification despite the language differences when choosing an appropriate machine learning approach. However, to maintain high classification accuracy, it is probably necessary that the subjects that the model classifies are from the language group that was included during the training process. Otherwise, the classification results decrease rapidly. We can observe this phenomenon in dropped MCC during cross-language validation (Table 7). The only improvement was an increase in MCC values from 0.42 to 0.55 in the classification of the US subjects when the model was trained on CZ data. Although the small size of the US dataset needs to be considered here, the value of MCC dropped from 0.42 to -0.24 when the US was classified by the model trained on CO data. It supports the hypothesis of the dependence of successful classification of PD on the chosen source and target language group during cross-language validation. This should be taken into account in the possible

Figure 5: Global normalized combination of importance coefficients of robust features **(left)** and description of the model trained by robust features of all languages together using SHAP values **(right)** (features ordered from the highest absolute SHAP value, negative values indicate a higher probability for the classification to HC and positive values to patients with PD).

use of transfer learning. Overall, the classifications were most successful when the model was trained on CZ data and worst when trained on CO data. Next to the combination of CO and any other language, the combination of IT and US seems also inappropriate. The leave-one-language-out validation approach results support the previous findings (Table 8) – classification of US subjects is the only one that shows better results; MCC increased from 0.42 to 0.48, which is a slightly lower improvement compared to the model trained on CZ data only. The drop in accuracy when classifying people speaking a different language to subjects used during training is consistent with previous studies' results [13, 25, 23], but it appears that data from another well-chosen language can be used to improve the model performance.

From the importance coefficients of each model (Figure 4), we can observe that in the classification, features are differently important for each scenario, which confirms the earlier findings [27]. After training the machine learning models with only the robust features, globally, the most important features are monopitch (Read-relF0SD), inappropriate silences (Read-SPIR) and articulatory decay (Read-RFA1) (Figure 5). From the SHAP values of the model trained on data of all languages, it is clear that some features of high importance in the individual models lose their ability to discriminate here (Read-#loc_max, [a]-NAQ, [a]-HRF). The features that have been most effective in this model are Read-relF0SD and Read-SPIR again. In third place, according to the absolute SHAP value, is the articulatory decay (Read-RFA2). Thus, reading text appears to be the crucial speech exercise for successful classification, which is consistent with the results of the study by Moro-Velazquez et al. [23].

4.3 Limitations of the study

This study has several limitations, the most major of which is the heterogeneity of the datasets. Each language dataset has a different number of subjects, with only 29 subjects in the American one, which is 185 fewer than in the Czech one. The US patients also have the disease for a much shorter time, which can explain why no feature discriminates significantly in this group. Moreover, since some groups' clinical

data (mainly LED and duration of the disease) are unknown, we cannot tell whether subjects in different datasets have the same level of disease progression. The fact that the voice and speech of the subjects in each dataset were recorded using a different acquisition protocol and hardware (microphones) cloud also play some role (even though we tried to remove this effect by feature transformation based on HC subjects in the specific language). Due to the different ways of performing speech tasks in each country, we did not test whether individual features from different language groups come from the same probability distribution. The different approach to the diadochokinetic task for the Italian acquisition also caused the inability to extract the feature quantifying the erratic articulation as the sum of the standard deviations of the duration of each syllable type (DDK-relSDSD). Furthermore, missing the monologue exercise in some datasets made it impossible to use this speech task in our multilingual study. We also need to consider that the classes (especially in the Israeli dataset) are unbalanced, which can mostly, along with the different sizes of datasets, affect the results of the cross-language validation of machine learning models. The robustness of the features and hypotheses arising from the classification results need to be verified in a follow-up study where data will be homogeneous and subjects preferably of the same age and level of PD progression.

5   Conclusion

We analyzed in detail the behaviour of acoustic speech features in different languages. The aim was to explore digital speech biomarkers of Parkinson's disease and determine which are independent of the speaker's language yet have high discrimination power.

Our statistical analysis found that approximately one-third of the significant features did not meet the conditions for language robustness. The most successful biomarkers in this sense include lower prominence of the second formant, monopitch, and a lower number of pauses detected during text reading. These biomarkers also performed best during classification using machine learning, both in the single language models and the model using all languages' features together. Classification accuracies ranged from 67% to 85%, depending on the language. The model trained with features of all languages together achieved a sensitivity of 73% and a specificity of 76%.

This work contributes insights into the field of objective assessment of speech affected by hypokinetic dysarthria and automated diagnosis of Parkinson's disease. It is the first study to use the concept of language robustness, and through the detailed exploration of speech features using both statistics and machine learning, it proposes several digital speech biomarkers that have the potential to be language independent and that could be potentially used in eHealth/mHealth applications.

Data Availability

Data are protected by privacy and security law. Code with used algorithms can be found in the GitHub repository [Multilingual_speech_analysis] [https://github.com/BDALab/multilingual_speech_analysis].

Acknowledgement

This work was supported by the Czech Ministry of Health under grant no. NU20-04-00294, by EU – Next Generation EU (project no. LX22NPO5107 (MEYS)), and by the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement no. 734718 (CoBeN).

Authorship contribution

Daniel Kovac: conceptualization, data processing, acous- tic features extraction, machine learning, GitHub repository management, article writing. **Jiri Mekyska**: conceptualization, acoustic features extraction, statistical analy- sis, machine learning, article writing. **Vered Aharonson**: conceptualization, data provision. **Pavol Harar**: statistical analysis, machine learning, GitHub repository management. **Zoltan Galaz**: machine learning. **Steven Rapcsak**: data provision. **Juan Rafael Orozco-Arroyave**: data provision. **Lubos Brabenec**: data provision. **Irena Rektorova**: conceptualization.

References

[1] S. G. Adams, A. Dykstra, M. Jenkins, and M. Jog. Speech-to-noise levels and conversational intelligibility in hypophonia and parkinson's disease. *Journal of medical speech-language pathology*, 16(4):165–173, 2008.

[2] L. Brabenec, P. Klobusiakova, P. Simko, M. Kostalova, J. Mekyska, and I. Rektorova. Non-invasive brain stimulation for speech in parkinson's disease: A randomized controlled trial. *Brain Stimulation*, 14(3):571–578, 2021.

[3] L. Brabenec, J. Mekyska, Z. Galaz, and I. Rektorova. Speech disorders in parkinson's disease: early diagnostics and effects of medication and brain stimulation. *Journal of neural transmission*, 124(3):303–334, 2017.

[4] M. Cernak, J. R. Orozco-Arroyave, F. Rudzicz, H. Christensen, J. C. Vásquez-Correa, and E. Nöth. Characterisation of voice quality of parkinson's disease using differential phonological posterior features. *Computer Speech & Language*, 46:196–208, 2017.

[5] P. Corcoran, A. Hensman, and B. Kirkpatrick. Glottal flow analysis in parkinsonian speech. In *BIOSIGNALS*, pages 116–123, 2019.

[6] F. L. Darley, A. E. Aronson, and J. R. Brown. Differential diagnostic patterns of dysarthria. *Journal of speech and hearing research*, 12(2):246–269, 1969.

[7] G. DeMaagd and A. Philip. Parkinson's disease and its management: part 1: disease entity, risk factors, pathophysiology, clinical presentation, and diagnosis. *Pharmacy and therapeutics*, 40(8):504, 2015.

[8] G. Dimauro and F. Girardi. Italian parkinson's voice and speech, 2019.

[9] J. R. Duffy. *Motor speech disorders e-book: Substrates, differential diagnosis, and management*. Elsevier Health Sciences, 2019.

[10] Z. Galaz, J. Mekyska, Z. Mzourek, Z. Smekal, I. Rektorova, I. Eliasova, M. Kostalova, M. Mrackova, and D. Berankova. Prosodic analysis of neutral, stress-modified and rhymed speech in patients with parkinson's disease. *Computer methods and programs in biomedicine*, 127:301–317, 2016.

[11] A. M. Goberman, M. Blomgren, and E. Metzger. Characteristics of speech disfluency in parkinson disease. *Journal of Neurolinguistics*, 23(5):470–478, 2010.

[12] V. L. Hammen and K. M. Yorkston. Speech and pause characteristics following speech rate reduction in hypokinetic dysarthria. *Journal of communication disorders*, 29(6):429–445, 1996.

[13] H. Hazan, D. Hilu, L. Manevitz, L. O. Ramig, and S. Sapir. Early diagnosis of Parkinson's disease via machine learning on speech data. In *2012 IEEE 27th Convention of Electrical and Electronics Engineers in Israel*, pages 1–4. IEEE, 2012.

[14] A. K. Ho, R. Iansek, C. Marigliani, J. L. Bradshaw, and S. Gates. Speech impairment in a large sample of patients with parkinson's disease. *Behavioural neurology*, 11(3):131–137, 1998.

[15] R. B. Hoodin and H. R. Gilbert. Nasal airflows in parkinsonian speakers. *Journal of Communication Disorders*, 22(3):169–180, 1989.

[16] O. Hornykiewicz. Biochemical aspects of Parkinson's disease. *Neurology*, 51(2 Suppl 2):S2–S9, 1998.

[17] F. S. Juste, F. C. Sassi, J. B. Costa, and C. R. F. de Andrade. Frequency of speech disruptions in parkinson's disease and developmental stuttering: A comparison among speech tasks. *Plos one*, 13(6):e0199054, 2018.

[18] Y. Kim and Y. Choi. A cross-language study of acoustic predictors of speech intelligibility in individuals with Parkinson's disease. *Journal of Speech, Language, and Hearing Research*, 60(9):2506–2518, 2017.

[19] D. Kovac, J. Mekyska, Z. Galaz, L. Brabenec, M. Kostalova, S. Z. Rapcsak, and I. Rektorova. Multilingual analysis of speech and voice disorders in patients with parkinson's disease. In *2021 44th International Conference on Telecommunications and Signal Processing (TSP)*, pages 273–277, 2021.

[20] C. McDonald, G. Gordon, A. Hand, R. W. Walker, and J. M. Fisher. 200 years of parkinson's disease: what have we learnt from james parkinson? *Age and ageing*, 47(2):209–214, 2018.

[21] L. Moro-Velazquez and N. Dehak. A review of the use of prosodic aspects of speech for the automatic detection and assessment of parkinson's disease. In *Automatic Assessment of Parkinsonian Speech Workshop*, pages 42–59. Springer, 2019.

[22] L. Moro-Velazquez, J. A. Gomez-Garcia, J. D. Arias-Londoño, N. Dehak, and J. I. Godino-Llorente. Advances in parkinson's disease detection and assessment using voice and speech: A review of the articulatory and phonatory aspects. *Biomedical Signal Processing and Control*, 66:102418, 2021.

[23] L. Moro-Velazquez, J. A. Gomez-Garcia, J. I. Godino-Llorente, F. Grandas-Perez, S. Shattuck-Hufnagel, V. Yagüe-Jimenez, and N. Dehak. Phonetic relevance and phonemic grouping of speech in the automatic detection of Parkinson's disease. *Scientific reports*, 9(1):1–16, 2019.

[24] N. Muñoz-Vigueras, E. Prados-Román, M. C. Valenza, M. Granados-Santiago, I. Cabrera-Martos, J. Rodríguez-Torres, and I. Torres-Sánchez. Speech and language therapy treatment on hypokinetic dysarthria in parkinson disease: Systematic review and meta-analysis. *Clinical Rehabilitation*, 35(5):639–655, 2021.

[25] J. Orozco-Arroyave, F. Hönig, J. Arias-Londoño, J. Vargas-Bonilla, K. Daqrouq, S. Skodda, J. Rusz, and E. Nöth. Automatic detection of Parkinson's disease in running speech spoken in three different languages. *The Journal of the Acoustical Society of America*, 139(1):481–500, 2016.

[26] J. R. Orozco-Arroyave, J. D. Arias-Londoño, J. F. Vargas-Bonilla, M. C. González-Rátiva, and E. Nöth. New Spanish speech corpus database for the analysis of people suffering from Parkinson's disease. In *Proceedings of the Ninth International Conference on Language Resources and Evaluation (LREC'14)*, pages 342–347, Reykjavik, Iceland, May 2014. European Language Resources Association (ELRA).

[27] A. S. Ozbolt, L. Moro-Velazquez, I. Lina, A. A. Butala, and N. Dehak. Things to consider when automatically detecting parkinson's disease using the phonation of sustained vowels: Analysis of methodological issues. *Applied Sciences*, 12(3):991, 2022.

[28] J. Parkinson. An essay on the shaky palsy. *London: Sherwood, Neely and Jones*, pages 1–6, 1817.

[29] W. Poewe. Global scales to stage disability in pd: the hoehn and yahr scale. *Rating Scales Parkinsons Dis*, pages 115–122, 2012.

[30] A. Rohl, S. Gutierrez, K. Johari, J. Greenlee, K. Tjaden, and A. Roberts. Chapter 7 - speech dysfunction, cognition, and parkinson's disease. In N. S. Narayanan and R. L. Albin, editors, *Cognition in Parkinson's Disease*, volume 269 of *Progress in Brain Research*, pages 153–173. Elsevier, 2022.

[31] J. Rusz, J. Hlavnička, M. Novotný, T. Tykalová, A. Pelletier, J. Montplaisir, J.-F. Gagnon, P. Dušek, A. Galbiati, S. Marelli, et al. Speech biomarkers in rapid eye movement sleep behavior disorder and parkinson disease. *Annals of neurology*, 90(1):62–75, 2021.

[32] J. Rusz, T. Tykalova, M. Novotny, D. Zogala, K. Sonka, E. Ruzicka, and P. Dusek. Defining speech subtypes in de novo parkinson disease: response to long-term levodopa therapy. *Neurology*, 97(21):e2124–e2135, 2021.

[33] Z. Thijs and C. R. Watts. Perceptual characterization of voice quality in nonadvanced stages of parkinson's disease. *Journal of Voice*, 2020.

[34] K. Tjaden and G. Wilding. Effects of speaking task on intelligibility in parkinson's disease. *Clinical Linguistics & Phonetics*, 25(2):155–168, 2011.

[35] T. Tykalová, J. Rusz, J. Švihlík, S. Bancone, A. Spezia, and M. T. Pellecchia. Speech disorder and vocal tremor in postural instability/gait difficulty and tremor dominant subtypes of parkinson's disease. *Journal of Neural Transmission*, 127(9):1295–1304, 2020.

[36] J. C. Vásquez-Correa, T. Arias-Vergara, C. D. Rios-Urrego, M. Schuster, J. Rusz, J. R. Orozco-Arroyave, and E. Nöth. Convolutional neural networks and a transfer learning strategy to classify Parkinson's disease from speech in three different languages. In *Iberoamerican Congress on Pattern Recognition*, pages 697–706. Springer, 2019.

[37] T. L. Whitehill. Studies of chinese speakers with dysarthria: informing theoretical models. *Folia Phoniatrica et Logopaedica*, 62(3):92–96, 2010.