Genomic epidemiology of SARS-CoV-2 within households in coastal Kenya: a case ascertained cohort study

Authors

Charles N. Agoti[1,2, *], Katherine E. Gallagher[1,3], Joyce Nyiro[1], Arnold W. Lambisia[1], Nickson Murunga[1], Khadija Said Mohammed[1], Leonard Ndwiga[1], John M. Morobe[1], Maureen W. Mburu[1], Edidah M. Ongera[1], Timothy O. Makori[1], My V.T. Phan[4], Matthew Cotten[4,5], Lynette Isabella Ochola-Oyier[1], Simon Dellicour[6,7], Philip Bejon[1, 8], George Githinji[1,2], and D. James Nokes[1,9]

Affiliations

[1] Kenya Medical Research Institute (KEMRI)-Wellcome Trust Research Programme (KWTRP), Kilifi, Kenya

[2] Pwani University, Kilifi, Kenya

[3] London School of Hygiene & Tropical Medicine, United Kingdom

[4] MRC/UVRI & London School of Hygiene and Tropical Medicine, Entebbe, Uganda.

[5] MRC-University of Glasgow, Centre for Virus Research Glasgow, United Kingdom

[6] Spatial Epidemiology Lab (SpELL), Université Libre de Bruxelles, Bruxelles, Belgium

[7] Department of Microbiology, Immunology and Transplantation, Rega Institute, Laboratory for Clinical and Epidemiological Virology, KU Leuven, University of Leuven, Leuven, Belgium

[8] Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom

[9] School of Life Sciences and Zeeman Institute for Systems Biology and Infectious Disease Epidemiology Research (SBIDER), University of Warwick, Coventry, United Kingdom

* Correspondence

Dr Charles N. Agoti,

Epidemiology and Demography Department,

KEMRI-Wellcome Trust Research Programme,

P.O. Box 230, Kilifi – 80108,

32   Kenya.

33   Email: cnyaigoti@kemri-wellcome.org

Article type

36   Research Article

Abstract (221 words)

Analysis of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) genomic sequences from household infections may provide useful epidemiological information for future control measures. Between December 2020 and July 2022, we conducted a case-ascertained household cohort study whereby households were recruited if a member was either a SARS-CoV-2 case or contact of a confirmed case. A total of 765 members of 214 households were prospectively monitored for SARS-CoV-2 infection and transmission. Follow-up visits collected a nasopharyngeal/oropharyngeal (NP/OP) swab on days 1, 4 and 7 for RT-PCR diagnosis. If any of these were positive, further swabs were collected on days 10, 14, 21 and 28. Of 2,780 NP/OP swabs collected, 540 (19.4%) tested SARS-CoV-2 positive and viral genome sequences were recovered for 288 (53.3%) positive samples. The genomes belonged to 23 different Pango lineages. Phylogenetic analysis including contemporaneous Coastal Kenya data estimated 233 putative transmission events involving 162 members of the 89 households, of which 60 (25%) were intra-household transmission events while 173 (75%) were infections that likely occurred outside the households. In 34 (38%) households, multiple virus introductions were observed (up to six) within the one-month follow-up period, in contrast to high-income settings, where a single introduction seemed to occur during epidemic waves. Our findings suggests that in this setting control of respiratory virus spread by household member isolation will be ineffective.

Introduction

Households are a fundamental unit of social structure and the frequent scene of
respiratory pathogen transmission including for severe acute respiratory syndrome
coronavirus 2 (SARS-CoV-2), the aetiological agent of coronavirus disease 2019
(COVID-19) [1,2]. The household secondary attack rate (i.e., the probability of infection of
susceptible household members directly from the first case in the household) for SARS-
CoV-2 has been estimated to be about 21.1% (95%CI: 17.4-24.8%) with considerable
heterogeneity observed over geographic regions and time periods [3-6]. An improved
understanding of SARS-CoV-2 household transmission, including the frequency of virus
transmission within a household compared to transmission from outside into the
household, may help refine local control measures.

As of August 2022, Kenya had experienced six waves of SARS-CoV-2 infections [7]. In
this period, Kenya's COVID-19 countermeasures as measured by the Oxford Stringency
Index (SI), a measure based on nine key indicators rescaled from 0-100, fluctuated
between 36 and 89[8] in response to the perceived threat. The different infection waves
were dominated by distinct virus variants: the first two by the early ancestral virus
lineage (mainly B.1), the third by the Alpha (B.1.1.7) variant of concern (VOC), the
fourth by Delta VOC (B.1.617.2), the fifth by Omicron VOC (BA.1), and the sixth by
Omicron sub-variants BA.4/5 [9]. Local epidemiological studies suggest many SARS-
CoV-2 infections have been asymptomatic or mild [10], as evidenced by the high
seroprevalence (48.5% by March 2021 [11]) without high numbers of hospitalisations or
excess mortality [12]. However, confirmed active SARS-CoV-2 infections by the Kenyan
Ministry of Health (MoH) have been reported in less than 1% of the Kenyan population
indicating inadequate testing. Consequently, gaps exist in our quantitative
documentation of SARS-CoV-2 circulation within the Kenyan population.

To date, SARS-CoV-2 genomic analysis has played a key role in elucidating COVID-19
pandemic transmission patterns [13-17]. For instance, genomic analysis has helped
uncover the nature of virus seeding into close living environments like, hospitals [18,19],
prisons[20], cruise ships [21], long-term care facilities [22], or learning institutions [23], and has

also uncovered a number of superspreading events [19,24]. Unlike many RNA viruses,

SARS-CoV-2 replication is believed to be under some level of proof-reading[25], limiting

its substitution rate (estimated at 9.90×10^{-4} nucleotide substitutions/site/year) [26], a

parameter critical in applying genomics to understand short-term epidemiological

dynamics.

Few studies have examined SARS-CoV-2 genomic diversity in putative within-

household transmission cases. In Ireland, Hare *et al* [27] found that most family members

testing positive had indistinguishable consensus genome sequences from other family

members and the early presumed index case had a divergent sub-lineage. In the

present study, we document SARS-CoV-2 transmission patterns within households in

coastal Kenya by analysing infections identified in a case-ascertained cohort (i.e.

households enrolled if a member was confirmed case of SARS-CoV-2 or a contact of

one) during successive local waves of infections [28]. We undertook detailed genomic

analyses to identify patterns of SARS-CoV-2 introductions into households and to

document the frequency and patterns of infection spread within households in coastal

Kenya.

Results

Baseline characteristics

A total of 765 participants from 214 households were recruited between 10th December

2020 and 29th July 2022. From these, 2,780 nasopharyngeal/oropharyngeal (NP/OP)

swabs were collected, 540 (19.4%) of which tested SARS-CoV-2 positive by RT-PCR

(**Table 1**). The positive swabs were from 254 infected participants in 119 households.

The temporal distribution of the swab collections for all the 214 households and their

RT-PCR testing results are shown in **S1 Fig**. The participants with a positive swab had

a median age of 27 years (IQR: 12.0-46.0; **Table 1)**, with 164 (64.6%) females.

Compared to participants who remained SARS-CoV-2 negative during the follow-up

period, positive cases were more likely to report at least one acute respiratory illness

(ARI) symptom (67.7% vs 32.5%; *p <0.001,* chi-squared test). The household

recruitments were predominantly coincidental with the national COVID-19 waves 3, 4, 5

and 6 (**Fig. 1A and B**) with only one household recruited during wave 2. The changes in

the national countermeasures during the study period as estimated by the Oxford SI is

provided in **Fig. 1C.**

Table 1. Demographic characteristics of study participants.

Characteristic	Positive	Negative	Total
NP/OP swabs	540	2240	2780
Recruited households	119	95	214
Recruited participants	254	511	765
Sex			
female (%)	164 (64.6)	297 (58.1)	461 (60.3)
male (%)	90 (35.4)	214 (41.9)	304 (39.7)
Age distribution (years)			
Median (IQR)	27.0 (12.0-46.0)	20.0 (10.0-40.0)	24.0 (11.0-44.0)
Age categories			
0-4 y (%)	31 (12.2)	54 (10.6)	85 (11.1)
5-9 y (%)	23 (9.1)	54 (10.6)	77 (10.1)
10-19 y (%)	48 (18.9)	133 (26.0)	181 (23.7)
20-39 y (%)	73 (28.7)	113 (28.6)	219 (28.6)
40-64 y (%)	63 (24.8)	92 (18.0)	155 (20.3)
65+ y (%)	14 (5.5)	31 (6.1)	45 (5.9)
Unknown	2 (0.8)	1 (0.2)	3 (0.4)
Symptom status			
asymptomatic	82 (32.3)	345 (67.5)	427 (55.8)
symptomatic	172 (67.7)	166 (32.5)	338 (44.2)

127   IQR: Interquartile Range, y: years

Fig. 1. The timeline of the household study and genomic sequencing results.
131 *Panel A: Reported SARS-CoV-2 infections observed in Kenya between March 2020 and*
132 *March 2022. Panel B: Temporal distribution of the collected NP/OPs and their RT-PCR*
133 *diagnostic results. Panel C: The level of government restrictions. Panel D: Temporal*
134 *distribution of the SARS-CoV-2 VOCs detected in the household study.*
135

136
137
138

Genomic sequencing and lineage/VOC classification

139
Near complete genomes (over 80% coverage) were obtained from 288 samples (53.3%
of positive samples) collected from 162 participants from 89 households. The samples
that failed sequencing (n = 252) had high Ct values (corresponding to low viral load;
>33.0; **S2. Fig**.) or yielded low quality PCR products during library preparation. The 288
genomes were classified into non-VOC (Pango lineage B.1; n = 11), Alpha VOC (n =
70), Beta VOC (n = 22), Delta VOC (n = 88) and Omicron VOC (n = 97; **Fig.1D**). All
Alpha and Beta sequences fell within lineage B.1.1.7 and B.1.351, respectively. Within
the Delta VOC, six Pango lineages were identified while within the Omicron VOC 14
Pango lineages were identified. The lineages within Omicron VOC included those that
classified under BA.1, BA.2, BA.4 and BA.5 sub-variants. A summary of the temporal
distribution of the 23 Pango lineages that were identified across the sequenced cases
and their history of global detection is presented in **S2 Fig. and S1 Table**.

Phylogenetic clustering of the household study genomes

153
To investigate the genetic diversity in the household study genome sequences, we
reconstructed a maximum likelihood (ML) phylogeny that included background coastal
Kenya co-circulating viruses (n = 2,555). As expected, the household genome
sequences clustered by VOC and Pango lineages with other Kenyan coastal sequences
(**S3 Fig.**). Notably, lineage B.1 sequences were found in multiple branches of the
phylogeny, including some at the base of branches leading to Beta and Delta VOCs. To
assess the genetic relatedness of the recovered genomes within and between
households, we reconstructed VOC-specific phylogenies with tips coloured by the
household of sampling (**Fig. 2**). Here we observed both intra- and inter-household
clustering (i.e., multiple sequences of a single clade identified in one household or
shared between two or more households). In a few households, sequences of a single
VOC but distinct clades were observed, indicating multiple distinct virus introductions
into the same household (**Fig. 2**).

Fig. 2. Phylogenetic patterns of variants of concern (VOC) in the household study using maximum-likelihood methods. These household study VOC phylogenies include other SARS-CoV-2 genome sequences generated from samples collected in six Kenyan coastal counties (Mombasa, Kwale, Kilifi, Taita Taveta, Tana River and Lamu) during the study period as background diversity (tips without symbols). On the phylogenies, household sample derived sequences are displayed as filled circles, coloured distinctly by household. In the Alpha, Beta, Delta, and Omicron phylogenies, 370, 175, 535 and 574 genome sequences were included, respectively.

Estimating the number of introductions into the households

SARS-CoV-2 has been reported to have a genomic evolutionary rate of ~2 nucleotide substitutions per genome per month [29]. A heterogenous distribution of the pairwise nucleotide differences of specimens identified in the same household was observed (range 0-63; median 0.0, mean 2.8; **S4 Fig.**). More than two nucleotide differences between genomes from the same household were observed in 25 households, implying

possible multiple introductions into these households. We further investigated the

number of virus introductions into each household using ancestral state reconstruction

(ASR) along the dated ML phylogeny [16,30]. A total of 155 virus introductions were

predicted into the 89 households where we recovered SARS-CoV-2 sequence data. On

classifying the transition events by sequence origin ("non-household" events - those

individuals not from same household - and "household" events - those where both

involved individuals were members of same household), we found that most transitions

were "non-household" compared to "household" transition events (75% vs. 25%; **Fig.**

3A). Overall, we estimated that a single virus introduction occurred for 55 households

(61%), two introductions for 16 households (17%), three introductions for nine

households (10%), four introductions for five households (6%), five introductions for

three households (3%) and six introductions in one household (**Fig. 3B**).

Fig. 3. *Patterns of SARS-CoV-2 introductions into studied households as determined by*
ancestral state reconstruction. Panel A: Alluvial plot showing the number of viral imports
and exports into and out of study households. Panel B: Frequency of distinct
introductions into the study households.

Discussion

We provide evidence of frequent multiple introductions of SARS-CoV-2 into rural coastal Kenyan households within a month, a finding that was unexpected. The conventional view has been that households with concurrently infected members most likely acquire the infection introduced through a single index case. This assumption has been repeatedly supported by several genomic studies. For instance, a phylogenetic analysis performed in a Dutch study following 85 households showed a single introduction into all the studied households [4]. By contrast, in the present study, about one-third of infected households (38%) had two or more introduction events within a distinct epidemic wave.

A variety of factors may explain the differences in household SARS-CoV-2 introduction patterns identified in our study compared to previous observations. First, in our setting, multiple families may live in one compound and eat from one kitchen and these were defined as a single household in this study [31]. Such living arrangement results in relatively larger household sizes that may be associated with higher chances of multiple virus introductions, especially at the peak of epidemic waves. Second, the dominance of informal sector jobs in this setting may have made options like working from home difficult to implement. As result the potential for acquiring infection outside the household setting was high relative to high-income settings.

Our study followed-up participants for a period of up to 1 month with serial sample collection and recovered genomes were analysed in the context of contemporaneous locally circulating diversity in coastal Kenya [32]. Despite observing minimal nucleotide variation between samples from members of the same household infection clusters, when we incorporated sampling dates through the ASR analysis, we were able to partially reconstruct potential within-household transmission events. This allowed the identification of multiple introductions into the households of closely related viruses, observation of putative within-household transmission and short-interval reinfections.

Few studies have examined SARS-CoV-2 households transmission dynamics within

Africa [33-35], and these have resulted in diverse findings. In rural Egypt, a 6-month study

reported a SAR of 89.8% [33], in South Africa a 13-month study reported a 25% infection

rate among vulnerable household contacts [34], and in Madagascar, a SAR of 38.8%

(CI:19.5-57.2) [36] was reported. None of these studies included a viral genomic analysis

to support any conclusions that the inferred household transmission clusters were

epidemiologically linked and arose from a single index case.

The Kenyan government countermeasures in place during the study period may have

had an impact on the spread of SARS-CoV-2 within the study households. In June

2020, the Kenyan government announced guidelines for home-based care for

asymptomatic or mildly symptomatic patients without co-morbidities [37]. Kenya started

vaccinating its population in March 2021, but the coverage was low (<15%) during the

study period [38], and thus it is unlikely that it affected transmission during our study. The

stringency index in the country during the study period fluctuated from 35% to 75%.

However, we did not detect variation in the pattern of introductions over time, which

could suggest that the various restrictions had minimal impact at the household level.

However, concluding on this would likely require more advanced epidemiological

modelling.

This study had several limitations. First, several positive NP/OP samples (46.7%) failed

to yield viral genome sequences or had large genome sequencing gaps due to PCR

amplicon drop-off. With such a high level of genome data missingness, the overall

phylogenetic signal was reduced in trying to establish who infected whom or

directionality of transmission. Second, we cannot rule out that a few of the sequence

changes observed could be sequencing or assembly artifacts [39]. Third, the case-

ascertained study design we used had the drawback that by the time the first sample

was collected, multiple positive cases had already occurred in most households. Most of

the index cases were recruited following presentation to a health facility with ARI. This

complicated our effort of fully determining who infected whom back in the household. To

overcome this challenge, future studies should observe members before entry of the

virus into households and genomic data co-analysed with other relevant epidemiological

data [40]. Fourth, our sampling interval, especially after week two, may have missed

persons who had been positive for less than the 7 days sample collection interval.

Higher density sampling has previously been associated with a higher attack rate[4].

Finally, we did not bring in other datatypes like, contact patterns, symptoms and shared

intra-host variation [41], which would have provided further insights into potential

transmission linkages. An analysis of secondary attack rates is being conducted that

includes all data, including serological evidence of infection, which is not included here.

In conclusion, we identified an unusually high number of independent virus introductions

into households in coastal Kenya during household temporally clustered infections. Our

findings suggests that control of SARS-CoV-2 transmission by household member

isolation alone may not stop community transmission in this setting. Our study further

highlights the importance of examining genomic data for accurate estimation and

interpretation of SARS-CoV-2 household epidemiological parameters.

Methods

Study design and recruitment

We conducted a case-ascertained study in coastal Kenya, where new households were

recruited into the study via five local health facilities or the Kilifi County Department of

Health rapid response team (RRTs). Households were defined as dwellings or groups of

dwellings that share the same kitchen or cooking space (**S2 Table**). Many of the

recruited households were from within the Kilifi Health and Demographic Surveillance

System (KHDSS) area located in Kilifi County, Coastal Kenya [42]. To get enrolled, a

household needed to have at least two occupants, to be accessible by road, and to

obtain permission from the household head. In the beginning, only households with a

member who was a contact of a confirmed case from a different household were

recruited, but due to slow enrolment, we revised the protocol to include households with

confirmed cases at first sampling. A household was exempted if at the time of

recruitment: two or more members had already developed COVID-19 symptoms (e.g.

fever, sore throat, cough etc), a member had been hospitalized due to COVID-19, or the

household had been enrolled in a trial of therapeutic COVID-19 product.

Follow-up

During each household visit, a NP/OP swab was obtained from all participants and

transported in cool boxes with ice packs to KEMRI-Wellcome Trust Research

Programme (KWTRP) laboratories within 48 hr for real-time RT-PCR testing. The study

had broadly two follow-up arms: "reduced follow-up" and "intense follow-up".

Households in the "reduced follow-up" arm were those where all the members tested

SARS-CoV-2 negative at day 1, 4, and 7. Therefore, in that case, follow-up was

discontinued henceforth with a few exceptions (**S1 Fig.**). The "intense follow-up arm"

was activated when a household member tested positive on day 1, 4, or 7. In that case,

the household was sampled again on day 10, 14, 21, and 28. Data on baseline

household and demographic characteristics were collected by the study team at

enrolment. During all households' visits, data on presence of ARI symptoms (e.g., fever,

cough, runny nose, sore throat, headache) were collected.

Laboratory procedures

SARS-CoV-2 diagnosis

SARS-CoV-2 testing of study samples was undertaken alongside samples collected in

six coastal counties of Kenya as part of the national COVID-19 testing as previously

described [10]. Four different viral RNA extraction kits (namely, QIAamp Viral RNA Mini

Kit, RNeasy ® QIAcube ® HT Kit, TIANamp Virus RNA Kit and Da An Gene Nucleic

acid Isolation and Purification Kit) were deployed in combination with five different RT-

PCR kits/protocol (namely, Da An Gene Co. detection Kit, European Virus Archive-

Global (EVAg) E gene protocol, Standard M Kit, Sansure Biotech Novel Coronavirus

(2019-nCoV) Nucleic Acid Diagnostic Real-time RT-PCR kit[10]). Positives were

determined using the kit/protocol-defined cycle thresholds (Ct). In kits where multiple

SARS-CoV-2 genomic regions were targeted, the average cycle threshold (Ct) was

calculated from the individual Cts.

Genome sequencing

We aimed to sequence whole genomes of all the RT-PCR positive samples with a cycle

threshold of < 33.0. Viral RNA was re-extracted from the specimens using QIAamp viral

RNA mini-Kit following the manufacturer's instructions and converted to cDNA using

Lunascript kit with ARTIC protocol primers [43]. Genome amplification was conducted

using Q5 PCR kit and ARTIC protocol primers (initially v3 and then v4). Sequencing

libraries were prepared using Oxford Nanopore Technologies (ONT) ligation sequencing

kit SQK-LSK109 and the ONT Native Barcoding Expansion kit as described in the

ARTIC protocol [43]. Sequencing was performed on Oxford Nanopore Technologies'

MinION or GridION devices using R9.4.1 flow cells.

Bioinformatic analysis

Genome assembly and lineage assignment

The ONT's raw sequencing reads (FAST5) were base-called and demultiplexed using

ONT's Guppy v3.5-4.2. The resultant files (FASTQ) were assembled into consensus

genomes using ARTIC bioinformatic pipeline reference-based approach

(https://artic.network/ncov-2019/ncov2019-bioinformatics-sop.html; last accessed 2022-

09-17). Only nucleotides with a read depth of more than × 20 were included into the

consensus sequence. Only sequences with >80% genome coverage were further

analysed. The genomes were assigned Pango lineages using the command-line

installation of pangolin v4.1.3, PUSHER-v1.3, scorpio v0.3.16 and constellation v0.1.6

[44,45].

Phylogenetic analysis

Multiple sequence alignments were generated using Nextalign v.1.10.1 referenced-

based aligner within the Nextclade tool v0.14.2 [46] using the command:

nextalign -r NC_045512.fasta -r <input_file.fasta>

Alignments were visualized using a custom Python script and "snipit" tool

(https://github.com/aineniamh/snipit; last accessed 2022-05-20). Pairwise distances

were calculated using pairsnp.py (https://github.com/gtonkinhill/pairsnp/; last accessed

2022-05-20) using the command:

359   *pairsnp -f <input_file.fasta> --csv --output <output_file.fasta>*

361   Phylogenetic relationships between all recovered genomes and between viruses

362   classified under the same VOC were obtained through the inference of maximum

363   likelihood (ML) trees performed with the program IQTREE v2.1.3 with a general time

364   reversible (GTR) substitution model using the command:

366   *Iqtree -s <input_alignment.fasta> -nt <no_of_cores> -m GTR*

We included contemporaneous genomes from the six coastal Kenya counties

(Mombasa, Kilifi, Kwale, Taita Taveta, Tana River, and Lamu) that were sequenced as

part of the national SARS-CoV-2 genomic surveillance to provide phylogenetic context

to the household study genomes. Each ML tree was subsequently time-calibrated using

the program TreeTime, assuming a constant evolutionary rate of SARS-CoV-2 genome

of 8.0×10^{-4} using the command:

treetime –tre <tree_file> --aln <alignment_file> --clock-rate 0.00084 --dates

376   *<metadata_file.csv>*

The phylogenetic trees were combined with metadata and visualized with the R

package "ggtree" v2.4.2 [47].

Virus introductions and putative transmissions

The number of independent virus introductions into the households was investigated

using two approaches.

i.   *Pairwise nucleotide substitution.*

Here, we compared observed nucleotide differences between pairs of household

genomes with the number of nucleotide mutations expected over the time interval

between the two sampling dates. The pairwise differences between the household study

genomes were computed using the program pairsnp

(https://github.com/gtonkinhill/pairsnp; accessed 06-Feb-2023).

ii.   Ancestral state reconstruction (ASR)

ASR approach was used to identify the introduction events into a household and count

the transitions between household members [16]. This was performed along the time-

scaled phylogeny obtained for each VOC. To infer the number of introduction events

into each household, a variable "source" was generated for which the sequences were

assigned the household ID or noted as "non-household". Using the dated phylogeny,

the nucleotide sequence alignment, the "source" metadata file, a mugration analysis

was run using TreeTime[48], using the command:

treetime mugration --tre <dated_tree_file> --states <metadata_file.csv> --attributes

402   *<source_attribute>*

To infer within household transition events, a variable "participant" was generated for

which the sequences were assigned the participant ID or noted as "non-household".

The mugration analysis was rerun using the dated phylogeny and transitions between

members of the same households noted using python scripts described in [16,30].

Statistical analysis

Summary statistics were computed for key demographic characteristics including mean,

median, standard deviation as appropriate. Infection prevalence was expressed using

proportions and comparison between groups included appropriate statistical tests (e.g.,

chi-square or Fisher's exact). All statistical analyses were performed in R version 4.0.5.

Ethical consideration

The study protocol was reviewed and approved by both the Scientific and Ethics

Research Unit (SERU) at Kenya Medical Research Institute (KEMRI), Nairobi, Kenya

(REF: SERU protocol # 4077) and the University of Warwick, Biomedical and Scientific

Research Ethics Committee, Coventry, United Kingdom (REF: BSREC 150/19-20

AM01)**.** Prior to baseline data and biospecimen collection, written informed consent was

obtained from all participants aged 18 years or older, while for participants aged less

than 18 years consent was obtained from their parents or legal guardians. Assent was

also sought for adolescents (13-17 years of age).

Data availability

The consensus genome sequences obtained in this study that passed our quality

control filters have been submitted to both GISAID and GenBank database (accession

numbers available in appendix pages of the supplementary material). The raw data files

have been prepared for deposition in Harvard DataVerse. For more detailed information

beyond the metadata used in the paper, there is a process of managed access requiring

submission of a request form for consideration by our Data Governance Committee

(http://kemri-wellcome.org/about-us/#ChildVerticalTab_15).

Code availability

The code for the analyses presented in this manuscript is available from Github using

the link https://github.com/cnyaigoti/Household_study_2021-2022.

Acknowledgements

We thank (a) the members of the Kilifi County rapid response team who worked with our

field study team in collecting the samples analysed here; (b) the members of the

COVID-19 KWTRP Testing Team who undertook real-time RT-PCR processing of the

samples received at KWTRP to identify positives (see full list of members below).

This work was supported by the National Institute for Health and Care Research (NIHR)

(project reference 17/63/82) using UK aid from the UK Government to support global

health research, The UK Foreign, Commonwealth and Development Office and

Wellcome Trust (grant# 220985). Members of COVID-19 Testing Team at KWTRP are

supported by multiple funding sources including UNITAD (BOHEMIA study received by

Dr Marta Maia funded UNITAID), EDCTP (Senior Fellowship and Research and

Innovation Action (RIA) grants received by Dr Francis Ndungu), GAVI (PCIVS grant

received by Prof. Anthony Scott). Dr Simon Dellicour acknowledges support from the

Fonds National de la Recherche Scientifique (F.R.S.-FNRS, Belgium; grant

n°F.4515.22), from the Research Foundation - Flanders (*Fonds voor Wetenschappelijk*

Onderzoek-Vlaanderen, FWO, Belgium; grant n°G098321N), and from the European

Union Horizon 2020 project MOOD (grant agreement n°874850). MP and MC were

supported by the Wellcome Trust and FCDO - Wellcome Epidemic Preparedness –

Coronavirus (AFRICO19, grant agreement number 220977/Z/20/Z). The views

expressed in this publication are those of the author (s) and not necessarily those of

NIHR, the Department of Health and Social Care, Foreign Commonwealth and

Development Office, Wellcome Trust or the UK government.

List of members of the COVID-19 Testing Team at KWTRP

Agnes Mutiso, Alfred Mwanzu, Angela Karani, Bonface M. Gichuki, Boniface Kaaria,

Brian Bartilol, Brian Tawa, Calleb Odundo, Caroline Ngetsa, Clement Lewa, Daisy

Mugo, David Amadi, David Ireri, Debra Riako, Domtila Kimani, Donwilliams Omuoyo,

Edwin Machanja, Elijah Gicheru, Elisha Omer, Faith Gambo, Horace Gumba, Isaac

Musungu, James Chemweno, Janet Thoya, Jedida Mwacharo, Jennifer Musyoki, John

Gitonga, Johnstone Makale, Justine Getonto, Kelly Ominde, Kelvias Keter, Lydia

Nyamako, Margaret Nunah, Martin Mutunga, Metrine Tendwa, Moses Mosobo, Nelson

Ouma, Nicole Achieng, Patience Kiyuka, Perpetual Wanjiku, Peter Mwaura, Rita Warui,

Robinson Cheruiyot, Salim Mwarumba, Shaban Mwangi, Shadrack Mutua, Susan

Njuguna, Victor Osoti, Wesley Cheruiyot, Wilfred Nyamu, Wilson Gumbi and Yiakon

Sein.

Role of funding source

478 The funders of the study had no role in study design, data collection, data analysis ,
479 data interpretation or writing of the report.

Author's contributions

482 The project was conceived and designed by CNA, KEG, JUN, MC, and DJN; Laboratory

483 processing of specimens was conducted by JUN, LIO, NM, AWL, KSM, LN, JMM,

484 MWM, EMO, and TOM; Management and analysis of data were handled by AWL, CNA,

485 NM, SD and GG. CNA wrote the first draft; MP, MC, LIO, SD, PB, and DJN critically

486 reviewed the manuscript to produce the final draft.

References

Lee, E. C., Wada, N. I., Grabowski, M. K., Gurley, E. S. & Lessler, J. The engines of SARS-CoV-2 spread. *Science* **370**, 406-407 (2020). https://doi.org:doi:10.1126/science.abd8755

Park, Y. J. *et al.* Contact Tracing during Coronavirus Disease Outbreak, South Korea, 2020. *Emerging Infectious Disease journal* **26**, 2465 (2020). https://doi.org:10.3201/eid2610.201315

Thompson, H. A. *et al.* Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Setting-specific Transmission Rates: A Systematic Review and Meta-analysis. *Clin Infect Dis* **73**, e754-e764 (2021). https://doi.org:10.1093/cid/ciab100

Kolodziej, L. M. *et al.* High SARS-CoV-2 household transmission rates detected by dense saliva sampling. *Clinical Infectious Diseases* (2022). https://doi.org:10.1093/cid/ciac261

Madewell, Z. J., Yang, Y., Longini, I. M., Jr, Halloran, M. E. & Dean, N. E. Household Secondary Attack Rates of SARS-CoV-2 by Variant and Vaccination Status: An Updated Systematic Review and Meta-analysis. *JAMA Network Open* **5**, e229317-e229317 (2022). https://doi.org:10.1001/jamanetworkopen.2022.9317

Jørgensen, S. B., Nygård, K., Kacelnik, O. & Telle, K. Secondary Attack Rates for Omicron and Delta Variants of SARS-CoV-2 in Norwegian Households. *JAMA* **327**, 1610-1611 (2022). https://doi.org:10.1001/jama.2022.3780

Tegally, H. *et al.* The evolving SARS-CoV-2 epidemic in Africa: Insights from rapidly expanding genomic surveillance. *Science*, eabq5358 (2022). https://doi.org:10.1126/science.abq5358

Hale, T. *et al.* A global panel database of pandemic policies (Oxford COVID-19 Government Response Tracker). *Nat Hum Behav* **5**, 529-538 (2021). https://doi.org:10.1038/s41562-021-01079-8

Nasimiyu, C. *et al.* Imported SARS-CoV-2 Variants of Concern Drove Spread of Infections across Kenya during the Second Year of the Pandemic. *COVID* **2**, 586-598 (2022).

Nyagwange, J. *et al.* Epidemiology of COVID-19 infections on routine polymerase chain reaction (PCR) and serology testing in Coastal Kenya. *Wellcome Open Res* **7**, 69 (2022). https://doi.org:10.12688/wellcomeopenres.17661.1

Uyoga, S. *et al.* Prevalence of SARS-CoV-2 Antibodies From a National Serosurveillance of Kenyan Blood Donors, January-March 2021. *Jama* **326**, 1436-1438 (2021). https://doi.org:10.1001/jama.2021.15265

Otiende, M. *et al.* Impact of COVID-19 on mortality in coastal Kenya: a longitudinal open cohort study. *medRxiv*, 2022.2010.2012.22281019 (2022). https://doi.org:10.1101/2022.10.12.22281019

Li, J., Lai, S., Gao, G. F. & Shi, W. The emergence, genomic diversity and global spread of SARS-CoV-2. *Nature* **600**, 408-418 (2021). https://doi.org:10.1038/s41586-021-04188-6

Bugembe, D. L. *et al.* Main Routes of Entry and Genomic Diversity of SARS-CoV-2, Uganda. *Emerg Infect Dis* **26**, 2411-2415 (2020). https://doi.org:10.3201/eid2610.202575

Githinji, G. *et al.* Tracking the introduction and spread of SARS-CoV-2 in coastal Kenya. *Nat Commun* **12**, 4809 (2021). https://doi.org:10.1038/s41467-021-25137-x

Wilkinson, E. *et al.* A year of genomic surveillance reveals how the SARS-CoV-2 pandemic unfolded in Africa. *Science* **374**, 423-431 (2021). https://doi.org:10.1126/science.abj4336

Vöhringer, H. S. *et al.* Genomic reconstruction of the SARS-CoV-2 epidemic in England. *Nature* **600**, 506-511 (2021). https://doi.org:10.1038/s41586-021-04069-y

Ellingford, J. M. *et al.* Genomic and healthcare dynamics of nosocomial SARS-CoV-2 transmission. *Elife* **10** (2021). https://doi.org:10.7554/eLife.65453

Illingworth, C. J. *et al.* Superspreaders drive the largest outbreaks of hospital onset COVID-19 infections. *Elife* **10** (2021). https://doi.org:10.7554/eLife.67308

20   Hershow, R. B. *et al.* Rapid Spread of SARS-CoV-2 in a State Prison After Introduction by Newly
Transferred Incarcerated Persons - Wisconsin, August 14-October 22, 2020. *MMWR Morb*
Mortal Wkly Rep **70**, 478-482 (2021). https://doi.org:10.15585/mmwr.mm7013a4

21   Hoshino, K. *et al.* Transmission dynamics of SARS-CoV-2 on the Diamond Princess uncovered
using viral genome sequence analysis. *Gene* **779**, 145496 (2021).
https://doi.org:10.1016/j.gene.2021.145496

22   Aggarwal, D. *et al.* The role of viral genomics in understanding COVID-19 outbreaks in long-term
care facilities. *Lancet Microbe* **3**, e151-e158 (2022). https://doi.org:10.1016/s2666-
5247(21)00208-1

23   Baumgarte, S. *et al.* Investigation of a Limited but Explosive COVID-19 Outbreak in a German
Secondary School. *Viruses* **14** (2022). https://doi.org:10.3390/v14010087

24   Popa, A. *et al.* Genomic epidemiology of superspreading events in Austria reveals mutational
dynamics and transmission properties of SARS-CoV-2. *Sci Transl Med* **12** (2020).
https://doi.org:10.1126/scitranslmed.abe2555

25   Smith, E. C., Blanc, H., Surdel, M. C., Vignuzzi, M. & Denison, M. R. Coronaviruses lacking
exoribonuclease activity are susceptible to lethal mutagenesis: evidence for proofreading and
potential therapeutics. *PLoS Pathog* **9**, e1003565 (2013).
https://doi.org:10.1371/journal.ppat.1003565

26   Nie, Q. *et al.* Phylogenetic and phylodynamic analyses of SARS-CoV-2. *Virus Res* **287**, 198098
(2020). https://doi.org:10.1016/j.virusres.2020.198098

27   Hare, D. *et al.* Genomic epidemiological analysis of SARS-CoV-2 household transmission. *Access*
Microbiol **3**, 000252 (2021). https://doi.org:10.1099/acmi.0.000252

28   Githinji, G. *et al.* The genomic epidemiology of SARS-CoV-2 variants of concern in Kenya.
medRxiv, 2022.2010.2026.22281446 (2022). https://doi.org:10.1101/2022.10.26.22281446

29   Duchene, S. *et al.* Temporal signal and the phylodynamic threshold of SARS-CoV-2. *Virus*
Evolution **6** (2020). https://doi.org:10.1093/ve/veaa061

30   Tegally, H. *et al.* Sixteen novel lineages of SARS-CoV-2 in South Africa. *Nat Med* **27**, 440-446
(2021). https://doi.org:10.1038/s41591-021-01255-3

31   Munywoki, P. K. *et al.* The source of respiratory syncytial virus infection in infants: a household
cohort study in rural Kenya. *J Infect Dis* **209**, 1685-1692 (2014).
https://doi.org:10.1093/infdis/jit828

32   Agoti, C. N. *et al.* Transmission networks of SARS-CoV-2 in Coastal Kenya during the first two
waves: a retrospective genomic study. *eLife* **11**, e71703 (2022).
https://doi.org:10.7554/eLife.71703

33   Gomaa, M. R. *et al.* Incidence, household transmission, and neutralizing antibody
seroprevalence of Coronavirus Disease 2019 in Egypt: Results of a community-based cohort.
PLoS Pathog **17**, e1009413 (2021). https://doi.org:10.1371/journal.ppat.1009413

34   Cohen, C. *et al.* SARS-CoV-2 incidence, transmission, and reinfection in a rural and an urban
setting: results of the PHIRST-C cohort study, South Africa, 2020-21. *Lancet Infect Dis* (2022).
https://doi.org:10.1016/s1473-3099(22)00069-x

35   Semakula, M. *et al.* The secondary transmission pattern of COVID-19 based on contact tracing in
Rwanda. *BMJ Global Health* **6**, e004885 (2021). https://doi.org:10.1136/bmjgh-2020-004885

36   Ratovoson, R. *et al.* Household transmission of COVID-19 among the earliest cases in
Antananarivo, Madagascar. *Influenza Other Respir Viruses* **16**, 48-55 (2022).
https://doi.org:10.1111/irv.12896

37   Herman-Roloff, A. *et al.* Adapting Longstanding Public Health Collaborations between
Government of Kenya and CDC Kenya in Response to the COVID-19 Pandemic, 2020-2021.
Emerg Infect Dis **28**, S159-s167 (2022). https://doi.org:10.3201/eid2813.211550

38 Orangi, S. *et al.* Epidemiological impact and cost-effectiveness analysis of COVID-19 vaccination
in Kenya. *BMJ Glob Health* **7** (2022). https://doi.org:10.1136/bmjgh-2022-009430
39 Bendall, E. E. *et al.* SARS-CoV-2 Genomic Diversity in Households Highlights the Challenges of
Sequence-Based Transmission Inference. *mSphere* **7**, e00400-00422 (2022).
https://doi.org:doi:10.1128/msphere.00400-22
40 Agoti, C. N. *et al.* Genomic analysis of respiratory syncytial virus infections in households and
utility in inferring who infects the infant. *Sci Rep* **9**, 10076 (2019).
https://doi.org:10.1038/s41598-019-46509-w
41 Gallego-García, P. *et al.* Limited genomic reconstruction of SARS-CoV-2 transmission history
within local epidemiological clusters. *Virus Evolution* **8** (2022).
https://doi.org:10.1093/ve/veac008
42 Scott, J. A. *et al.* Profile: The Kilifi Health and Demographic Surveillance System (KHDSS). *Int J*
Epidemiol **41**, 650-657 (2012). https://doi.org:dys062 [pii]
10.1093/ije/dys062
43 Tyson, J. R. *et al.* Improvements to the ARTIC multiplex PCR method for SARS-CoV-2 genome
sequencing using nanopore. *bioRxiv* (2020). https://doi.org:10.1101/2020.09.04.283077
44 O'Toole, Á. *et al.* Assignment of epidemiological lineages in an emerging pandemic using the
pangolin tool. *Virus Evol* **7**, veab064 (2021). https://doi.org:10.1093/ve/veab064
45 Rambaut, A. *et al.* A dynamic nomenclature proposal for SARS-CoV-2 lineages to assist genomic
epidemiology. *Nat Microbiol* **5**, 1403-1407 (2020). https://doi.org:10.1038/s41564-020-0770-5
46 Hadfield, J. *et al.* Nextstrain: real-time tracking of pathogen evolution. *Bioinformatics* **34**, 4121-
4123 (2018). https://doi.org:10.1093/bioinformatics/bty407
47 Yu, G., Smith, D. K., Zhu, H., Guan, Y. & Lam, T. T.-Y. ggtree: an r package for visualization and
annotation of phylogenetic trees with their covariates and other associated data. *Methods in*
Ecology and Evolution **8**, 28-36 (2017). https://doi.org:https://doi.org/10.1111/2041-
210X.12628
48 Sagulenko, P., Puller, V. & Neher, R. A. TreeTime: Maximum-likelihood phylodynamic analysis.
Virus Evol **4**, vex042 (2018). https://doi.org:10.1093/ve/vex042
611