

1 Childhood deprivation and Health related quality of life and
2 associated factors among Pediatric cancer patients at National
3 Hospital, Tanzania
4

5 Authors: Mwanaheri Chubi^{1*}, Stephen Kibusi², Lulu Chirande³, Shakiru Juma⁴

6 *Corresponding author

7
8 ¹Department of Clinical Nursing, School of Nursing and Public Health, The University
9 of Dodoma, Dodoma Tanzania

10 Email: chubimwana@gmail.com

11 ²Department of Clinical Nursing, School of Nursing and Public Health, The University
12 of Dodoma, Dodoma Tanzania

13 Email: skibusi@gmail.com

14 ³Department of Pediatric and child health, School of Medicine, Muhimbili Univeristy of
15 Health and Allied Sciences, Dar es salaam, Tanzania

16 Email: chirandelulu@yahoo.com

17 ⁴Department of Pediatric and child health, School of Medicine, The University of
18 Dodoma, Dodoma Tanzania

19 Email: shakiluj@gmail.com

20

21

22

23

24

25 **Abstract**

26 **Introduction**

27 Pediatrics cancer is one of the most important life-threatening, non-communicable diseases
28 worldwide. However, pediatric cancer patients suffer from physical disabilities associated with
29 cancer treatment. However, there is unclear information about level of health-related quality of
30 life and associated factors. Aimed of this study was to assess level of Health-Related Quality
31 of Life among pediatric cancer patients.

32 **Methods**

33 A hospital-based Analytical cross-sectional study design with a quantitative approach was
34 employed among 91 conveniently sampled pediatric cancer patients with their
35 parents/caregivers. Self-administered questionnaires were used to collect data. Data were
36 analyzed using SPSS computer software program version 25. One-way ANOVA and linear
37 regression were used to quantify and establish an association between Childhood Deprivation,
38 Health-Related Quality of Life, and associated factors among Pediatric Cancer Patients at 95%
39 CI, and a 5% level of significance.

40 **Results**

41 Results, the response rate was 92.3% (n = 84). The most prominent kind of pediatric cancer was
42 Acute Lymphoblastic Leukemia 30 (35.7%) and Hodgkin lymphoma 11(13%). The mean age
43 of pediatric cancer patients was 10.26±3.90 years while 59 (70.2%) were male. In linear
44 regression, Childhood Deprivation ($\beta=-1.640$; P-value< 0.05) on PedsQL™ 4.0 and ($\beta= -2.175$;
45 P-value< 0.01) on PedsQL™ 3.0. Findings imply that decreases the level of Childhood
46 Deprivation, the chance of improving Quality of life increases.

47 **Conclusion**

48 The magnitude of acute lymphoblastic leukemia is higher among pediatric cancer patients.
49 There was a significant link between pediatric cancers and childhood deprivation and thus,
50 compromised quality of life. Innovative pediatric cancer care policies, guidelines, and or
51 strategies may need to be advocated to address the problem accordingly.

52 Introduction

53 Pediatrics cancer is one of the most important life-threatening non-communicable
54 diseases worldwide. It has been predicted that more than 1,000 children will be diagnosed with
55 cancer every day and 400,000 children will be diagnosed with cancer every year, most living in
56 low- and middle-income countries [1]. Pediatric cancer patients are individuals who have been
57 diagnosed with cancer from 0 to 18 years [2] with a survival rate of more than 80% in high-
58 income countries and 5 to 25% in low and middle-income countries respectively [3].

59 However, little is documented about pediatric cancer patients in sub-Saharan Africa as
60 there is no population database of pediatric cancers in Africa, making it difficult to know the
61 true prevalence of pediatric cancers [4]. estimated a mortality rate of 4.6/100,000, from
62 pediatric cancer with an incidence rate of 0.5 as high, with the mortality rate being twice as high
63 for low- and middle-income countries where they are disadvantaged in terms of education,
64 health, and standard of living [5,6].

65 Although evidence shows that the survival rate has significantly increased following the
66 advancement of pediatric cancer therapies such as multiple chemotherapies, surgery,
67 radiotherapy, supportive care, and bone marrow transplantation [7]. Furthermore, studies found
68 that childhood cancer has the best 5 years relative survival rate after diagnosis and a high rate
69 in children with retinoblastoma and ALL tumor types had more than 50% survival in Nairobi
70 compared to Zimbabwe and Uganda, (8), and less than 18 months survival rate to Burkitt
71 lymphoma in Tanzania [9]. Nevertheless, the prompt number of a male has been observed in
72 many countries compared to female pediatric cancer patients most of them spend much time
73 receiving cancer treatment approximately six months to 3 years [10], after diagnosis.

74 In Tanzania, 18.8% of pediatric cancer patients were diagnosed in 2019, with an
75 incidence of 1.42/100,000 whereby Wilms tumor was 17.2%, retinoblastoma 16.8%, and acute
76 lymphoblastic leukemia was 15.9%, furthermore, Burkett lymphoma and Wilm's tumor has
77 been presented mostly in lake zone region [11]. The World Health Organization has launched
78 the Global Initiative for Childhood Cancer (GICC) to maximize the survival rate by up to 60%
79 by 2030, worldwide [12].

80 Despite the substantial gains in recent years, due to major support rendered by developed
81 countries and various international organizations, to support the government in the eradication

82 of child poverty, evidence shows that human basic needs are deprived and significantly remain
83 a major health problem among children [13]. However, there is limited published data about
84 childhood deprivation in pediatric cancer patients worldwide, few studies reveal that in pediatric
85 cancer patients living in the most deprived areas at diagnosis [14], and associated with poor
86 quality of life. Furthermore, evidence shows that most pediatric patients who were diagnosed
87 with cancer disease are hard to reach cancer treatment centers early as they should travel a long
88 distance for care [11].

89 In Tanzania, about 88.9% of children suffer from more than three deprivations and half
90 of them suffer from more than five deprivation including insufficient nutrition, health,
91 protection, education, information, sanitation, water, and housing [15]. Furthermore, studies
92 found that children living in poverty face many deprivations to their human rights, survival,
93 health and nutrition, education, and participation [16]. In addition, a study done in Malaysia
94 revealed that children living in poor (housing, and living conditions) and scarcity health care
95 delivery are significantly vulnerable to physical, social, school, and emotional problems [17]
96 which is among of the factors associated to lower mean scores of health related quality of life
97 of pediatric cancer patients.

98 Health-Related Quality of life is defined as perceived physical, emotional, mental, and
99 social well-being that is affected by a disease condition and its treatment [18]. WHO, Global
100 Initiative Childhood Cancer on its strategies approach to cure all, based on improving health-
101 related quality of life by maximizing the abilities of pediatric cancer patients to achieve
102 individual goals while reaching productive fellows of society [4]. Furthermore, it is expected
103 that the pediatric cancer patient should be free from the major physical and psychological
104 challenges which will improve health related quality of life by alleviating those effects of
105 cancer treatments such as learning difficulties, growth, and cognitive problems [19], by
106 maximizing early interventions

107 Studies found that age at diagnosis, sex, place of residence, type of cancer disease, medical
108 treatment, [20,21], level of education, marital status of the parents/caregiver, employment, poor
109 housing and living standards [22], impact the perceived Health Related Quality of Life of
110 pediatric cancer patients and their parents/caregivers in both generic core and and specific
111 cancer module. Where Pediatrics Quality of Life Inventory (PedsQLTM) Generic core 4.0

112 measures all pediatrics with various chronic diseases and Cancer module 3.0 is the specific for
113 pediatric cancer patients and measure the severity of the disease [18,23,24].

114 So far, some study shows there is a large disparity in quality of life between high and low-
115 income countries. Evidence found that there is significant variation in HRQOL perceived by
116 pediatric cancer patients and their parents on PedsQL scores between countries, whereby the
117 US reported high scores for physical, emotional, and school functioning [25]. Despite the
118 increased incidence of pediatric cancer, childhood deprivation and Health-Related Quality of
119 Life (HRQoL) among pediatrics diagnosed with cancer are not known in Tanzania. This implies
120 that there is a need to know the status of childhood deprivation and Health-related quality of
121 life and associated factors to pediatric cancer patients in Tanzania. The importance of this study
122 is to highlight Sustainable Development Goal number 3 (SDG3), sub-goal 3.1 (Health and well-
123 being), which is to ensure healthy living and promote well-being for all people of all ages.

124 Tanzania is one of the developing countries with an increasing number of pediatric cancer
125 patients in a situation MOHCDGEC has launched several initiatives to address the burden.
126 These initiatives include modern and accurate diagnostic technologies and multimodal therapy
127 including chemotherapy, radiotherapy, surgery and rehabilitation services to improve the
128 quality of life of all pediatric cancer patients to ensure all requirements are easily accessed.
129 Currently, 9 Hospitals in Tanzania provide cancer therapy including Muhimbili National
130 Hospital which provides all modalities that are subject to increases in the survival rate of
131 pediatric cancer patients from 0 to 5 years, where about 50% of all children diagnosed with
132 cancer in 2019 survive, [26,27].

133 However, challenges related to diagnosis and treatment are critical, pediatric cancer
134 patients experience stressful physical and emotional suffering that could greatly affect their
135 well-being including lack of self-esteem, physical incapacity, and poor educational progress
136 [28]. Furthermore, they are subjected to fear, anxiety, and depression and also weakening of
137 immunity [28,29], related to pain, nausea and vomiting, fatigue, weakness, loss of limb
138 function, attention deficit problems and loss of hair while on the cancer treatment period [28].
139 Hence if the problem is left unattended, increases long-term effects and mortality. Furthermore,
140 most of the previous studies have been conducted to establish the prevalence, incidence, needs,
141 and concerns of childhood cancer [3,9,11,29].

142 They helped to expose early, late and long-term effects of cancer treatment, risk of secondary
143 malignancies, to risk of recurrence and also they confront many issues during the treatment
144 period [30], which include childhood deprivation and social problems. Furthermore, their
145 experience has been associated with an increased risk of detrimental psychosocial effects
146 impacting emotional health, socialization, educational and vocational achievement, and health
147 care access [31].

148 Many questions remain unanswered regarding interrelationship issues related to Deprivation
149 and Quality of Life in this unique group. Therefore, this study aims to assess Childhood
150 Deprivation and Health-Related Quality of Life among pediatric cancer patients, which could
151 help the health profession to initiate comprehensive and consistent early intervention that could
152 improve Quality of Life.

153 **Methods and Materials**

154 **Research design**

155 It is a hospital-based, analytical cross-sectional study design with a quantitative research
156 approach that was held to find the association between variables including childhood
157 deprivation and HRQoL among pediatric cancer patients with their parents/caregivers

158 **Study setting**

159 The study was conducted in the pediatric oncology unit at Muhimbili National Hospital,
160 which is a National referral hospital located at Ilala, Dar es Salaam Tanzania (Eastern Indian
161 Ocean coast of Africa), with 1,500 beds capacity, attending 1,000 to 1,200 outpatients per week
162 and admitting 1,000 to 1,200 inpatients per week. The actual data was collected at the pediatric
163 oncology unit whereby pediatric cancer patients were found from all over the country ranging
164 from 0 to 18 years old. It is a hub of pediatric oncology in Tanzania, with the unit attending 300
165 to 350 new cancer children each year [26].

166 Furthermore, approximately more than 150 pediatric cancer patients attend follow-up visits
167 at pediatric oncology units per month, this includes all from cancer Hostels and homes with a
168 revisit plan either weekly or monthly depending on their progress and treatment schedule. The
169 unit runs its clinic on 3 alternative days per week (Monday, Wednesday, and Friday). The study
170 were conducted from 6th may to 30th June 2022.

171 **Study population**

172 The primary subjects in this study were confirmed pediatric cancer patients aged 5 to 18
173 years who received cancer treatment at the oncology unit (clinic) after being reviewed by an
174 oncologist and planned for treatment.

175 **Inclusion and exclusion criteria**

176 All pediatric patients who confirmed they had cancer disease and were on cancer
177 treatment at the oncology clinic were included in the study aged from 5 to 18 years, and
178 understood and communicate verbal instructions.

179 **Sampling and Sampling procedure**

180 The following formula adopted from Cochran (1977) was used to determine the minimum
181 sample size (n) by mean that was required for this study. A convenient sampling technique was
182 used to recruit pediatric cancer patients aged 5 to 18 years who already completed an induction
183 course of cancer therapy (Tembe et al, 2021). The patient's files were used to find the diagnosis,
184 phase, and types of cancer treatment of the child before recruitment. As shown in Fig 1.

185 **Variables**

186 **Outcome variables**

187 The two dependent variables were assessed including Health-Related Quality of Life and
188 Childhood Deprivation among Pediatric cancer patients.

189 **Predictors**

190 The variables show the association on effect to dependent variable including social
191 demographic data of Pediatric cancer patients and their Parents/caregivers and the disease
192 profile

193 **Data source**

194 The study used and modified the Social demographic characteristics from Tanzania
195 malaria indicator survey of 2017 [32] and disease status from previous source [24] [18,24,34]
196 and Multidimensional Poverty Index(MPI) was developed by Sabina Alkire and Foster in 2007,
197 by using demographic and health survey data from USAID and from UNICEF (Multiple
198 indicator cluster surveys (MICS) [16] and Pediatrics quality of Life inventory (PedsQL™ 4.0

199 Generic core scale and PedsQLTM 3.0 Cancer module) from American Pediatric Cancer
200 Association [35]

201 **Measures and instrumentation**

202 **Social demographic characteristics;** the data were collected using designed
203 structured questionnaire, including pediatric cancer patients characteristic: Age, sex, level of
204 education, place of residence and disease status. Parents/caregivers information including age,
205 sex, level of education, employment status, marital status and commenced caring the child

206 **Childhood deprivation;** defined as lack of essential needs such as nutrition, water
207 and sanitation facilities, shelter, access to basic health care services, education, participation,
208 and protection leaving children not capable to enjoy the right to reach their full potentials as
209 members of the community [32]. In this study, Global Multidimensional Poverty Index 2014
210 (MPI) scale used to measure poverty for catching multiple childhood deprivation including
211 education, health and standard of living. Lack of essential needs among pediatric cancer patients
212 resulting on unable to fulfill and participate various activities as a member of community.

213 The Global Multidimensional Poverty Index2014 (MPI) scale was used to examine each
214 person's deprivations across 10 with indicators in three equally weighted dimensions such
215 health, education and standard of living (Nutrition, Child mortality, Cooking fuel, Sanitation,
216 Drinking water, Electricity, Housing, Assets, Years of schooling and school attendance)
217 Whereas each indicator is equally evaluated within its dimension, health and education
218 indicators are weighted 1/6 each and the standard of living indicators are weighted 1/18 each.
219 However, the normal MPI ranges from 0 to 1 (the domain was weight by using scale of 0 and
220 1) where 0 if the pediatric cancer patient fulfils the wealth criteria and 1 when does not fulfill
221 wealth criteria. However, the standard of living involved 2 subscales one on item 6 and second
222 on item 7 making a total of 12 questions.

223 **Health Related Quality of life;** defined as perceived physical, emotional, mental
224 and social well-being is affected by a disease condition and its treatment, [18]. The Pediatric
225 Quality of Life Inventory Version 4.0. (PedsQL™ 4.0) Generic core scale, used to assess the
226 Quality of life of pediatric patients with cancer diseases. The scale comprises with 23-item with
227 4 components such as Physical Functioning score (8 items), Emotional Functioning score (5
228 items), Social Functioning score (5 items), and School Functioning score (5 items). The scales

229 contained in both child report design include 5– 7 (young child), 8–12 (child), and 13–18
230 (adolescent).

231 The child report form for children completed as self-administered interview questions.
232 (24,37). Items are scored on a 5-point response scale where by (0 = never a problem, 1 = almost
233 never a problem, 2 = sometimes a problem, 3 = often a problem, 4 = almost always a problem).
234 Whereas the specific domain and entire score is calculated from the equivalent questions,
235 ranging from 0 to 100 (0=100, 1=75, 2=50, 3=25, 4=0). Furthermore, for ages 5-7 years' items
236 are scored on a 3-point response scale (0 = not at all, 2 = sometimes a problem, 4 = a lot of
237 problems).

238 The PedsQL™ 3.0 cancer module will be used to obtain the subject's information
239 regarding HRQL. The module is a multidimensional instrument having eight domains which
240 consist of 27 items that are covered by five scales: 1 (pain and hurt-two items),2 nausea (5
241 items), 3 procedural anxiety (3 items), 4 treatment anxiety (3 items), 5 worries (3 items), 6
242 cognitive problems (5 items), 7 perceived physical appearance (3 items) and 8 communication
243 (3 items). Items are scored on a 5-point Likert scale whereby (0 = never a problem, 1 = almost
244 never a problem, 2 = sometimes a problem, 3 = often a problem, 4 = almost always a problem
245 for the children aged 8 to 18 self-report and parent proxy report. children aged 5 to 7 years have
246 a 3-point Likert scale, where 0=not at all, 2= sometimes and 4 =a lot. The data was collected
247 by using a visual aid (happy face, neutral face, and sad face) for ages 5-7 years, as a tool
248 direction. Whereas the specific domain and entire score are calculated the same as in PedsQL™
249 4.0 [35].

250 However, both Likert was transformed and shared with all age groups to 3 Likert scales
251 0=never a problem, 2 sometimes, and 4 almost always a problem where (0=100, 2=50, and
252 4=0). Furthermore, scores ranged from 27 to 135 for PedsQL 3.0 cancer module and 23 to 115
253 for PedsQL 4.0 Generic scale. Therefore, for this study, those who scored higher means have a
254 better Quality of Life (above 69.7), and below are considered a poor quality of life [35,38],
255 since there is no median severity of the disease

256 **Reliability of the study**

257 On MPI, the Exploratory factor analysis was performed for reduction and determine
258 weights of each item set at 0.3 as recommended by previous studies [37,38]. The correlation

259 coefficient was set at a cut-off point of ≥ 0.30 , whereas, the Kaiser-MeyerOklin (KMO) value
260 was set at 50% with a 5% level of significance as shown in table 3.1. 11 items were subjected
261 to an initial solution of which two items 1 and 4 were extracted as they weighed < 0.3 . The
262 second exploratory factor analysis involved 9 items that weighed > 0.3 and were for further
263 analysis. The total variance was explained by factor analysis 3 (initial Eugen value = 13.317 and
264 the cumulative percent = 63.646)

265 The descriptive analysis was done which was subjected to establish mean, standard
266 deviation, percent, variance, range, and normality statistics including skewness and kurtosis.
267 However, the data for childhood deprivation were approximately normally distributed, whereby
268 Kolmogorov statistical value was 0.112 (DF-84), P-value < 0.05 with skewness of 1.407, and
269 kurtosis 1.381. Therefore, the non-parametric measurement was obtained and the mean was
270 used as a central tendency to define the endpoint of childhood deprivation scores above the
271 mean were treated as deprived or otherwise not deprived. As shown in table 3.1 mean score and
272 standard of error for the cut point of level of childhood deprivation among pediatric cancer
273 patients was 3.32 (0.215), with a minimum of 0 and a maximum of 8.

274 The Pediatric Quality of life, the actual Cronbach alpha for the PedsQL™ 4.0 Generic
275 core scale was 0-0.7 and the PedsQL™ cancer module inventory were ranging from 0- 0.7. The
276 tool was tested before conducting the study to assess it consistently, by providing a -pre-testing
277 questionnaire in a geographical location outside of the study area, which has been sampled in
278 the study. 16 (10%) pediatric patients with sickle cell disease were interviewed by using MPI,
279 PedsQL™ 4.0 and PedsQL™ 3.0, was tested by using the Cronbach alpha values, which
280 measured the internal consistency. The place of residence was used to test the tools. Both
281 Health-related Qualities of life were excellent at Cronbach alpha 0.7. However, the MPI was
282 excellent with Cronbach alpha ranging from 0.3-0.72 points in factor analysis after increases of
283 illitaration to 75% in the first round, because the sample size was below 300 participants.

284 **Sample size**

285 The following formula adopted from Cochran 1977 was used to determine the minimum sample
286 size (n) that was required for this study

$$287 \quad n = \frac{\sigma^2 Z^2}{e^2}$$

$$288 \quad e^2$$

289 Whereas

290 n = Minimum sample size

291 σ = standard deviation (13.965 for the children aged ≥ 5 years (18)

292 z = significant level 1.96

293 e = acceptable sampling error (3%)

294 Therefore

295 $N = (13.965 \times 1.96 / 3)^2$

296 N was 83 with an attrition rate of 10% of no response, therefore the final calculation was 91.

297 **Analytical strategy**

298 Obtained data was managed by manually cross checked for completeness and accuracy
299 to detect any missing information and was entered to SPSS version 25 so as to clean the data
300 by running frequency, and to view if it is skewed or not skewed to detect any extreme value and
301 opt the type of data analysis model. Descriptive statistics were performed to establish social
302 demographic characteristics of pediatric cancer patients and their parents/caregivers and also
303 were used to find the level of childhood deprivation and health related quality of life.

304 Cross tabulation of frequency and percentage of socio demographic characteristics and
305 disease profile were performed. One-way ANOVA and independent t-test two tailed was used
306 to find the difference between variables of sociodemographic characteristics and outcome
307 variables. Linear regression was used to investigate the association between outcome variable
308 and variable of interest including socio demographic characteristics and disease profile.

309 **Ethics approval and consent to participate**

310 Permission to conduct the study was brought from the University of Dodoma through the
311 Institution Research Review Committee (IRRC), and the Executive director of MNH to allow
312 the study to be conducted at the oncology unit. The informed consent form was obtained from
313 parents/caregivers and assent was obtained from pediatric cancer patients by the Principle
314 Investigator before being interviewed. No names of participants was appearing on the
315 questionnaire, only numbers were coded to ensure confidentiality. The data collection
316 procedure was conducted in an unoccupied separate room to ensure privacy. The participants

317 had the right to withdraw from the study. However, during the study, no challenge or any change
318 arises regarding Health Related or childhood deprivation among pediatric cancer patients. Data
319 was handled and managed securely by using a folder with a key in the IRRC computer and not
320 shared with other users for any purpose out of the intention of this study.

321 **Results**

322 **Socio-demographic information of the pediatric cancer patients**

323 Pediatric cancer patients and their parents/carers qualified and participated in the study
324 with a 92.3% (n=84) response rate of 91. As shown in Table 1, the mean age of 84 pediatric
325 cancer patients was 10,263,90 with a minimum of 5 and a maximum of 18 years The most
326 prominent age group ranged between 8 and 12 years 40.5% (n=34), as most of them were male
327 70.2% (n=59) compared to females 29.8% (n=25). Most of pediatric cancer patients living in
328 urban 38.1% (n=32), Rural 36.9% (n=31) and Peri-Urban 25% (n=21). Only 47.6% (n=40) of
329 them attended school during the study compared to 52.4% (n=44) of them are not going to
330 school, while about 63.1% (n=53) had primary school level.

331 **Table 1. Socio-demographic information of the pediatric cancer patients**
332 **(n=84)**

Characteristics	Frequency(n)	Percent (%)
Age in years		
Mean(SD) 10.26±3.90		
Min 5		
Max 18		
Age of group in years		
5 to 7	22	26.2
8 to 12	34	40.5
13 to 18	28	33.3
Gender of child		
Male	59	70.2
Female	25	29.8
Place of residence		
Urban	32	38.1
Peri-urban	21	25.0
Rural	31	36.9
School attendance		
Attending	40	47.6

Not attending	44	52.4
Level of education of the child		
Not went to school	18	21.4
Kindergarten	8	9.5
Primary school	53	63.1
Secondary school	5	6.0

333 **Source:** Field Data (2022)

334 **Socio-demographic characteristics of parents/caregivers**

335 Table 2 shows that the mean age of 84 qualified parents/carers who participated in the
 336 study was 38.62±10.87 years, with a minimum of 18 and a maximum of 68 years. The most
 337 conspicuous 58.3% (n=49) of parents/caregivers aged 20-39, 52% (n=44) were female and
 338 47.6% (n=40) were male, 41.7% (n =35) were mothers. Also whom 71.6% (n=61) were married,
 339 58.3% (n=49) had elementary school, 85.7% (n= 72) were informally employed and about
 340 78.6% (n=66) started caring for the child before the illness. The results may indicate that the
 341 majority of caregivers are young and capable of caring for pediatric cancer patients and
 342 influencing the patient's good health.

343 **Table 2. Socio-demographic characteristics of Parents/caregivers (n=84)**

Caregiver characteristics	Frequency	Percent
Age in years		
Mean(SD) 38.62±10.87		
Min 18		
Max 68		
Age by the group in years		
10 to19	1	1.2
20 to 39	49	58.3
40 to 59	31	36.9
60 to 79	3	3.6
Gender		
Male	40	47.6
Female	44	52.4
Marital status of the caregiver		
Single	16	19.0
Married	61	72.6
Divorced	6	7.1
Widow	1	1.2
Level of education of the caregiver		

Never went to school	6	7.1
Primary school	49	58.3
Secondary school	21	25.0
College/university	8	9.5
Employment status of the parent/caregiver		
Formal employment	12	14.3
Informal employment	72	85.7
Commenced caring for the child		
Before illness	66	78.6
During illness	18	21.4

344 **Source:** Field Data (2022)

345 **Disease profile of pediatric cancer patients**

346 As shown in Table 3, the results show that the most prominent pediatric cancer was 35.7%
 347 (n=30) acute lymphoblastic leukemia, followed by 13.7% (n=11) Hodgkin's lymphoma, and the
 348 last one for rhabdomyosarcoma, one for primitive neuroectodermal tumor, one for endometrial
 349 carcinoma and one for esophageal carcinoma. Tumor while most of them confirmed cancer and
 350 started cancer treatment in MNH 89.3% (n=75). The median year of diagnosis was 2019,772.47
 351 with a minimum of 2009 and a maximum of 2022, with the majority of pediatric cancer patients
 352 63.1% (n=53) diagnosing cancer in 2021 and early 2022.

353 The mean time since diagnosis was 1,890.99 with a minimum of 1 and a maximum of 6
 354 years in pediatric cancer patients, while most of them were 1 year 45.2% (n=38), 1 to 2 years
 355 29.8% (n= 25), 3 to 4 years 15.5% (n = 13) and 5 years are only 9.5% (n = 8) since confirmed
 356 disease. About 35.7% (n=30) of them were treated in 4 hospitals before diagnosis, 76.2% (n=64)
 357 started cancer treatment 2 weeks after initial diagnosis, 78.6% (n=66) had received
 358 chemotherapy, 73.8% (n=66) had received chemotherapy, 73.8% (n=64) had received cancer
 359 treatment =62) psychological support. However, 54.8% (n=46) of pediatric cancer patients
 360 hospitalized 1 to 3 times before 3 months, 42.9% (n=36) had stomach pain, and 36.9% (n=31)
 361 had fever, while receiving cancer treatment. Only 44.0% (n=37) of pediatric cancer patients
 362 attended school/university in the last 3 months.

363 The results imply that the number of children diagnosed with cancer is currently still
 364 increasing, while the majority of them delayed the start of cancer treatment, as most of them
 365 confirmed cancer and started cancer treatment at the National Referral Hospital, where all

366 modalities of cancer therapy are available. However, most pediatric cancer patients experience
367 stomach pain as the main problem while receiving cancer therapy, especially chemotherapy.

368 **Table 3. Disease profile of pediatric cancer patients (n = 84)**

Characteristics	Frequency	Percent
Year of diagnosis		
Mean(SD); 2019.77 ±2.47		
Min 2009		
Max 2022		
Year of cancer disease diagnosed		
2009	1	1.2
2010-2015	2	2.4
2016-2020	28	33.3
2021-2022	53	63.1
Time since diagnosis in years		
Mean(SD); 1.89 ±0.99		
Min 1		
Max 6		
Time since diagnosis by group(years)		
<1 year	38	45.2
1 to 2 years	25	29.8
3 to 4 years	13	15.5
≥5 years	8	9.5
No hospitals treated the child before diagnosis		
1	9	10.7
2	24	28.6
3	21	25.0
≥4	30	35.7
Diagnosis		
Acute Lymphoblastic Leukemia	30	35.7
Hodgkins Lymphoma	11	13.1
Cellular carcinoma with Xeroderma Pigmentosum	10	11.9
Wilm's Tumor	9	10.7
Burkitt Lymphoma	8	9.5
Retinoblastoma	5	6
Acute Amyloid Lymphoblastic	4	4.8
Osteosarcoma	3	3.6
Rhabdomyosarcoma	1	1.2

Primitive Neuroectodema tumor	1	1.2
Endometrial Carcinoma	1	1.2
Oesophageal tumor	1	1.2
Cancer medication started after diagnosis		
≥ 2 weeks	60	71.4
< 2 weeks	24	28.6
Treatment		
Chemotherapy	66	78.6
Combination (chemotherapy, radiotherapy, and surgery)	7	8.3
Combination (chemotherapy and radiotherapy)	2	2.4
Combination (chemotherapy and surgery)	9	10.7
Hospitalization in prior 3 months		
1 to 3 times	46	54.8
>3 times	35	41.7
No hospitalization	3	3.6
Psychosocial counseling		
Received	62	73.8
Not received	22	26.2
Activities done in the last 3 months		
School/studying	37	44.0
Playing/exercises	23	27.4
Hospitalized	12	14.3
Too ill	9	10.7
Hand capped/ cannot work	3	3.6
Complications related to cancer treatment		
Stomach pain	36	42.9
Fever	31	36.9
Heart palpitation	7	8.3
Anemia	3	3.6
Combination; Fever, stomach pain, heart palpitation, loss of appetite, and body malaise	2	2.4
Lack of appetite and fatigue	2	2.4
Diarrhea	1	1.2
Pneumonia	1	1.2
None	1	1.2

369 **Source:** Field Data (2022)

370 **The proportion of the type of pediatric cancer by sex,age and**
371 **hospitalization prior to 3 months**

372 Descriptive analysis was performed to determine the proportion of cancer type by gender,
373 age group and hospitalization prior to 3 months in pediatric cancer patients as shown in table
374 4-6. The findings show that males were most affected at 70.2% (n=59) compared to females. In
375 addition, ALL was 45.8% (n=27) in males and 12.0% (n=3) and Hodgkin's lymphoma 15.3%
376 (n=9) in males and 8.0% (n=2) most prominent in female pediatric cancer patients. In addition,
377 the least cancer type was affected only in female pediatric cancer patients, including 4.0% (n =
378 1) Primitive Neuro-Ectodermal tumors (PNET), 4.0% (n = 1) endometrial cancer, and 4.0% (n
379 = 1) Esophageal tumor.

380 The proportion of cancers by age group. Results showed that pediatric cancer patients
381 aged 8-12 and 13-18 years were most affected by ALL compared to other cancers, at 38.2%
382 (n=13) and 42.9% (n=12), respectively. However, Hodgkin's lymphoma and Wilms' tumor
383 affected most pediatric cancers at ages 5-7 and 8-12 years, with the same proportions of 22.7%
384 (n=5) and 11.8% (n=4), respectively. Therefore, the most affected age group with pediatric
385 cancer was between 8 and 12 years. In terms of hospitalization prior to 3 months, most pediatric
386 cancer patients with ALL were admitted 1 to 3 times and more than 3 times, 39.1% (n=18) and
387 31.4% (n=11), respectively Hodgkin lymphoma and cellular carcinoma with xerodermic
388 pigmentosa as shown in Table .

389 **Table 4. The proportion of diagnosis by age group (years) of pediatric cancer patients n=84**

Variable	ALL	Retino brastobl astoma	wilm's tumor	hodgikins lymphom a	Burkitt lympho ma	Osteo sarcoma	Rhabd o myosar coma	Cellular carcinoma with xeroderma pigmentosa	AML	PNET	Endom etrial carcino ma	oesophage al tumor	Total
5 to 7	5(22.7%)	2(9.1%)	5(22.7%)	5(22.7%)	2(9.1%)	0(0.0%)	0(0.0%)	2(9.1%)	1(4.5%)	0(0.0%)	0(0.0%)	0(0.0%)	22(26.2%)
8 to 12	13(38.2%)	1(2.9%)	4(11.8%)	4(11.8%)	2(5.9%)	0(0.0%)	1(2.9%)	5(14.7%)	2(5.9%)	1(2.9%)	1(2.9%)	0(0.0%)	34(40.5%)
13 to 18	12(42.9%)	2(7.1%)	0(0.0%)	2(7.1%)	4(14.3%)	3(10.7%)	0(0.0%)	3(10.7%)	1(3.6%)	0(0.0%)	0(0.0%)	1(3.6%)	28(33.3%)
Total	30(35.7%)	5(6.0%)	9(10.7%)	11(13.1%)	8(9.5%)	3(3.6%)	1(1.2%)	10(11.9%)	4(4.8%)	1(1.2%)	1(1.2%)	1(1.2%)	84(100%)

390

391

392 **Table 5. The proportion of diagnosis by age group (years) of pediatric cancer patients n=84**

Variable	ALL	Retino brastobla stoma	wilm's tumor	hodgikins lymphom a	Burkitt lymphom a	Osteo sarcoma	Rhabdo myosar coma	Cellular carcinoma with xeroderma pigmentosa	AML	PNET	Endometri al carcinoma	oesophage al tumor	Total
5 to 7	5(22.7%)	2(9.1%)	5(22.7%)	5(22.7%)	2(9.1%)	0(0.0%)	0(0.0%)	2(9.1%)	1(4.5%)	0(0.0%)	0(0.0%)	0(0.0%)	22(26.2%)
8 to 12	13(38.2%)	1(2.9%)	4(11.8%)	4(11.8%)	2(5.9%)	0(0.0%)	1(2.9%)	5(14.7%)	2(5.9%)	1(2.9%)	1(2.9%)	0(0.0%)	34(40.5%)
13 to 18	12(42.9%)	2(7.1%)	0(0.0%)	2(7.1%)	4(14.3%)	3(10.7%)	0(0.0%)	3(10.7%)	1(3.6%)	0(0.0%)	0(0.0%)	1(3.6%)	28(33.3%)
Total	30(35.7%)	5(6.0%)	9(10.7%)	11(13.1%)	8(9.5%)	3(3.6%)	1(1.2%)	10(11.9%)	4(4.8%)	1(1.2%)	1(1.2%)	1(1.2%)	84(100%)

393

394

395

396

397

398 **Table 6. The proportion of diagnosis by hospitalization before 3 months of pediatric cancer patients n=84**

Variable	ALL	Retino brastobla stoma	wilm's tumor	hodgikins lymphoma	Burkitt lymphom a	osteosar coma	Rhabdom yo sarcoma	Cellular carcinoma with xeroderma pigmentosa	AML	PNET	Endometria l carcinoma	oesophag eal tumor	TOTAL
No Hospitalization	1(33.3%)	0(0.0%)	1(33.3%)	1(33.3%)	0(0.0%)	0(0.0%)	0(0.0%)	0(0.0%)	0(0.0%)	0(0.0%)	0(0.0%)	0(0.0%)	3(3.6%)
1 to 3 times	18(39.1%)	3(6.5%)	4(8.7%)	5(10.9%)	5(10.9%)	2(4.3%)	1(2.2%)	6(13.0%)	1(2.2%)	0(0.0%)	0(0.0%)	1(2.2%)	46(54.8%)
More than 3 times	11(31.4%)	2(5.7)	4(11.4%)	5(14.3%)	3(8.6%)	1(2.9%)	0(0%)	4(11.4%)	3(8.6%)	1(2.9%)	1(2.9%)	0(0.0%)	35(41.7%)
TOTAL	30(35.7%)	5(6.0%)	8(9.5%)	11(13.1%)	8(9.5%)	3(3.6%)	1(1.2%)	10(11.9%)	4(4.8%)	1(1.2%)	1(1.2%)	1(1.2%)	84(100%)

400 **Level of childhood deprivation**

401 A descriptive analysis was carried out, in which mean, frequency, and percentage were
402 determined. As shown in the Fig 1, the data for the degree of childhood deprivation were
403 approximately normally distributed, since all values above the mean were considered
404 disadvantaged in pediatric cancer patients, since the cut-off value was 3.32. The results show
405 that approximately 56.0% (n=47) of pediatric cancer patients were deprived.

406 **Mean scores of childhood deprivation by age**

407 The overall mean values are below the limit in both age categories. However, the
408 observed differences in standard of living where the mean is above the threshold are at ages 5-
409 7 years (3.64±1.81) compared to ages 8-12 and 13-18 years (2.44±1.54) and (2.71±1.76) or as
410 shown in Table 7. The results suggest that most young pediatric cancer patients have a lower
411 standard of living compared to older patients.

412 **Table 7. Mean scores of childhood deprivation by age n=84**

Variable	5-7 yrs.		8-12 yrs.		13-18 yrs.		P-value
	Mean	SD	Mean	SD	Mean	SD	
Total mean score	1.39	0.91	0.97	0.79	1.04	0.86	0.331
Education	0.32	0.48	0.15	0.36	0.14	0.36	0.212
Health	0.23	0.43	0.32	0.48	0.29	0.46	0.746
Standard of living	3.64	1.81	2.44	1.54	2.71	1.76	0.036

413 **Bonferroni alpha Post hoc test of childhood deprivation among** 414 **pediatric cancer patients for age n=84**

415 Multiple comparison tests in pediatric cancer patients showed a difference in significance
416 between those aged 5-7 years and 8-12 years for standard of living, with the p-value being
417 <0.05. However, in other domains, the observed mean difference was not statistically
418 significant, as shown in Table 8. The results reflect that young pediatric cancer patients may
419 deprived on standard of living.

420 **Table 8. Bonferroni alpha Post hoc test of childhood deprivation among**
421 **pediatric cancer patients for age n=84**

Domain	Comparison (years)	MD	S.E	95% CI		P-value
				Lower	Upper	

Education	5 to 7- 13 to 18	0.171	0.107	-0.09	0.43	0.344
	8 to 12-13 to 18	0.004	0.100	-0.24	0.25	1.000
	5 to7-8 to 12	-0.004	0.100	-0.25	0.24	1.000
Health	5 to 7- 13 to 18	-0.096	0.125	-0.40	0.21	1.000
	8 to 12- 13 to 18	-0.058	0.131	-0.38	0.26	1.000
	5 to7-8 to 12	-0.038	0.117	-0.32	0.25	1.000
Standard of living	5 to 7- 13 to 18	1.195	0.462	0.07	2.33	0.035
	8to 12- 13 to 18	0.922	0.481	-0.25	2.10	0.177
	5 to 7- 13 to 18	0.273	0.431	-0.78	1.33	1.000

422 **Source:** Field Data (2022)

423 **Mean score of childhood deprivation by place of residence**

424 As shown in Table 9, the mean total childhood deprivation scores by pediatric cancer
425 patients' city of residence. The results show that the overall mean is below the cut-off (3.32).
426 However, mean living standard scores for those living in rural areas were higher (3.77 ± 1.75 ; p-
427 value < 0.01) than those in urban and peri-urban areas. The finding implicates that living in rural
428 areas could affect pediatric cancer patients, which resulted in poor quality of life.

429 **Table 9. Mean scores of Childhood Deprivation by place of residence n=84**

Variable	Urban		Peri-Urban		Rural		P-value
	Mean	SD	Mean	SD	Mean	SD	
Total mean score	0.77	0.74	0.89	0.75	1.48	0.89	0.257
Education	0.16	0.37	0.24	0.44	0.19	0.40	0.765
Health	0.19	0.39	0.14	0.36	0.48	0.51	0.007
Standard of living	1.97	1.47	2.81	1.44	3.77	1.75	0.000

430 **Source:** Field Data (2022)

431 **Bonferroni alpha Post hoc test of childhood deprivation among** 432 **pediatric cancer patients for place of residence**

433 Multiple comparison tests in pediatric cancer patients showed a significant difference
434 between those living in urban-rural and peri-urban-rural areas in terms of health functioning by
435 the Bonferroni alpha post-hoc test with an adjusted p-value of < 0.024 or 0.020 . In terms of
436 standard of living, the observed difference was between urban and rural. Therefore, the results
437 suggest that pediatric cancer patients living in rural areas are disadvantaged in terms of health

438 and standard of living. In addition, no difference was observed in education as shown in Table
439 10.

440 **Table 10. Bonferroni alpha Post hoc test of childhood deprivation among**
441 **pediatric cancer patients for place of residence n=84**

Variable	Comparison	MD	S.E	95% CI		P-value
				Lower	Upper	
Education	Urban-peri urban	-0.082	0.112	-0.36	0.19	1.000
	Urban-rural	-0.037	0.100	-0.28	0.21	1.000
	Peri urban-rural	-0.045	0.113	-0.32	0.23	1.000
Health	Urban-peri urban	0.045	0.122	-0.25	0.34	1.000
	Urban-rural	-0.296	0.109	-0.56	-0.03	0.024
	Peri urban-rural	0.341	0.122	0.04	0.64	0.020
Standard of living	Urban-peri urban	-0.841	0.441	-1.92	0.24	0.180
	Urban-rural	-1.805	0.396	-2.77	-0.84	0.000
	Peri urban-rural	0.965	0.444	-0.12	2.05	0.098

442 **Source:** Field Data (2022)

443 **Level of Health-Related Quality of Life and associated factors**
444 **among pediatric cancer patients**

445 The descriptive analysis was performed to determine the level of health-related quality of
446 life (HRQoL), with the endpoint being the mean score from the tool such as the Pediatric
447 Quality of Life Inventory, both the generic core scale (PedsQL™ 4.0) child report, while mean
448 for child report (69.7) and parent proxy report (65.4). Therefore, the mean scores above the
449 endpoint were treated with better quality of life and below the mean with poor quality of life

450 **Level of health-related quality of life among pediatric cancer**
451 **patients measured by (PedsQL™ 4.0) Generic core scale**

452 As shown in table 11, the results of PedsQL™ 4.0 show that the total mean scores were
453 lower (67.79±21.63). This finding implies that pediatric cancer patients have a poor quality of
454 life, as the overall mean scores were below the cut-off point (69.7).

455 **Table 11. Total mean scores of the Level of Health-Related Quality Life**
456 **measured by (PedsQL™ 4.0) Generic core scale (n = 84)**

Domain	Mean score	Median	SD	Min	Max
--------	------------	--------	----	-----	-----

Total mean score	67.79	75.94	21.63	18.44	100.00
Physical functioning	66.74	78.75	21.06	18.75	100.00
Emotional functioning	77.44	80.00	17.83	25.00	100.00
Social functioning	86.73	100.00	18.97	30.00	100.00
School functioning	40.25	45.00	28.67	0.00	100.00

457 **Source:** Field Data (2022)

458 **Mean scores of Health-related quality of life measured by**
 459 **(PedsQL™ 4.0) Generic core scale by sex**

460 Table 13 shows that the average total scores of males compared to females were below
 461 the cutoff point (69.7) in both the Child Report. However, only physical functioning was
 462 statistically significant (p-value 0.039). In addition, academic performance in both males and
 463 females was below the threshold but was not statistically significant. This finding suggests that
 464 the quality of life of female pediatric cancer patients improved and got better.

465 **Table 13. Mean scores of Health-related quality of life measured by**
 466 **(PedsQL™ 4.0) Generic core scale by sex (n = 84)**

Domain	Male		Female		P-value
	Mean score	SD	Mean score	SD	
Total mean score	65.46	21.85	72.08	21.35	0.313
Physical functioning	63.51	19.98	74.38	21.96	0.039
Emotional functioning	76.52	18.53	79.50	16.19	0.473
Social functioning	85.68	19.90	89.20	16.69	0.440
School functioning	38.14	27.82	45.24	30.57	0.302

467 **Source:** Field Data (2022)

468 **Mean scores of Health-related quality of life measured by**
 469 **(PedsQL™ 4.0) Generic core scale for age**

470 Table 14 One-way ANOVA analysis was used to establish the relationship through mean
 471 results of the PedsQL 4.0 Generic Core Scale for Child report by age group. The results show
 472 that age groups (13-18 and 8-12 years) had a significant impact (M=70.00±15.32 and
 473 73.33±10.31, respectively; p<0.01) on the health-related quality of life of pediatric cancer
 474 patients. However, the high mean was statistically significant for physical functioning at ages
 475 8–12 and 13–18 years (M=71.59±19.09; 71.14±19.99; p-value 0.001) compared to the 5–7

476 year-olds who reported a low (M=53.12±22.36) in children. However, with respect to school
 477 functioning, the mean was observed to be low for all age groups at ages 5–7, 8–12, and 13–18,
 478 respectively (M=15.45±23.03; 46.50±25.98 and 52.142±4.32; p-value 0.01). The finding may
 479 suggest that the child's perceived physical functioning affects the quality of life of pediatric
 480 cancer patients.

481 **Table 14. Mean Scores of PedsQL™ 4.0 Generic Core Scale for Child and**
 482 **Parents proxy by age group (n=84)**

Domain	5-7 yrs.		8-12 yrs.		13-18 yrs.		P-value
	Mean score	SD	Mean score	SD	Mean score	SD	
Total mean score	57.32	12.34	70.00	15.32	73.33	10.31	<0.01
Physical functioning	53.12	22.36	71.59	19.09	71.14	19.99	<0.001
Emotional functioning	76.81	18.09	78.38	16.45	76.78	19.73	0.925
Social functioning	83.86	24.39	83.52	18.93	92.85	12.12	0.111
School functioning	15.45	23.03	46.50	25.98	52.14	24.32	<0.006

483 **Source:** Field Data (2022)

484 **Bonferroni alpha Post hoc test for health related quality of**
 485 **pediatric cancer patients for age**

486 Multiple comparison tests in pediatric cancer patients showed a difference in significance
 487 between those aged 5-7 years and 8-12;13-18 years for physical and academic functioning, with
 488 the p-value being <0.01. While in the parents' report, only the school function shows significant
 489 differences. However, in other domains, the observed mean difference was not statistically
 490 significant, as shown in Table 15. The results reflect that young pediatric cancer patients may
 491 face physical problems that can lead to developmental delays and limited school attendance and
 492 other problems.

493 **Table 15. Bonferroni alpha Post hoc test for health related quality measured**
 494 **by (PedsQL™ 4.0) Generic Core Scale of pediatric cancer patients**
 495 **for age (n = 84)**

Variable	Comparison (year)	MD	S.E	95% CI		P-value
				Lower	Upper	
Physical functioning	5 to7-8 to 12	-18.47	5.38	-31.62	-5.33	0.003
	5 to 7- 13 to 18	-18.42	5.60	-32.10	-4.73	0.004

	8 to 12-13 to 18	-0.06	5.02	-12.32	12.20	1.000
Emotion functioning	5 to 7-8 to 12	-1.56	4.93	-13.62	10.49	1.000
	5 to 7- 13 to 18	0.03	5.14	-12.52	12.59	1.000
	8 to 12- 13 to 18	-1.60	4.60	-12.84	9.65	1.000
Social functioning	5 to 7-8 to 12	0.33	5.11	-12.17	12.84	1.000
	5 to 7- 13 to 18	-8.99	5.32	-22.01	4.02	0.285
	8to 12- 13 to 18	9.33	4.77	-2.33	20.99	0.162
School functioning	5 to 7-8 to 12	-31.05	6.76	-47.57	-14.53	0.000
	5 to 7- 13 to 18	-36.69	7.04	-53.89	-19.49	0.000
	8 to 12- 13 to 18	-36.69	6.30	-9.76	21.05	1.000

496 **Source:** Field Data (2022)

497 MD-Mean Difference

498 **Mean scores of Health-related quality of life measured by (PedsQL 4.0)**
 499 **Generic core scale among pediatric cancer patients by hospitalization prior**
 500 **to 3 months**

501 Pediatric cancer patients were hospitalized more than 3 times prior to 3 months, where the total
 502 mean scores were below 62.23 ± 20.15 . The overall trend physical functioning and school
 503 functioning decreased quality of life, as shown in Table 16.

504 **Table 16. Mean scores of Health-related quality of life (PedsQL 4.0 Generic**
 505 **core scale) pediatric cancer patients by hospitalization before 3**
 506 **months (n = 84)**

Domain	No Hospitalization		1-3 times		➤ 3 times		P-value
	Mean score	SD	Mean score	SD	Mean score	SD	
Total mean score	79.71	30.47	72.33	21.16	62.23	20.15	0.192
Physical functioning	54.17	39.57	68.55	19.60	65.45	21.49	0.468
Emotional functioning	73.33	23.09	80.76	18.82	73.43	15.57	0.172
Social functioning	76.67	25.17	90.43	18.28	82.71	18.83	0.124
School functioning	48.33b	34.03	49.57	27.93	27.31	24.71	0.002

507 **Source:** Field Data (2022)

508 Multiple comparison tests in pediatric cancer patients showed a difference in significance
 509 between those admitted 1-3 times and more than 3 times for school function 3 months before

510 function by Bonferroni Alpha Post-Hoc Test with an adjusted p-value of <0.001 as shown in
511 Table 17

512 **Table 17. Bonferroni alpha Post hoc test of PedsQL™ 4.0 cancer generic**
513 **scale with pediatric cancer patients for hospitalization prior in 3 months (n**
514 **= 84)**

Variable		Mean D	S.E	95% CI		P-value
				Lower	Upper	
Total mean score	1-2	-9.20	12.56	-39.97	21.49	0.858
	1-3	0.9	12.68	30.09	31.89	0.897
	2-3	-10.09	4.73	-21.66	1.46	0.353
Physical functioning	1-2	-14.38	12.58	-45.14	16.39	0.770
	1-3	-11.28	12.70	-42.33	19.78	1.000
	2-3	-3.09	4.74	-14.68	8.48	1.000
Emotion functioning	1-2	-7.43	10.52	-33.15	18.29	1.000
	1-3	-0.09	10.62	-26.06	25.87	1.000
	2-3	-7.33	3.96	-17.01	2.35	0.203
Social functioning	1-2	-13.77	11.15	-41.03	13.49	0.662
	1-3	-6.05	11.26	-33.57	21.48	1.000
	2-3	-7.72	4.19	-17.98	2.54	0.209
School functioning	1-2	-1.23	15.97	-40.27	37.80	1.000
	1-3	21.02	16.12	-18.39	60.43	0.588
	2-3	-22.25	6.01	-36.95	-7.55	0.001

515 **Source:** Field Data (2022)

516 1- No hospitalization, 2- hospitalization 1-3 times, 3-hospitalization >3 times

517 **Health-Related Quality of Life and associated factors among** 518 **Pediatric cancer patients**

519 As shown in Table 18, regression analysis was performed to find the association between
520 associated factors and HRQoL of pediatric cancer patients and their caregivers using the
521 Pediatrics Quality of Life Inventory (PedsQL™ 4.0) Generic Core Scale. The results show a
522 significant correlation between level of education, gender and age (=5.675; P-value <0.01:
523 =3.685; P-value <0.01: =7.719; P-value <0.01). However, childhood deprivation (= -1.640; P-
524 value <0.05), place of residence (= -3.131; P-value <0.05), work done in the last 7 days (= -6.476;
525 P- value <0.01). hence, the finding implies that higher education, older age, and female gender

526 increase the chance of an improvement in the quality of life. In addition, the chance of better
527 health in pediatric cancer patients increases when childhood deprivation decreases, activities
528 are reduced, and live in urban areas.

529

530 **Table 18: Health related quality of life and associated factors measured by**
531 **PedsQL™ 4.0 (n = 84)**

Variable	B	T	95% CI		P-value
			Lower	Upper	
Child factors					
Childhood deprivation	-1.640	-2.211	-3.116	-0.165	0.030
Place of residence	-3.131	-0.219	-5.797	-0.464	0.022
Age of a child	7.719	5.105	4.710	10.727	0.001
Sex of child	3.685	1.451	-1.368	8.737	0.001
Level of education of the child	5.675	4.395	3.103	8.246	0.000
Work done in the past 7 days	-6.476	-2.530	-11.574	-1.378	0.013
Psychosocial counseling	-7.707	-2.818	-13.155	-2.259	0.006
Types of treatment received	-1.842	-2.529	-3.292	-0.391	0.014
Hospitalization in prior 3 months	-2.197	-0.621	-9.350	4.957	0.538
Most activity done in the last 3 months	1.611	1.247	-1.001	4.223	0.220
Caregiver factors					
Level of education of the caregiver	-3.528	-1.556	-8.109	1.054	0.128
Caregiver employment	5.763	1.184	-4.076	15.603	0.243
Commence caring	-0.417	-0.061	-14.117	13.284	0.951

532 **Source:** Field Data (2022)

533 **Level of health-related quality of life among pediatric cancer**
 534 **patients measured by(PedsQL™ 3.0) Generic core scale**

535 The results in Table 19 show that the mean total score (69.86±25.90) perceived by
 536 pediatric cancer patients corresponded to the cut-off point (69.7). The results suggest that
 537 pediatric cancer patients may have a poor quality of life.

538 **Table 19 Total mean scores of Level of Health-Related Quality of Life**
 539 **(PedsQL 3.0) cancer module (n = 84)**

Domain	Mean	median	SD	Min	Max
Total mean score	69.86	67.29	25.90	4.17	100.00
Pain and hurt	70.98	75.00	21.97	25	100.00
Nausea	76.49	80.00	18.54	0.00	100.00
Procedural anxiety	59.92	58.33	31.49	0.00	100.00
Treatment anxiety	84.92	100.00	23.33	0.00	100.00
Worry	77.48	75.00	22.73	0.00	100.00
Cognitive problem	62.20	77.50	37.29	0.00	100.00
Perceived physical appearance	83.63	100.00	24.42	8.33	100.00
Communication	48.81	50.00	27.48	0.00	100.00

540 **Source:** Field Data (2022)

541 **Mean scores of Health-related quality of life measured by PedsQL**
 542 **3.0 cancer module perceived by pediatric cancer patients for sex**

543 As shown in Table 20 independent t-test two tailed, the overall means are above the cut-
 544 off point and are not statistically significant. However, for procedural anxiety, cognitive
 545 problems, and communication, both male and female pediatric cancer patients had mean scores
 546 below their self-perceived limits. In addition, in the parent report, the mean score is lower in
 547 males (65.59±43.18) on cognitive function, although this was not significant. The finding
 548 implies that most male patients have a poor quality of life compared to female patients.

549 **Table 20. Mean scores of Health-related quality of life (PedsQL 3.0 Generic**
 550 **core scale) perceived by pediatric cancer patients for sex (n = 84)**

Domain	Male		Female		P-value
	Mean	SD	Mean	SD	
Total mean score	70.27	25.48	71.20	27.02	0.521

Pain and hurt	75.68	17.77	78.40	20.50	0.542
Nausea	71.39	22.27	70.00	21.65	0.814
Procedural anxiety	60.45	31.28	58.67	32.84	0.541
Treatment anxiety	83.89	24.84	87.33	19.56	0.901
Worry	77.68	20.56	77.00	27.67	0.485
Cognitive problem	60.33	37.53	66.60	37.13	0.329
Perceived physical appearance	85.88	23.53	78.33	26.13	0.197
Communication	46.89	26.03	53.33	30.71	0.364

551 **Source:** Field Data (2022)

552 **Mean scores of Health-related quality of life measured by PedsQL**
 553 **3.0 cancer module perceived by pediatric cancer patients for age**

554 As shown in Table 21 One-Way ANOVA Analysis of PedsQL Cancer Module 3.0. The
 555 finding shows that there is an association between health-related quality of life by age group
 556 where pediatric cancer patients ages 8-12 and 13-18 years have a high overall mean
 557 (74.31±12.32 and 75.15±9.74, respectively) that is statistically significant (p-value 0.01). The
 558 difference was seen in nausea (p-value 0.020), procedure anxiety (p-value 0.01), and cognitive
 559 problem (p-value 0.000) in the child report. Therefore, the results may suggest that aging may
 560 improve health-related quality of life in pediatric cancer.

561 **Table 21. Mean scores of Health-related quality of life measured by PedsQL**
 562 **3.0 cancer module for age (n=84)**

Domain	5-7 yrs.		8-12 yrs.		13-18yrs.		P-value
	Mean	SD	Mean	SD	Mean	SD	
Total mean score	58.90	11.37	74.31	12.32	75.15	9.74	<0.01
Pain and hurt	64.77	23.98	73.89	22.05	72.32	19.94	0.296
Nausea	67.27	22.50	80.88	16.89	78.39	14.72	<0.020
Procedural anxiety	71.96	33.09	87.99	16.30	91.36	17.49	<0.01
Treatment anxiety	32.19	33.95	71.81	24.36	67.26	23.77	<0.007
Worry	87.87	21.93	78.18	21.12	68.45	22.26	<0.009
Cognitive problem	23.63	33.45	71.32	30.28	81.42	24.18	<0.01
Perceived physical appearance	90.90	18.34	78.18	27.29	84.52	24.08	0.159
Communication	32.57	24.92	52.20	26.29	57.44	26.18	<0.003

563

564 **Bonferroni alpha Post hoc test of PedsQL™ 3.0 cancer module**
565 **scale of pediatric cancer patients for age**

566 Table 22 Bonferroni alpha post hoc test for the PedsQL 3.0 cancer module in pediatric
567 cancer patients aged 5-7, 8-12, and 13-18 years. The multiple comparison test revealed
568 statistical differences between ages 5-7 and 8-13:5-7 and 13-18 for all domains except concern,
569 with significance observed only between ages 5-7 and 13-18, with a p -score of <0.001 for
570 procedural anxiety and cognitive problems with a p-value of <0.05 for nausea, worry, and
571 communication. In parents/carers, post hoc tests show statistically significant anxiety, worry,
572 cognitive, and communicative emotional functioning between children enrolled once or twice
573 and children enrolled more than three times in relation to anxiety, worry, and communication.

574 **Table 22: Bonferroni alpha Post hoc test of PedsQL™ 3.0 cancer module for**
575 **age 5-7, 8-12, and 13-18 (n = 84)**

Variable				95% CI		P-value
	Comparison (years)	MD	S.E	Lower	Upper	
Pain and hurt	5 to7-8 to 12	-9.124	5.993	-23.776	5.528	0.395
	5 to 7- 13 to 18	-7.549	6.241	-22.805	7.708	0.690
	8 to 12-13 to 18	1.576	5.590	-12.090	15.241	1.000
Nausea	5 to7-8 to 12	-13.609	4.894	-25.574	-1.645	0.020
	5 to 7- 13 to 18	-11.120	5.096	-23.578	1.338	0.096
	8 to 12- 13 to 18	2.489	4.564	-8.669	13.648	1.000
Procedural anxiety	5 to7-8 to 12	-39.617	7.388	-57.679	-21.554	<0.001
	5 to 7- 13 to 18	-35.065	7.693	-53.872	-16.257	<0.001
	8to 12- 13 to 18	4.552	6.891	-12.295	21.398	1.000
Treatment anxiety	5 to7-8 to 12	-16.021	6.083	-30.891	-1.150	0.030
	5 to 7- 13 to 18	-19.399	6.334	-34.883	-3.916	<0.009
	8 to 12- 13 to 18	-3.379	5.673	-17.248	10.491	1.000
Worry	5 to7-8 to 12	9.693	5.943	-4.836	24.221	0.320
	5 to 7- 13 to 18	19.426	6.188	4.299	34.554	<0.007
	8 to 12-13 to 18	9.734	5.543	-3.816	23.284	0.249
Perceived physical appearance	5 to7-8 to 12	12.723	6.613	-3.443	28.889	0.174
	5 to 7- 13 to 18	6.385	6.885	-10.448	23.218	1.000
	8 to 12- 13 to 18	-6.338	6.168	-21.416	8.740	0.922
Cognitive	5 to7-8 to 12	-47.687	8.019	-67.291	-28.084	<0.001

	5 to 7- 13 to 18	-57.792	8.349	-78.204	-37.380	<0.001
	8to 12- 13 to 18	-10.105	7.479	-28.389	8.179	0.541
Communication	5 to7-8 to 12	-19.630	7.090	-36.963	-2.298	<0.021
	5 to 7- 13 to 18	-24.865	7.382	-42.912	-6.817	<0.003
	8 to 12- 13 to 18	-5.235	6.613	-21.400	10.931	1.000

576 **Source:** Field Data (2022)

577 **Associated factors and Health-Related Quality of Life among**
578 **Pediatric cancer patients(PedsQL™ 3.0)**

579 As shown in Table 23 a regression model to determine the effect of individual
580 characteristics on the PedsQL 3.0 cancer module reports of children and parent proxy reports
581 of quality of life. About PedsQL as perceived by pediatric cancer patients (child report). The
582 finding indicates that the educational level of pediatric cancer patients is significantly associated
583 with quality of life (=7.902 p<0.01), length of time since onset of disease (=2.165; p-value
584 <0.01), and the number of siblings in the family (=2.488; p-value<0.05). The results suggest
585 that a higher level of education, a longer survival time with cancer, and a large number of
586 siblings increase the chance of a better quality of life. However, childhood deprivation included
587 (= -2.175; p-value <0.01), complications related to cancer treatment (= -0.977; p-value <0.01),
588 hospitalization in the past 3 months (= -0.747; p-value < 0.01).

589 In addition, results related to PedsQL perceived by parents/carers (parent report) show
590 that childhood deprivation (= -1.589; p-value < 0.05), complications related to cancer treatment
591 (= -1.159; p-value < 0.01). , hospitalization in the last 3 months (= -5.589; p-value <0.01).
592 Therefore, the finding shows that reducing childhood deprivation, complications related to
593 cancer, and the number of hospitalizations before 3 months increases the chance of an
594 improvement in quality of life.

595 **Table 23. The health-related quality of life and associated factors (PedsQL™**
596 **3.0 cancer module) (n = 84)**

Variable	B	t	95% CI		P-Value
			Lower	Upper	
Child report					
Childhood Deprivation	-2.175	-3.122	-3.561	-0.789	0.002

Place of residence	-4.573	-2.658	-8.051	-1.096	0.011
Age of the child	1.796	0.770	-2.920	6.512	0.446
Sex of child	3.506	1.150	-2.658	9.670	0.257
Level of education of the child	7.902	7.302	5.747	10.056	0.000
Work done in the past 7 days	-7.305	-3.491	-11.472	-3.138	0.001
Hospitalization in prior 3 months	-5.474	-2.450	-9.924	-1.024	0.017
Types of treatment the child received	-0.454	-0.584	-2.025	1.117	0.563
Psychosocial counseling	-1.760	-0.662	-7.135	3.615	0.512
Most activity is done in the last 3 months	0.847	0.946	-0.963	2.657	0.350
Parent/caregiver factors					
Level of education of the parent/caregiver	-0.589	-0.375	-3.764	2.585	0.709
Employment	-0.823	-0.244	-7.640	5.995	0.809
Commence caring	15.677	3.338	6.185	25.169	0.002

597 **Source:** Field Data (2022)

598 **Discussion**

599 The present study found that there is association between childhood deprivation and
600 health related quality of life among pediatric cancer patients, while both level of childhood
601 deprivation and health related quality were higher. However male pediatric cancer patients were
602 many and suffering from Acute Lymphoblastic Leukemia.

603 The study results showed that acute lymphoblastic leukemia (ALL) and Hodgkin's
604 lymphoma were the most prominent leading cancers than other types of pediatric cancers. The
605 most affected pediatric patients were males over 7 years of age compared to female patients.
606 This is evident as many male pediatric cancer patients of school and adolescent age were
607 compared with females in this study. The higher proportion of ALL and Hodgkin's lymphoma
608 observed in this study was similar to previous studies in South Africa which showed that various
609 cancers were observed in which the majority of children diagnosed with acute lymphoblastic
610 leukemia 12% and males were most affected [24]. However, the differences were found in
611 latest study, the leading pediatric cancers were Wilms tumor 17.2%, retinoblastoma 16.8%,

612 and acute lymphoblastic leukemia 15.9% [11]. There was a higher incidence of male subjects,
613 according to international data reporting that males are more likely to have cancer than females,
614 with a 6:5 ratio for males and females [18,39]. In addition, ALL was the most common cancer
615 observed among all types of pediatric cancers [8,24].

616 In addition, many pediatric cancer patients were diagnosed in 2021 and early 2022
617 compared to previous years. This reflects that interventions such as childhood cancer awareness
618 and community awareness programs were more widely implemented during this period than in
619 previous years. In addition, priority was given to empowering healthcare providers through
620 (short and long cancer training), in-service training, conferences, and seminars against pediatric
621 cancer. Nervous was probably a time when there were advances in the diagnosis of pediatric
622 cancer, resulting in a rising number of children being diagnosed and confirmed with cancer in
623 Tanzania. However, only a few conducted studies reported that the mean time since diagnosis
624 increased significantly after the advancement of pediatric cancer therapies such as multiple
625 chemotherapies, surgery, radiotherapy, and psychosocial therapy [7]. Pediatric cancer patients
626 were treated in more than 4 hospitals seeking treatment while the cancer symptoms were pre-
627 diagnosis, with the majority of them, referred to MNH for diagnosis confirmation and treatment,
628 resulting in late diagnosis and treatment as most of them took more than 2 weeks to arrive at
629 the treatment center. This finding, also shown in previous studies, showed that about 74.5% of
630 pediatric cancer patients live more than 4 hours to reach the cancer centers [11]. This is one of
631 the factors contributing to being late in diagnosis, delaying cancer treatment can lead to early
632 cancer complications, and early infant mortality since most pediatric cancer patients came from
633 disadvantaged areas. [3,11], because of onset at a late stage of cancer.

634 Regarding the frequency of hospitalizations, most adolescent pediatric cancer patients were
635 admitted 1 to 3 times prior to 3 months, which may be due to complications related to cancer
636 itself and treatments. The result was consistent with a few studies conducted in which frequency
637 of admission is related to concern [18]. Despite advances in cancer therapy that maximize
638 survival time, pediatric cancer patients experience abdominal pain, fever, and palpitations
639 during cancer treatment. As reported by many studies, most children with cancer face problems
640 related to cancer and its therapy, including pain, nausea, vomiting, and physical incapacity
641 leading to weakened immunity [28,29]. School/study events were the most activities performed
642 by pediatric cancer patients in the last 3 months after play/excise and the few could not do any

643 activities because they have a physical disability (handicap) due to cancer and they Delay in
644 treatment and intervention.

645 In this study, the Multidimensional Poverty Index (MPI) was used to assess the level of
646 childhood disadvantage in 84 of 91 pediatric cancer patients. The results of this study show that
647 childhood deprivation was significantly higher in pediatric cancer patients as the proportion of
648 deprived was larger than non-deprived, which is attributed to most of them missing their
649 educational development while going to the hospital for treatment, frequency of admission, and
650 informal employment of their parents and caregivers continued. The results were similar to
651 previous demographic surveys: 50.2% of all children worldwide suffered from extreme poverty
652 in terms of education, health, and nutrition, while 31% in sub-Saharan Africa were
653 multidimensionally poor [15]. However, no significant difference was observed regarding the
654 sex of the pediatric cancer patients, but the overall mean of the female patients was slightly
655 higher compared to the male pediatric cancer patients, which may be due to the small number
656 of female podiatry cancer patients participating in this study. However, about 12% of male
657 children in Tanzania were disadvantaged in more than three essential needs, including early
658 school dropping, health, and nutrition [15]

659 This study found that childhood cancer occurs in rural areas, has low levels of education,
660 visits more than 4 hospitals while seeking treatment, and is significantly disadvantaged in single
661 parent/carer living. However, the results showed a significant association between the standard
662 of living and health. The difference was observed in place of residence between urban and rural
663 and peri-urban and rural, while mean total scores were high for pediatric cancer patients living
664 in rural areas compared to urban and peri-urban areas. Similar results of other studies from the
665 2016 NBS and UNICEF survey showed that about 81% of children living in rural areas are 3
666 and more disadvantaged, while few of them are in poverty (rural) areas compared to urban areas
667 life [13]. In addition, children living in rural areas suffering from chronic diseases have been
668 disadvantaged in more than basic needs compared to those living in urban [16]. Furthermore,
669 few studies documented that most pediatric cancer patients living in non-deprived areas had
670 more likely to have better health and improve their disease status to increase the survivors
671 [40], compared to those coming from deprived areas or rural areas.

672 Time to start medication was significantly associated with childhood deprivation, as most
673 pediatric cancer patients were delayed in arriving at cancer treatment centers. they took more

674 than 2 weeks to start cancer medication after diagnosis of the types of cancer compared with
675 previous finding access to cancer treatment centers and improving treatment modalities in
676 national hospitals are important in delaying cancer treatment [3]. However, the findings
677 suggested that there was no statistical difference either in gender or age among pediatric cancer
678 patients as also reported in a previous study that there is no association between deprivation
679 and gender or age group [41].

680 In this PedsQL study, the generic core module 4.0 was used to assess the health-related
681 quality of life of 84 of 91 pediatric cancer patients aged 5 to 18 years whose parents/carers
682 attended oncology clinics for treatment and other interventions. The results showed that the
683 mean HRQoL scores perceived by pediatric cancer patients were lower in PedsQL inventory.
684 Cancer is among the factors that have been perceived to contribute to the impairment of physical
685 and school functioning in pediatric cancer patients. Male pediatric cancer patients are the most
686 vulnerable group affected in both physical and school functioning, as their numbers were higher
687 than female pediatric cancer patients in this study. Physical disability of pediatric cancer
688 patients accompanied by school affection. Similar results were reported in previous studies,
689 male pediatric cancer patients were the most prominent group discharged because of physical
690 problems, seeking cancer treatment, and frequent admission to their school [18,24]

691 Furthermore, pediatric cancer patients aged 5-7 years were perceived as having physical
692 and school functioning problems, while other groups were perceived only as having school
693 functioning problems. HRQoL corresponding to age group implies that pediatric cancers are
694 associated with better quality of life in older than younger pediatric cancer patients. In addition,
695 most of them had not attended school because they did not feel well connected to cancer and
696 its treatments and went to the doctor or hospital. However, participated in various activities in
697 the last three months, including school/study, which is reliable with the current study not
698 attending school for < months and continued in the last 30 days [24].

699 The similarity is also suggested by a study conducted in the US and Chile, in which pediatric
700 cancer patients in Chile during the active phase of treatment and returning to school during their
701 cancer treatment had a lower mean score. For all US school cancer treatment courses School
702 functioning were the greatest impact on affecting the quality of life in pediatric cancer patients
703 and the dissimilarity observed in the US, where there was a higher mean school functioning
704 score throughout the treatment course [25]. The result suggests that there is a difference

705 between countries' scores on quality of life improvement and the impact of educational support.
706 In addition, starting school and regular school attendance are believed to affect growth and
707 development [10].

708 In this study, a linear regression analysis was performed to determine the impact of
709 individual characteristics on HRQoL in pediatric cancer patients. Pediatric cancer patients, how
710 were they disadvantaged, place of residence, participation in work/any activities in the last 7
711 days, their HRQoL were impaired resulting in poor quality of life (QoL). Poor health, the
712 standard of living, and living in rural areas are the factors that contribute most to Poor QoL in
713 pediatric cancer patients in this study. However, the results are consistent with recent studies
714 suggesting that all children with cancer living in disadvantaged and rural areas are likely to
715 impact poor QoL and experience a better quality of life if their socioeconomic status is
716 improved [22,43].

717 However, the results obtained show that pediatric cancer patients who were female, school
718 age, and adolescents (over 7 years) with higher education levels were perceived as having a
719 better quality of life in both the child report and the parent proxy report. The results suggest
720 that the age of pediatric cancer patients is more likely to affect the quality of life if it contributes
721 to the normal developmental milestone even during the disease. However, younger age (5-7)
722 years is associated with poor quality of life, which the child perceives in terms of physical
723 functioning. This finding is the same as previous studies, which found that most pediatric
724 patients diagnosed with cancer at a younger age had a poor observed quality of life [18,44].
725 This finding reinforces the importance of preventing the physical disability and dysfunction
726 associated with cancer treatment and the disease itself.

727 In addition, the study was statistically significant in terms of perceived disease severity
728 in pediatric cancer patients, who had lower overall mean. Procedural anxiety, cognitive
729 problems, and communication problems were the critical problems perceived by pediatric
730 cancer patients. Pediatric cancer patients rely on their parents/caregivers to provide their health
731 information on their behalf. However, differences in three subscales, including nausea,
732 procedural anxiety, and cognitive problems, were seen in the reports of children aged 8-12 and
733 13-18 compared to 5-7 years in this study. The observed outcome is no different from the
734 current study, which found that younger-aged pediatric cancer patients were statistically
735 associated with poor health-related quality of life on the fear of procedures subscale [18]. This

736 finding underscores the importance of preventing the anxiety associated with medical
737 intervention during cancer treatment. Procedural anxiety can be distressing pediatric cancer
738 patients, making treatment difficult

739 Therefore, the results are likely contributed by other causes such as dissimilarity in
740 developmental factors. In addition, the association was observed in the communication range,
741 which was below the cutoff mean (69.7) in each age group. The same result was found in other
742 studies that the communication range of pediatric cancer patients was considered difficult to
743 interact with healthcare providers (Tembe, 2021). Additionally, most research on
744 communication in childhood cancer has focused on parents/caregivers or healthcare providers
745 rather than the patients themselves. Therefore, studies have suggested that understanding
746 pediatric cancer patients require a healthcare provider to facilitate effective communication
747 (patient-provider) which could improve quality of life and client satisfaction (Lin et al., 2020)
748 through early intervention and reduce the impact of cancer therapies complication

749 The lower mean was found for pain and injury, nausea, and treatment anxiety perceived
750 by pediatric cancer patients aged 5-7 years. In addition, a dissimilarity was found in the
751 cognitive domain of the children's report, where the mean score was lower at ages 5-7 than at
752 ages 8-12 and 13-18, which is similar to other studies (Bishop et al., 2018; Tembe, 2021). . In
753 addition, another analysis was performed (Bonferroni alpha post hoc test) to find out the
754 difference between the age groups of the pediatric cancer patients, with the finding being
755 consistent between the age groups (5-7 and 13-18; 5-7 and 13-18) significant was treatment
756 anxiety. This may be related to a lack of patient orientation by healthcare providers and the
757 developmental characteristics of children. In addition, cognitive problems have been observed
758 between the ages of 5 and 7 years that being younger and being diagnosed with cancer can
759 affect the quality of life, which is because most pediatric cancer patients do not attend school
760 and spend most of their time in hospital spending.

761 In this study, a linear regression analysis was performed to determine the impact of
762 individual characteristics on HRQoL in pediatric cancer patients. Pediatric cancer patients, how
763 were they disadvantaged, place of residence, participation in work/any activities in the last 7
764 days, their HRQoL were impaired resulting in poor quality of life (QoL). Poor health, the
765 standard of living, and living in rural areas are the factors that contribute most to Poor QoL in
766 pediatric cancer patients in this study. However, the results are consistent with recent studies

767 suggesting that all children with cancer living in disadvantaged and rural areas are likely to
768 impact poor QoL and experience a better quality of life if their socioeconomic status is
769 improved [22,44].

770 However, the results obtained show that pediatric cancer patients who were female,
771 school age, and adolescents (over 7 years) with higher education levels were perceived as
772 having a better quality of life. The results suggest that the age of pediatric cancer patients is
773 more likely to affect the quality of life if it contributes to the normal developmental milestone
774 even during the disease. However, younger age (5-7) years is associated with poor quality of
775 life, which the child perceives in terms of physical functioning. This finding is the same as
776 previous studies, which found that most pediatric patients diagnosed with cancer at a younger
777 age had a poor observed quality of life [10,18]. This finding reinforces the importance of
778 preventing the physical disability and dysfunction associated with cancer treatment and the
779 disease itself. In addition, the regression model showed a significant association with quality of
780 life (PedsQLTM 3.0 inventory) in both the child's report and the parent's proxy report.

781 The finding shows that the child's age, hospitalization within the last 3 months, childhood
782 deprivation, and school attendance were perceived as serious, and in the parent representative's
783 report the observed outcome was significant for hospitalization within the last 3 months and
784 school attendance. The results may suggest that age, frequent hospitalizations, childhood
785 deprivation, and irregular school attendance are important determinants of improved quality of
786 life in pediatric cancer patients. However, most of the deprived pediatric cancer patients failed
787 to attend school after frequent hospitalizations [44]. However, this study has discover that the
788 childhood deprivation is higher which lead to lower quality of life among pediatric cancer
789 patients related to delay treatment due to living in rural areas where there is limited of cancer
790 centers and poor standard of living,

791 **Strength and limitation of the study**

792 This study has discover that the childhood deprivation is higher which lead to lower
793 quality of life among pediatric cancer patients related to delay treatment due to living in rural
794 areas where there is limited of cancer centers and poor standard of living. The study did not
795 rely on examining the clinical presentation factors associated with HRQoL. Also, the generality
796 of this study may be limited by inclusion and exclusion criteria, particularly with regard to age,
797 which did not involve pediatric cancer patients younger than 5 years of age and not able to

798 understand and communicate verbally, and those who have been diagnosed with cancer and
799 started or completed cancer treatment. Information bias can be observed after the Parent Proxy
800 Report on Childhood Deprivation and Health-Related Quality of Life.

801 **Conclusion**

802 The magnitude of acute lymphoblastic leukemia is higher among pediatric cancer
803 patients. There was a significant link between pediatric cancers and childhood deprivation and
804 thus, compromised quality of life among pediatric cancer patients. Innovative pediatric cancer
805 care policies, guidelines, and or strategies may need to be advocated to address the problem
806 accordingly.

807 **Suggestions**

808 Future research to maximize the size of the study by recruiting a large sample is needed
809 to investigate routine clinical factors that may affect improving QOL outcomes using sensitivity
810 measures during the treatment of pediatric cancer patients that provide important information
811 for early information of healthcare providers about treatment decisions represent intervention.

812 **Supporting information**

813 Dataset available, shared upon request

814 **Conflicts of Interest**

815 The authors has no conflicts of interest relevant to this article.

816 **Acknowledgments**

817 The authors express gratitude to the University of Dodoma for providing ethical clearance to
818 conduct this study. We also thank the Muhimbili National Hospital administration for
819 allowing us to conduct the study, also we thank the pediatric cancer patients and their
820 parents/caregivers for volunteering to participate in this study fully till the end.

821 **Author Contributions**

822 Conceptualization: Mwanaheri Chubi.

823 Data curation: Mwanaheri Chubi

824 Formal analysis: Mwanaheri Chubi.

- 825 Methodology: Mwanaheri Chubi.
- 826 Supervision: Stephen Kibusi, Lulu Chirande and Shakiru Jumanne.
- 827 Varidation : Mwanaheri Chubi
- 828 Writing: original draft: Mwanaheri Chubi.
- 829 Writing: review & editing: Stephen Kibusi, Lulu Chirande and Shakiru Jumanne.
- 830 Funding: This research received no external funding
- 831

832 **REFERENCE**

- 833 1. WHO. CureAll framework. WHO global initiative for childhood cancer: increasing
834 access, advancing quality, saving lives. World Health Organization. 2021.
- 835 2. Zhang A, Wang K, Zebrack B, Tan CY, Walling E, Chugh R. Psychosocial, behavioral,
836 and supportive interventions for pediatric, adolescent, and young adult cancer survivors:
837 A systematic review and meta-analysis. *Crit Rev Oncol Hematol* [Internet].
838 2021;160(October 2020):103291. Available from:
839 <https://doi.org/10.1016/j.critrevonc.2021.103291>
- 840 3. Schroeder K, Saxton A, McDade J, Chao C, Masalu N, Chao C, et al. Pediatric cancer in
841 northern Tanzania: Evaluation of diagnosis, treatment, and outcomes. *J Glob Oncol*.
842 2018;2018(4).
- 843 4. Geel JA, Challinor J, Ranasinghe N, Myezo KH, Eyal KC, Aderounmu W, et al. Pediatric
844 cancer care in Africa: SIOP Global Mapping Program report on economic and
845 population indicators. *Pediatr Blood Cancer*. 2021;68(11):1–9.
- 846 5. Boyle P, Ngoma T, Sullivan R, Ndlovu N, Autier P, Stefan C, et al. State of Oncology
847 in Africa 2015 [Internet]. 2016. 1–298 p. Available from: [http://www.i-pri.org/wp-](http://www.i-pri.org/wp-content/uploads/2017/02/STATE-OF-ONCOLOGY-IN-AFRICA-2015-WEB-VERSION.pdf)
848 [content/uploads/2017/02/STATE-OF-ONCOLOGY-IN-AFRICA-2015-WEB-](http://www.i-pri.org/wp-content/uploads/2017/02/STATE-OF-ONCOLOGY-IN-AFRICA-2015-WEB-VERSION.pdf)
849 [VERSION.pdf](http://www.i-pri.org/wp-content/uploads/2017/02/STATE-OF-ONCOLOGY-IN-AFRICA-2015-WEB-VERSION.pdf)
- 850 6. Geel JA, Ranasinghe N, Myezo KH, Davidson A, Howard SC, Hessissen L, et al.
851 Pediatric cancer care in Africa: SIOP Global Mapping process. *Pediatr Blood Cancer*.
852 2021;68(11).
- 853 7. Karimi M, Cox AD, White S V., Karlson CW. Fatigue, Physical and Functional Mobility,
854 and Obesity in Pediatric Cancer Survivors. *Cancer Nurs*. 2020;43(4):E239–45.
- 855 8. Joko-Fru WY, Parkin DM, Borok M, Chokunonga E, Korir A, Nambooze S, et al.
856 Survival from childhood cancers in Eastern Africa: A population-based registry study.
857 *Int J Cancer*. 2018 Nov 15;143(10):2409–15.
- 858 9. Kayungu SJ. Finding an optimal treatment regimen for endemic Burkitt’s lymphoma in
859 a limited resource setting: A historical comparison of an INCTR-03-06 protocol for
860 children treated at a tertiary pediatric oncology unit in Tanzania. *J Clin Oncol*.

- 861 2020;38(15_suppl):e22507–e22507.
- 862 10. Pan HT, Wu LM, Wen SH. Quality of Life and Its Predictors among Children and
863 Adolescents with Cancer. *Cancer Nurs.* 2017;40(5):343–51.
- 864 11. Maillie L, Mutagonda A, Chirande L, Rehema L, Jumanne S, Meleki E, et al. Abstract
865 101: The Tanzanian Pediatric Cancer Network: A Comprehensive Evaluation of the
866 Incidence of Presenting Patients and Access to Pediatric Cancer Care in Tanzania. In
867 2021. p. 101–101.
- 868 12. WHO. Global initiative for childhood cancer [Internet]. 2020. Available from:
869 <https://www.who.int/publications/m/item/global-initiative-for-childhood-cancer>
- 870 13. NBS & UNICEF. Child Poverty in Tanzania National Bureau of Statistics and United
871 Nations Children’s Fund THE UNITED REPUBLIC OF TANZANIA [Internet]. 2016.
872 Available from: [https://www.unicef.org/esa/sites/unicef.org.esa/files/2018-09/UNICEF-](https://www.unicef.org/esa/sites/unicef.org.esa/files/2018-09/UNICEF-Tanzania-2016-Child-Poverty.pdf)
873 [Tanzania-2016-Child-Poverty.pdf](https://www.unicef.org/esa/sites/unicef.org.esa/files/2018-09/UNICEF-Tanzania-2016-Child-Poverty.pdf)
- 874 14. Poulalhon C, Goujon S, Marquant F, Faure L, Guissou S, Bonaventure A, et al. Factors
875 associated with 5- and 10-year survival among a recent cohort of childhood cancer
876 survivors (France, 2000–2015). *Cancer Epidemiol.* 2021;73(June).
- 877 15. NBS & UNICEF. Child Poverty in TANZANIA [Internet]. 2019. Available from:
878 [https://www.unicef.org/tanzania/media/2106/file/UNICEF-Tanzania-Child-Poverty-](https://www.unicef.org/tanzania/media/2106/file/UNICEF-Tanzania-Child-Poverty-Report-2019.pdf)
879 [Report-2019.pdf](https://www.unicef.org/tanzania/media/2106/file/UNICEF-Tanzania-Child-Poverty-Report-2019.pdf)
- 880 16. Lwidiko A, Kibusi SM, Nyundo A, Mpondo BCT. Association between HIV status and
881 depressive symptoms among children and adolescents in the Southern Highlands Zone,
882 Tanzania: A case-control study. *PLoS One.* 2018 Feb 1;13(2).
- 883 17. Zainal NR, Kaur G, Ahmad NA, Khalili JM. Housing Conditions and Quality of Life of
884 the Urban Poor in Malaysia. *Procedia - Soc Behav Sci.* 2012;50(July 2012):827–38.
- 885 18. Alabbas F, Elyamany G, Alkhayat N, AlShahrani M, Ibrahim W, Elborai Y, et al.
886 Assessment of Health-related Quality of Life in Saudi Children with Cancer. *Heal Serv*
887 *Insights.* 2021;14:117863292110293.
- 888 19. MoHSW. National Cancer Control Strategy (NCCS) 2013 - 2022. 2013. 2013;72.

- 889 20. Ahmad AA, Saburi E, Rajaei T, Mortazavi Y, Thomas X, Chelghoum Y, et al. Types of
890 cancer that develop in children. *Anticancer Res* [Internet]. 2018;9(1):1–2. Available
891 from:
892 <https://link.springer.com/content/pdf/10.1023/A:1010062929140.pdf>
893 [http://0-ovidsp.ovid.com.wam.city.ac.uk/ovidweb.cgi?T=JS&PAGE=reference&D=emed10&](http://0-ovidsp.ovid.com.wam.city.ac.uk/ovidweb.cgi?T=JS&PAGE=reference&D=emed10&NEWS=N&AN=40807300)
894 [NEWS=N&AN=40807300](https://www.ledsmagazine.com/articles/print/volume-15/issue-5/features/developer-foru)
895 [https://www.ledsmagazine.com/articles/print/volume-](https://www.ledsmagazine.com/articles/print/volume-15/issue-5/features/developer-foru)
- 896 21. Sung L, Klaassen RJ, Dix D, Pritchard S, Yanofsky R, Dzolganovski B, et al.
897 Identification of paediatric cancer patients with poor quality of life. *Br J Cancer*.
898 2009;82–8.
- 899 22. Kim J, Park E. Impact of socioeconomic status and subjective social class on overall and
900 health-related quality of life. *BMC Public Health* [Internet]. 2015; Available from:
901 <http://dx.doi.org/10.1186/s12889-015-2014-9>
- 902 23. Varni JW, Panepinto JA. Cognitive functioning, patient health communication, and
903 worry mediate pain predictive effects on health-related quality of life in youth with sickle
904 cell disease. *Pediatr Blood Cancer*. 2020 Nov 1;67(11).
- 905 24. Tembe SC. The Health-Related Quality of Life of South African Paediatric In-patients
906 with Cancer. *South African J Occup Ther*. 2021;51(1):44–53.
- 907 25. Bishop MW, Billups CA, Gattuso JS, Bishop MW, Advani SM, Daw NC, et al. Health-
908 related quality of life and survival outcomes of pediatric patients with nonmetastatic
909 osteosarcoma treated in countries with different resources. *J Glob Oncol*.
910 2018;2018(4):1–11.
- 911 26. Tumaini la Maisha. Curing Children’s cancer in Tanzania: 2015-2020 [Internet]. 2020.
912 Available from: [https://www.ucd.ie/t4cms/TLM PDF.pdf](https://www.ucd.ie/t4cms/TLM%20PDF.pdf)
- 913 27. Directrelief. In Sub-Saharan Africa, Childhood Cancer Is Often Deadly. A New Initiative
914 Is Working to Change That. 2020.
- 915 28. Kohi TW, Von Essen L, Masika GM, Gottvall M, Dol J. Cancer-related concerns and
916 needs among young adults and children on cancer treatment in Tanzania: A qualitative
917 study 11 Medical and Health Sciences 1117 Public Health and Health Services. *BMC*

- 918 Cancer. 2019 Jan 17;19(1).
- 919 29. Masika GM, Gottvall M, Kohi TW, Von Essen L, Dol JS. Concerns and Needs of
920 Support among Guardians of Children on Cancer Treatment in Dar es Salaam: A
921 Qualitative Study. *Cancer Nurs.* 2020;43(6):E342–8.
- 922 30. Hydeman JA, Uwazurike OC, Adeyemi EI, Beaupin LK. Survivorship needs of
923 adolescent and young adult cancer survivors: a concept mapping analysis. *J Cancer*
924 *Surviv.* 2019 Feb 15;13(1):34–42.
- 925 31. Riaz N, Wolden SL, Gelblum DY, Eric J. Survivors of Childhood/Adolescent Cancer:
926 Life-long Risks and Responsibilities. 2016;118(24):6072–8.
- 927 32. Ministry of Health and Social Welfare (MoHSW) [Tanzania Mainland], Ministry of
928 Health (MoH) [Zanzibar], (NBS) NB of S, Statistician(OCGS) O of the CG, International
929 I. Tanzania Service Provision Assessment. 2015;502. Available from:
930 <https://dhsprogram.com/pubs/pdf/spa22/spa22.pdf>
- 931 33. National Malaria Control Programme -Ministry of Health. Tanzania Malaria Indicator
932 Survey 2017 - Final Report. 2017; Available from:
933 <https://dhsprogram.com/publications/publication-mis31-mis-final-reports.cfm>
- 934 34. Abraham ZS, Massawe ER, Kahinga AA, Mapondella KB, Massawe WA, Swai H, et al.
935 Clinical Profile of Paediatric Head and Neck Cancers at a Tertiary Hospital in Tanzania.
936 *Rwanda J Med Heal Sci.* 2020 Apr 14;3(1):11–20.
- 937 35. Varni JW, Limbers CA, Burwinkle TM. Parent proxy-report of their children’s health-
938 related quality of life: An analysis of 13,878 parents’ reliability and validity across age
939 subgroups using the PedsQL™ 4.0 Generic Core Scales. *Health Qual Life Outcomes.*
940 2007;5:1–10.
- 941 36. UN General Assembly. A/RES/61/146 Rights of the child. Vol. 1974. 2007.
- 942 37. Limbers CA, Ripperger-Suhler J, Heffer RW, Varni JW. Patient-reported pediatric
943 quality of life inventory™ 4.0 generic core scales in pediatric patients with attention-
944 deficit/hyperactivity disorder and comorbid psychiatric disorders: Feasibility, reliability,
945 and validity. *Value Heal.* 2011;14(4):521–30.
- 946 38. López León A, Carreño Moreno S, Arias-Rojas M. Relationship Between Quality of Life

- 947 of Children With Cancer and Caregiving Competence of Main Family Caregivers. *J*
948 *Pediatr Oncol Nurs*. 2021;38(2):105–15.
- 949 39. EL Hajjar ST, Alkhanaizi MS. Exploring the Factors That Affect Employee Training
950 Effectiveness: A Case Study in Bahrain. *SAGE Open* [Internet]. 2018;8(2). Available
951 from: <https://journals.sagepub.com/doi/epub/10.1177/2158244018783033>
- 952 40. Ul Hadia N, Abdullah N, Sentosa I. An Easy Approach to Exploratory Factor Analysis:
953 Marketing Perspective. *J Educ Soc Res* [Internet]. 2016;6(1):215–23. Available from:
954 https://www.researchgate.net/publication/290042054_An_Easy_Approach_to_Exploratory_Factor_Analysis_Marketing_Perspective_Noor_Ul_Hadi
- 956 41. American Cancer Society. Treating Children with Cancer. *Am Cancer Soc* [Internet].
957 2016;1–4. Available from: <https://www.cancer.org/cancer/cancer-in-children/how-are-childhood-cancers-treated.html>(accessed 19/7/2022)
- 959 42. Régnier Denois V, Bourmaud A, Nekaa M, Bezzaz C, Bousser V, Kalecinski J, et al.
960 The impact of social inequalities on children’s knowledge and representation of health
961 and cancer Véronique. *Eur J Pediatr* [Internet]. 2018;177(8):1219–30. Available from:
962 <https://pubmed.ncbi.nlm.nih.gov/29808238/>
- 963 43. Taralee Hamner, Robert D. Latzman, Natasha E. Latzman, T. David Elkin and SM.
964 Quality of Life Among Pediatric Patients With Cancer: Contributions of Time Since
965 Diagnosis and Parental Chronic Stress. *Pediatr Blood Cancer* [Internet].
966 2015;62(7):1232–1236. Available from:
967 <https://onlinelibrary.wiley.com/doi/abs/10.1002/pbc.25468>
- 968 44. H Adıgüzel, I Ergin, Z Ocek RD. Influence of deprivation on health-related quality of
969 life in children: Cross sectional study, Turkey. *Eur J Public Health* [Internet].
970 2018;28(4):485–6. Available from:
971 https://academic.oup.com/eurpub/article/28/suppl_4/cky218.224/5192239 accessed on
972 7/20/2022
- 973 45. Pan H, Wu L, Wen S. Quality of Life and Its Predictors Among Children and Adolescents
974 With Cancer. *Cancer Nurs* [Internet]. 2016;00(0):1–9. Available from:
975 https://journals.lww.com/cancernursingonline/Fulltext/2017/09000/Quality_of_Life_and_Its_Predictors_Among_Children.2.aspx




Fig 1: Flow diagram representing sampling procedure and sample size

Level of childhood deprivation




Fig 1: Histogram showing the level of childhood deprivation by mean score among pediatric cancer patients