Title: Low back pain care pathways – is the last provider seen more important than the first: A

retrospective cohort study

Authors:

David Elton, DC, Optum Labs, Eden Prairie, Minnesota USA

Meng Zhang, PhD, Optum Labs, Eden Prairie, Minnesota USA

Corresponding author:

David Elton

Optum Labs

11000 Optum Circle, Eden Prairie, MN 55343

Phone: (952) 562-1349

david.elton@optum.com

Declarations

Ethics approval and consent to participate

Because the data was de-identified or a Limited Data Set in compliance with the Health Insurance Portability and Accountability Act and customer requirements, the UnitedHealth Group Office of Human Research Affairs determined that this study was exempt from Institutional Review Board review.

Consent for publication

Not applicable

Availability of data and materials

The data are proprietary and are not available for public use but, under certain conditions, may be made available to editors and their approved auditors under a data-use agreement to confirm the findings of the current study.

Competing interests

At the time of manuscript submission **DE and MZ** are UnitedHealth Group employees and UNH stockholders. No other potential conflicts of interest or competing interests exist.

Funding

None

Authors' contributions

Study conception and design; **DE**. Data acquisition; **DE, MZ.** Data analysis and interpretation; **DE, MZ.** Draft or substantially revise manuscript; **DE, MZ.**

Acknowledgements

Abstract

Background: The associations between the type of health care provider (HCP) initially contacted by an individual with low back pain (LBP), guideline concordance, and cost of care are well understood. Less is known about types of HCP that become involved after the initial HCP, particularly the last HCP. The purpose of this retrospective cohort study was to examine sequence patterns in the types of HCP involved in an episode of LBP, exposure to 13 types of services, and total cost.

Methods: A US national sample of commercially insured individuals with a single episode of non-surgical LBP occurring in 2017-2019 was analyzed using episode of care unit of analysis. The primary independent variable was the type of initial contact HCP. Dependent measures included types of subsequent HCPs involved in an episode of LBP, rate of use of 13 types of health care services, and total cost.

Results: 503,958 individuals aged 18 years and older initially contacted 196,522 different HCPs for an episode of non-surgical LBP. 30.2% of individuals saw a second HCP, 11.8% a third, 5.2% a fourth and 2.5% a fifth. Compared to primary care providers (PCP) (71.0% of episodes), when chiropractors (DC) were the initial contact HCP they were the most likely to be the only and last HCP seen (84.0% of episodes) (risk ratio 1.18, 95% confidence interval 1.18-1.19). DCs were also the most likely to be the last HCP seen when the second HCP in an episode (72.7%) (1.10, 1.09-1.12), the third HCP (67.7%) (1.10, 1.08-1.13), or the fourth HCP (61.6%) (1.13, 1.08-1.17). Physical therapists (PT), while less likely than PCPs to be the only and last HCP seen (60.0%) (0.84, 0.83-0.86), were more likely than PCPs to be the last HCP when the second (70.9%) (1.08, 1.06-1.09), third (63.2%) (1.03, 1.00-1.06), or fourth HCP (60.7%) (1.11, 1.06-1.16) seen during an episode. PTs were the only type of HCP involved in more episodes as the second HCP (7,344) than as the initial HCP (5,114).

Conclusions: DCs are the most likely to be the last HCP seen for an episode of non-surgical LBP. PTs are as likely as DCs to be the last HCP seen except when initially contacted by an individual with LBP. There are numerous potential confounders and limitations to consider when interpreting this novel finding.

Keywords: Low back pain; pathway; guideline; initial contact; first provider; last provider; utilization; cost; value

Background

Low back pain (LBP) clinical practice guidelines (CPG) recommended a stepped approach to
management with an emphasis on non-pharmaceutical and non-interventional first-line approaches for
most LBP.[1,2] The type of health care provider (HCP) initially contacted by an individual with LBP, and
associated subsequent service use and CPG concordance, is the subject of a growing body of research. [3-8] A high proportion of individuals with LBP initially seek treatment from primary care providers (PCP) and
physician specialists with management emphasizing second- and third-line approaches.[3] Non-prescribing
HCPs, like chiropractors (DC), physical therapists (PT), or licensed acupuncturists (LAc) are more likely to
have episodes associated with an emphasis on first-line therapies, with less use of second- and third-line
services.[3,6] A variety of confounders impact the interpretation and translation of findings associated with
the type of HCP initially contacted by an individual with LBP.[4,6,7]

Relatively little is known about longitudinal patterns of all types of HCPs involved in an episode of LBP
and association with guideline concordance and total cost. Research into the sequence of types of HCPs
involved in an episode of LBP has tended to focus on service use rather than types of HCPs involved, and
has often focused on a single type of initial contact HCP such as PCP [9,10], physiatry [11], emergency
medicine (EM)[12,13], and PT.[5,8,14] These studies identified variation in approaches to LBP, yet have not
provide comprehensive insights into patterns of HCP involvement in LBP. A PubMed.gov search for key
terms "last provider seen" and "back pain" returned no results.

The aim of this study was to examine patterns in the types of HCP involved in a single episode of non-
surgical LBP and associated total episodic use of 13 types of health care services and total cost, with a
specific focus on the type of HCP seen last. The hypothesis was types of HCP emphasizing CPG

concordant first-line approaches, like DCs, PTs, LAcs, and doctors of osteopathy (DO), would be commonly seen as the last HCP during an episode of LBP even if becoming involved later in an episode.

Methods

Study design, population, setting and data sources

This is a retrospective cohort study of individuals with a single episode of non-surgical LBP during the 2017-2019 study period. An analytic database was created which included de-identified enrollment records, administrative claims data for all inpatient and outpatient services, and pharmacy prescriptions, for commercially insured enrollees from a single national commercial insurer. De-identified HCP demographic information and professional licensure status was included in an HCP database. ZIP code level population race and ethnicity data was extracted from the US Census Bureau [15], socioeconomic status (SES) Area Deprivation Index (ADI) data, from the University of Wisconsin Neighborhood Atlas® [16], and household adjusted gross income (AGI) from the Internal Revenue Service.[17]

A review was performed to assess compliance with de-identification requirements and as it was determined the data was de-identified or a Limited Data Set in compliance with the Health Insurance Portability and Accountability Act and customer requirements, the UnitedHealth Group Office of Human Research Affairs determined that this study was exempt from Institutional Review Board review. The study was conducted and reported based on the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) guidelines.[18] [Supplement 1 – STROBE Checklist]

The analysis does not include an adjustment for typical confounders such as individual age, sex and co-morbidities [19,20] using common yet potentially inadequate approaches such as propensity score matching [21] due to the inability to control for important considerations of individuals with LBP initially contacting different types of HCP, and subsequent HCPs involved in an episode. Examples of considerations not available include: HCP options convenient to an individual's home, workplace or daily travel routes including public transportation if used, individual preference for specific services or type of HCP including gender or racial concordance, recommendations from family or friends and influence of HCP marketing efforts, perceived LBP severity, anticipated potential out of pocket costs and willingness to pay for services, time availability to participate in multiple visits, and appointment availability within an individual's timing expectations meeting these and other criteria.[22] As an alternative to blurring the line between association and causality through a process that simultaneously introduces distortion and complexity into results, actual measures of individual demographic and episodic characteristics, and unadjusted associations, are reported for each type of HCP seen initially, or subsequently, during an episode of LBP.

To partially address potential confounders, sub analyses were performed for episodes where an individual initially contacted a PCP, orthopedic surgeon (OS), or EM physician. To examine the influence of population factors on results, the flow of episodes between types of HCP, and the of % of episodes where each type of HCP was the last HCP seen were calculated for individuals living in zip codes with 75-100 ADI and 0-25% non-Hispanic white (NHW) population, 50-75 ADI and 50-75% NHW population, and 0-25 ADI and 75-100% NWH population.

Cohort selection and unit of analysis

The cohort included individuals aged 18 years and older with a single complete episode of non-surgical LBP commencing and ending during the calendar years 2017-2019. This timeframe was selected to follow the release of the American College of Physicians (ACP) LBP CPG [2] in 2017 and before the influence of the COVID-19 epidemic on care patterns in early 2020. All individuals had continuous medical and pharmacy commercial insurance coverage during the entire study period.

Episode of care was selected as the unit of analysis. This approach has been shown to be a valid way to organize all administrative claims data associated with a condition.[23] The *Symmetry® Episode Treatment Groups® (ETG®)* and *Episode Risk Groups® (ERG®)* version 9.5 methodologies and definitions were used to translate administrative claims data into episodes of care, which have been reported as a valid measurement for comparison of HCPs based on cost of care.[24] A previous analysis of the same underlying data indicates a low risk of misclassification bias associated with using episode of care as the unit of analysis.[3] Using episode of care unit of measurement has potential translational benefits in supporting the transition from fee for service to value-based episodic bundled payment arrangements, something particularly relevant when examining episodes with multiple HCPs involved.

The analysis included complete episodes defined as having at least 91-day pre- and 61-day post-episode clean periods during which no services were provided by any HCP for any LBP diagnosis. LBP episodes including a surgical procedure, or associated with diagnoses of malignant and non-malignant neoplasms, fractures and other spinal trauma, infection, congenital deformities and scoliosis, autoimmune disorders, osteoporosis, and advanced arthritis were excluded from the analysis. Individuals with multiple LBP episodes during the study period were also excluded. These exclusions were made to address a potential study limitation of individuals with more complex conditions confounding study

results examining the type of HCP initially contacted, and subsequent types of HCP seen during an episode of LBP.

Variables

Data preprocessing, table generation, and initial analyses were performed using Python (*Python Language Reference, Version 3.7.5.*, n.d.). To evaluate whether measures were derived from a normally distributed sample we conducted a goodness-of-fit measure using D'Agostino's K-squared test. Non-normally distributed data are reported using the median and interquartile range (IQR). Odds (OR) and risk (RR) ratios, and associated 95% confidence intervals (CI), were calculated for select measures. RR were reported as this is the measure more widely understood in associational analyses and due to the tendency for ORs to exaggerate risk in situations where an outcome is relatively common.[25]

The primary independent variable was 17 types of HCP initially contacted by an individual experiencing a single episode of non-surgical LBP during the study period. All types of HCP could be accessed directly without a referral. The primary dependent variables were types of HCP involved after the initial HCP, the number of days into an episode when subsequent HCP types were involved, the rate of use of 13 types of health care services and total cost. Service utilization reflects services provided by any type of HCP an individual saw during the complete episode of LBP. Total cost included costs associated with all services provided for LBP during an episode, including those not specifically identified in the 13 categories used in the analyses. Costs for services for which an insurance claim was not submitted were not available.

With 97.5% of episodes involving five or fewer HCPs, the analyses focused on the first five types of HCP seen during an episode. This resulted in 1,419,925 possible HCP sequence combinations. [Figure 1] Rate

(%) and timing (days), median and IQR, and RR and 95% CI results for cohort demographics, HCP

involvement, episodic service use, and total cost were calculated for all HCP sequence combinations.

Total episode cost and percent of episodes including an MRI or prescription opioid were calculated for

episodes where a PCP, DC, PT or OS was the last HCP seen.

Results

The cohort consisted of 503,958 individuals, associated with 503,958 complete non-surgical LBP

episodes for which 196,522 different HCPs were initially contacted. There were $387,867,096 in

reimbursed health care expenditures. The median pre- (646 days) and post-episode (405 days) clean

periods were substantially longer than the *ETG*® clean period definitions. Primary care physicians (PCP)

(35.2% of episodes), DCs (25.9%), nurse practitioners (7.5%), radiologists (6.0%) and OSs (5.5%) where

the most common types of HCP initially contacted. [Supplement 1] [Supplement 2] There were

potentially meaningful differences in individual and population factors between types of HCP seen at

different points in an episode. [Supplement 3]

30.2% of episodes involved 2 HCPs, 11.8% involved 3 HCPs, 5.2% involved 4 HCPs and 2.5% of episodes

involved 5 HCPs. All types of HCP were involved in progressively fewer episodes when moving from

initial to second, third, fourth or fifth HCP, except PTs who participated in more episodes as a second

HCP (7,344) than as an initial HCP (5,114). [Table 1]

When initially contacted by individuals with LBP, and compared to the PCP reference (71.0%), DCs

(84.0%, risk ratio 1.18, 95% confidence interval 1.18-1.19), LAcs (81.7%, 1.15, 1.13-1.17) and DOs

(80.2%, 1.13, 1.08, 1.18) were most likely to be the last and only HCP seen during an episode. [Table 2]

[Figure 2] When seen as a second, third or fourth HCP during an episode, DCs and PTs were the only HCPs more likely than a PCP to be the last HCP seen. Neurosurgeons and radiologists were the least likely to be the last HCP seen. Sub analyses of the 75-100 ADI 0-25% NHW, 50-75 50-75% NHW, and 0-25 ADI 75-100% NHW segments revealed nearly identical results. [Supplement 4] When the initial HCP contacted, DCs were more likely than PCPs to be the last HCP seen in each segment. LAcs were more likely than PCPs to be the last HCP seen in the 75-100 ADI 0-25% NHW and 0-25 ADI 75-100% NHW segments. [Figure 3] When positioned as the second HCP involved in an episode, DCs were more likely than PCPs to be the last HCP seen in each segment. PTs were more likely than PCPs to be the last HCP seen in the 50-75 50-75% NHW and 0-25 ADI 75-100% NHW segments. [Figure 4]

The sub-analyses of individuals initially contacting a PCP, OS, or EM had similar last HCP seen results. For individuals initially contacting a PCP, and when seen as a second HCP, DCs (76.2%, risk ratio 1.15, 95% confidence interval 1.13-1.18) and PTs (76.1%, 1.15, 1.12-1.18) were the most likely to be the last only HCP seen. For individuals initially contacting an OS, and when seen as a second HCP, DCs (69.5%, 1.12, 1.05-1.19) and PTs (71.2%, 1.14, 1.09-1.20) were the most likely to be the last only HCP seen. For individuals initially contacting EM, and when seen as a second HCP DCs (76.5%, 1.12, 1.05-1.20) were the most likely to be the last only HCP seen. [Figure 5] This was also the case when a DC or PT was the third or fourth provider seen for individuals initially contacting a PCP. DCs and PTs were also the most likely to be the last HCP seen when the second, third or fourth HCP for individuals initially contacting an OS or EM. [Table 3]

Whether seen as the initial, second, third or fourth HCP, when the last HCP seen during an episode the rate of MRI and opioid exposure progressively increased the later in an episode a PCP, DC, PT or OS became involved. Compared to both PCPs and OSs, DCs and PTs were associated with a lower rate of

opioid exposure at any time during an episode. DCs were associated with a lower rate of MRI exposure than PCPs, PTs, or OSs. [Figure 6] Similarly, when the last HCP seen, total episode cost progressively increased later in an episode a PCP, DC, PT or OS became involved. When the last HCP seen, PCPs and DCs were associated with lower median total episode cost with PTs and OSs associated with higher median total episode cost. [Figure 7] [Table 4]

Discussion

As a novel measure with many potential confounders and in the absence of clinical outcomes data it is important not to over interpret the finding that non-prescribing HCPs, particularly DCs and PTs, are most likely to be the last HCP seen for an episode of LBP. When coupled with the growing body of evidence demonstrating high levels of CPG concordance when these same types of HCP are initially contacted by individuals with LBP, additional exploration of attributes of the last HCP seen is warranted to more thoroughly understand the role these types of HCP play in managing LBP. While non-prescribing HCPs are most likely to be the last HCP seen for an episode of LBP, the rate of use of second- and third-line services and total episode cost increases the later non-prescribing HCPs are involved in an episode. This may provide additional support for positioning non-prescribing HCPs early in an episode of LBP as a strategy for improving CPG concordance and value.

This study has several limitations including its retrospective design and those associated with the use of administrative databases. The cohort had continuous highly uniform commercial insurance coverage and the processing of administrative claims data included extensive quality and actuarial control measures, nonetheless, data errors, variability in benefit plan design, variability in enrollee cost-sharing responsibility, and missing information were potential sources of confounding or bias. Although the

commercial insurer HCP database is under continual validation it may have included errors or missing information. Summarizing total episode cost has potential limitations associated with insurance coverage, nature of network participation, and alternative reimbursement models. While individuals from all 50 states and most US territories were included, providing a measure of generalizability, the cohort did not describe a U.S representative sample.

Another important limitation was the risk of confounding and bias associated with the limited ability to control for individual preference for types of HCP seen during an episode of LBP, individual expectations, or requests for specific health care services, and potentially meaningful differences in clinical complexity of individuals contacting or being referred to different types of HCP. This limitation was partially addressed by narrowing the study population with several exclusions. The study excluded LBP associated with serious pathology, individuals with multiple episodes of LBP, and episodes involving a surgical procedure. Sub analyses were conducted to provide insight into to the degree to which individual and population factors may have influenced results.

While this study focused on individuals with a single episode of LBP, without the availability of clinical outcomes or patient self-report data, the last HCP seen cannot be assumed to be associated with, or a proxy for, resolution of LBP. Individuals may have been dissatisfied with care and discontinued all treatment, may have paid for additional treatment on a self-pay basis, or elected to pursue self-management. With the cohort having continuous insurance coverage during the study period and a median post-episode clean period of 401 days (Q1 248, Q3 638, minimum 61) during which there was no treatment of any type for LBP, it is highly unlikely that after the last HCP seen there was a worsening of LBP symptoms. In future analyses, expanding the cohort to include individuals with multiple episodes of LBP will be important.

The volume of data involved in exploring associations in the 1.4 million possible HCP combinations presented another limitation. This will be addressed through subsequent analyses of cohort characteristics and differing rates of exposure to the types of health care services.

With this study of the last HCP seen for LBP being novel, it is challenging to corroborate previous studies. When involved in an episode of LBP as either the initial or subsequent HCP, the finding that DCs are more likely to be the last HCP seen, and associated with the lowest total cost, of any type of HCP is consistent with DCs emphasis on CPG concordant non-pharmaceutical and non-interventional services. [3] The finding that PTs, are more frequently contacted as a second HCP than as an initial HCP is consistent with PTs historically being positioned as referral recipients rather than initial contact HCPs. [26,27] When contacted as a second, third or fourth HCP PTs are as likely as a DC to be the last HCP seen which makes sense given the similar emphasis on CPG recommended first-line services. [3] The finding of PTs being infrequently contacted as the initial HCP is consistent with a previous study [3], however when contacted as the initial HCP, PTs are less likely than PCPs to be the last HCP seen which indicates the potential for less practice autonomy than DCs and warrants further study. A possible interpretation of these findings is that DCs and PTs possess similar knowledge, skills and training to effectively resolve LBP, with DCs positioned as portal of entry autonomous HCPs and PTs positioned as a referral recipient from and referral source to other types of HCPs.

Conclusions

The last HCP seen by an individual with LBP is a novel area of investigation and further exploration is warranted to understand whether this study's findings can be replicated and whether the approach

provides meaningful insights into the types of HCP that are most helpful in resolving LBP. Whether seen

initially, or later in an episode, DCs are most likely to be the last HCP seen by an individual with LBP. PTs

are similar, except when they are the initial HCP contacted by an individual with LBP. While DCs and PTs

are most likely to be the last HCP seen during an episode of LBP the later in an episode a DC or PT is seen

associated beneficial attributes such as opioid avoidance are lost, and total episode cost increases. This

adds additional support to the growing body of literature advocating for positioning these types of HCPs

earlier in an episode of LBP.

References

1.   Meroni R, Piscitelli D, Ravasio C, et al. Evidence for managing chronic low back pain in primary care: a review of recommendations from high-quality clinical practice guidelines. *Disabil Rehabil*. Apr 2021;43(7):1029-1043. doi:10.1080/09638288.2019.1645888

2.   Qaseem A, Wilt TJ, McLean RM, et al. Noninvasive Treatments for Acute, Subacute, and Chronic Low Back Pain: A Clinical Practice Guideline From the American College of Physicians. *Ann Intern Med*. Apr 4 2017;166(7):514-530. doi:10.7326/M16-2367

3.   Elton D, Kosloff TM, Zhang M, et al. Low back pain care pathways and costs: association with the type of initial contact health care provider. A retrospective cohort study. *medRxiv*. 2022:2022.06.17.22276443. doi:10.1101/2022.06.17.22276443

4.   Fritz JM, Kim J, Dorius J. Importance of the type of provider seen to begin health care for a new episode low back pain: associations with future utilization and costs. *Journal of Evaluation in Clinical Practice*. 2016;22(2):247-252. doi:https://doi.org/10.1111/jep.12464

5.   Frogner BK, Harwood K, Andrilla CHA, Schwartz M, Pines JM. Physical Therapy as the First Point of Care to Treat Low Back Pain: An Instrumental Variables Approach to Estimate Impact on Opioid Prescription, Health Care Utilization, and Costs. *Health Services Research*. 2018;53(6):4629-4646. doi:https://doi.org/10.1111/1475-6773.12984

6.   Harwood KJ, Pines JM, Andrilla CHA, Frogner BK. Where to start? A two stage residual inclusion approach to estimating influence of the initial provider on health care utilization and costs for low back pain in the US. *BMC Health Serv Res*. May 23 2022;22(1):694. doi:10.1186/s12913-022-08092-1

7.   Kazis LE, Ameli O, Rothendler J, et al. Observational retrospective study of the association of initial healthcare provider for new-onset low back pain with early and long-term opioid use. *BMJ Open*. Sep 20 2019;9(9):e028633. doi:10.1136/bmjopen-2018-028633

8.   Liu X, Hanney WJ, Masaracchio M, et al. Immediate Physical Therapy Initiation in Patients With Acute Low Back Pain Is Associated With a Reduction in Downstream Health Care Utilization and Costs. *Physical Therapy*. 2018;98(5):336-347. doi:10.1093/ptj/pzy023

9.   Murphy C, French H, McCarthy G, Cunningham C. Clinical pathways for the management of low back pain from primary to specialised care: a systematic review. *Eur Spine J*. Jul 2022;31(7):1846-1865. doi:10.1007/s00586-022-07180-4

10.   Coeckelberghs E, Verbeke H, Desomer A, et al. International comparative study of low back pain care pathways and analysis of key interventions. *Eur Spine J*. Apr 2021;30(4):1043-1052. doi:10.1007/s00586-020-06675-2

11.   Magel J, Hansen P, Meier W, et al. Implementation of an Alternative Pathway for Patients Seeking Care for Low Back Pain: A Prospective Observational Cohort Study. *Phys Ther*. Dec 1 2018;98(12):1000-1009. doi:10.1093/ptj/pzy105

12.   Magel J, Suslavich K, Roper K, Fritz J, Madsen T. Emergency department evaluation, treatment, and functional outcomes among patients presenting with low back pain. *Am J Emerg Med*. Sep 2022;59:37-41. doi:10.1016/j.ajem.2022.06.048

13.   Davidson SR, Bolsewicz K, Kamper SJ, et al. Perspectives of emergency department clinicians on the challenges of addressing low back pain in the emergency setting: A qualitative study. *Emerg Med Australas*. Apr 2022;34(2):199-208. doi:10.1111/1742-6723.13854

14.   Ojha HA, Wyrsta NJ, Davenport TE, Egan WE, Gellhorn AC. Timing of Physical Therapy Initiation for Nonsurgical Management of Musculoskeletal Disorders and Effects on Patient Outcomes: A Systematic Review. *J Orthop Sports Phys Ther*. Feb 2016;46(2):56-70. doi:10.2519/jospt.2016.6138

15.   Bureau USC. ACS Demographic and Housing Estimates. Accessed July 8, 2020. https://data.census.gov/cedsci/table?q=United%20States&g=0100000US&y=2018&tid=ACSDP1Y2018.DP05&hidePreview=true.

16.   Kind AJH, Buckingham WR. Making Neighborhood-Disadvantage Metrics Accessible - The Neighborhood Atlas. *N Engl J Med*. Jun 28 2018;378(26):2456-2458. doi:10.1056/NEJMp1802313

17.   Service IR. US Department of Treasury SOI tax stats - individual income tax statistics - 2017 ZIP code data. Accessed July 20, 2020. https://www.irs.gov/statistics/soi-tax-stats-individual-income-tax-statistics-2017-zip-code-data-soi.

18.   von Elm E, Altman DG, Egger M, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: guidelines for reporting observational studies. *Int J Surg*. Dec 2014;12(12):1495-9. doi:10.1016/j.ijsu.2014.07.013

19.   Hernan MA. The C-Word: Scientific Euphemisms Do Not Improve Causal Inference From Observational Data. *Am J Public Health*. May 2018;108(5):616-619. doi:10.2105/AJPH.2018.304337

20.   Smith GD, Ebrahim S. Data dredging, bias, or confounding. *BMJ*. Dec 21 2002;325(7378):1437-8. doi:10.1136/bmj.325.7378.1437

21.   King G NR. Why Propensity Scores Should Not Be Used for Matching. *Political Analysis*. 2019;27(4):435-454. doi:10.1017/pan.2019.11

22.   National Academies of Sciences E, and Medicine. *Health-Care Utilization as a Proxy in Disability Determination*. 2 Factors That Affect Health-Care Utilization. National Academies Press 2018.

23.   Peterson C, Grosse SD, Dunn A. A practical guide to episode groupers for cost-of-illness analysis in health services research. *SAGE Open Med*. 2019;7:2050312119840200. doi:10.1177/2050312119840200

24.   Insight O. Symmetry episode treatment groups: measuring health care with meaningful episodes of care. 2012.

25.   Viera AJ. Odds ratios and risk ratios: what's the difference and why does it matter? *South Med J*. Jul 2008;101(7):730-4. doi:10.1097/SMJ.0b013e31817a7ee4

26.   Clark B, Clark L, Showalter C, Stoner T. A call to action: direct access to physical therapy is highly successful in the US military. When will professional bodies, legislatures, and payors provide the same advantages to all US civilian physical therapists? *J Man Manip Ther*. Aug 2022;30(4):199-206. doi:10.1080/10669817.2022.2099893

27.   Hon S, Ritter R, Allen DD. Cost-Effectiveness and Outcomes of Direct Access to Physical Therapy for Musculoskeletal Disorders Compared to Physician-First Access in the United States: Systematic Review and Meta-Analysis. *Phys Ther*. Jan 4 2021;101(1)doi:10.1093/ptj/pzaa201

List of Abbreviations:

LBP – Low back pain

US – United States

CPG – Clinical practice guideline

PCP – Primary care provider

DC – Doctor of Chiropractic

PT – Physical Therapist

LAc – Licensed acupuncturist

HCP – Health care provider

ADI – Area Deprivation Index

WTP – Willingness to pay

STROBE – Strengthening the Reporting of Observational Studies in Epidemiology

ETG® – Episode Treatment Group®

ERG® – Episode Risk Group®

ACP – American College of Physicians

RR – Risk ratio

OR – Odds ratio

CI – Confidence interval

CMT – Chiropractic manipulative treatment

AC – Active care

MT – Manual therapy

Table 1 - Non-surgical low back pain episode count by type of health care provider (HCP) and episode sequence position																
		# of Unique HCPs					# of Episodes					% of Episodes				
HCP type		First HCP	Second HCP	Third HCP	Fourth HCP	Fifth HCP	First HCP	Second HCP	Third HCP	Fourth HCP	Fifth HCP	First HCP	Second HCP	Third HCP	Fourth HCP	Fifth HCP
Primary Care	PCP	68888	28730	11787	4889	2301	177265	40767	13766	5239	2364	35.2%	26.8%	23.2%	20.0%	18.5%
	Nurse	21226	9897	4101	1919	1049	37857	12820	4749	2108	1031	7.5%	8.4%	8.0%	8.0%	8.1%
	PA	12787	6746	2955	1373	781	25350	9418	3542	1545	718	5.0%	6.2%	6.0%	5.9%	5.6%
	DO	121	45	30	9	7	469	64	42	11	10	0.1%	0.0%	0.1%	0.0%	0.1%
Non-Prescriber	DC	21644	9953	4396	1735	739	130520	18586	5823	2011	797	25.9%	12.2%	9.8%	7.7%	6.2%
	PT	3019	4143	2769	1601	776	5114	7344	4076	2077	1135	1.0%	4.8%	6.9%	7.9%	8.9%
	LAc	1430	397	159	67	36	2634	486	175	73	30	0.5%	0.3%	0.3%	0.3%	0.2%
Specialist	OS	8590	5241	3048	1740	1120	27799	10867	4971	2362	1152	5.5%	7.1%	8.4%	9.0%	9.0%
	PMR	2966	2127	1622	1120	767	12496	5501	3236	1817	1009	2.5%	3.6%	5.5%	6.9%	7.9%
	PM	2497	1888	1561	1126	782	9013	4280	2884	1757	1031	1.8%	2.8%	4.9%	6.7%	8.1%
	NS	1373	1164	894	650	441	3058	2084	1322	817	432	0.6%	1.4%	2.2%	3.1%	3.4%
	Neuro	2352	1321	709	371	194	3917	1729	843	410	224	0.8%	1.1%	1.4%	1.6%	1.8%
	Rheum	1642	998	558	304	158	3354	1463	667	332	166	0.7%	1.0%	1.1%	1.3%	1.3%
	MD Oth	9230	5122	2332	1259	723	13877	6716	2963	1539	840	2.8%	4.4%	5.0%	5.9%	6.6%
Emergency /Urgent Care	EM	8365	3658	1313	511	301	15760	4838	1463	544	233	3.1%	3.2%	2.5%	2.1%	1.8%
	Rad	10098	8613	4763	2468	2039	30393	23828	8454	3475	1551	6.0%	15.7%	14.3%	13.3%	12.2%
	UC	1120	517	186	73	43	5082	1358	272	87	39	1.0%	0.9%	0.5%	0.3%	0.3%
Total		177348	90560	43183	21215	12257	503958	152149	59248	26204	12762	100.0%	100.0%	100.0%	100.0%	100.0%
% of Total Episodes												100.0%	30.2%	11.8%	5.2%	2.5%

PCP=primary care provider, DO=doctor of osteopathy, DC=doctor of chiropractic, PT=physical therapist, LAc=licensed acupuncturist, OS=orthopedic surgeon, PMR=physical medicine & rebilitation, PM=pain medicine, NS=neurosurgeon, Neuro=neurologist, Rheum=rheumatologist, MD Oth = other medical physician, EM=emergency medicine, Rad=radiologist, UC=urgent care

Table 2 - % of non-surgical low back pain episodes where health care provider (HCP) type was the last HCP seen									
HCP type		% where no follow-up HCP				Risk Ratio and 95% confidence interval compared to PCP			
		First No Second	Second No Third	Third No Fourth	Fourth No Fifth	First No Second	Second No Third	Third No Fourth	Fourth No Fifth
Primary Care	PCP	71.0%	65.9%	61.4%	54.7%	Reference			
	Nurse	62.1%	60.2%	53.4%	48.3%	0.87 (0.87, 0.88)	0.91 (0.90, 0.93)	0.87 (0.84, 0.90)	0.88 (0.84, 0.93)
	PA	58.8%	56.4%	51.8%	47.5%	0.83 (0.82, 0.84)	0.86 (0.84, 0.87)	0.84 (0.81, 0.87)	0.87 (0.82, 0.92)
	DO	80.2%	64.1%	66.7%	36.4%	**1.13 (1.08, 1.18)**	0.97 (0.81, 1.17)	1.09 (0.88, 1.35)	0.67 (0.30, 1.45)
Non-Prescriber	DC	84.0%	72.7%	67.7%	61.6%	**1.18 (1.18, 1.19)**	**1.10 (1.09, 1.12)**	**1.10 (1.08, 1.13)**	**1.13 (1.08, 1.17)**
	PT	60.0%	70.9%	63.2%	60.7%	0.84 (0.83, 0.86)	**1.08 (1.06, 1.09)**	**1.03 (1.00, 1.06)**	**1.11 (1.06, 1.16)**
	LAc	81.7%	62.3%	62.9%	50.7%	**1.15 (1.13, 1.17)**	0.95 (0.88, 1.01)	1.02 (0.91, 1.15)	0.93 (0.74, 1.16)
Specialist	OS	63.2%	56.8%	52.8%	49.8%	0.89 (0.88, 0.90)	0.86 (0.85, 0.88)	0.86 (0.83, 0.88)	0.91 (0.87, 0.95)
	PMR	64.0%	58.5%	56.5%	55.7%	0.90 (0.89, 0.91)	0.89 (0.87, 0.91)	0.92 (0.89, 0.95)	1.02 (0.97, 1.07)
	PM	61.5%	54.7%	55.3%	52.9%	0.87 (0.85, 0.88)	0.83 (0.81, 0.85)	0.90 (0.87, 0.93)	0.97 (0.92, 1.02)
	NS	51.6%	47.6%	43.9%	45.4%	0.73 (0.70, 0.75)	0.72 (0.69, 0.76)	0.71 (0.67, 0.76)	0.83 (0.77, 0.90)
	Neuro	57.1%	56.6%	55.2%	51.7%	0.80 (0.78, 0.83)	0.86 (0.82, 0.90)	0.90 (0.84, 0.96)	0.95 (0.86, 1.04)
	Rheum	62.0%	57.8%	54.1%	52.1%	0.87 (0.85, 0.90)	0.88 (0.84, 0.92)	0.88 (0.82, 0.95)	0.95 (0.86, 1.06)
	MD Oth	55.9%	59.8%	55.3%	52.0%	0.79 (0.78, 0.80)	0.91 (0.89, 0.93)	0.90 (0.87, 0.93)	0.95 (0.90, 1.00)
Emergency /Urgent Care	EM	60.4%	56.0%	52.4%	43.2%	0.85 (0.84, 0.86)	0.85 (0.83, 0.87)	0.85 (0.81, 0.90)	0.79 (0.72, 0.87)
	Rad	49.0%	49.5%	42.0%	38.3%	0.69 (0.68, 0.70)	0.75 (0.74, 0.76)	0.68 (0.67, 0.70)	0.70 (0.67, 0.74)
	UC	62.6%	65.3%	55.5%	50.6%	0.88 (0.86, 0.90)	0.99 (0.95, 1.03)	0.90 (0.81, 1.01)	0.93 (0.75, 1.14)

Cells in red have risk ratio 95% confidence interval crossing the PCP reference of 1
Cells highlighted with bold font have at least 20 episodes and risk ratio 95% plus confidence interval greater than the PCP reference of 1 indicating greater likelihood to be the last HCP seen

PCP=primary care provider, DO=doctor of osteopathy, DC=doctor of chiropractic, PT=physical therapist, LAc=licensed acupuncturist, OS=orthopedic surgeon, PMR=physical medicine & rebilitation, PM=pain medicine, NS=neurosurgeon, Neuro=neurologist, Rheum=rheumatologist, MD Oth = other medical physician, EM=emergency medicine, Rad=radiologist, UC=urgent care

Table 3 - Non-surgical low back pain episodes initially contacting a primary care provider (PCP), orthopedic surgeon (OS) or emergency medicine physician (EM), subsequent health care providers (HCP) seen, and % of episodes where the subsequent HCP was the last HCP seen																		
	Initial HCP Contacted For First Episode - PCP						Initial HCP Contacted For First Episode - OS						Initial HCP Contacted For First Episode - EM					
	Second HCP		Third HCP		Fourth HCP		Second HCP		Third HCP		Fourth HCP		Second HCP		Third HCP		Fourth HCP	
	Episodes	No 3rd	Episodes	No 4th	Episodes	No 5th	Episodes	No 3rd	Episodes	No 4th	Episodes	No 5th	Episodes	No 3rd	Episodes	No 4th	Episodes	No 5th
PCP	11709	66.1%	6762	63.4%	2055	53.7%	1645	62.2%	564	52.3%	232	48.3%	1614	68.4%	571	66.7%	256	56.2%
Nurse	4787	62.3%	1147	52.5%	570	50.9%	441	51.0%	168	50.0%	81	43.2%	541	65.1%	199	60.8%	72	41.7%
PA	3340	60.7%	933	49.8%	456	45.0%	755	48.6%	187	49.7%	104	44.2%	541	59.9%	166	54.8%	66	59.1%
DO	16	68.8%	9	100.0%	1	100.0%	6	83.3%	3	0.0%	0		3	100.0%	3	66.7%	1	100.0%
DC	6824	76.2%	1189	69.6%	529	62.2%	699	69.5%	200	64.5%	98	54.1%	362	76.5%	132	73.5%	51	64.7%
PT	2660	76.1%	1270	65.2%	683	61.1%	1573	71.2%	491	62.3%	238	63.9%	78	62.8%	113	70.8%	56	71.4%
LAc	128	69.5%	49	51.0%	6	66.7%	18	66.7%	4	75.0%	4	100.0%	11	45.5%	6	50.0%	3	33.3%
OS	3589	57.7%	1329	52.7%	689	50.2%	1148	54.1%	1273	54.2%	437	51.3%	274	60.9%	164	62.2%	80	57.5%
PMR	1772	60.4%	836	53.9%	553	50.8%	564	60.6%	258	59.3%	155	56.1%	126	65.1%	93	58.1%	43	65.1%
PM	1293	59.7%	604	57.5%	446	59.4%	400	60.5%	194	60.3%	160	56.2%	62	58.1%	40	57.5%	31	54.8%
NS	701	48.1%	437	43.9%	272	44.1%	99	45.5%	67	41.8%	42	57.1%	63	58.7%	49	51.0%	38	50.0%
Neuro	580	58.4%	221	55.2%	144	44.4%	138	53.6%	51	31.4%	23	60.9%	31	61.3%	22	54.5%	7	57.1%
Rheum	471	56.9%	151	57.6%	74	55.4%	90	55.6%	39	66.7%	23	47.8%	20	50.0%	7	42.9%	7	71.4%
MD Oth	2009	63.4%	753	56.0%	396	54.5%	487	60.4%	206	58.7%	120	46.7%	317	64.0%	98	58.2%	46	47.8%
EM	1224	53.1%	433	49.9%	175	40.0%	109	53.2%	38	60.5%	27	37.0%	535	56.6%	284	58.1%	70	54.3%
Rad	9814	52.8%	2863	41.5%	1188	40.1%	2039	47.4%	538	42.9%	221	35.3%	1606	51.6%	442	43.2%	159	34.6%
UC	555	66.3%	88	50.0%	22	45.5%	20	45.0%	10	70.0%	4	75.0%	58	58.6%	19	47.4%	6	33.3%
Risk ratio and 95% confidence interval compared to the PCP reference																		
PCP	1						1						1					
Nurse	0.94 (0.92, 0.97)		0.83 (0.78, 0.88)		0.95 (0.87, 1.04)		0.82 (0.74, 0.90)		0.96 (0.81, 1.13)		0.90 (0.67, 1.19)		0.95 (0.89, 1.02)		0.91 (0.80, 1.03)		0.74 (0.55, 0.99)	
PA	0.92 (0.89, 0.95)		0.79 (0.73, 0.84)		0.84 (0.75, 0.93)		0.78 (0.72, 0.85)		0.95 (0.81, 1.12)		0.92 (0.71, 1.18)		0.88 (0.81, 0.95)		0.82 (0.71, 0.95)		1.05 (0.84, 1.32)	
DO	1.04 (0.75, 1.45)		1.58 (1.55, 1.61)		1.86 (1.79, 1.94)		1.34 (0.93, 1.92)		N/A		N/A		1.46 (1.41, 1.51)		1.00 (0.45, 2.23)		1.78 (1.60, 1.98)	
DC	**1.15 (1.13, 1.18)**		**1.10 (1.05, 1.14)**		**1.16 (1.07, 1.25)**		**1.12 (1.05, 1.19)**		**1.23 (1.08, 1.40)**		1.12 (0.89, 1.40)		**1.12 (1.05, 1.20)**		1.10 (0.98, 1.24)		1.15 (0.91, 1.45)	
PT	**1.15 (1.12, 1.18)**		1.03 (0.98, 1.07)		**1.14 (1.06, 1.22)**		**1.14 (1.09, 1.20)**		**1.19 (1.07, 1.32)**		**1.32 (1.12, 1.56)**		0.92 (0.77, 1.09)		1.06 (0.93, 1.21)		**1.27 (1.04, 1.55)**	
LAc	1.05 (0.94, 1.18)		0.80 (0.61, 1.06)		1.24 (0.70, 2.19)		1.07 (0.77, 1.49)		1.43 (0.81, 2.54)		2.07 (1.81, 2.37)		0.66 (0.35, 1.27)		0.75 (0.34, 1.67)		0.59 (0.12, 2.95)	
OS	0.87 (0.85, 0.90)		0.83 (0.79, 0.88)		0.94 (0.86, 1.02)		0.87 (0.81, 0.93)		1.04 (0.94, 1.14)		1.06 (0.90, 1.25)		0.89 (0.81, 0.99)		0.93 (0.82, 1.06)		1.02 (0.82, 1.27)	
PMR	0.91 (0.88, 0.95)		0.85 (0.80, 0.91)		0.95 (0.86, 1.04)		0.97 (0.90, 1.05)		1.13 (1.00, 1.29)		1.16 (0.96, 1.41)		0.95 (0.83, 1.09)		0.87 (0.73, 1.04)		1.16 (0.91, 1.48)	
PM	0.90 (0.86, 0.95)		0.91 (0.84, 0.97)		**1.11 (1.02, 1.21)**		0.97 (0.89, 1.06)		**1.15 (1.00, 1.32)**		1.17 (0.96, 1.41)		0.85 (0.69, 1.05)		0.86 (0.66, 1.13)		0.97 (0.70, 1.37)	
NS	0.73 (0.67, 0.79)		0.69 (0.62, 0.77)		0.82 (0.71, 0.95)		0.73 (0.59, 0.91)		0.80 (0.60, 1.07)		1.18 (0.88, 1.59)		0.86 (0.70, 1.06)		0.76 (0.58, 1.01)		0.89 (0.64, 1.24)	
Neuro	0.88 (0.83, 0.95)		0.87 (0.77, 0.98)		0.83 (0.69, 1.00)		0.86 (0.73, 1.01)		0.60 (0.40, 0.91)		1.26 (0.89, 1.80)		0.90 (0.68, 1.19)		0.82 (0.56, 1.20)		1.02 (0.53, 1.95)	
Rheum	0.86 (0.80, 0.93)		0.91 (0.79, 1.04)		1.03 (0.84, 1.27)		0.89 (0.74, 1.08)		**1.27 (1.01, 1.61)**		0.99 (0.63, 1.55)		0.73 (0.47, 1.13)		0.64 (0.27, 1.51)		1.27 (0.79, 2.05)	
MD Oth	0.96 (0.93, 0.99)		0.88 (0.83, 0.94)		1.02 (0.92, 1.12)		0.97 (0.89, 1.05)		1.12 (0.98, 1.29)		0.97 (0.77, 1.22)		0.94 (0.86, 1.02)		0.87 (0.73, 1.04)		0.85 (0.62, 1.17)	
EM	0.80 (0.76, 0.85)		0.79 (0.71, 0.87)		0.75 (0.62, 0.90)		0.85 (0.71, 1.02)		1.16 (0.88, 1.51)		0.77 (0.46, 1.28)		0.83 (0.76, 0.90)		0.87 (0.78, 0.98)		0.97 (0.76, 1.23)	
Rad	0.80 (0.78, 0.82)		0.65 (0.62, 0.69)		0.75 (0.69, 0.81)		0.76 (0.72, 0.81)		0.82 (0.72, 0.93)		0.73 (0.59, 0.91)		0.75 (0.71, 0.80)		0.65 (0.57, 0.73)		0.61 (0.48, 0.78)	
UC	1.00 (0.94, 1.07)		0.79 (0.64, 0.97)		0.85 (0.53, 1.34)		0.72 (0.44, 1.18)		1.34 (0.89, 2.02)		1.55 (0.87, 2.78)		0.86 (0.69, 1.07)		0.71 (0.44, 1.14)		0.59 (0.19, 1.85)	

Cells in red have risk ratio 95% confidence interval crossing the PCP reference of 1

Cells highlighted with bold font have at least 20 episodes and risk ratio 95% plus confidence interval greater than the PCP reference of 1 indicating greater likelihood to be the last HCP seen

PCP=primary care provider, DO=doctor of osteopathy, DC=doctor of chiropractic, PT=physical therapist, LAc=licensed acupuncturist, OS=orthopedic surgeon, PMR=physical medicine & rebilitation, PM=pain medicine, NS=neurosurgeon, Neuro=neurologist, Rheum=rheumatologist, MD Oth = other medical physician, EM=emergency medicine, Rad=radiologist, UC=urgent care

Table 4 - Non-surgical low back pain service use and total cost for select health care providers (HCP) positioned as the last HCP seen																
	Primary care provider (PCP)				Doctor of Chiropractic (DC)				Physical Therapist (PT)				Orthopedic Surgeon (OS)			
Episode Sequence	1st	2nd	3rd	4th	1st	2nd	3rd	4th	1st	2nd	3rd	4th	1st	2nd	3rd	4th
	No 2nd	No 3rd	No 4th	No 5th	No 2nd	No 3rd	No 4th	No 5th	No 2nd	No 3rd	No 4th	No 5th	No 2nd	No 3rd	No 4th	No 5th
Episodes	88980	24766	8251	2835	107340	13409	3928	1235	3030	5177	2572	1261	15314	5965	2593	1172
% of episodes including service																
Manipulation- Chiropractic	4.1%	20.7%	18.4%	20.9%	93.6%	91.6%	94.8%	93.9%	7.8%	6.6%	10.1%	12.5%	3.1%	11.6%	13.5%	14.9%
Active Care	9.8%	15.1%	21.2%	28.3%	30.9%	35.5%	43.4%	47.9%	96.6%	97.2%	97.3%	97.9%	20.1%	27.6%	35.2%	42.2%
Passive Therapy	4.5%	10.3%	11.9%	14.8%	36.8%	38.9%	46.3%	50.3%	28.7%	26.5%	28.5%	30.3%	6.7%	11.0%	14.3%	14.2%
Manual Therapy	7.3%	10.3%	14.0%	18.8%	17.9%	21.0%	27.3%	33.2%	77.4%	74.2%	73.8%	74.8%	14.1%	18.8%	24.0%	28.5%
Acupuncture	0.3%	0.6%	0.7%	0.7%	0.5%	1.2%	1.9%	2.3%	1.2%	0.6%	0.9%	1.0%	0.2%	0.5%	0.6%	0.7%
Manipulation - Osteopathic	2.5%	1.3%	1.6%	1.5%	0.1%	0.4%	0.5%	0.4%	0.3%	0.4%	0.7%	0.6%	0.7%	0.4%	0.8%	0.5%
Imaging - Radiography	21.1%	31.6%	45.7%	51.5%	18.8%	29.5%	43.0%	53.6%	16.3%	42.7%	57.7%	63.4%	73.6%	68.7%	71.9%	73.5%
Rx - NSAID	44.2%	42.7%	49.5%	52.6%	4.0%	20.1%	24.0%	30.9%	9.9%	22.4%	30.8%	33.0%	25.8%	39.0%	44.2%	50.1%
Rx - Skeletal Muscle Relaxant	47.5%	43.3%	46.7%	47.9%	2.3%	16.1%	22.3%	27.0%	9.3%	22.8%	29.2%	32.8%	15.8%	26.6%	33.2%	36.3%
Imaging - MRI	4.9%	6.5%	13.3%	24.8%	1.2%	3.7%	8.8%	16.8%	9.6%	11.4%	22.2%	33.0%	18.5%	27.8%	40.2%	50.5%
Rx-Opioid	26.4%	26.9%	35.9%	39.9%	3.4%	11.4%	15.6%	21.1%	6.6%	8.6%	13.7%	19.2%	12.5%	20.0%	28.1%	33.4%
Spinal Injection	2.5%	2.7%	4.6%	7.9%	0.6%	2.0%	3.8%	5.4%	3.4%	3.3%	5.4%	10.2%	7.0%	8.8%	12.3%	18.2%
Imaging-CT	1.1%	2.2%	4.3%	6.0%	0.2%	0.6%	1.4%	3.4%	0.8%	0.9%	1.4%	2.5%	1.3%	2.7%	4.3%	5.5%
Total Episode Cost - Median (Q1, Q3)																
Total Episode Cost	$106	$218	$455	$835	$165	$320	$573	$877	$528	$688	$1,012	$1,333	$229	$504	$920	$1,343
	(30,	(97,	(202,	(364,	(68,	(149,	(293,	(488,	(250,	(418,	(592,	(818,	(121,	(236,	(429,	(719,
	220)	537)	1131)	1926)	397)	658)	1071)	1661)	1041)	1172)	1693)	2297)	597)	1205)	1867)	2473)

Figure 1. Flow of episodes from initial (1) to second (2) to third (3) to fourth (4) to fifth (5) type of health care provider (HCP) seen during an episode. PCP=primary care provider, DC=doctor of chiropractic, Rad=radiologist, OS=orthopedic surgeon, PA=physician's assistant, EM=emergency medicine, Oth=other, PMR=physical medicine & rehabilitation, PM=pain medicine, PT=physical therapist, UC=urgent care, Neuro=neurologist, Rheum=rheumatologist, NS=neurosurgeon, LAc=license acupuncturist, DO=doctor of osteopathy, None=no next HCP seen (previous HCP was the last HCP)

Figure 2. Risk ratio and 95% confidence interval compared to primary care provider (PCP) reference for % of episodes where HCPs with at least 20 episodes were the initial and last or 2nd and last HCP . PA=physician's assistant, DO=doctor of osteopathy, DC=doctor of chiropractic, PT=physical therapist, LAc=licensed acupuncturist, OS=orthopedic surgeon, PMR=physical medicine & rehabilitation, PM=pain management, NS=neurosurgeon, Neuro=neurologist, Rheum=rheumatologist, Oth=other, EM=emergency medicine, Rad=radiologist, UC=urgent care

Figure 3. Risk ratio and 95% confidence interval compared to primary care provider (PCP) reference for % of episodes where HCPs with at least 20 episodes were the initial and last HCP seen segmented by population Area Deprivation Index (ADI) and % of the population that is non-Hispanic white (NHW). PA=physician's assistant, DO=doctor of osteopathy, DC=doctor of chiropractic, PT=physical therapist, LAc=licensed acupuncturist, OS=orthopedic surgeon, PMR=physical medicine & rehabilitation, PM=pain management, NS=neurosurgeon, Neuro=neurologist, Rheum=rheumatologist, Oth=other, EM=emergency medicine, Rad=radiologist, UC=urgent care

Figure 4. Risk ratio and 95% confidence interval compared to primary care provider (PCP) reference for % of episodes where HCPs with at least 20 episodes were the second and last HCP seen segmented by population Area Deprivation Index (ADI) and % of the population that is non-Hispanic white (NHW). PA=physician's assistant, DO=doctor of osteopathy, DC=doctor of chiropractic, PT=physical therapist, LAc=licensed acupuncturist, OS=orthopedic surgeon, PMR=physical medicine & rehabilitation, PM=pain management, NS=neurosurgeon, Neuro=neurologist, Rheum=rheumatologist, Oth=other, EM=emergency medicine, Rad=radiologist, UC=urgent care

Figure 5. - Risk ratio and 95% confidence interval compared to primary care provider (PCP) reference for % of episodes where for health care providers with at least 20 episodes the type of second HCP seen was also the last HCP seen. PA=physician's assistant, DO=doctor of osteopathy, DC=doctor of chiropractic, PT=physical therapist, LAc=licensed acupuncturist, OS=orthopedic surgeon, PMR=physical medicine & rehabilitation, PM=pain management, NS=neurosurgeon, Neuro=neurologist, Rheum=rheumatologist, Oth=other, EM=emergency medicine, Rad=radiologist, UC=urgent care

Figure 6. % of episodes with an MRI scan or prescription opioid for episodes where a primary care physician (PCP), chiropractor (DC), physical therapist (PT) or orthopedic surgeon (OS) was the last health care provider seen based on whether the PCP, DC, PT, or OS where the 1st, 2nd, 3rd or 4th HCP seen during an episode

Figure 7. Median total episode cost and interquartile range (Q1, Q3) for episodes where a primary care physician (PCP), chiropractor (DC), physical therapist (PT) or orthopedic surgeon (OS) was the last health care provider seen based on whether the PCP, DC, PT, or OS where the 1st, 2nd, 3rd or 4th HCP seen during an episode