Overweight and Obesity among Women at Reproductive Age in Cambodia: Data Analysis of Cambodia Demographic and Health Survey 2014

Samnang Um[1*], An Yom[1], Jonathan A. Muir[2], Heng Sopheab[1]

1- School of Public Health at the National Institute of Public Health, Cambodia
2- The Global Health Institute, Emory University, Atlanta, Georgia, United States

*Corresponding authors: Samnang Um, email: umsamnang56@gmail.com,

[ORCID: https://orcid.org/0000-0002-1521-1260],

An Yom, email: anyomniph@gmail.com,

Jonathan A. Muir, email: jamuir@emory.edu

[ORCID: https://orcid.org/0000-0002-8039-7145]

Heng Sopheab, email: hsopheab@niph.org.kh,

[ORCID: https://orcid.org/0000-0002-6553-1334]

Address: Lot#80, 289 Samdach Penn Nouth St. (289), Phnom Penh

ABSTRACT

Introduction

Overweight and obesity increase the risk of cardiovascular disease, type 2 diabetes mellitus,

hypertension, stroke, and some type of cancers, and maternal health globally. In Cambodia, the

prevalence of overweight and obesity among women aged 15-49 years increased from 6% in

2000 to 18% in 2014, becoming a public health burden. We examined socio-demographic and

behavioral factors associated with overweight and/or obesity among women of reproductive age

in Cambodia.

Methods

We analyzed existing data from the 2014 Cambodia Demographic and Health Survey (CDHS)

that used a two-stage stratified cluster sampling design. Data analysis was restricted to non-

pregnant women, resulting in an analytic sample of 10,818 women. Multivariable logistic

regressions that accounted for the CDHS sampling design were performed using STATA V16 to

examine factors associated with overweight and obesity.

Results

Prevalence of overweight and obesity among non-pregnant women of reproductive age were

15.2% and 2.8% respectively. Factors independently associated with increased odds of

overweight and/or obesity including women aged 20-29 years with adjusted odds ratio

[AOR=2.4; 95% CI: 1.6-3.6], 30-39 years [AOR=4.6; 95% CI: 3.0-6.9], and 40-49 years

[AOR=6.6; 95% CI: 4.3-10.1], married women [AOR=1.8; 95% CI: 1.3-2.7], urban residence

[AOR=1.3; 95% CI: 1.1-1.5], and women having at least 4 children [AOR=1.7; 95% CI: 1.2-

2.5]. The following factors were associated with decreased odds of overweight and obesity:

completed at least secondary education [AOR=0.7; 95% CI: 0.6-0.8], agricultural work

[AOR=0.7; 95% CI: 0.5-0.8], and manual labor work [AOR=0.7; 95% CI: 0.6-0.9].

Conclusions

Increased age, married women, living in urban residence, and having at least four children were
the main risk factors associated with overweight and/or obesity. Conversely, higher education,
working in agriculture, and working in manual labor were negatively associated with overweight
and/or obesity. Cambodia's non-communicable disease (NCD) public health programs should
consider these characteristic for targeting interventions to further reduce overweight and/or
obesity in the coming years.
54
55 **Keywords:** Overweight, obesity, non-communicable diseases, NCDs, women at reproductive
56 age, CDHS, Cambodia
57

Introduction

Obesity or being overweight is associated with increased risk of cardiovascular disease, type 2 diabetes mellitus, systemic hypertension, obstructive sleep apnea, stroke, and some type of cancers, and maternal disorders among women of reproductive age (WRA) [1, 2]. According to the WHO, overweight and obesity are leading risks for global deaths with at least 2.8 million adults dying annually due to these conditions [3]. Globally, 2 billion adults aged 18 years and older were overweight, and 650 million were obese in 2016 [3]. In both developed and developing countries, obesity or being overweight were higher in women than men; for instance, obesity was 15% in women compared to 11% in men [2, 3]. The Global Nutrition Report 2019 showed that 26.1% of women and 20.4% of men were overweight, while obesity was 6.3% in women compared to 3.5% in men [4]. The final report for the 2104 Cambodia Demographic and Health Survey (CDHS) indicated that overweight and obesity increased from 6% in 2000 to 18% in 2014 [5]. It was estimated that overweight and obesity contribute to rising healthcare costs in Cambodia (approximately 1.7% of its annual gross domestic product (GDP) per capita) and is a major contributor to mortality and decreased general health and productivity [6].

Factors commonly associated with overweight and obesity among women include higher socioeconomic status, older age, marriage, living in an urban residence, and lack of education [7, 8]. Women with formal employment had higher odds of being overweight or obese than informally employed women [9]. Also, women who used hormonal contraceptives such as oral contraceptive pills, implants, patches, and rings were at higher risk of being overweight or obese [10]. Globally, 30% of daily smokers are overweight or obese [11], with women smokers at greater risk for overweight or obesity than men smokers [12, 13]. While health scholars have examined these factors internationally, to our knowledge, factors associated with overweight and obesity specifically among WRA in Cambodia have not been explored. Therefore, we examined socio-demographic and behavioral factors associated with overweight and/or obesity among WRA.

Materials and Methods

Ethics Statement

The data used in this study were extracted from CDHS 2014 data [14], which are publicly available with all personal identifiers of study participants removed. Permission to analyze the data was granted through registering with the DHS program website and submitting an application outlining the intended use of the datasets [14]. Informed consent was obtained from the study participants before data collection. The data collection tools and procedures for CHDS 2014 was approved by the Cambodia National Ethics Committee for Health Research (Ref: # 056 NECHR) and the Institutional Review Board (IRB) of ICF in Rockville, Maryland, USA.

Data sources and procedures

We used the women's dataset from CDHS 2014, which was a nationally representative population-based household survey. Survey data were obtained through face-to-face survey interviews using a formal survey instrument by trained interviewers. Two-stage stratified cluster sampling was used to collect the samples. In the first stage, enumeration areas (EAs) stratified by urban-rural were selected from the sampling frame using probability proportional to cluster size. In the second stage, a defined number of households, generally 25-30, were selected from the list of households in each EA using the systematic sampling [5]. Eligible WRA in selected households were invited to complete a survey interview. In total, 17,578 women aged 15-49 years were interviewed. Weight and height measurements for women were taken by trained female field staff using standardized instruments and procedures [5]. Height in centimeters (cm), measured using standardized measuring boards with accuracy to 0.1 cm, and weight in kilograms (kg), measured using solar-powered scales (UNICEF electronic scale or Uniscale) with accuracy to 0.1 kg [5]. Data quality assurance and cleaning followed standard operating procedures, which cover every step of the research process, from data collection to data entry [15]. The data were restricted to only include non-pregnant women aged 15-49 years with available body mass index (BMI) data, resulting in the exclusion of 974 pregnant women and 5,786 women with missing BMI data. Data restriction resulted in a final analytic sample of 10,818 WRA.

Outcome variable

The outcome variable of this study was **Overweight/Obese** for WRA. For the outcome variable **Overweight/Obese**, BMI was first calculated as weight in kilograms divided by the square of height in meters (kg/m^2) [16]. BMI was then measured as follows: underweight $(< 18.0\ kg/m^2)$, normal weight $(18.5\text{-}24.9\ kg/m^2)$, overweight $(25.0\text{-}29.9\ kg/m^2)$ and obesity $(\geq 30.0\ kg/m^2)$ [5]. **Overweight/Obese** was defined as a binary outcome for which women with a BMI $\geq 25.0\ kg/m^2$ were classified as overweight and obese (coded = 1), while women with a BMI $< 25.0\ kg/m^2$ were coded as other (coded = 0).

Explanatory variables

Explanatory variables included socio-demographic characteristics and behavioral factors: Women's age in years (15-19, 20-29, 30-39, and 40-49), marital status (not married, married or living together, and divorced or widowed or separated), education level (no education, primary, secondary and higher), occupation (not working, agriculture, manual labor or unskilled, professional or sealer or services). Households' wealth status was represented by a wealth index calculated via principal component analysis (PCA) and using variables for household assets and dwelling characteristics. Weighted scores divided into five wealth quintiles (poorest, poorer, medium, richer, and richest) each comprising 20% of the population [5, 15, 16]. Then, the original variable was then recombined into three categories including richer/richest, middle, and poorer/poorest. Residence (urban vs. rural), number of children ever born (no children, one child, two-three children, four and more children), smoking (non-smoker vs smoker), contraceptive usage (not used, hormonal methods (using a pill, Norplant, injection, and vaginal rings), non-hormonal methods (condoms, the diaphragm, the IUD, spermicides, lactational amenorrhea, and sterilization), and traditional methods (periodic abstinence and withdrawal).

Statistical analysis

Data analyses were performed using STATA version 16 (Stata Corp 2017, College Station, TX) and accounted for the CDHS sampling weights and complex survey design. Socio-demographic characteristics and behavioral factors were described in weighted frequency and percentage. Descriptive analyses that coupled cross-tabular frequency distributions with chi-square tests were

used to assess associations between explanatory variables and the outcome variable **Overweight/Obese**. Variables associated with the outcome variable with a significance level of p-value $\leq$ 0.10 or that had theoretical justification (for example, women's age, wealth index, and residence) were included in the unadjusted and adjusted logistic regression analyses [17-19].

Unadjusted logistic regression was used to determine the magnitude effect of associations between overweight and obesity with socio-demographic characteristics and behavioral factors reported as odds ratios (OR) with 95% confidence intervals (CI). Then, adjusted logistics regression was used to assess independent associations, reported as adjusted odds ratios (AOR), with overweight and obesity after adjusting for other independent variables included in the model. Multicollinearity was checked for some original variables including women's age and the number of children ever born, education, and wealth index.

Results

Characteristics of the study samples

The mean age of women was 30 years old (SD = 9.9 years) in which age group 20-39 years old accounted for 60.4%. Almost 66.0% were married; 48.4% had completed primary education and 13.6% did not receive formal schooling. A total of 34.2% of women did agricultural work, 21.4% were unemployed. Of the total sample, 36.4% women were from the poor households. The majority (80.7%) of women resided in rural areas. More than one third had two to three children while 31% had no children. Only 2.2% of women reported cigarette smoking, 7.2% reported using hormonal contraceptives. Mean women's BMI was 22 kg/m^2 (SD = 3.6 kg/m^2) and 15.2% and 2.8% were overweight and obese respectively (**Table 1**).

183 **Table 1**. Socio-demographic and behaviors characteristics of the weighted samples of women
184 aged 15-49 years old in Cambodia, 2014 (n □=□10,765, weighted count)

Variables	Freq.	%
Women's mean age in years (SD)		30.0 (9.9)
15-19	1741	16.2
20-29	3553	**33.0**
30-39	2946	27.4
40-49	2526	23.5
Marital status		
Not married	2846	26.4
Married or living together	7096	**65.9**
Widowed/divorced/separated	823	7.6
Education		
No education	1459	**13.6**
Primary	5214	48.4
Secondary and higher	4092	38.0
Current work status		
Not working	2307	21.4
Agricultural	3681	**34.2**
Professional	2585	24.0
Manual labor and unskilled	2190	20.3
Wealth index		
Poor	3915	**36.4**
Middle	2064	19.2
Rich	4786	44.5
Place of residence		
Rural	8687	**80.7**
Urban	2078	19.3
Number of children born		
0	3321	30.9
1	1721	16.0
2-3	3553	33.0
4+	2170	**20.2**
Smoking status		
Non smoker	10530	97.8
Smoker	235	**2.2**
Ever report of contraceptive use		
No method	6399	**59.4**
Traditional method	2223	20.6
Non-hormonal method	1370	12.7
Hormonal method	774	7.2
BMI mean in kg/m^2 (SD)		22.0 (3.6)
Underweight	1501	13.9
Normal weight	7327	68.1
Overweight	1637	**15.2**
Obese	300	**2.8**

Factors associated with overweight and obesity in bivariate analysis

Socio-demographic characteristics were significantly associated with overweight and obesity (**Table 2**). Women had higher prevalence of overweight/obesity if they were aged 40-49 years (31%) compared to younger age groups, (p value <0.001), married (22.9%) compared to other relationship status (p value <0.001), had no education (22.4%) compared to higher education (p value <0.001), were employed in a professional job (28.4%) compared to other jobs, or were from the rich households (21.4%) compared to poor households (p value <0.001). In addition, women who lived in urban areas had a higher prevalence of overweight and obesity than those living in rural areas (22.4% vs 16.9%, p value <0.001). Overweight/obesity increased with parity. Women reported at least four children had a significantly highest overweight and obesity (29.3%), (p value < 0.001). Likewise, women used non-hormonal contraceptives methods had a significant higher proportion of overweight and obesity (27.7%) than women using other methods (p value <0.019).

Factors associated with overweight and obesity in adjusted logistic regression

As shown in **Table 3**, several factors were independently associated with increased odds of having overweight and obesity among women. These factors included age group 20-29 years [AOR=2.4; 95% CI: 1.6-3.6], 30-39 years [AOR=4.6; 95% CI: 3.0-6.9] and 40-49 years [AOR=6.6; 95% CI: 4.3-10.1] married [AOR=1.8; 95% CI: 1.3-2.7], women who had professional job [AOR=1.3; 95% CI: 1.0-1.5], living in urban residence [AOR=1.3; 95% CI: 1.1-1.5], middle wealth quintile [AOR=1.4; 95% CI :1.1-1.8], and rich wealth quintile [AOR=1.8 ; 95% CI: 1.5-2.1], having at least two-three children or more. On the contrary, following factors were independently associated with decreased odds of having overweight and obese: women with higher education [AOR=0.7; 95% CI: 0.6-0.8], working in agricultural jobs [AOR=0.7; 95% CI: 0.5-0.8], and in manual labor jobs [AOR=0.7; 95% CI: 0.6-0.9].

 Table 2. Factors associated with overweight and obesity among women aged 15-49 years in Chi^2

 analysis (n□=□10,765, weighted count)

Characteristics	Overweight/Obese		Normal/Underweight		p-value
	Freq.	%	Freq.	%	
Women's age group in years					
15-19	51	2.9	1690	97.0	
20-29	390	11.0	3162	89.0	**<0.001**
30-39	705	24.0	2241	76.0	
40-49	790	**31.0**	1735	69.0	
Marital status					
Not married	159	5.6	2688	94.4	
Married or living together	1628	**22.9**	5468	77.1	**<0.001**
Widowed/divorced/separated	151	18.3	672	81.7	
Education					
No education	327	**22.4**	1132	77.6	
Primary	1051	20.2	4164	79.8	**<0.001**
Secondary and higher	559	13.7	3532	86.3	
Current work status					
Not working	381	16.5	1926	83.5	
Agricultural	596	16.2	3085	83.8	
Professional	675	**26.1**	1910	73.9	**<0.001**
Manual labor and unskilled	285	13.0	1905	87.0	
Wealth index					
Poor	543	13.9	3372	86.1	
Middle	368	17.8	1696	82.2	**<0.001**
Rich	1026	**21.4**	3760	78.6	
Place of residence					
Rural	1470	16.9	7217	83.1	
Urban	466	**22.4**	1612	77.6	**<0.001**
Number of children ever born					
0	215	6.5	3106	93.5	
1	238	13.8	1483	86.2	**<0.001**
2-3	847	23.8	2706	76.2	
4+	636	**29.3**	1533	70.7	
Smoking status					
Non smoker	1907	18.1	8623	81.9	0.108
Smoker	30	12.7	205	87.3	
Ever report of contraceptive use					

	964	15.1	5435	84.9	
No method	964	15.1	5435	84.9	
Traditional method	294	21.5	1076	78.6	
Non-hormonal method	214	**27.7**	559.2	72.3	**<0.001**
Hormonal method	465	20.9	1758	79.1	

Table 3. Risk factors associated with overweight and obesity in unadjusted and adjusted logistic regression analysis

Characteristics	Total (N=10,765)			Total (N=10,761)		
	OR	**95% CI**	**P value**	**AOR**	**95% CI**	**P value**
Women's age group in years						
15-19	1.0	Ref.		1.0	Ref.	
20-29	4.1	2.8-5.8	<0.001	**2.4**	**1.6-3.6**	**<0.001**
30-39	10.4	7.3-14.6	<0.001	**4.6**	**3.0-6.9**	**<0.001**
40-49	15.0	10.6-21.2	<0.001	**6.6**	**4.3-10.1**	**<0.001**
Marital status						
Not married	1.0	Ref.		1.0	Ref.	
Married or living together	5.0	4.2-6.1	<0.001	**1.8**	**1.3-2.7**	**0.002**
Widowed/divorced/separated	3.8	2.9-5.0	<0.001	1.1	0.7-1.7	0.638
Education level						
No education	1.0	Ref.		1.0	Ref.	
Primary education	0.9	0.7-1.0	0.150	0.9	0.8-1.1	0.495
Secondary and higher	0.6	0.5-0.7	<0.001	**0.7**	**0.6-0.8**	**0.001**
Current work status						
Not working	1.0	Ref.		1.0	Ref.	
Professional	1.8	1.5-2.2	<0.001	**1.3**	**1.0-1.5**	**0.025**
Agricultural	1.0	0.8-1.2	0.827	**0.7**	**0.5-0.8**	**0.001**
Manual labor and unskilled	0.8	0.6-1.0	0.025	**0.7**	**0.6-0.9**	**0.011**
Wealth index						
Poor	1.0	Ref		1.0		
Middle	1.4	1.1-1.7	0.009	**1.4**	**1.1-1.8**	**0.003**
Rich	1.7	1.5-2.0	<0.001	**1.8**	**1.5-2.1**	**<0.001**
Place of residence						
Rural	1.0	Ref.		1.0	Ref.	
Urban	1.4	1.2-1.6	<0.001	**1.3**	**1.1-1.5**	**0.006**
Number of children ever born						
0	1.0	Ref.		1.0	Ref.	

	OR	95% CI	p	AOR	95% CI	p
1	2.3	1.9-2.9	<0.001	1.1	0.8-1.7	0.505
2-3	4.5	3.8-5.4	<0.001	**1.5**	**1.1-2.2**	**0.026**
4+	6.0	4.9-7.3	<0.001	**1.7**	**1.1-2.5**	**0.008**
Smoking status						
Non smoker	1.0	Ref		-	-	
Smoker	0.7	0.4-1.1	0.111	-	-	
Ever report of contraceptive						
No method	1.0	Ref		1.0	Ref	
Traditional method	1.5	1.3-1.9	<0.001	0.8	0.6-1.1	0.052
Non-hormonal method	2.2	1.7-2.7	<0.001	1.0	0.8-1.3	0.769
Hormonal method	1.5	1.3-1.7	<0.001	1.0	0.8-1.1	0.554

223 *OR: Unadjusted odds ratios, **AOR**: Adjusted odds ratios, **95% CI**: 95% confidence interval*

Discussion

Factors associated with being overweight or obese for non-pregnant women include older age,

increased parity, marriage, urban residence, higher wealth, and working in a professional

occupation. Women with higher education as well as those working in agricultural or manual

labor were less likely to be overweight or obese.

Older aged women were more likely overweight or obese as compared to younger women aged

15-19 years. This is consistent with other studies that showed obesity was more prevalent in

older WRA [20]. The risk of women becoming overweight or obese rises with age, possibly due

to unhealthy food consumption and/or a lack of physical activity [21]. Women with higher

education are less likely to be overweight or obesity than women with no school due to increased

knowledge and awareness; for example, studies in Nigeria and South Korea reported that

educated women had a lower risk of being overweight or obese and that this may be linked to

education influencing healthy behavior [22-25]. Education is a critical predictor of women's

healthy behaviors and health outcomes, including diet, physical activity [26]. Women with better

socioeconomic status and who live in urban areas had higher odds of being overweight and

obese, which is consistent with prior research [27, 28]. Women living in urban areas with high

income may choose to use cars and other fuel-based vehicles; this convenience results in reduced
physical activity and may also increase consumption of fast food. Women with higher
socioeconomic status tend to use more improved technologies for a more comfortable lifestyle
[29, 30]. Women have at least two and more children at risk of being overweight and obese,
similar other studies [31]. During pregnancy, factors such as stress, depression, and/or anxiety
may play a role in hypothalamic-pituitary-adrenal hyperactivity [32, 33]. Women with several
children may also have gained weight as a result of their reduced physical activity and have less
time to focus on health behaviors including weight management.

This study has limitations. It was a cross-sectional study that does not indicate causality.
Additionally, we used to exit secondary data that may result in omitted variable bias as other
factors that may be relevant to overweight and obesity such as level of physical activity, dietary
intake such as alcohol, sweet food, or soft drink consumption, family history, and awareness
about overweight and obesity were not included in the analysis due to data limitations. Finally,
these data were gathered in 2014 and may no longer reflect the present conditions of non-
pregnant women in Cambodia. A subsequent study should further evaluate these relationships
and the general prevalence of overweight and obesity among non-pregnant Cambodian women
once additional data becomes available (the next round of data gathering of Cambodia DHS data
is currently underway, but has experienced delays due to COVID-19). However, this study has
several strengths. First, it draws upon a large, nationally representative sample with a high
response rate of 97%. Data were collected using validated survey methods including calibrated
measurement tools and highly trained data collectors, which contribute to improved data quality
[15]. Formally incorporating the complex survey design and sampling weights into the analysis
bolsters the rigor of the analysis and enables generalizing our findings to the population of non-
pregnant women in Cambodia.

Our results point to socio-demographic characteristics including older age, urban residence,
higher education levels, higher income and wealth, and behavioral factors such as type of
employment as risk factors associated with overweight and obesity among women of
reproductive age in Cambodia. It is crucial to design intervention programs that target these
socio-demographic factors and to raise awareness on the importance of consuming healthy food

as well as the benefits of regular physical activity, especially among older women and those with

professional   jobs.

Acknowledgments

The authors would like to thank DHS-ICF, who approved the data used for this paper.

Authors' Contributions

Um S. contributed to conceptualization, methods, data analysis, writing the original draft, and reviewing/editing the manuscript; **An Y.** contributed to writing and reviewing/editing the manuscript; **Sopheab H.** contributed to reviewing/editing the manuscript; **Jonathan M**. contributed to reviewing/editing the manuscript. All authors read and approved the final manuscript.

Data Availability

Our study used the 2014 Cambodia Demographic and Health Survey (CDHS) datasets. The DHS data are publicly available from the website: https://www.dhsprogram.com/data/dataset_admin

Funding

The authors received no funding support for this study.

Competing interests

The authors have declared that no competing interests exist.

Abbreviations

BMI, Body mass index; NCDs, Non-communicable diseases; WHO, World Health Organization; CDHS, Cambodia Demographic Health Survey; WRA, Women reproductive age; AOR, Adjusted odds ratio; PPS, Probability proportional to size. EA, Enumeration areas.

References

1. Leddy MA, Power ML, Schulkin J: **The impact of maternal obesity on maternal and fetal health**. *Reviews in obstetrics and gynecology* 2008, **1**(4):170.

2. WHO: **World Health Organization, Noncommunicable diseases: Mortality; [https://www.who.int/data/gho/data/themes/topics/topic-details/GHO/ncd-mortality]: Accessed on July 12, 2022.**

3. WHO: **World Health Organization, Obesity and overweight; [https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight]: Accessed on July 12, 2022.**

4. Report GN: **Country Nutrition Profiles: [https://globalnutritionreport.org/resources/nutrition-profiles/asia/south-eastern-asia/cambodia/]; Accessed on July 12, 2022. 2022**.

5. NIS: **National Institute of Statistics. Directorate General for HC. International ICF . Cambodia Demographic and Health Survey 2014. Phnom Penh, Cambodia: National Institute of Statistics/Cambodia, Directorate General for Health/Cambodia, and ICF International; 2015.** 2014.

6. **World Food Programme in Cambodia; [https://www.wfp.org/countries/cambodia]: Accessed on July 13, 2022.**

7. Abrha S, Shiferaw S, Ahmed KY: **Overweight and obesity and its socio-demographic correlates among urban Ethiopian women: evidence from the 2011 EDHS**. *BMC Public Health* 2016, **16**(1):636.

8. Balarajan Y, Villamor E: **Nationally representative surveys show recent increases in the prevalence of overweight and obesity among women of reproductive age in Bangladesh, Nepal, and India**. *The journal of nutrition* 2009, **139**(11):2139-2144.

9. Oddo VM, Bleich SN, Pollack KM, Surkan PJ, Mueller NT, Jones-Smith JC: **The weight of work: the association between maternal employment and overweight in low-and middle-income countries**. *International Journal of Behavioral Nutrition and Physical Activity* 2017, **14**(1):1-10.

10. Lopez LM, Bernholc A, Chen M, Grey TW, Otterness C, Westhoff C, Edelman A, Helmerhorst FM: **Hormonal contraceptives for contraception in overweight or obese women**. *The Cochrane database of systematic reviews* 2016, **2016**(8):Cd008452.

11. de Munter JS, Tynelius P, Magnusson C, Rasmussen F: **Longitudinal analysis of lifestyle habits in relation to body mass index, onset of overweight and obesity: results from a large population-based cohort in Sweden**. *Scandinavian journal of public health* 2015, **43**(3):236-245.

12. Piirtola M, Jelenkovic A, Latvala A, Sund R, Honda C, Inui F, Watanabe M, Tomizawa R, Iwatani Y, Ordonana JR: **Association of current and former smoking with body mass index: A study of smoking discordant twin pairs from 21 twin cohorts**. *PloS one* 2018, **13**(7):e0200140.

13. Chiolero A, Jacot Sadowski I, Faeh D, Paccaud F, Cornuz J: **Association of cigarettes smoked daily with obesity in a general adult population**. *Obesity* 2007, **15**(5):1311-1318.

14. **DHS Program. The Demographic and Health Surveys (DHS) Program; ICF: Rockville,**
MD, USA, 2020; Available online: https://dhsprogram.com/data/available-datasets.cfm
(accessed on 28 March 2022).

15. BOWLER P: **CUNNINGHAM, A. and JARDINE, N.(editors). Romanticism and the**
sciences. Cambridge University Press, Cambridge: 1990. Pp xxii, 345; illustrated. Price:£
15, US $19.95. ISBN: 0-521-35685-7 (paperback). In.; 1992.

16. Organization WH: **Physical status: The use of and interpretation of anthropometry, Report**
of a WHO Expert Committee: World Health Organization; 1995.

17. Um S, Chan YZC, Tol B, Sopheab H: **Determinants of exclusive breastfeeding of infants**
under six months among Cambodian mothers. *Journal of pregnancy* 2020, **2020**.

18. Bursac Z, Gauss CH, Williams DK, Hosmer DW: **Purposeful selection of variables in logistic**
regression. *Source code for biology and medicine* 2008, **3**(1):1-8.

19. Um S, Sopheab H, Yom A, Muir JA: **Trends and factors associated with anemia among**
pregnant women aged 15-49 years old in Cambodia: Data analysis of the Cambodia
Demographic and Health Surveys. *medRxiv* 2022:2022.2009.2008.22279710.

20. Biswas T, Uddin MJ, Mamun AA, Pervin S, P Garnett S: **Increasing prevalence of overweight**
and obesity in Bangladeshi women of reproductive age: Findings from 2004 to 2014. *PloS*
one 2017, **12**(7):e0181080.

21. Alemu E, Atnafu A, Yitayal M, Yimam K: **Prevalence of overweight and/or obesity and**
associated factors among high school adolescents in Arada Sub city, Addis Ababa, Ethiopia.
Journal of nutrition & food sciences 2014, **4**(2):1.

22. Hamad R, Elser H, Tran DC, Rehkopf DH, Goodman SN: **How and why studies disagree about**
the effects of education on health: A systematic review and meta-analysis of studies of
compulsory schooling laws. *Social Science & Medicine* 2018, **212**:168-178.

23. Chung A, Backholer K, Wong E, Palermo C, Keating C, Peeters A: **Trends in child and**
adolescent obesity prevalence in economically advanced countries according to
socioeconomic position: a systematic review. *Obesity reviews* 2016, **17**(3):276-295.

24. Barlow P: **The effect of schooling on women's overweight and obesity: A natural experiment**
in Nigeria. *Demography* 2021, **58**(2):685-710.

25. McLaren L: **Socioeconomic status and obesity**. *Epidemiologic reviews* 2007, **29**(1):29-48.

26. Liu H, Guo G: **Lifetime socioeconomic status, historical context, and genetic inheritance in**
shaping body mass in middle and late adulthood. *American sociological review* 2015,
80(4):705-737.

27. Mangemba NT, San Sebastian M: **Societal risk factors for overweight and obesity in women**
in Zimbabwe: a cross-sectional study. *BMC public health* 2020, **20**(1):1-8.

28. Kamal SM, Hassan CH, Alam GM: **Dual burden of underweight and overweight among**
women in Bangladesh: patterns, prevalence, and sociodemographic correlates. *Journal of*
health, population, and nutrition 2015, **33**(1):92.

29. Ghose B: **Frequency of TV viewing and prevalence of overweight and obesity among adult**
women in Bangladesh: a cross-sectional study. *BMJ open* 2017, **7**(1):e014399.

30. Islam M, Sathi NJ, Abdullah HM, Naime J, Butt ZA: **Factors Affecting the Utilization of**
Antenatal Care Services During Pregnancy in Bangladesh and 28 Other Low-and Middle-
income Countries: A Meta-analysis of Demographic and Health Survey Data. *Dr Sulaiman*
Al Habib Medical Journal 2022, **4**(1):19-31.

31.   Mbochi RW, Kuria E, Kimiywe J, Ochola S, Steyn NP: **Predictors of overweight and obesity in**
389   **adult women in Nairobi Province, Kenya**. *BMC public health* 2012, **12**(1):1-9.
390   32.   Li W, Wang Y, Shen L, Song L, Li H, Liu B, Yuan J, Wang Y: **Association between parity and**
391   **obesity patterns in a middle-aged and older Chinese population: a cross-sectional analysis**
392   **in the Tongji-Dongfeng cohort study**. *Nutrition & metabolism* 2016, **13**(1):1-8.
393   33.   Pasquali R, Vicennati V, Cacciari M, Pagotto U: **The hypothalamic□pituitary□adrenal axis**
394   **activity in obesity and the metabolic syndrome**. *Annals of the New York Academy of Sciences*
395   2006, **1083**(1):111-128.