Circulating proteome of hospitalized patients uncovers six endophenotypes of COVID-19 and points to FGFR and SHC4-signaling in acute respiratory distress syndrome

William Ma[1,*], Antoine Soulé[1,*], Katelyn Yixiu Liu[2], Catherine Allard[3], Karine Tremblay[4, †], Simon Rousseau[2, †], and Amin Emad[1, 5, †]

[*] These authors contributed equally.

[1] Department of Electrical and Computer Engineering, McGill University, Montréal, QC, Canada

[2] The Meakins-Christie Laboratories at the Research Institute of the McGill University Heath Centre Research Institute, & Department of Medicine, Faculty of Medicine, McGill University, Montréal, QC, Canada

[3] Statistical department, Centre de recherche du Centre hospitalier universitaire de Sherbrooke (CRCHUS), Sherbrooke, Canada.

[4] Pharmacology-physiology Department, Faculty of Medicine and Health Sciences, Université de Sherbrooke, Saguenay, QC, Canada; Centre intégré universitaire de santé et de services sociaux du Saguenay-Lac-Saint-Jean, Saguenay, QC, Canada; CRCHUS, Sherbrooke, Canada.

[5] Mila, Quebec AI Institute, Montréal, QC, Canada

† Corresponding Authors:
Amin Emad,
755 McConnell Engineering Building, 3480 University Street, Montreal H3A 0E9, Canada
Email: amin.emad@mcgill.ca

Simon Rousseau,
RI-MUHC, E M3.2244, 1001 Décarie, Montréal H4A 3J1, Canada,
Email: simon.rousseau@mcgill.ca

Karine Tremblay,
Pavillon des Augustines, local AUG-5-01A, 225 St-Vallier street, Chicoutimi G7H 7P2, Canada
Email: karine.tremblay@usherbrooke.ca

Abstract

Defining the molecular mechanisms of novel diseases such as COVID-19 is crucial to identify

treatable traits to improve patient care. To circumvent *a priori* bias and the lack of in-depth

knowledge of this new disease, we opted for an unsupervised stratification approach, followed

by detailed multi-modal characterization using proteome, metabolomic, genomic, and clinical

features. Using the detailed circulating proteome, as measured by 4985 aptamers (SOMAmers),

robust consensus clustering identified six endophenotypes (EPs) present among 731 SARS-CoV-2

PCR-positive hospitalized participants to *Biobanque québécoise de la COVID-19 (BQC19)*, with

varying degrees of disease severity and times to intensive care unit (ICU) admission. In particular,

one endophenotype, EP6, was associated with a greater proportion of ICU admission, mechanical

ventilation, acute respiratory distress syndrome (ARDS) and death. Clinical features of this

endophenotype showed increased levels of C-reactive protein, D-dimers, elevated neutrophils,

and depleted lymphocytes. Moreover, metabolomic analysis supported a role for immuno-

thrombosis in severe COVID-19 ARDS. Furthermore, Fibroblast Growth Factor Receptor (FGFR)

and SH2-containing transforming protein 4 (SHC4) signaling were identified as molecular features

associated with severe COVID-19. Importantly, a predictive model was developed and validated

on an additional set of 631 SARS-CoV-2 PCR-positive patients to enable prediction of these

endophenotypes, which reflect patients' likelihood of admission to ICU, solely based on clinical

laboratory measurements. This suggests the use of blood markers as surrogates for generalizing

these EPs to new patients and automating identification of high-risk groups in the clinic.

Introduction

The coronavirus disease 2019 (COVID-19) is a new human disease caused by the coronavirus SARS-CoV-2 infection that has been assessed pandemic by the World Health Organization in March 2020. As SARS-CoV-2 infection spread, a breath of outcomes of the infection became apparent, from asymptomatic individuals to severely ill and dying, from complete recovery to long-lasting symptoms (Marshall et al., 2020; Rubin, 2020). The emergence of novel diseases such as COVID-19 presents the medical and scientific community with numerous challenges. Among them, defining the molecular mechanisms of disease related to specific outcomes is important to identify treatable traits and improve the performance of healthcare systems facing the challenges brought by the pandemic.

Successfully reaching this precision medicine goal requires a more granular definition of the pathology. A symptom-based method to discover molecular mechanisms of the disease may result in a challenge emerging from the fact that the same higher-level phenomenon, such as COVID-19 severity, can be produced by several different molecular mechanisms, a phenomenon termed the «many-one» limitation (Hull, 1974). Recent advances in computing strategies, such as machine learning, has enabled the development of methods that help overcome this limitation by starting from molecular profiles instead of symptoms to define endophenotypes, *i.e.* subgroups of individuals who are inapparent to traditional methods but share a common set of molecular factors that can lead to treatable traits (Berrettini, 2005). Establishing successful treatment strategies requires a tailored approach to the underlying molecular mechanisms that can help predict and alter disease trajectories (Russell and Baillie, 2017). Endophenotypes can

become apparent using extensive molecular phenotyping combined with machine learning

algorithms (Blatti III et al., 2020; Emad et al., 2020; Te Pas et al., 2017). Current investigations of

endophenotypes in COVID-19 have mainly relied on supervised approaches using fixed outcomes

(such as disease severity) and integrating clinical variables at the onset (Al-Hadrawi et al., 2022).

We hypothesize that using an unsupervised approach exploiting a rich molecular dataset can

provide novel mechanistic insights into the pathobiology of severe COVID-19 that can help

physicians improve diagnosis and prognosis.

Therefore, this study aims to identify endophenotypes linked to diverse clinical trajectories of

COVID-19 using the extensive molecular phenotyping of a cohort of 731 SARS-CoV-2 positive

hospitalized patients from the *Biobanque québécoise de la COVID-19* (BQC19,

www.quebeccovidbiobank.ca) (Tremblay et al., 2021), a prospective observational cohort of

SARS-CoV-2 positive and negative participants recruited in the province of Québec, Canada, to

improve our understanding of COVID-19 pathobiology and our capacity to alter disease

outcomes.

In this manuscript, we report the identification of six endophenotypes in hospitalized SARS-CoV-

2 positive participants to BQC19, associated with different clinical trajectories. The molecular

information underpinning these endophenotypes was used to increase our understanding of

pathobiology and predict the likelihood of patients admitted to the hospital to be admitted to

the ICU using clinical blood workups.

Results

Unsupervised clustering of SARS-CoV-2-positive hospitalized BQC19 participants reveal endophenotypes associated with varying disease severity

In this study, we aimed to identify endophenotypes of COVID-19 based on the circulating proteome of patients in our cohort of SARS-CoV-2 positive hospitalized participants to BQC19 (n = 1,362, Table 1), using an unsupervised approach. Figure S1 shows the distribution of the time of hospital admission of the patients and the corresponding waves as defined by National Institute of Public Health of Quebec (https://www.inspq.qc.ca/covid-19). For this purpose, we performed consensus agglomerative clustering of the subset of patients (n = 731, Table S1) for whom data corresponding to circulating proteome measured by a multiplex SOMAmer affinity array (Somalogic, ~5,000 aptamers) (Gold et al., 2010) was available in BQC19. The remaining samples were kept aside for follow-up analysis. First, the optimal number of clusters (k = 6) was identified using two criteria, Akaike's Information Criteria (AIC) and Bayesian Information Criteria (BIC) (Figure 1A); then, consensus agglomerative clustering (Euclidean distance and Ward linkage) (Murtagh and Legendre, 2014; Ward Jr, 1963) using bootstrap subsampling was performed to obtain six robust clusters (see Methods for details) (Figure 1, Figures S2 and S3).

The clinical and pathological characteristics of patients in each endophenotype is provided in Table S1. To characterize the identified endophenotypes (EPs) with respect to disease severity, we performed two-sided Fisher's exact test to assess their enrichment (or depletion) in "severe" or "dead" outcome. EP6 was significantly enriched in the severe/dead outcomes (Benjamini-Hochberg false discovery rate (FDR) = 1.72E-18) with these outcomes observed in 66.1% of EP6

patients. Meanwhile, EP1 was significantly depleted in severe/dead outcomes (FDR = 8.23E-11)

(Figure 2A, Table S1) with these outcomes only observed in 11.6% of EP1 patients. In addition,

EP6 was enriched in participants 1) receiving oxygen therapy (FDR = 6.24E-12), 2) receiving

ventilatory support (FDR = 6.63E-12), and 3) being admitted to intensive care unit (ICU) (FDR=

1.26E-22) (Figure 2A, Table S1). Kaplan Meier analysis (Kaplan and Meier, 1958) also confirmed

that the identified EPs have a distinct temporal pattern of admission to ICU (multivariate logrank

test P = 5.06E-36), with EP1 (EP6) having the highest (lowest) chance of not being admitted to

ICU (or die prior to that) in a 40-day span since their admission to the hospital (Figure 2B). A

similar pattern can be observed when patients that died before admission to ICU were excluded

(Figure S4, multivariate logrank test P = 6.55E-36). A two-sided Mann-Whitney U (MWU) test

showed that patients in EP5 were generally older than other EPs (FDR = 6.94E-5), while EP3

included younger patients (FDR = 1.82E-4). However, EP6 (which had the most severe patients)

did not show enrichment in older patients or individuals with high BMI (two-sided MWU

FDR>0.05) (Figure 2C, Table S1).

These analyses revealed that the unsupervised approach was able to identify endophenotypes

with distinct disease characteristics and outcomes using the circulating proteome of the patients.

We identified EP6 as a group of participants with an increase in many measures of COVID-19

disease severity.

EP6 is enriched with BQC19's participants having acute respiratory distress syndromes

In accordance with increase disease severity, EP6 was also enriched in COVID-19 medical

complications (two-sided Fisher's exact test): acute respiratory distress syndrome (ARDS) (FDR =

1.43E-10), acute kidney injury (FDR = 6.37E-7), bacterial pneumonia (FDR=2.13E-6), liver

dysfunction (FDR = 9.32E-3), and hyperglycemia (FDR = 1.56E-2) (Figure 3, Table S2). The

frequency of ARDS was just below 8% in EP1 compared to greater than 44% in EP6, making this

complication a key feature of this cluster (Figure 3, Table S2).

EP6 is enriched in blood metabolites associated with severe COVID-19.

To further characterize each EP and gain insight into mechanisms of disease, metabolomic

profiling of plasma samples was done in parallel to the SOMAmer analysis. The results yielded

data on 1,435 metabolites, of which 576 were found significantly altered in EP6 (two-sided MWU

FDR < 0.01). Moreover, the metabolomic characterization of the plasma samples supported the

distinction in blood composition at the levels of metabolomic sub-pathways and individual

metabolites between the different EPs (Figure 4 and S3).

Pathway enrichment analysis identifies FGFR-signaling in severe COVID-19 acute respiratory

distress syndrome

To gain insight into the molecular mechanisms underlying the severity in EP6, we performed

pathway enrichment analysis using the Knowledge Engine for Genomics (KnowEnG) (Blatti III *et*

al., 2020) for the aptamers associated with EP6. Since aptamers were used to identify the EPs, it

is expected that many of them would be significantly associated with the EPs. As a result, we

selected a very strict threshold of FDR < 1E-20 (two-sided MWU test) to identify top aptamers

associated with EP6 (Table S4). We then used the gene set characterization pipeline of KnowEnG

with Reactome (Gillespie et al., 2021) pathway collection. This analysis showed that while EP6 is

characterized by pathways associated with Interleukins and Cytokine Signaling in Immune

system, multiple instances linked it with Fibroblast Growth Factor Receptor (FGFR) signaling,

identifying this pathway as a potential driver of severe pathology that was not present in other

EPs (Tables 2 and S4).

SHC4 genotype and protein expression levels are associated with higher odds of belonging to

EP6

To further improve our understanding of the molecular mechanisms underlying EP6, we

leveraged an additional dataset of Genome Wide association Study (GWAS) corresponding to

these patients. We identified 25 single nucleotide variations (SNVs) distributed in 13 annotated

genes, that were below a p-value threshold of 1E-4 differentiating EP6 versus the rest (Table 3).

We then investigated each of the SNVs to which we could assign a gene and an aptamer, to assess

whether their protein product in circulation was differentially regulated by the genotype (Table

4). We discovered two genes, *SHC4* (encoding SHC adaptor protein 4) and *CACNA2D3* (encoding

calcium voltage-gated channel auxiliary subunit alpha2 delta3) for which there was a significant

association between genotype and protein expression levels (p-values < 0.05). While *CACNA2D3*

may have mild impact on EP6 membership (odd ratio = 0.61, Table 4), SHC4 was one of the top-

enriched aptamers (position 32 out of 4,985), with odds ratio of 11.98 and 2.00 for the protein

product and SNV, respectively of belonging to EP6 for the alternative allele. Therefore, the GWAS

analysis revealed that the signaling adaptor protein SHC4 may play an important mechanistic role

in contributing to severe disease pathology. To gain further insight into the potential role of SHC4,

we investigated the metabolites associated with SHC4 in EP6. The only significantly correlated

metabolite was 6-oxopiperidine-2-carboxylate (Spearman's rank correlation = 0.40, FDR = 0.012),

where the p-values were corrected for the number of metabolites using Benjamini-Hochberg

FDR.

A predictive model based on blood markers predicts EPs in a separate validation cohort

To further characterize each EP, we assessed the clinical laboratory results obtained from blood

draws and compared them between the groups. We focused on 21 markers that were measured

in at least 50% of the patients used for consensus clustering (Figure 5A and Table S5) and used

the summary value reported in the BQC19 database corresponding to the most extreme

measurement among multiple blood draws (Table S5 provides this information for each blood

marker). Figure 5A shows the elevation and depletion of these markers in the identified EPs. EP6

is characterized by abnormal values in markers of inflammation (lymphopenia, total white blood

cells count, neutrophilia, C-reactive protein (CRP)), liver damage (alanine aminotransferase (ALT),

albumin, lactate deshydrogenase (LDH)), blood clotting disorder (D-dimers, low hemoglobin,

International Normalized Ratio (INR) and hyperglycemia (glucose).

To identify relationships that may shed light on factors influencing clinical laboratory results

defining EP6, we also performed Spearman's rank correlation analyses between each blood test

values and metabolites (Table S5).

Since EPs and particularly EP6, which we identified as the EP with worst outcome, showed a clear

and distinct clinical laboratory result signature compared to other EPs, we sought to develop a

predictive model based on these signatures. Due to the presence of missing values for these

markers in our cohort, we developed a nearest-centroid classifier that is capable of dealing with

missing values and can predict EPs based on blood markers (see Methods for details). To test the

ability of this model on prediction of EPs on an independent yet similar dataset, we used data

corresponding to 631 SARS-CoV-2 positive hospitalized BQC19's participants that did not have

circulating proteome data and hence were not used to identify the endophenotypes. The clinical

and pathological characteristics of patients in each predicted endophenotype (PEP) is provided

in Figure 5B-5E and Table S6.

Our predictive model identified 116 of these patients to belong to EP6. Fisher's exact test showed

a significant enrichment of the predicted EP6 (PEP6) in severity/dead (FDR = 1.77E-22), while

PEP1 and PEP2 were significantly depleted in these outcomes (FDR = 1.61E-4 and FDR = 1.30E-8,

respectively), as shown in Figure 5B and Table S6. Similar to EP6, PEP6 was also significantly

enriched in participants 1) receiving oxygen therapy (FDR = 2.38E-8), 2) receiving ventilatory

support (FDR = 4.40E-8), and 3) being admitted to ICU (FDR = 1.23E-24) (Table S6). Kaplan Meier

analysis also confirmed that these PEPs have a distinct temporal pattern of admission to ICU

(multivariate logrank test P = 1.05E-34), with PEP6 having the lowest chance of not being

admitted to ICU (or die prior to that) in a 40-day span since their admission to the hospital (Figure

5E).

These results suggest that our predictive model can use these 21 blood markers to generalize the

definition of endophenotypes to patients for whom the proteome data is unavailable.

Discussion

The results presented herein came from the unsupervised clustering of the circulating proteome

from a large cohort of deeply-phenotyped SARS-CoV-2 positive hospitalized participants, that

provided both expected and novel findings into the molecular mechanisms regulating COVID-19.

They led to the identification of six endophenotypes with different clinical trajectories and

distinguishing characteristics summarized in Table 5. One endophenotype, (EP6) was associated

with worst clinical outcome of COVID-19 (enriched in acute respiratory distress syndrome)

reflected by a greater proportion of ICU admission, mechanical ventilation, and severe/death

outcomes (Figure 1). Clinical features of this endophenotype were consistent with published

literature with increased levels of CRP, D-dimers, elevated neutrophils, and depleted

lymphocytes (Figure 5A, Table S5). Our approach enabled the identification of interleukins, FGFR

and SHC4 signaling as cardinal features of the molecular pathways associated with severe COVID-

19. Importantly, this information was sufficient to train an accurate predictive model that could

in the near future support clinical care, since it only requires data from routine clinical laboratory

results for prognosis.

The approach: unsupervised clustering capacity at identifying clinically meaningful

*subpopulation*s

Our unsupervised clustering approach in conjunction with a rich molecular dataset enabled us to

identify endophenotypes that could not be captured using traditional methods classifying the

population in two bins solely based on severity. This is in part because of the «many-one»

limitation: the same higher-level phenomenon (COVID-19 severity) can be produced by several

different molecular mechanisms. Determining endophenotypes using an unsupervised method

provides a higher granularity and increases the chance to identify distinct molecular mechanisms

and pathways resulting in similar COVID-19 severity. Accordingly, we identified two

endophenotypes with more favorable outcomes (EP1 and EP2), three endophenotypes with

intermediate outcomes in terms of severity (EP3, EP4 and EP5) and one endophenotype which

led to worst outcomes compared to all others (EP6).

The identification of endophenotypes was done systematically using robust consensus clustering

of aptamer expression levels in which the optimum number of clusters was determined

congruently using two well-established measures: AIC and BIC. The consensus clustering using

bootstrap sampling (1000 times) ensured identification of robust clusters that are not sensitive

to exclusion of some of the samples (20% randomly selected and excluded at each cycle).

Moreover, identifying the best number of clusters using AIC/BIC (both of which agreed with each

other) allowed us to reveal the patterns of the EPs directly from the data, instead of imposing a

pattern onto it through human supervision. This is an important strength of the study that

enabled us to identify distinct molecular patterns of patients that could have remained

undetected using other traditional approaches.

Moreover, to improve the translational applicability of EPs, we developed a predictive model

based only on laboratory measured blood markers to generalize the definition of these

endophenotypes to unseen samples without measured aptamer expression levels.

Characteristics of EPs predicted solely based on their blood markers were consistent with the

original EPs, suggesting the use of blood markers as surrogates for generalizing these EPs to new

patients and automating identification of high-risk groups in the clinic. This approach takes into

account the effect of multiple blood variables simultaneously and incorporates the full

distribution of each variable. This is in contrast to the clinical laboratory results that are

automatically flagged as within or outside normal range, one variable at a time, therefore

increasing the clinical applicability of our model by leveraging a wider spectrum of information

to prognose patient outcome.

COVID-19 molecular pathology

The datasets used in this study carry rich molecular information on mechanisms of disease for

COVID-19. EP6 is characterized by its enrichment in ARDS (44% vs 8% in EP1, Figure 3), acute

kidney injury (45%) and liver dysfunction (19%) (Table S2). Low levels of Sphingosine 1-phosphate

(S1P) are associated with ARDS and were shown to be associated with greater ICU admission and

decrease survival in COVID-19 (Marfia et al., 2021). Accordingly, we found that S1P levels are

depleted in EP6 while they are enriched in EP1 (Figure 4A). Interestingly, the aptamers detecting

neutral ceramidase, an enzyme converting ceramides into sphingosine, is enriched in EP1

(although changes in protein abundance as detected by aptamers may not necessarily reflect

changes in enzymatic activity). Accordingly, dihydroceramides and ceramides are depleted in

cluster EP1. Conversely, dihydroceramides and ceramides are significantly enriched in EP6,

suggesting that there is a shunting of the pathway away from sphingosine towards more pro-

inflammatory ceramides in EP6 associated with metabolic disorders (Lachkar et al., 2021).

Moreover, one of the top aptamers found enriched in EP6, is the enzyme Serine

palmitoyltransferase 2 (SPTLC2) (Table S4) (Han et al., 2009; Lone et al., 2020). These results

suggest a counterbalance between ceramides and sphingosine, where the former is associated

with poorer outcomes during critical illness, whereas higher levels of the latter is associated with

more favorable outcomes in ARDS.

Metabolomic profile of EP6 supports a role for immuno-thrombosis-mediated organ damage in

COVID-19

To obtain a more comprehensive understanding of the alteration in metabolic profiles in EP6, we

investigated which metabolic sub-pathways were significantly enriched in EP6 (Figure 4, Table

S3). The two top pathways (FDR < 1E-2) are "Methionine, Cysteine, SAM and Taurine

Metabolism" and "phosphatidinylethanolamine (PE)". Interestingly, these two pathways are

known to interact (Blachier et al., 2020), with PE methylation a major consumer of S-

Adenosylmethionine (SAM) leading to the synthesis of S-Adenosylhomocysteine (SAH) and

cystathionine (Ye et al., 2017), which itself has been found upstream of 2-hydroxybutyrate and

2-aminobutyrate in a model of hepatoxicity (Parman et al., 2011). SAH, cysthathionine, 2-

hydroxybutyrate and 2-aminobutyrate are significantly enriched in EP6. Congruently, EP6 is

enriched (FDR < 1E-2) in liver dysfunction (Figure 3, Table S2), with markers of liver dysfunction

all significantly altered in clinical blood works: ALT, albumin, bilirubin and LDH (Figure 5A, Table

S5).

PEs become exposed at the surface of cell membranes upon exposure to stress, inflammation,

and cell death (Ran et al., 2002; Stafford and Thorpe, 2011). In a Syrian hamster model, infection

with SARS-CoV-2 had markedly increased PE expression in the animals that were fed a high salt,

high fat diet, demonstrating the interaction between infection and metabolic disorder with the

abundance of circulating PE (Port et al., 2021). Phospholipids-containing microparticles from

platelet activation contribute to Tissue Factor activation and pro-thrombinase activity (Ataga,

2009). Platelets-derived microparticles have a much greater procoagulant activity than activated

platelets (Sinauridze et al., 2007). Exposure of glycerophospholipids in conjunction with

phosphatidinylserine (PS) enhances factor X activation and increases pro-thrombinase activities

(Majumder et al., 2011; Tavoosi et al., 2011). Interestingly, red blood cells exposed to paclitaxel,

PS were exposed to the surface by protein kinase C (PKC) zeta activation of scramblase (Kim et

al., 2018). The aptamer detecting PKC zeta is one of the top aptamers associated with EP6 (Table

S4). PKC zeta can be activated by testosterone, Dehydroepiandrosterone(DHEA) (Sato et al.,

2008) and dexamethasone (Kajita et al., 2001), signals of relevance to EP6. The membranes of

azurophilic granules, which contains Cathepsin G representing the most enriched aptamer in EP6,

are enriched in PE relative to PC (MacDonald and Sprecher, 1989). Accordingly, EP6 in addition

to significantly enhanced PE abundance, showed decrease platelets count, increased D-Dimers,

INR and activated partial thromboplastin time (aPTT) (Figure 5A, Table S5), hallmarks of

Disseminated Intravascular Coagulation (DIC), a serious and often lethal complication of sepsis

(Papageorgiou et al., 2018). Liver lesions are frequently observed in DIC (Esaki et al., 1984), where

liver damage can cause DIC, or exacerbate its manifestation due to its function in clearing

activated products of the coagulation cascade. Taken together, the metabolomic profile of EP6

supports pro-coagulation activity in circulation that can be linked to organ damage.

Many early reports suggested a role for immunothrombosis involving neutrophil-mediated

release of NETs contributing to endothelial dysfunction as a mechanism of microthrombosis in

COVID-19 associated ARDS (Barnes et al., 2020; Bonaventura et al., 2021; Ding et al., 2021). These

findings have been supported by several studies carried out in humans (Blasco et al., 2021;

Desilles et al., 2022; Englert et al., 2021; Leppkes et al., 2020; Middleton et al., 2020; Obermayer

et al., 2021; Ouwendijk et al., 2021; Petito et al., 2021; Skendros et al., 2020). All of these studies

were performed with 7 to 77 participants. Our study supports these findings in two important

ways: 1) we used a much greater sample size (n = 731) and, 2) the identification of molecular

factors associated with immuno-thrombosis emerged from an *unsupervised* analysis of deep

phenotyping of the participant population. The strength of the extensive characterization

performed in this study has enabled the finer definition of molecular mechanisms of disease by

providing associations between the circulating proteome, metabolome and clinical laboratories

results.

Tryptophan and polyamine metabolisms in COVID-19 acute kidney injury

SARS-CoV-2 infection has been shown to be associated with altered kynurenin levels associated

with increased IL-6 and kidney injury (Thomas et al., 2020). Interestingly, we also observed

Kynurenin to be enriched in EP6 but depleted in EP1, supporting the association of increased

kynurenin with COVID-19 severity and the enrichment in acute kidney injury complication.

Spermidine/spermine N(1)-acetyltransferase (SSAT) contributes to polyamine synthesis. In

addition, its extracellular metabolite, N1,N12-diacetylspermine, is one of the top 15 metabolites

enriched in EP6 (Figure 4A), is positively correlated with Urea and creatinine, and is negatively

correlated with lymphocyte numbers (Table S5). Deletion of SSAT in mice is protective against

LPS-induced kidney injury (Zahedi et al., 2010). SSAT activity is associated with white blood cell

count in Acute Myeloid Leukemia and Chronic Myeloid Leukemia patients (Pirnes-Karhu et al.,

2014). This suggests that the tryptophan and polyamine metabolisms are associated with acute

kidney injury in COVID-19 and identifies potential pathways of disease progression.

Novel molecular markers of COVID-19 pathology: FGFR and SHC4

Two of the outstanding novel molecular factors associated with COVID-19 ARDS identified by our

study are FGFR and SHC4. Circulating levels of the pro-angiogenic FGF-2 has been associated with

COVID-19 severity and creatine levels in a study of 208 SARS-CoV-2 positive participants (Smadja

et al., 2021). It is noteworthy that the use of Nintedanib, an inhibitor of FGFR, Vascular

Endothelial Growth Factor Receptor (VEGFR) and Platelet-Derived Growth Factor Receptor

(PDGF-R) approved for use in interstitial lung disease, improved pulmonary inflammation and

helped wean off mechanical ventilation of three middle-aged obese COVID-19 patients where

lung function restoration has been challenging (Bussolari et al., 2021). While the adaptor protein

SHC4 has not been experimentally demonstrated to modulate FGFR signaling, a 12-gene

biomarker signature associated with melanoma contains FGFR2, FGFR3 and SHC4 (Liu et al.,

2013). It is attractive to speculate that SHC4 may act downstream of FGFR or other associated

growth factor receptors, favoring immuno-thrombosis associated with COVID-19 ARDS. In view

of the limited knowledge of this understudied member of the SHC family, we looked at the

metabolites correlated with SHC4 to gain insights into its possible functions. Interestingly, 6-

oxopiperidine-2-carboxylate (the only significantly correlated metabolite to SHC4 in EP6) was

found to be negatively associated with glomerular filtration rate in a GWAS study of kidney

disease and hypertension in African-Americans (Luo et al., 2021). Acute kidney injury is a frequent

complication of acute liver failure (Agarwal et al., 2013) and liver dysfunction is associated with

abnormal coagulation (Williams et al., 1993). The overall molecular information coming from the

multi-modal analysis of EP6 points to a triad between liver function, kidney function and

hemostasis that becomes dysfunctional following ARDS-associated immuno-thrombosis.

Whether the presence of SHC4 in circulation is a marker of dysfunction of this triad, or an active

actor immuno-thrombosis remains to be determined. Moreover, the identity of the cells

expressing SHC4, leading to its presence in the circulation is not known. Taken together, our

identification of FGFR and SHC4 signaling pathways distinguishing EP6 from other

endophenotypes, supports further investigation of antagonists of those pathways to treat severe

manifestations of COVID-19 and their potential use as biomarkers of severe disease activity.

Limitations and considerations

The data presented in this study come from individuals participating to BQC19, a prospective

observational cohort built to study COVID-19 in Québec (Canada) with its specific population

profile as reported previously (Tremblay *et al.*, 2021). While the number of participants was

sufficient to establish the endophenotypes using the extensive proteomic profile available in

BQC19, it was insufficient for traditional genome-wide association study (GWAS) to identify

relations between SNVs and the identified endophenotypes (Zhou et al., 2021). Instead, we

exploited the top SNVs that distinguished EP6 from other EPs in a pQTL analysis. Because these

results show a potential genetic functional causality, it gives us the confidence that these

associations are likely not due to random chance; however, the robustness of this approach

needs to be further tested in other studies. A chronological bias may also be present, as most of

the participants used for endophenotyping in this study were recruited during the first two waves

of the pandemic (Figure S1), prior to widespread vaccination in Québec and prior to the

appearance of the Omicron variant and sub-variants. Therefore, some of the features of the

identified endophenotypes may change over the course of the pandemic. It will be essential to

continue to assess the molecular profiles longitudinally to better understand the dynamic nature

of host-pathogen interactions. It will also be interesting to compare the profiles of COVID-19

ARDS to other viral-induced ARDS, to identify commonalities as well as distinguishing features.

In this study, we used circulating proteome as determined by aptamers to identify

endophenotypes, a task for which it is well suited as it can capture a dynamic landscape.

However, several important considerations need to be mentioned. First, raw (unnormalized)

expression values of the same aptamer can be used to compare different samples, but these

values cannot be used to compare different aptamers against each other in the same sample,

since they only show the relative abundance of expression and not absolute expression. As such,

one needs to first normalize these values across samples (one aptamer at a time) and then

subject them to follow-up analysis such as clustering, an approach that we adopted in this study.

Second, since information only shows relative abundance, focusing on an individual aptamer and

analyzing it will require additional measurements to establish absolute abundance. Moreover, if

one wants to analyze aptamers individually (instead of the collective approach that we used in

this study), they need to consider the effect of complexes and non-specific binding that may

result in noisy data. As such, we suggest that the aptamer expression data be used collectively

and after proper normalization, which enabled us to identify various EPs and important molecular

mechanisms discussed in this study.

In this work, we developed a predictive model based on blood markers that enables us to

generalize the definition of EPs to scenarios where data on circulating proteome is not available.

This significantly increases the applicability of this approach and may enable automating

identification of high-risk individuals in the clinic. However, time and follow-up studies are

required to move this predictive model and the definitions of EPs from research to clinic and

develop it as an acceptable clinical practice. Nonetheless, our approach can be used to study

COVID-19 in different cohorts and identify characteristics that can guide the treatment of the

disease.

Future directions

There is enormous amount of data within this study and BQC19 that can and should be exploited

by the scientific and medical community to improve our understating of COVID-19 and novel

emerging acute respiratory illnesses. Analyzing in more details the molecular profiles of the other

EPs, in particular EP3-5, which led to similar outcomes from distinct molecular pathways (Table

5) should further yield important insights into mechanisms of the disease. For example, EP5 is

enriched in males and cardiovascular disease complications, while EP4 is enriched in female (like

EP1) but with different distinct clinical trajectories pointing to sex-dependent and independent

molecular mechanisms of disease.

Conclusion

The major strength of this study is its starting point: unsupervised analysis of a large and deeply

phenotyped cohort. We showed that this approach can address both fundamental scientific

questions pertaining to mechanisms of disease and help the medical community improve patient

outcomes through early identification of patient that may follow a severe clinical course during

COVID-19.

Methods:

Datasets and preprocessing

The Biobanque québécoise de la COVID-19 (BQC19; www.quebeccovidbiobank.ca) is aimed at

coordinating the collection of patients' data and samples for COVID-19 related research. Data

and samples were collected from ten sites across the province of Québec (Canada) (Tremblay *et*

al., 2021). BQC19 organizes the collected data, including clinical information and multi-omics

experimental data, before making it available in successive releases. For this study, we used

release #5 of the clinical data published in December 2021, the circulating proteome determined

using SOMAmers (Zhou *et al.*, 2021) and Metabolomics data (Ford et al., 2020). BQC19 GWAS

imputation data was generated by Tomoko Nakanishi at Brent Richards lab, Jewish General

Hospital and McGill University. Detailed codes used for generating the data can be found in:

https://github.com/richardslab/BQC19_genotype_pipeline

Our main corpus of analysis consisted of n = 1,362 hospitalized and SARS-CoV-2 positive patients

(based on qRT-PCR) of BQC19. This included n = 731 patients for which both clinical and

proteomic data were available as well as n = 631 patients for which proteomic data was not

available, but their clinical data contained measurements for more than half (at least 11 out of

21) of the blood markers that we used as a validation set for the predictive model developed in

this study.

We also obtained data (n = 731) corresponding to the circulating proteome measured between

April 2, 2020 and April 20, 2021 by a multiplex SOMAmer affinity array (Somalogic, 4,985

aptamers) from BQC19 (release #3 Sep. 2021, associated patients' data updated in release #5

Dec. 2021). When measurements of the same patients but at different time points were

available, we used the one corresponding to the first time point. SomaScan is a biotechnological

protocol commercialized by the Somalogic company. It relies on a set of artificial aptamers linked

to a fluorophore and each designed to bind a single protein. Once added to the sample, the

activity of each aptamer is measured through fluorescence and used to approximate the

expression level of the targeted protein. SomaScan protocol comprises several levels of

calibration and normalization to correct technical biases (Gold *et al.*, 2010). Log2 and Z-score

normalization were performed on each aptamer separately in addition to the manufacturer's

provided normalized data (hybridization control normalization, intraplate median signal

normalization, and median signal normalization). Since the data was analyzed by Somalogic in

two separate batches, we applied the z-score transformation separately to each batch, to reduce

batch effects. These additional transformations ensure that the measured values of different

aptamers are comparable and can be used in cluster analysis.

We obtained metabolomic data (1,435 metabolites) from BQC19. We used the batch-normalized,

missing values imputed and log-transformed version of the data.

Analyses of the GWAS dataset

For the GWAS analyses, annotation of SNVs were done using the biomaRt package (Durinck et

al., 2009) from R (R Core Team, 2022) and all analyses were done using R version 4.1.3. Quality

control steps were derived in majority from a 2017 QC tutorial article (Marees et al., 2018). At

the beginning, we had 867,450 markers and 2,429 samples. We import Plink format data into R

using the "read_plink" function from genio package (Ochoa, 2022) from R. We removed 103,592

non ACGT bi-allelic markers. We calculated the predicted sex by looking at the rate of

homozygote markers on chromosome 23. We removed 3,588 markers with call rates < 98%,

448,932 monomorphic markers and markers with MAF <0.05, and 28,092 markers with Hardy

Weinberg equilibrium < 1E-6 (calculated by the "HWE.exact" function from the genetics package

(Warnes et al., 2021) from R. For the EP6 cluster group analyses, we removed in addition 1,747

samples that were not in the cluster analysis, 8 samples with a sex discrepancy (based on

predicted sex calculated earlier and reported sex), and 3 samples with a heterozygosity rate > 3

standard deviations. We finally removed a pair of samples who had approximately the same

genome probably due to an error of manipulation. We couldn't know which one was the right

sample, so we removed both of them. We also had 2 pairs of individuals who had a pi-hat of ~0.5

(meaning first degree relatives), we decided to keep one sample per pair, the one with the higher

call rate. All other pairs of individuals had a pi-hat < 0.21 that is judge acceptable considering our

population. Pi-hat were calculated with the "snpgdsIBDMoM" function from SNPRelate package

(Zheng et al., 2012) from R. At the end of quality controls, 283,246 markers on 655 samples have

been used to perform association analyses.

To perform the principal component analyses (PCA), we took a subsample of independent

markers (pruning) with a maximum sliding window of 500,000 base pairs and a linkage

disequilibrium (LD) threshold of 0.2 using the "snpgdsLDpruning" function from the SNPRelate

package from R. We ran the PCA with the "snpgdsPCA" function from SNPRelate package from

R. The first 2 principal components (PCs) were considered significant.

For the GWAS analyses of the 283,246 remaining markers between EP6 cluster compared to all

others, we modeled a logistic regression with the dichotomous variable indicating if the

participants belong to the EP6 cluster as the outcome variable. We used the additive model for

markers as an independent variable and we adjusted the models with the first two PCs. Odds

ratio and p-values were calculated on each model. QQ-plots have been performed as quality

control of the models; p-values were plotted using "qqplot.pvalues" function from gaston

package (Perdry and Dandine-Roulland, 2020) from R (data not shown). We compared the

aptamers' normalized level of expression (based on normalization described earlier) between the

three groups of genotypes for each studied genes by performing standard ANOVA analyses

followed by Tukey *post hoc* tests (referred to in this study as pQTL analysis). Since aptamers

tested are limited compared to the SNVs, we have fixed significance p-values threshold below

1E-4 to report more SNVs instead of the more common 1E-5 suggestive threshold. Finally, to

identify if the cluster EP6 may be explained by the aptamers and the SNVs, we performed

multiple logistic regression analyses models include aptamers expression values, SNV genotypes

(additive model) and the two principal components. OR are reported with 95% confidence

intervals.

Consensus agglomerative clustering

Patients were clustered using agglomerative clustering, with Euclidean distance and Ward's

linkage (Murtagh and Legendre, 2014; Ward Jr, 1963). To identify number of clusters k, we used

the elbow method based on the Akaike Information Criterion (AIC) and Bayesian Information

Criterion (BIC). More specifically, we calculated the AIC and BIC for clustering using k = 2, 3, …, 20

and used the Kneedle algorithm (Satopaa et al., 2011) to identify the value of k corresponding to

the "elbow", where increasing the value of k does not provide much better modeling of the data.

Kneedle identified k = 6 as the number of clusters based on both AIC and BIC (Figure 1A).

Given the number of clusters in the data, we then used consensus clustering with sub-sampling

to obtain robust endophenotypes. We randomly sampled 80% of the patients 1000 times. Each

time, we used agglomerative clustering above with k = 6 to identify clusters. Given these 1000

clusterings, we calculated the frequency of two patients appearing in the same cluster, when

both were present in the randomly formed dataset. We then performed one final agglomerative

clustering of these frequency scores to identify the six endophenotypes (Figure S2A and Figure

1B). The distribution of Rand-Index, showing the concordance between each one of the 1000

clusterings and the final consensus clustering is provided in Figure S2B (mean Rand-Index =

0.823), reflecting a high degree of consistency.

Metabolomic pathway characterization of EP6

The 1435 metabolites measured were organized into 122 sub-pathways in the original dataset

(denoted as "SUB_PATHWAYS"). We first identified metabolites whose values were significantly

higher or lower in EP6 compared to other EPs (two-sided MWU test, FDR<0.01). Then, we used

these metabolites to perform pathway enrichment analysis (one-sided Fisher's exact test) based

on 122 pathways. The resulting p-values (Table S3) were then corrected for multiple tests using

Benjamini-Hochberg FDR.

Nearest-centroid predictor based on blood markers

In order to predict endophenotypes from blood tests, we developed a missing-value resilient

nearest-centroid classifier. We used the set of patients that were used to form the original EPs

(n = 731) as the training set and the set of patients that did not have proteome data as the

validation set (n = 631). First, we z-score normalized each blood marker across all the patients in

the training set, one marker at a time. We then formed a blood marker signature (a vector of

length 21) for each EP. Each element of an EP's signature corresponds to the mean of the

corresponding marker across all patients of that EP.

To predict the EP label of each patient in the test set, we first z-score normalized their blood

marker measurements using the mean and standard deviation of the blood markers calculated

from the training set. Then, we calculated the cosine distance between each test patient's blood

marker profile and the centroids (excluding missing values) and identified the nearest EP as the

predicted EP (PEP) label of the patient.

Acknowledgements

This work was made possible through open sharing of data and samples from the Biobanque

québécoise de la COVID-19, funded by the Fonds de recherche du Québec - Santé, Génome

Québec, the Public Health Agency of Canada and, as of March 2022, the Ministère de la Santé et

des Services Sociaux du Québec. We thank all participants to BQC19 for their contribution. This

study was supported by the Fonds de recherche du Québec - Santé (FRQS)- Cardiometabolic

Health, Diabetes and Obesity Research Network (CMDO)- Initiative. This work was also supported

by Natural Sciences and Engineering Research Council of Canada (NSERC) grant RGPIN-2019-

04460 (AE).

Permissions

The study was approved by the Institutional Ethics Review Board of the "Centre intégré

universitaire de santé et de services sociaux du Saguenay-Lac-Saint-Jean" (CIUSSS-SLSJ) affiliated

to Université de Sherbrooke [protocol #2021-369, 2021-014 CMDO – COVID19].

Figures

Figure 1: Unsupervised consensus clustering of SARS-CoV-2 positive patients.
A) The elbow points (circles in red) of Akaike's Information Criteria (AIC) and Bayesian
Information Criteria (BIC) curves versus number of clusters consistently corresponded to k=6 as
the optimal number of clusters. B) The heatmap shows the expression of aptamers (rows) in each
sample (columns). The dendogram shows the identified endophenotypes. C) Characterization of
samples based on sex at birth, highest world health organization (WHO) severity level achieved,
intensive care unit (ICU) admission, ventilatory support, and oxygen therapy. For the last three
rows, a sample colored "black" reflects a label of "yes".

Figure 2: Characterization of endophenotypes (EPs).
A) Enrichment or depletion of each EP in clinical variables (one cluster versus rest). Two-sided
Fisher's exact tests are used to calculate the p-values, which are corrected for multiple tests using
Benjamini-Hochberg false discovery rate (FDR). Gradients of blue show depletion, while gradients
of red show enrichment. FDR values above 0.05 are depicted as white. B) The number of patients
in each EP and the colors used to represent them in panels C, D, and E. C) Kaplan-Meier analysis
of the time between patients' admission to the hospital and their admission to intensive care unit
(ICU) (or death if earlier) for each EP (Delta). D) Distribution of age in each EP. E) Distribution of
BMI in each EP. F) COVID-19 severity in each EP.

Figure 3: Frequency and significance of complications in different EPs.
The value in each cell shows the percentage of patients of that EP (column) that suffered from the complication (row). The colors represent two-sided Fisher's exact test false discovery rate (FDR, corrected for multiple tests). Red represents enrichment, while blue represents depletion. FDR values below 0.05 are shown as white.

594
Figure 4: Metabolite characteristics of endophenotypes (EPs).
A) The heatmap shows the over-expression (red) and under-expression (blue) of metabolites in
different EPs (two-sided Mann–Whitney U test). Row names show metabolites followed by the
sub-pathway to which they belong in parentheses. Only top 15 metabolites (based on false
discovery rate for EP6) for which a definite name and sub-pathway was available are shown. Full
list is provided in Table S3. B) The heatmap shows the enrichment (one-sided Fisher's exact test)
of EPs in differen   e Table S3 for the full list.

Figure 5: Differential expression patterns of blood markers in the identified endophenotypes (EPs) and characterization of predicted endophenotypes (PEPs) based on the predictive model. A) Heatmaps of false discovery rates (FDR) values for two-sided one-vs-rest Mann–Whitney U tests for 21 blood markers for each EP. Abbreviations used: WBC = white blood cells, LDH = lactate deshydrogenase, ALT = alanine aminotransferase, aPTT = activated partial thromboplastin time, INR = International Normalized Ratio. FDR values below 0.05 are shown as white. B) World health organization COVID-19 severity assessed in each PEP. C) Distribution of age in each PEP. D) Distribution of BMI in each PEP. E) Kaplan-Meier analysis of the time between patients' admission to the hospital and their admission to ICU (or death if earlier) for each PEP (Delta). The colormap

612   in panel E shows the number of patients in each PEP group and the colors used to represent them

613   in panels C, D, and E. PEPs were predicted based on blood markers and were not used to originally

614   identify the EPs.

 Tables

616 **Table 1:** Clinical and pathological characteristics of the BQC19's participants used in this study.

		Cohort (n = 1,362) No. (%)
Age in years	<45	17.5
	45-65	33.6
	>65	48.7
	Unknown	0.3
Body mass index in kg/m²	<20	3.8
	20-25	13.0
	25-35	27.8
	>35	7.4
	Unknow	47.9
Sex at birth	Female	44.8
	Male	55.2
Severity	Dead	4.4
	Severe	21.8
	Moderate	51.0
	Mild	11.8
	Unknown	10.9
Oxygen therapy	Yes	48.2
	No	21.0
	Unknown	30.8
Ventilatory support	Yes	47.6
	No	14.0
	Unknown	38.5
Admission to intensive care unit	Yes	17.5
	No	33.6
	Unknown	48.7

618 **Table 2:** Reactome pathways associated with EP6, based on expression of aptamers. KnowEnG
619 analytical platform was used. The p-values were calculated using a one-sided Fisher's exact test
620 and were corrected for multiple tests using Benjamini-Hochberg method. Only signaling
621 pathways with FDR<5E-4 are shown in this table (see Table S4 for the full list).
622

Pathway	FDR
Signaling by Interleukins	2.75E-12
Cytokine Signaling in Immune system	6.25E-10
Immune System	2.28E-06
Signaling by FGFR	4.75E-06
Constitutive Signaling by Aberrant PI3K in Cancer	2.94E-05
FGFR2 ligand binding and activation	5.77E-05
Extracellular matrix organization	1.85E-04
FGFR3 ligand binding and activation	1.85E-04
FGFR3c ligand binding and activation	1.85E-04
FGFRL1 modulation of FGFR1 signaling	1.85E-04
TNFs bind their physiological receptors	2.58E-04
PI3K/AKT Signaling in Cancer	3.28E-04
PI5P, PP2A and IER3 Regulate PI3K/AKT Signaling	4.14E-04
Interleukin-4 and 13 signaling	4.91E-04

623
624

625   **Table 3**: SNVs differentiating EP6 against all other endophenotype clusters.

SNV ID	Position	Gene Symbol	SNV [1]	MAF	HWE	OR [2]	p-value
rs1394671	chr5:7348567	-	G > A	0.240	0.095	2.28	2.41E-06
rs12186698	chr5:168880668	*SLIT3*	T > C	0.087	0.247	2.93	3.72E-06
rs11625406	chr14:50049474	-	C > A	0.438	0.137	0.47	1.03E-05
rs11862889	chr16:83828886	-	T > C	0.104	0.446	2.62	1.07E-05
rs2995918	chr4:37905775	*TBC1D1*	C > T	0.494	0.776	0.49	1.20E-05
rs16897810	chr5:68415296	-	G > A	0.093	0.001	2.67	1.43E-05
rs2376263	chr17:35436659	*SLFN13*	G > A	0.261	0.073	2.01	2.48E-05
rs4790712	chr17:1614620	*SLC43A2*	G > A	0.441	0.003	1.97	2.50E-05
rs2294566	chr20:41472939	*CHD6*	C > A	0.239	0.265	2.04	2.78E-05
rs7164451	chr15:48921859	*SHC4*	G > A	0.386	0.053	1.89	3.01E-05
rs57664621	chr8:22808443	*PEBP4*	G > C	0.174	0.016	2.16	3.07E-05
rs28482919	chr3:14903094	*FGD5*	T > C	0.187	0.053	0.32	3.20E-05
rs6559283	chr9:89712560	-	T > C	0.375	<0.001	1.96	3.27E-05
rs657075	chr5:132094425	-	A > G	0.104	0.328	2.49	4.28E-05
rs56235109	chr15:62424001	*TLN2*	A > G	0.235	0.366	0.38	4.67E-05
rs7620057	chr3:179473377	*GNB4*	T > C	0.099	1.000	2.51	4.67E-05
rs10466868	chr12:131455375	-	T > G	0.108	0.401	2.51	5.94E-05
rs2236798	chr1:18735127	*PAX7*	A > G	0.054	0.114	2.90	6.11E-05
rs3774814	chr4:5464702	*STK32B*	C > G	0.205	<0.001	0.36	6.75E-05
rs4497815	chr19:22903215	-	G > A	0.216	0.205	2.07	7.52E-05
rs6765694	chr3:54601810	*CACNA2D3*	G > A	0.404	0.833	0.51	7.66E-05
rs2797773	chr6:37559045	-	C > T	0.422	0.000	0.53	8.06E-05
rs17014760	chr4:129419725	-	A > A	0.316	0.000	1.89	9.33E-05
rs10948260	chr6:45835559	-	G > A	0.366	0.189	1.81	9.52E-05
rs12035677	chr1:232391209	-	A > G	0.063	0.000	3.02	9.95E-05

626   Abbreviations used: SNV = Single nucleotide variation, HWE = Hardy Weinberg Equilibrium, MAF
627   = Minor allele frequency, OR = Odd ratio.
628
629   [1] SNV are described following GWAS annotations: refence allele > alternative allele (e.g., G > A).
630   [2] Logistic regression analyses using additive model adjusted for the 2 principal components.
631

632 **Table 4:** Association between genotypes and aptamer expression levels

SNV	Gene Symbol	Nearest gene	Aptamers normalized expression [1]				Multiple logistic regression analyses [3]			
			HM_{ref}	HTZ	HM_{alt}	p-val [2]	**Aptamers** OR	p-val	**SNV** OR	p-val
rs6765694	*CACNA2D3*	-	-0.09 (±0.89)	0.01 (±1.10)	0.24 * (±0.99)	**0.011**	0.61 (0.46-0.79)	2.5E-4	0.52 (0.37-0.74)	2.4E-4
rs10948260	-	*CLIC5*	0.02 (±0.92)	0.02 (±1.09)	-0.02 (±0.97)	0.796	0.68 (0.55-0.83)	1.7E-4	1.83 (1.35-2.48)	8.8E-5
rs657075	-	*IL3*	-0.01 (±1.01)	0.05 (±0.99)	0.32 (±0.46)	0.328	1.37 (1.14-11.64)	8.2E-4	2.49 (1.60-3.88)	5.2E-5
rs7164451	*SHC4*	-	-0.07 (±0.90)	0.01 (±1.05)	0.23 * (±1.16)	**0.017**	11.98 (7.61-18.87)	8.4E-27	2.00 (1.30-3.09)	1.8E-3
rs12186698	*SLIT3*	-	0.00 (±1.02)	0.07 (±1.06)	0.15 (±0.89)	0.509	0.82 (0.64-1.06)	0.135	2.99 (1.89-4.73)	2.7E-6
rs56235109	*TLN2*	-	-0.03 (±0.98)	0.06 (±1.03)	0.04 (±1.27)	0.353	0.58 (0.45-0.75)	2.7E-5	0.38 (0.23-0.60)	4.7E-5

633 Abbreviations used: SNV = Single nucleotide variation, HM_{ref} = Homozygotes for the reference
634 allele, HM_{alt} = Homozygotes for the alternative allele, HTZ = Heterozygotes, OR = Odd ratio.
635
[1] Aptamers' normalized levels of expression are reported as mean (± standard deviation).
Normalization steps for aptamer expressions are described in Methods.
[2] p-value, standard ANOVA analyses followed by Tukey *post hoc* analyses. Asterisks (*) identify
difference between HM_{ref} and HM_{alt} genotypes.
[3] Multiple logistic regression analyses models include aptamers expression values, SNV
genotypes (additive model) and the two principal components. OR are reported with 95%
confidence intervals in parentheses.
643
644

645 **Table 5:** Summary of the characteristics of each endophenotype.
646 In this table, High (Low), denoted as H (L) implies that the average value of the variable in the
647 corresponding EP was significantly higher (lower) than the other EPs (considered together), while
648 N (Nondescript) implies that it was not significantly different. The table is formed based on results
649 described in Tables S1, S3 and S5.

Endophenotype	Age	Sex at birth	BMI	Metabolic sub-pathways	Blood markers
EP1	H	F	N	Leucine, Isoleucine and Valine Metabolism	High lymphocyte, Low neutrophil
EP2	N	N	L	N	High albumin, Low white blood cells
EP3	L	M	N	Phospholipid Metabolism	High platelet, Low creatinine
EP4	N	N	N	Ascorbate and Aldarate Metabolism	High lactate deshydrogenase
EP5	H	N	N	Benzoate Metabolism	High creatinine, Low haemoglobin
EP6	N	M	N	Methionine, Cysteine, SAM and Taurine Metabolism	High white blood cells, Low lymphocyte

650 Abbreviations used: H = High, L = Low, N = Nondescript, F = Female, M = Male
651

References

Agarwal, B., Gatt, A., Riddell, A., Wright, G., Chowdary, P., Jalan, R., Burroughs, A.K., and
Davenport, A. (2013). Hemostasis in patients with acute kidney injury secondary to acute liver
failure. Kidney International *84*, 158-163. https://doi.org/10.1038/ki.2013.92.

Al-Hadrawi, D.S., Al-Rubaye, H.T., Almulla, A.F., Al-Hakeim, H.K., and Maes, M. (2022). Lowered
oxygen saturation and increased body temperature in acute COVID-19 largely predict chronic
fatigue syndrome and affective symptoms due to Long COVID: A precision nomothetic approach.
Acta Neuropsychiatr, 1-12. 10.1017/neu.2022.21.

Ataga, K.I. (2009). Hypercoagulability and thrombotic complications in hemolytic anemias.
Haematologica *94*, 1481-1484. 10.3324/haematol.2009.013672.

Barnes, B.J., Adrover, J.M., Baxter-Stoltzfus, A., Borczuk, A., Cools-Lartigue, J., Crawford, J.M.,
Daßler-Plenker, J., Guerci, P., Huynh, C., and Knight, J.S. (2020). Targeting potential drivers of
COVID-19: Neutrophil extracellular traps. Journal of Experimental Medicine *217*.

Berrettini, W.H. (2005). Genetic bases for endophenotypes in psychiatric disorders. Dialogues
Clin Neurosci *7*, 95-101. 10.31887/DCNS.2005.7.2/wberrettini.

Blachier, F., Andriamihaja, M., and Blais, A. (2020). Sulfur-Containing Amino Acids and Lipid
Metabolism. The Journal of Nutrition *150*, 2524S-2531S. 10.1093/jn/nxaa243.

Blasco, A., Coronado, M.-J., Hernández-Terciado, F., Martín, P., Royuela, A., Ramil, E., García, D.,
Goicolea, J., Del Trigo, M., and Ortega, J. (2021). Assessment of neutrophil extracellular traps in
coronary thrombus of a case series of patients with COVID-19 and myocardial infarction. JAMA
cardiology *6*, 469-474.

Blatti III, C., Emad, A., Berry, M.J., Gatzke, L., Epstein, M., Lanier, D., Rizal, P., Ge, J., Liao, X., and
Sobh, O. (2020). Knowledge-guided analysis of "omics" data using the KnowEnG cloud platform.
PLoS biology *18*, e3000583. 10.1371/journal.pbio.3000583.

Bonaventura, A., Vecchié, A., Dagna, L., Martinod, K., Dixon, D.L., Van Tassell, B.W., Dentali, F.,
Montecucco, F., Massberg, S., and Levi, M. (2021). Endothelial dysfunction and
immunothrombosis as key pathogenic mechanisms in COVID-19. Nature Reviews Immunology
21, 319-329.

Bussolari, C., Palumbo, D., Fominsky, E., Nardelli, P., De Lorenzo, R., Vitali, G., De Cobelli, F.,
Rovere-Querini, P., and Scandroglio, A.M. (2021). Case Report: Nintedaninb May Accelerate Lung
Recovery in Critical Coronavirus Disease 2019. Frontiers in Medicine *8*, 766486.

Desilles, J.P., Solo Nomenjanahary, M., Consoli, A., Ollivier, V., Faille, D., Bourrienne, M.C.,
Hamdani, M., Dupont, S., Di Meglio, L., and Escalard, S. (2022). Impact of COVID-19 on thrombus
composition and response to thrombolysis: Insights from a monocentric cohort population of

COVID-19 patients with acute ischemic stroke. Journal of Thrombosis and Haemostasis *20*, 919-
928.

Ding, J., Hostallero, D.E., El Khili, M.R., Fonseca, G.J., Milette, S., Noorah, N., Guay-Belzile, M.,
Spicer, J., Daneshtalab, N., and Sirois, M. (2021). A network-informed analysis of SARS-CoV-2 and
hemophagocytic lymphohistiocytosis genes' interactions points to Neutrophil extracellular traps
as mediators of thrombosis in COVID-19. PLoS Computational Biology *17*, e1008810.
10.1371/journal.pcbi.1008810.

Durinck, S., Spellman, P.T., Birney, E., and Huber, W. (2009). Mapping identifiers for the
integration of genomic datasets with the R/Bioconductor package biomaRt. Nature Protocols *4*,
1184-1191. 10.1038/nprot.2009.97.

Emad, A., Ray, T., Jensen, T.W., Parat, M., Natrajan, R., Sinha, S., and Ray, P.S. (2020). Superior
breast cancer metastasis risk stratification using an epithelial-mesenchymal-amoeboid transition
gene signature. Breast Cancer Research *22*, 74. 10.1186/s13058-020-01304-8.

Englert, H., Rangaswamy, C., Deppermann, C., Sperhake, J.-P., Krisp, C., Schreier, D., Gordon, E.,
Konrath, S., Haddad, M., and Pula, G. (2021). Defective NET clearance contributes to sustained
FXII activation in COVID-19-associated pulmonary thrombo-inflammation. EBioMedicine *67*,
103382.

Esaki, Y., Hirokawa, K., Fukazawa, T., and Matsuda, T. (1984). Immunohistochemical study on the
liver in autopsy cases with disseminated intravascular coagulation (DIC) with reference to
clinicopathological analysis. Virchows Archiv A *404*, 229-241.

Ford, L., Kennedy, A.D., Goodman, K.D., Pappan, K.L., Evans, A.M., Miller, L.A.D., Wulff, J.E.,
Wiggs, B.R., III,, Lennon, J.J., Elsea, S., and Toal, D.R. (2020). Precision of a Clinical Metabolomics
Profiling Platform for Use in the Identification of Inborn Errors of Metabolism. The Journal of
Applied Laboratory Medicine *5*, 342-356. 10.1093/jalm/jfz026.

Gillespie, M., Jassal, B., Stephan, R., Milacic, M., Rothfels, K., Senff-Ribeiro, A., Griss, J., Sevilla, C.,
Matthews, L., Gong, C., et al. (2021). The reactome pathway knowledgebase 2022. Nucleic Acids
Research *50*, D687-D692. 10.1093/nar/gkab1028.

Gold, L., Ayers, D., Bertino, J., Bock, C., Bock, A., Brody, E., Carter, J., Cunningham, V., Dalby, A.,
and Eaton, B. (2010). Aptamer-based multiplexed proteomic technology for biomarker discovery.
Nature Precedings, 1-1.

Han, G., Gupta, S.D., Gable, K., Niranjanakumari, S., Moitra, P., Eichler, F., Brown Jr, R.H., Harmon,
J.M., and Dunn, T.M. (2009). Identification of small subunits of mammalian serine
palmitoyltransferase that confer distinct acyl-CoA substrate specificities. Proceedings of the
National Academy of Sciences *106*, 8186-8191.

Hull, D.L. (1974). Informal aspects of theory reduction. (Cambridge University Press), pp. 653-
670.

Kajita, K., Ishizuka, T., Miura, A., Kanoh, Y., Ishizawa, M., Kimura, M., Muto, N., and Yasuda, K.
(2001). Glucocorticoid-induced insulin resistance associates with activation of protein kinase C
isoforms. Cellular signalling *13*, 169-175.

Kaplan, E.L., and Meier, P. (1958). Nonparametric Estimation from Incomplete Observations.
Journal of the American Statistical Association *53*, 457-481. 10.1080/01621459.1958.10501452.

Kim, K., Chang, Y.-K., Bian, Y., Bae, O.-N., Lim, K.-M., and Chung, J.-H. (2018). Pro-coagulant and
pro-thrombotic effects of paclitaxel mediated by red blood cells. Thrombosis and Haemostasis
118, 1765-1775.

Lachkar, F., Ferré, P., Foufelle, F., and Papaioannou, A. (2021). Dihydroceramides: their emerging
physiological roles and functions in cancer and metabolic diseases. American Journal of
Physiology-Endocrinology and Metabolism *320*, E122-E130.

Leppkes, M., Knopf, J., Naschberger, E., Lindemann, A., Singh, J., Herrmann, I., Stürzl, M., Staats,
L., Mahajan, A., and Schauer, C. (2020). Vascular occlusion by neutrophil extracellular traps in
COVID-19. EBioMedicine *58*, 102925.

Liu, W., Peng, Y., and Tobin, D.J. (2013). A new 12-gene diagnostic biomarker signature of
melanoma revealed by integrated microarray analysis. PeerJ *1*, e49. 10.7717/peerj.49.

Lone, M.A., Hülsmeier, A.J., Saied, E.M., Karsai, G., Arenz, C., von Eckardstein, A., and Hornemann,
T. (2020). Subunit composition of the mammalian serine-palmitoyltransferase defines the
spectrum of straight and methyl-branched long-chain bases. Proceedings of the National
Academy of Sciences *117*, 15591-15598.

Luo, S., Feofanova, E.V., Tin, A., Tung, S., Rhee, E.P., Coresh, J., Arking, D.E., Surapaneni, A.,
Schlosser, P., Li, Y., et al. (2021). Genome-wide association study of serum metabolites in the
African American Study of Kidney Disease and Hypertension. Kidney International *100*, 430-439.
https://doi.org/10.1016/j.kint.2021.03.026.

MacDonald, J., and Sprecher, H. (1989). Distribution of arachidonic acid in choline-and
ethanolamine-containing phosphoglycerides in subfractionated human neutrophils. Journal of
Biological Chemistry *264*, 17718-17726.

Majumder, R., Liang, X., Quinn-Allen, M.A., Kane, W.H., and Lentz, B.R. (2011). Modulation of
prothrombinase assembly and activity by phosphatidylethanolamine. Journal of Biological
Chemistry *286*, 35535-35542.

Marees, A.T., de Kluiver, H., Stringer, S., Vorspan, F., Curis, E., Marie-Claire, C., and Derks, E.M.
(2018). A tutorial on conducting genome-wide association studies: Quality control and statistical
analysis. Int J Methods Psychiatr Res *27*, e1608. 10.1002/mpr.1608.

Marfia, G., Navone, S., Guarnaccia, L., Campanella, R., Mondoni, M., Locatelli, M., Barassi, A.,
Fontana, L., Palumbo, F., and Garzia, E. (2021). Decreased serum level of sphingosine-1-

phosphate: a novel predictor of clinical severity in COVID-19. EMBO molecular medicine *13*,
e13424.

Marshall, J.C., Murthy, S., Diaz, J., Adhikari, N., Angus, D.C., Arabi, Y.M., Baillie, K., Bauer, M.,
Berry, S., and Blackwood, B. (2020). A minimal common outcome measure set for COVID-19
clinical research. The Lancet Infectious Diseases *20*, e192-e197.

Middleton, E.A., He, X.-Y., Denorme, F., Campbell, R.A., Ng, D., Salvatore, S.P., Mostyka, M.,
Baxter-Stoltzfus, A., Borczuk, A.C., and Loda, M. (2020). Neutrophil extracellular traps contribute
to immunothrombosis in COVID-19 acute respiratory distress syndrome. Blood *136*, 1169-1179.

Murtagh, F., and Legendre, P. (2014). Ward's Hierarchical Agglomerative Clustering Method:
Which Algorithms Implement Ward's Criterion? Journal of Classification *31*, 274-295.
10.1007/s00357-014-9161-z.

Obermayer, A., Jakob, L.-M., Haslbauer, J.D., Matter, M.S., Tzankov, A., and Stoiber, W. (2021).
Neutrophil extracellular traps in fatal COVID-19-associated lung injury. Disease markers *2021*.

Ochoa, A. (2022). genio: Genetics Input/Output Functions. R package version 1.1.1.
https://CRAN.R-project.org/package=genio.

Ouwendijk, W.J., Raadsen, M.P., Van Kampen, J.J., Verdijk, R.M., Von Der Thusen, J.H., Guo, L.,
Hoek, R.A., Van Den Akker, J.P., Endeman, H., and Langerak, T. (2021). High levels of neutrophil
extracellular traps persist in the lower respiratory tract of critically ill patients with coronavirus
disease 2019. The Journal of infectious diseases *223*, 1512-1521.

Papageorgiou, C., Jourdi, G., Adjambri, E., Walborn, A., Patel, P., Fareed, J., Elalamy, I.,
Hoppensteadt, D., and Gerotziafas, G.T. (2018). Disseminated intravascular coagulation: an
update on pathogenesis, diagnosis, and therapeutic strategies. Clinical and Applied
Thrombosis/Hemostasis *24*, 8S-28S.

Parman, T., Bunin, D.I., Ng, H.H., McDunn, J.E., Wulff, J.E., Wang, A., Swezey, R., Rasay, L.,
Fairchild, D.G., and Kapetanovic, I.M. (2011). Toxicogenomics and metabolomics of
pentamethylchromanol (PMCol)-induced hepatotoxicity. Toxicological Sciences *124*, 487-501.

Perdry, H., and Dandine-Roulland, C. (2020). gaston: Genetic Data Handling (QC, GRM, LD, PCA)
& Linear Mixed Models. R package version 1.5.7. https://CRAN.R-project.org/package=gaston.

Petito, E., Falcinelli, E., Paliani, U., Cesari, E., Vaudo, G., Sebastiano, M., Cerotto, V., Guglielmini,
G., Gori, F., and Malvestiti, M. (2021). Association of neutrophil activation, more than platelet
activation, with thrombotic complications in coronavirus disease 2019. The Journal of infectious
diseases *223*, 933-944.

Pirnes-Karhu, S., Jantunen, E., Mäntymaa, P., Mustjoki, S., Alhonen, L., and Uimari, A. (2014).
Spermidine/spermine N1-acetyltransferase activity associates with white blood cell count in
myeloid leukemias. Experimental Hematology *42*, 574-580.

Port, J.R., Adney, D.R., Schwarz, B., Schulz, J.E., Sturdevant, D.E., Smith, B.J., Avanzato, V.A.,
Holbrook, M.G., Purushotham, J.N., and Stromberg, K.A. (2021). High-fat high-sugar diet-induced
changes in the lipid metabolism are associated with mildly increased COVID-19 severity and
delayed recovery in the Syrian hamster. Viruses *13*, 2506.

R Core Team (2022). R: A Language and Environment for Statistical Computing. (R Foundation for
Statistical Computing). https://www.R-project.org/.

Ran, S., Downes, A., and Thorpe, P.E. (2002). Increased exposure of anionic phospholipids on the
surface of tumor blood vessels. Cancer research *62*, 6132-6140.

Rubin, R. (2020). As their numbers grow, COVID-19 "long haulers" stump experts. Jama *324*,
1381-1383.

Russell, C.D., and Baillie, J.K. (2017). Treatable traits and therapeutic targets: goals for systems
biology in infectious disease. Current opinion in systems biology *2*, 140-146.

Sato, K., Iemitsu, M., Aizawa, K., and Ajisaka, R. (2008). Testosterone and DHEA activate the
glucose metabolism-related signaling pathway in skeletal muscle. American Journal of
Physiology-Endocrinology and Metabolism *294*, E961-E968.

Satopaa, V., Albrecht, J., Irwin, D., and Raghavan, B. (2011). Finding a" kneedle" in a haystack:
Detecting knee points in system behavior. (IEEE), pp. 166-171.

Sinauridze, E.I., Kireev, D.A., Popenko, N.Y., Pichugin, A.V., Panteleev, M.A., Krymskaya, O.V., and
Ataullakhanov, F.I. (2007). Platelet microparticle membranes have 50-to 100-fold higher specific
procoagulant activity than activated platelets. Thrombosis and haemostasis *97*, 425-434.

Skendros, P., Mitsios, A., Chrysanthopoulou, A., Mastellos, D.C., Metallidis, S., Rafailidis, P.,
Ntinopoulou, M., Sertaridou, E., Tsironidou, V., and Tsigalou, C. (2020). Complement and tissue
factor–enriched neutrophil extracellular traps are key drivers in COVID-19 immunothrombosis.
The Journal of clinical investigation *130*, 6151-6157.

Smadja, D.M., Philippe, A., Bory, O., Gendron, N., Beauvais, A., Gruest, M., Peron, N., Khider, L.,
Guerin, C.L., and Goudot, G. (2021). Placental growth factor level in plasma predicts COVID-19
severity and in-hospital mortality. Journal of Thrombosis and Haemostasis *19*, 1823-1830.

Stafford, J.H., and Thorpe, P.E. (2011). Increased exposure of phosphatidylethanolamine on the
surface of tumor vascular endothelium. Neoplasia *13*, 299-IN292.

Tavoosi, N., Davis-Harrison, R.L., Pogorelov, T.V., Ohkubo, Y.Z., Arcario, M.J., Clay, M.C., Rienstra,
C.M., Tajkhorshid, E., and Morrissey, J.H. (2011). Molecular Determinants of Phospholipid
Synergy in Blood Clotting *. Journal of Biological Chemistry *286*, 23247-23253.
10.1074/jbc.M111.251769.

Te Pas, M.F.W., Madsen, O., Calus, M.P.L., and Smits, M.A. (2017). The Importance of
Endophenotypes to Evaluate the Relationship between Genotype and External Phenotype.
International Journal of Molecular Sciences *18*, 472.

Thomas, T., Stefanoni, D., Reisz, J.A., Nemkov, T., Bertolone, L., Francis, R.O., Hudson, K.E.,
Zimring, J.C., Hansen, K.C., and Hod, E.A. (2020). COVID-19 infection alters kynurenine and fatty
acid metabolism, correlating with IL-6 levels and renal status. JCI insight *5*.

Tremblay, K., Rousseau, S., Zawati, M.n.H., Auld, D., Chassé, M., Coderre, D., Falcone, E.L.,
Gauthier, N., Grandvaux, N., and Gros-Louis, F. (2021). The Biobanque québécoise de la COVID-
19 (BQC19)—A cohort to prospectively study the clinical and biological determinants of COVID-
19 clinical trajectories. PloS one *16*, e0245031. 10.1371/journal.pone.0245031.

Ward Jr, J.H. (1963). Hierarchical grouping to optimize an objective function. Journal of the
American statistical association *58*, 236-244.

Warnes, G., Gorjanc, G., Leisch, F., and Man, M. (2021). genetics: Population Genetics. R package
version 1.3.8.1.3. https://CRAN.R-project.org/package=genetics.

Williams, R., Schalm, S.W., and O'Grady, J.G. (1993). Acute liver failure: redefining the syndromes.
The Lancet *342*, 273-275. https://doi.org/10.1016/0140-6736(93)91818-7.

Ye, C., Sutter, B.M., Wang, Y., Kuang, Z., and Tu, B.P. (2017). A metabolic function for phospholipid
and histone methylation. Molecular cell *66*, 180-193. e188.

Zahedi, K., Barone, S., Kramer, D.L., Amlal, H., Alhonen, L., Jänne, J., Porter, C.W., and Soleimani,
M. (2010). The role of spermidine/spermine N 1-acetyltransferase in endotoxin-induced acute
kidney injury. American Journal of Physiology-Cell Physiology *299*, C164-C174.

Zheng, X., Levine, D., Shen, J., Gogarten, S.M., Laurie, C., and Weir, B.S. (2012). A high-
performance computing toolset for relatedness and principal component analysis of SNP data.
Bioinformatics *28*, 3326-3328. 10.1093/bioinformatics/bts606.

Zhou, S., Butler-Laporte, G., Nakanishi, T., Morrison, D.R., Afilalo, J., Afilalo, M., Laurent, L.,
Pietzner, M., Kerrison, N., Zhao, K., et al. (2021). A Neanderthal OAS1 isoform protects individuals
of European ancestry against COVID-19 susceptibility and severity. Nature Medicine *27*, 659-667.
10.1038/s41591-021-01281-1.