Chikungunya intra-vector dynamics in *Aedes albopictus* from Lyon (France) upon exposure to a human viremia-like dose range reveals vector barrier's permissiveness and supports local epidemic potential

Barbara Viginier[1], Lucie Cappuccio[1], Céline Garnier[1], Edwige Martin[2], Carine Maisse[1], Claire Valiente Moro[2], Guillaume Minard[2], Albin Fontaine[3,4,5], Sébastian Lequime[6], Maxime Ratinier[1], Frédérick Arnaud[1] & Vincent Raquin*[1]

IVPC UMR754, INRAE, Universite Claude Bernard Lyon 1, EPHE, PSL Research University, F-69007 Lyon, France.

Universite Claude Bernard Lyon 1, CNRS, INRAE, VetAgro Sup, UMR Ecologie Microbienne, F-69622 Villeurbanne, France.

Unité Parasitologie et Entomologie, Département Microbiologie et maladies infectieuses, Institut de Recherche Biomédicale des Armées (IRBA), Marseille, France.

Aix Marseille Univ, IRD, SSA, AP-HM, UMR Vecteurs–Infections Tropicales et Méditerranéennes (VITROME), Marseille, France.

IHU Méditerranée Infection, Marseille, France.

Cluster of Microbial Ecology, Groningen Institute for Evolutionary Life Sciences, University of Groningen, Groningen, The Netherlands.

*Corresponding author

Correspondence: vincent.raquin@ephe.psl.eu

ABSTRACT

Arbovirus emergence and epidemic potential, as approximated by the vectorial capacity formula, depends on host and vector parameters, including the vector's intrinsic ability to replicate then transmit the pathogen known as vector competence. Vector competence is a complex, time-dependent, quantitative phenotype influenced by biotic and abiotic factors. A combination of experimental and modelling approaches is required to assess arbovirus intra-vector dynamics and estimate epidemic potential. In this study, we measured infection, dissemination, and transmission dynamics of chikungunya virus (CHIKV) in a field-derived *Aedes albopictus* population (Lyon, France) after oral exposure to a range of virus doses spanning human viraemia. Statistical modelling indicates rapid and efficient CHIKV progression in the vector mainly due to an absence of a dissemination barrier, with 100% of the infected mosquitoes ultimately exhibiting a disseminated infection, regardless of the virus dose. Transmission rate data revealed a time-dependent, but overall weak, transmission barrier, with individuals transmitting as soon as 2 days post-exposure (dpe) and >50% infectious mosquitoes at 6 dpe for the highest dose. Based on these experimental intra-vector dynamics data, epidemiological simulations conducted with an agent-based model showed that even at low mosquito biting rates, CHIKV could trigger outbreaks locally. Together, this reveals the epidemic potential of CHIKV upon transmission by *Aedes albopictus* in mainland France.

Keywords: Arbovirus ; vector ; mosquito ; *Aedes albopictus* ; chikungunya virus ; epidemiology ; vector competence ; modelisation

Introduction

Arthropod-borne viruses (arboviruses) are pathogens transmitted to vertebrate hosts by hematophagous arthropods, mainly mosquitoes. Arbovirus spread is a multi-factorial, dynamic process that can be estimated using the vectorial capacity (VCap) model, which aims to determine the average number of infectious vector bites that arise per day from one infected host in a susceptible human population (Smith et al., 2012). The vector-centric component of VCap integrates mosquito ecological (density per host, survival) and behavioural (daily biting rate per human) factors along with the vector's proxies of virus transmission efficiency such as vector competence (VComp) and its time-related expression, the extrinsic incubation period (EIP). VComp represents the ability of mosquitoes to : i) allow midgut infection following an infectious blood meal, ii) disseminate the virus beyond the midgut barrier, and iii) retransmit the virus through the saliva during the next bite. In VCap models, VComp and EIP are often simplified under the EIP_{50}, the time required to reach 50% of infectious mosquitoes. Effectively, each individual mosquito has a given EIP leading to a range of EIPs in the population. EIP distribution can be assessed experimentally by measuring the time between the initial mosquito infection and the mosquito infectiousness using an adequate experimental design (transmission assay for a relevant number of individual mosquitoes and time points). Taking into account the time-dependency of Vcomp improves VCap estimation and therefore allows to capture the full epidemic potential of arboviruses (Lequime et al., 2020). VComp is impacted by biotic (*e.g.*, mosquito and virus genotype, virus dose, mosquito microbiota) and abiotic (*e.g.*, temperature) factors (Viglietta et al., 2021), but how these factors shape VComp dynamics has still to be determined.

Dengue virus (DENV), yellow fever virus (YFV), Zika virus (ZIKV), and chikungunya virus (CHIKV) pose a major sanitary threat as they are responsible for hundreds of millions of human infections each year worldwide, leading to severe morbidity and mortality (Labeaud et al., 2011; Bhatt et al., 2013). These arboviruses are primarily transmitted to humans by *Aedes aegypti* mosquitoes, although the Asian tiger mosquito, *Aedes albopictus,* is often incriminated as a vector. Indeed, *Ae. albopictus* is an important vector of arboviruses as evidenced by vector competence laboratory assays and the detection of infected field specimens (Gratz, 2004; Paupy et al., 2009). Notably, *Ae. albopictus* was identified as the main vector during CHIKV outbreaks in Gabon (2007), Congo (2011) as well as in a major outbreak at La Réunion island (2006) (Schuffenecker et al., 2006; Bonilauri et al., 2008; Pagès et al., 2009; Paupy et al., 2012; Mombouli et al., 2013). In Europe, this vector species is incriminated for autochthonous circulation of CHIKV for instance in Italy (Venturi et al., 2017) and mainland France (Delisle et al., 2015). Vector competence studies established that vector competence of *Ae. albopictus* for CHIKV depends on genetic (*e.g.*, mosquito and virus genotype) and environmental (*e.g.*, temperature) factors (Tsetsarkin et al., 2007; Vazeille et al., 2007; Zouache et al., 2014; Sanchez-Vargas et al., 2019). Host viremia, approxymated by virus dose in the blood meal during artifical mosquito infectious feeding experiments, is another major factor that drives mosquito vector competence (Nguyet et al., 2013; Aubry et al., 2020). Vertebrate host viremia for CHIKV last 4 to 12 days with an increase in blood viral titer prior to symptoms appearance, up to a peak around 8 $\log_{10}$ infectious particles/mL followed by a decrease until virus clearance for most of the cases (Schwartz & Albert, 2010). Beyond non-human primates, an estimate of CHIKV human viremia dynamics is lacking due to limited longitudinal monitoring of infected patients, despite it could help to decipher the duration and magnitude of human infectiousness for mosquitoes (Labadie et al., 2010). In *Ae. albopictus*, two studies exposed mosquitoes to a range of CHIKV doses in the blood meal with varying outcomes on vector competence as estimated by mosquito infection and dissemination rate (Pesko et al., 2009; Hurk et al., 2010). Vector competence studies on *Ae. albopictus* from mainland France measured CHIKV transmission rate although upon exposure to a single dose, always above 6.5 $\log_{10}$ FFU/mL range which does not cover the range of human vireamia (Moutailler et al., 2009; Vega-Rua et al., 2013; Zouache et al., 2014; Vega-Rúa et al., 2015). In addition, these studies focused on a limited number of time points after CHIKV exposure that prevent to capture the dynamics of vector competence. A study monitoring intra-vector dynamics of CHIKV and its epidemiological relevance is still lacking, notably upon variations of vector competence major drivers such as virus dose.

Here, we studied intra-vector infection dynamics of a field-derived population of *Ae. albopictus* from Lyon metropolis exposed by artificial membrane feeding to a range of human viraemia-like CHIKV (La Réunion 06.21 isolate, East Central South Africa (ECSA) clade) doses, based on our model of human CHIKV viremia in the blood. Strains from ECSA clade carrying the same A226V mutation on enveloppe *E1* gene than 06.21 were identified from autochtonous cases in mainland France, supporting the choice of the 06.21 strain for this study

(Franke et al., 2019). Individual mosquitoes were analyzed from day 2 to day 20 post-exposure (dpe) to
determine infection, dissemination, and transmission rates by infectious titration in addition to the
quantification of CHIKV RNA load in the saliva. This allowed us to estimate CHIKV intra-vector dynamics and
the strength of vector infection, dissemination, and transmission barriers as well as the distribution of EIP
according to the virus dose in the blood meal. These data were implemented in the agent-based model *nosoi*
(Lequime et al., 2020) to estimate, using realistic vectorial capacity parameters, the epidemic potential of
CHIKV in a French population of *Ae. albopictus*. Our results improve our understanding on vector-virus
interactions and provides key informations to better anticipate and prevent CHIKV emergence in mainland
France.

Methods

Modelling chikungunya viraemia in humans

Chikungunya virus (CHIKV) loads in human blood along with the time course of infection in patients were
recovered from two studies. The first study monitored blood CHIKV viraemia from a retrospective cohort of
102 febrile patients in Bandung, West Java, Indonesia, between 2005 and 2009 (Riswari et al., 2015). The
second study assessed CHIKV RNA viremic profile from 36 sera from day 1 to day 7 of illness during a CHIKV
epidemic in 2009 in Thepa and Chana districts of Songkhla province, Thailand (Appassakij et al., 2013). For the
second study, the median value of the blood CHIKV RNA loads from a group of patients (n=2 to 21) per day of
illness was used. The viraemia data from RT-qPCR was expressed on the logarithmic scale to the base 10 before
model fitting. Wood's gamma-type function was used to model the viraemia dynamic. The function is given in
the following equation:

$$y(t) = at^b e^{-ct} \text{ (Eq. 1)}$$

where *y(t)* represents the level of viraemia in the blood at *t* days post-infection, with *a*, *b,* and *c*
representing constants linked to the viraemia dynamics (Islam et al., 2013). Viraemia data were originally
expressed in time pre- or post-symptom onset, while the model represents viraemia as a function of time post-
infection. A fixed arbitrary median intrinsic incubation period of 6 days was added to each viraemia time to
standardize the time scale between the data and the model. This fixed incubation period falls into the
estimated 2-10 days incubation range (Moloney et al., 2014) and was chosen to ensure that all observed
viraemia data occurred after infection. The model was fitted to the data using non-linear least-squares
regression implemented in the *nls* function in the R environment (RCoreTeam, 2022). This method proposed a
possible intra-human CHIKV viraemia dynamic with 95% confidence intervals.

CHIKV stock production and titration

The CHIKV strain 06.21 from the Indian Ocean lineage was isolated from a newborn serum sample with
neonatal encephalopathy in La Réunion island in 2005 (Schuffenecker et al., 2006). This strain was amplified in
Aedes albopictus cell line C6/36 as previously described (Raquin et al., 2015). CHIKV was inoculated at a
multiplicity of infection of 0.01 on *Ae. albopictus* C6/36 cells cultivated in Leibovitz's L-15 media (Gibco) with
10 % (v:v) 1X Tryptose Phosphate Broth (Gibco), 10 % (v:v) foetal bovine serum and 0.1 % (v:v) 10,000 units/mL
penicillin/streptomycin (Gibco). Cells were incubated for 3 days at 28 °C before the cell supernatant was
clarified by centrifugation for 5 min at 500 g and stored at -80 °C as aliquots. CHIKV infectious titer was
measured on C6/36 using fluorescent focus assay (Raquin et al., 2015). Briefly, 3x10^5 cells/well were inoculated
in 96-well plates (TPP) with 40 µL/well of viral inoculum (after culture media removal) and incubated for 1 h at
28 °C. 150 µL/well of a mix 1:1 L-15 media and 3.2% medium viscosity carboxymethyl cellulose (Sigma) were
added as an overlay before incubation of the cells for 3 days at 28 °C. After incubation, cells were fixed in 150
µL/well of 4 % paraformaldehyde for 20 min at room temperature (RT) and then rinsed 3 times in 100 µL/well
of 1X Dulbecco's phosphate-buffered saline (DPBS) (Gibco) prior to immune labelling. Cells were permeabilized
for 30 min in 50 µL/well of 0.3 % (v:v) Triton X-100 (Sigma) in 1X DPBS + 1 % Bovine Serum Albumin (BSA,
Sigma) at RT then rinsed 3 times in 100 µL/well of 1X DPBS. A Semliki Forest virus anticapsid antibody cross-
reacting with CHIKV was used as a primary antibody, diluted 1:600 in 1X DPBS + 1 % BSA (Greiser-Wilke et al.,
1989). Cells were incubated in 40 µL/well of primary antibody for 1 h at 37°C, rinsed 3 times in 100 µL/well of
1X DPBS then incubated in 40 µL/well of anti-mouse Alexa488 secondary antibody (Life Technologies) at 1:500

in 1X DPBS + 1 % BSA for 30 min at 37 °C. Cells were rinsed 3 times in 100 μL/well 1X DPBS, then once in 100
μL/well tap water, stored at 4 °C overnight before the enumeration of fluorescent foci under Zeiss Colibri 7
fluorescence microscope at 10X objective. The CHIKV infectious titer was expressed as the $\log_{10}$ fluorescent
focus unit (FFU) per mL. Plates were then stored at 4 °C protected from light to allow further reading. The
infectious titer of the neat CHIVK 06.21 stock was 8.63 $\log_{10}$ FFU/mL.

Mosquito colony maintenance

The Lyon metropolis population of *Aedes albopictus* originates from a field sampling of larvae in 2018 that
were brought back to insectary for rearing (Microbial Ecology lab, Lyon, France). Sampling locations included
Villeurbanne (N : 45°46'18990'' E : 4°53'24615'') and Pierre-Bénite (N :45°42'11534'' E : 4°49'28743'') in Lyon
metropolis area, mainland France. Mass rearing of the population under standard laboratory conditions (28 °C,
80% relative humidity, 16:8 hours light:dark cycles) using mice feeding (*Mus musculus*) allowed to maintain
genetic diversity, in accordance with the Institutional Animal Care and Use Committee from Lyon1 University
and the French Ministry for Higher Education and Research (Apafis #31807-2021052715018315). Prior to
infectious blood feeding, eggs were hatched for 1 h in dechlorinated tap water, and larvae were reared at 26
°C (12:12h light:dark cycle) at a density of 200 larvae in 23 x 34 x 7 cm plastic trays (Gilac) in 1.5 L of
dechlorinated tap water supplemented with 0.1 g of a 3:1 (TetraMin tropical fish food:Biover yeast) powder
every two days. Adults were maintained in 32.5 x 32.5 x 32.5 cm mesh cages (Bugdorm) at 28 °C, 80 % relative
humidity, 12:12h light:dark cycle with permanent access to 10% sugar solution.

Experimental mosquito exposure to CHIKV

Female mosquitoes (4 to 8-day old) from F_{10} generation were confined in 136 x 81 mm plastic feeding
boxes (Corning-Gosselin) with ~60 individual per box then transferred to the level 3 biosafety facility (SFR
Biosciences, AniRA-L3, Lyon Gerland) at 26 °C, 12:12 h light:dark cycle deprived from sugar solution 16 h before
the infectious blood meal. The blood meal was composed of a 2:1 (v:v) mixture of washed human erythrocytes
(from multiple anonymous donors collected by EFS AURA under the CODECOH agreement DC-2019-3507) and
viral suspension at several doses, and supplemented with 2 % (v:v) of 0.5 M ATP, pH 7 in water (Sigma). Feeders
(Hemotek) were covered with pig small intestine and filled with 3 mL of infectious blood mixture. Females
were allowed to feed for 1 h at 26 °C and blood aliquots were taken before (T0) and after (1h) the feeding and
stored at -80 °C for virus titration (Figure S1). Mosquitoes were anaesthetized on ice and fully engorged
females were transferred in 1-pint cardboard containers (10-25 females/container) and maintained with 10 %
sucrose. Cardboard containers were placed in 18 x 18 x 18 inches cages (BioQuip) and kept in climatic chambers
at 26 °C, 70 % humidity. Two independent vector competence experiments were conducted with 370 and 418
individuals mosquitoes per experiment, respectively. In a first experiment (n=370), mosquito body and head
infection were tested for the presence of infectious virus at 4 time points while in the second experiment
(n=418), mosquito head and saliva were analyzed at 10 time points.

Mosquito dissection and CHIKV detection

At the selected day post-exposure (dpe) to CHIKV, individual saliva were collected then the heads and
bodies were recovered. Prior to saliva collection, mosquitoes were anaesthetized on ice then legs and wings
were removed under a stereomicroscope. Individuals were placed on plastic plates maintained by double-
sided adhesive tape. The proboscis was inserted in a trimmed 10 μL filtered tip containing 10 μL of foetal
bovine serum (FBS) held above the mosquito by modelling clay (Heitmann et al., 2018). Two μL of 1 %
pilocarpine hydrochloride (Sigma) supplemented with 0.1 % Tween-20 (Sigma) in water were added on the
thorax of each mosquito to enhance salivation. Mosquitoes were allowed to salivate at 26 °C, 80 % relative
humidity for 1 h. The FBS that contains the saliva was expelled in an ice-cold tube filled with 150 μL of DMEM
media (Gibco) supplemented with antibiotics solution (Amphotericin B 2.5 μg/ml, Nystatin 1/100, Gentamicin
50 μg/ml, Penicillin 5 U/ml and Streptomycin 5 μg/ml (Gibco)). Following salivation, each mosquito's head and
body were separated using a pin holder with 0.15 mm minutien pins (FST). Heads and bodies were transferred
in individual grinding tubes (Qiagen) containing 500 μL of DMEM supplemented with antibiotics (see above)
and one 3-mm diameter tungsten bead (Qiagen). Samples were ground on a 96-well adapter set for 2 x 1 min,
30 Hz using a TissueLyser II (Qiagen), then stored at -80 °C. CHIKV detection was performed once on 40 μL of
undiluted (raw) saliva, head and body samples using fluorescent focus assay on C6/36 cells (see above). Each
mosquito sample was declared positive or negative for CHIKV in the presence or absence of a fluorescent

signal, respectively. Each 96-well plate contained positive (virus stock) and negative (raw grinding media) controls. Two independent persons examined each plate. Of note, saliva samples were deposited immediately (no freezing step) on C6/36 cells to maximize CHIKV detection. 30 µL of saliva sample were immediately mixed with 70 µL of TRIzol (Life Technologies) and stored at -80°C before RNA isolation. The rest of the samples were stored at -80 °C as a backup.

RNA isolation from saliva

Total RNA was isolated from 30 µL of saliva mixed with 70 µL TRIzol and then stored at -80°C, as described (Raquin et al., 2017). After thawing samples on ice, 20 µL chloroform (Sigma) were added. The tubes were mixed vigorously, incubated at 4 °C for 5 min and centrifuged at 17,000 G for 15 min, 4 °C. The upper phase was transferred in a new tube containing 60 µL isopropanol supplemented with 1 µL GlycoBlue (Life Technologies). Samples were mixed vigorously and stored at -80 °C overnight to allow RNA precipitation. After 15 min at 17,000 G, 4 °C, the supernatant was discarded, and the blue pellet was rinsed with 500 µL ice-cold 70 % ethanol in water. The samples were centrifuged at 17,000 G for 15 min, 4 °C, the supernatant was discarded, and the RNA pellet was allowed to dry for 10 min at room temperature. Ten µL RNAse-free water (Gibco) were added, and samples were incubated at 37 °C for 10 min to solubilize RNA prior to transfer in RNAse-free 96-well plates and storage at -80 °C.

CHIKV RNA load quantification in saliva

Total RNA (2 µL) isolated from individual mosquito saliva were used as template in a one-step TaqMan RT-qPCR assay. The QuantiTect Virus kit (Qiagen) was used to prepare the reaction mix in a final volume of 30 µL. The reaction solution consisted of 6 µL 5X master mix, 1.5 µL primers (forward 5'-CCCGGTAAGAGCGGTGAA-3' and reverse 5'-CTTCCGGTATGTCGATGGAGAT-3') and TaqMan probe (5'-6FAM-TGCGCCGTAGGGAACATGCC-BHQ1-3') (Hurk et al., 2010) mixed at 0.4 µM and 0.2 µM final concentration respectively, 0.3 µL 100X RT mix, 20.2 µL RNAse-free water (Gibco) and 2 µL template saliva RNA. RT-qPCR reaction was conducted on a Step One Plus machine (Applied) for 20 min at 50 °C (RT step), 5 min at 95°C (initial denaturation) and 40 cycles with 15 s at 95 °C and 45 s at 60 °C. Serial dilutions of CHIKV 06.21 synthetic RNA from 8 to 1 $\log_{10}$ copies/µL were used as an external standard to estimate CHIKV RNA copies in saliva samples. Each plate contained duplicates of standard synthetic RNA samples as well as negative controls and random saliva samples without reverse transcriptase (RT-) also in duplicate. Aliquots from the same standard RNA (thawed only once) were used for all the plates, and samples from a single time-point were measured on the same plate to allow sample comparison.

Statistical analyses

Mosquito infection (number of CHIKV-positive mosquito bodies / number engorged mosquitoes), dissemination (number of CHIKV-positive heads / number of CHIKV-positive bodies) and transmission rate (number of positive CHIKV-saliva / number of CHIKV-positive heads) were analysed by logistical regression and considered as binary response variables. The time (dpe) and virus dose ($\log_{10}$ FFU/mL) were considered continuous explanatory variables in a full factorial generalized linear model with a binomial error and a logit link function. Logistic regression assumes a saturation level of 100% and could not be used to model the relationship between the probability of transmission (response variable) and the time post-infection, the dose and their interaction (predictors). Therefore, we first estimated the saturation level (K) for each dose and subtracted the value N = number of mosquitoes with CHIKV dissemination x (100% - K) to the number of mosquitoes without virus in their saliva at each time post virus exposure to artificially remove mosquitoes that would never ultimately transmit the virus from the dataset. Logistic regression was then used on these transformed data to predict transmission rates across time post virus exposure and the virus dose (Figure S2). The statistical significance of the predictors' effects was assessed by comparing nested models using deviance analysis based on a chi-squared distribution. All the statistical analyses were performed in R Studio (Posit), and figures were created with the package *ggplot2* within the *Tidyverse* environment (Wickham et al., 2019). The R script used for this study is available, and supplementary table 1 summarizes the proportion of infected, disseminated and infectious mosquitoes for all the experiments conducted.

Epidemiological modelling using *nosoi*

A series of stochastic agent-based model simulations were performed using the R package *nosoi* and a specific branch available on *nosoi*'s GitHub page (https://github.com/slequime/nosoi/tree/fontaine) as previously done (Lequime et al., 2020). Briefly, 100 independent simulations were run in replicates for each condition. Each simulation started with one infected human and was run for 365 days or until the allowed number of infected individuals (100,000 humans or 1,000,000 mosquitoes, respectively) was reached. We considered transmission only between an infected mosquito and an uninfected human or between an infected human and an uninfected mosquito. Vertical and sexual transmission, and the impact of potential superinfection were ignored during the simulations. We assumed no particular structure within host and vector populations.

It was also considered that humans do not die from infection and leave the simulation after they clear the infection (here 12 days). Each human agent experienced a Poisson-like distribution of bites per day with a mean value manually set at 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or 60 based on field measurement of *Aedes albopictus* blood-feeding behaviour (Delatte et al., 2010). Human-to-mosquito transmission followed a time-post-infection-dependent probability function, identical for all human agents, computed from the human viraemic profile (Eq. 1, see above) and dose-response experiments (Eq. 2) :

$$p_{trans\ H \to M}(t) = \frac{1}{1+\exp\left(-\left(-11.1678+1.9929*\left(a*t^b*\exp(-c*t)\right)\right)\right)} \text{(Eq. 2)}$$

where based on our model, a = 0.01479, b = 7.21809 and c = 1.11915.

The daily survival probability of infected mosquito agents was set empirically at 0.85 (Favier et al., 2005; Fontaine et al., 2018). Human biting (only one per mosquito agent) was set at fixed dates depending on a gonotrophic cycle duration drawn for each mosquito in a truncated Poisson distribution with a mean of 4 days (no draws below 3). The mosquito-to-human transmission was determined for each mosquito agent based on its individual EIP value acting as a threshold for transmission (if time post-infection is greater or equal to the EIP value, the mosquito can transmit). However, a certain proportion (based on saturation parameter K, see above) of mosquitoes never transmitted. The individual EIP value was dependent on the virus dose that initiated the infection based on this equation (Lequime et al., 2020):

$$DD50 = \frac{\left(\log\left(\frac{-P}{P-1.001}\right) - \beta_0 - \beta_1 \times X1\right)}{\beta_2 + \beta_3 \times X1} \text{(Eq. 3)}$$

where P = 0.5 (i.e., the median transmission probability), β_0 is the Y-intercept value (-2.328973), β_1 (0.278953), β_2 (0.136746) and β_3 (0.003276) are model coefficients associated to the virus dose, time post virus exposure and their interaction, respectively. X1 represents the virus dose value.

Results

Estimating CHIKV viraemia in humans by modelisation of clinical data

Intra-human dynamic of CHIKV viraemia over time post-infection was approximated using time course of human CHIKV viraemia in individual patients from two studies (Appassakij et al., 2013; Riswari et al., 2015). CHIKV loads were assessed at 3 to 6 different time points, prior or post symptoms onset from the blood of 5 patients. The range of CHIKV viraemia duration among the 5 patients was 4-12 days, with a minimal and a maximum CHIKV load of 1 and 8.78 $\log_{10}$ PFU equivalent/mL, respectively. Of note, two patients displayed 1.04 and 3.25 $\log_{10}$ PFU equivalent/mL before symptoms onset, respectively. Modelling CHIKV viraemic profile using a Wood's gamma-type function indicates that mean viral load rapidly increases to peak after 6.45 days (*i.e.* within 24 h after symptoms onset) at 7.55 $\log_{10}$ PFU equivalent/mL (7.01-8.78 $\log_{10}$ equivalent PFU/mL depending on the patient) (Figure 1).

313
314
Figure 1 - Estimated time course of CHIKV load in human blood as a function of days
post-infection. A Wood's gamma-type function was used to model CHIKV viraemia
dynamics based on the time course of human viraemia data in 5 patients. The black line
represents model prediction using mean fit parameter values. Each dot represents a single
experimental measurement with colours corresponding to different patients. The vertical
gold line indicates the day of symptoms onset. The grey ribbon represents upper and lower
predicted values. Refers to raw data table "ChikV_Viremia_dynamic.txt" (see data
availability section).
323

The dose, but not the time, modulates mosquito infection rate

Female *Ae. albopictus* were seprately exposed to a human erythrocytes suspension containing three CHIKV
doses (3.94, 6.07 and 8.63 $\log_{10}$ FFU/mL) that span the estimated range of human viraemia as estimated above
(Figure 1). The mortality rate was very low regardless of time or oral dose, remaining below 5%. Mosquito
infection rate (IR) remained below 7 % (n=35 to 53 individuals tested) at 3.94 $\log_{10}$ FFU/mL, ranged from 65 to
85 % at 6.07 $\log_{10}$ FFU/mL (n=20 to 35) and raised above 94 % at 8.63 $\log_{10}$ FFU/mL (n=12 to 24) (Figure 2A). A
detail table of these data is presented in supplementary table S1. IR significantly increases with the dose but
it does not depend on the time post-exposure (Wald χ^2, P_{dose} = 1.1 x 10^{-6}, P_{time} = 0.9 and $P_{dose*time}$ = 0.17). As IR
depends on the virus dose but not on the time post-exposure, we fitted a logistic model to the data considering
CHIKV titer in the blood meal as a unique explanatory variable (Figure 2B). Dose-dependent IR describes a
sigmoid with a median oral infection dose ($OID_{50\%}$) of 5.6 $\log_{10}$ FFU/mL and an $OID_{25\%}$ and $OID_{75\%}$ of 5.05 $\log_{10}$
FFU/mL and 6.15 $\log_{10}$ FFU/mL, respectively. The oral infection saturation level was reached at about 7.5 $\log_{10}$
FFU/mL.
337

Figure 2 - Dose-dependent infection rate of *Ae. albopictus* mosquitoes exposed to CHIKV 06.21. (A) The mosquito infection rate corresponds to the proportion (in %) of mosquito bodies positive for CHIKV infection out of the total of engorged mosquitoes, measured at 2, 6, 9 and 14 days post-exposure for three CHIKV doses (3.94, 6.07 and 8.63 $\log_{10}$ FFU/mL) in the blood meal. The number of individuals analysed at each time point is indicated above the bars that represents the 95% confidence interval. (B) Mosquito infection rate as a function of CHIKV dose in the blood meal. Blue dots correspond to the observed infection rate upon the three CHIKV doses tested. Dot size is proportional to the number of mosquitoes tested. The black line was obtained by fitting a logistic model to the data. The grey ribbon indicates the 95 % confidence interval. The oral infectious dose (OID) to infect 25 %, 50 % and 75 % of the mosquitoes exposed to CHIKV is indicated with the associated standard error (in $\log_{10}$ FFU/mL). Refers to raw data table "Data_titer_EIPdyna_body_head_final.txt" (see data availability section).

Dose- and time-dependent mosquito dissemination dynamics

The proportion of CHIKV-positive heads among positive bodies (*i.e.,* mosquito dissemination rate, DIR) was analysed using virus dose, time post-exposure and their interaction as explanatory variables. At 2 days post-exposure, <50 % of the mosquitoes presented a disseminated infection for the doses 3.94 (n=1 individual tested) and 6.07 (n=18) $\log_{10}$ FFU/mL, whereas DIR was already above 80 % after 2 days in mosquitoes exposed to 8.63 $\log_{10}$ FFU/mL of CHIKV (n=23) (Figure 3A). Notably, DIR increases >80 % for all the three doses after 6 dpe (n=1 to 29 individual tested per time point and dose) (Figure 3A and Table S1). Although they are not in interaction, both time and dose impact DIR (Wald χ^2, P_{dose} = 8.29 x 10^{-6}, P_{time} = 1.75 x 10^{-6} and $P_{dose*time}$ = 0.83). The plateau was 100 % at doses 3.94 and 8.63 $\log_{10}$ FFU/mL and 95.6% at the dose 6.07 $\log_{10}$ FFU/mL (n=22/23). The time to reach 50 % dissemination in *Ae. albopictus* exposed to CHIKV was 7.5 days, 2.2 days and <1 day for 3.94, 6.07 and 8.63 $\log_{10}$ FFU/mL CHIKV doses in the blood meal, respectively. DIR was inferred from experimental data for a larger set of CHIKV dose ranging from 3 to 8 $\log_{10}$ FFU/mL (Figure 3B). All the CHIKV doses tested led to 100% dissemination within the 40 days range used for predictions, although a longer time is required to reach this plateau at the lowest dose.

369
370
Figure 3 - Dose-dependent dissemination rate of *Ae. albopictus* mosquitoes exposed
to CHIKV 06.21. (A) The mosquito dissemination rate corresponds to the number of CHIKV-
positive heads out of the infected (CHIKV-positive bodies) individuals, measured at 2, 6, 9
and 14 days post-exposure for three virus doses (3.94, 6.07 and 8.63 $\log_{10}$ FFU/mL) in the
blood meal. Dot size is proportional to the number of mosquitoes tested. No disseminated
females were detected at day 14 post-exposure at the 3.94 $\log_{10}$ FFU/mL dose. Logistic
regression was used to model the time-dependent effect of the virus dose on mosquito
dissemination rate. Lines correspond to fit values with their 95% confidence intervals
displayed as ribbons. The time needed to reach 50 % dissemination is 7.5, 2.2 and <1 day
for the 3.94, 6.07 and 8.63 $\log_{10}$ FFU/mL CHIKV doses, respectively, as indicated within
each facet label. (B) Predicted dissemination dynamics according to virus dose and time
post-exposure for a range of CHIKV blood meal titers (3 to 8 $\log_{10}$ FFU/mL). Refers to raw
data table "Data_titer_EIPdyna_final.txt" (see data availability section).

Time post-exposure modulates transmission rate and viral load in the saliva
The presence of infectious CHIKV particles in individual mosquito saliva collected by forced salivation
technique was monitored at a fine time scale. This allowed us to measure the transmission rate (TR) and
quantify individual viral load in saliva over time, and to estimate the extrinsic incubation period (EIP). Two virus
doses (5.68 and 8.06 $\log_{10}$ FFU/mL) were used to obtain a workable proportion of infectious mosquitoes at a
high number of time points that covers mosquito expected lifespan while remaining in the range of human
viraemia. From day 2 post-exposure, TR increases following a sigmoid shape, reaching a plateau of around 60%
for both doses (Figure 4A). TR was analysed using virus dose, time post-exposure and their interaction as
explanatory variables, and only time was significant (Wald χ^2, P_{time} = 0.0037, P_{dose} = 0.18, and $P_{dose*time}$ = 0.8).
Infectious saliva samples were detected as soon as day 2 post-exposure to CHIKV, with a 33 % TR at dose 5.68
$\log_{10}$ FFU/mL (n=1/3 individuals tested) and 14 % at dose 8.06 $\log_{10}$ FFU/mL (n=2/14). The time needed to
reach 50 % infectious mosquitoes (*i.e.* Extrinsic Incubation Period 50%, EIP$_{50\%}$) was 7.5 and 3.5 days for doses
5.68 and 8.06 $\log_{10}$ FFU/mL, respectively. A TR saturation at 100 % is a prerequisite to applying logistic
regression analysis to the data. Therefore, the proportion of mosquitoes that would not ultimately transmit
the virus was artificially removed from the dataset based on the predicted saturation level at each dose (*i.e.*,
40 % of mosquitoes without the virus in their saliva were removed at each time post-infection). Logistic
regression was used on these transformed data to predict TR across a range of doses over time (Figure S2).
To decipher if CHIKV load in the saliva could be associated with the virus dose mosquitoes were challenged
with, total RNA was isolated at each time point from the saliva of individual mosquitoes exposed to 5.68 or
8.06 $\log_{10}$ FFU/mL. Viral load was measured by TaqMan RT-qPCR assay and analysed according to virus dose
and time post-exposure. A high individual variation is noticed (up to 10,000 fold difference between
individuals), although CHIKV load in the saliva seems to decrease over time (Figure 4B). Only the time post-
exposure significantly affected viral load when considering saliva samples that were CHIKV-positive both in
qRT-PCR and in infectious titration (Anova, P_{time} = 0.006, P_{dose} = 0.66 and $P_{dose*time}$ = 0.94). Of note, both time

and virus dose affected viral load in the saliva when considering RNA-positive samples regardless of the
presence of infectious virus (Figure S3).
411

412
413
Figure 4 - Transmission dynamics of CHIKV 06.21 by *Ae. albopictus*. (A) Mosquito
transmission rate corresponds to the number of CHIKV-positive saliva out of the number
of CHIKV-positive heads collected at 2, 3, 4, 5, 6, 7, 10, 12, 14, and 20 days post-exposure
for two virus doses (5.68 and 8.06 $\log_{10}$ FFU/mL) in the blood meal. Dot size is proportional
to the number of saliva tested. (B) CHIKV RNA load of each saliva scored positive for
infectious CHIKV was measured by TaqMan RT-qPCR assay using a synthetic RNA as
standard, then expressed in $\log_{10}$ CHIKV RNA copies/saliva. Each dot represents a saliva
sample from a mosquito exposed to the indicated dose. Refers to raw data tables
"Data_titer_EIPdyna_final.txt" and "Data_CHIKV_RNA_load_saliva.txt" (see data
availability section).
424
Simulation of CHIKV epidemic upon dose-dependent intra-vector dynamics
A stochastic agent-based model was used to assess the epidemiological impact of within-host CHIKV
dynamics using the R package *nosoi*, as done previously for ZIKV (Lequime et al., 2020). Starting with one
infected human in a population of susceptible humans and mosquitoes, the model simulates CHIKV
transmissions according to human viraemia, its derived probability of mosquito infection, and virus
transmission timeliness (EIP). The model was run 100 independent times for a maximum of 365 days for a
range of eleven mean individual mosquito biting rates (1, 2, 3, 4, 5, 6, 7, 8, 9, 10 and 60 independent
mosquitoes biting per person per day). Simulations led to large outbreaks (>100 secondary infections) even
under a low mosquito biting rate (Figure 5A). The maximum threshold of mosquito infection was reached
regardless of biting intensity, although the time needed to reach this threshold and the inter-simulation
variation were higher at the lowest biting intensity (*i.e.* 1 bite per day) compare to other conditions (Figure
5B). Accordingly, secondary cases values distributions across simulations were narrow for all conditions except
at 1 bite per day reflecting the explosive nature of the outbreak. The mean secondary case values increases as
a function of the mosquito biting intensity with a mean (± SD) of 3.68 (± 1.92), 7.37 (± 2.71), 11.06 (± 3.32),
14.75 (± 3.84), 18.43 (± 4.29), 22.12 (± 4.7), 25.81 (± 5.07), 29.5 (± 5.43), 33.18 (± 5.75), 36.87 (± 6.07) and
221.15 (± 6.09) for 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 and 60 mosquito bites per person per day, respectively (Figure 5C).
441

442
443
Figure 5 - Influence of dose-dependent intra-mosquito CHIKV dynamics on outbreak
simulations with various levels of mosquito bites. Stochastic agent-based epidemiological
simulations considering within-vector infection dynamics on transmission probability
during mosquito-human infectious contacts were performed in 100 independent
replicates. A total of 11 mosquito bite intensity levels were tested: 1, 2, 3, 4, 5, 6, 7, 8, 9,
10 and 60 bites per human per day. (A) Stacked proportions of outbreak simulations
resulting in no secondary infected human host, < 100 and ≧ 100 infected human hosts. (B)
Cumulative number of infected humans over time. Each curve represents a simulation run.
(C) Violin plots showing the number of secondary cases densities for each intensity of
mosquito exposure. Refers to raw data tables "Compiled_results_run5.csv",
"cumulative_run5.csv" and "R0dist_run5.csv" (see data availability section).

Discussion

Vector competence (VComp) of *Aedes albopictus* for chikungunya virus (CHIKV) has been widely
studied, notably since the La Réunion outbreak in 2006. Studies outlined a strong impact of mosquito and
CHIKV genotype as well as temperature on *Ae. albopictus* potential for CHIKV transmission, including large
VComp variations among worldwide *Ae. albopictus* populations exposed to the highly transmissible La
Réunion 2006 CHIKV 06.21 isolate (Zouache et al., 2014; Mariconti et al., 2019; Gloria-Soria et al., 2020;
Vega-Rúa et al., 2020). However, knowledge gaps remain regarding intra-vector virus dynamics and its
impact on CHIKV epidemic potential notably regarding the virus dose. Our work contributes to fill these
gaps, providing important data on the interplay between CHIKV and *Ae. albopictus* as well as on the
vireamia-dependent human infectiousness for mosquitoes.

Dose and time-dependent barriers to CHIKV intra-vector dynamics
Ae. albopictus midgut infection is strongly influenced by CHIKV dose in the blood meal. Previous studies
documented an oral infectious dose for 50 % of the mosquitoes ($OID_{50\%}$) ranging from 1.7 to 3.52 $\log_{10}$
infectious particles per mL of blood for CHIKV 06.21 and thus lower than the 5.60 $\log_{10}$ FFU/mL $OID_{50\%}$
estimated in our *Ae. albopictus* population (Tsetsarkin et al., 2007; Pesko et al., 2009; Hurk et al., 2010).
ZIKV $OID_{50\%}$ was 5.62 $\log_{10}$ FFU/mL in *Ae. albopictus* while in *Ae. aegypti*, ZIKV and DENV $OID_{50\%}$ ranged
from 4.73 to 8.10 $\log_{10}$ FFU/mL and from 3.95 to 5.5 $\log_{10}$ PFU/mL, respectively (Nguyet et al., 2013; Aubry
et al., 2020; Lequime et al., 2020). The CHIKV time-independent infection rate and relatively low $OID_{50\%}$ in
Ae. albopictus suggests that primary infection of midgut cells is rapid and efficient. Such CHIKV midgut
infection pattern could be promoted by the presence of several potential midgut receptors for *Alphavirus*
entry in mosquitoes (Franz et al., 2015). Importantly, a 0.5 $\log_{10}$ FFU/mL increase in $OID_{50\%}$ during the
exponential phase results in twice as much infected mosquitoes, which could exacerbate outbreaks,
notably upon large vector densities. Therefore, dose-response experiments at a small dose range for each
virus and mosquito genotype of interest would significantly improve our understanding of vector
competence.
According to our results, CHIKV dissemination from the *Ae. albopictus* midgut depends on the
interaction between time post-exposure and virus dose. A previous study showed that at day 6 post-
exposure, CHIKV dissemination rate in *Ae. albopictus* increases with virus dose, being ~10%, ~50% and
>80% upon 3.6, 4.4 and 5.2 $\log_{10}$ PFU/mL in the blood meal respectively (Pesko et al., 2009). Here, we show
that if at least 6 days are needed to reach ~30 % dissemination upon 3.94 $\log_{10}$ FFU/mL, virus doses of 6.07
and 8.63 $\log_{10}$ FFU/mL led overall to ~90 % dissemination regardless of the time point (except at day 2 for
the 6.07 $\log_{10}$ FFU/mL dose where only 40 % dissemination was observed). Of note, CHIKV dissemination
rate at 3.94 $\log_{10}$ FFU/mL shall be interpreted with caution due to the low sample size that arise directly
from the low infection rate (1.9 to 6.5%, n = 37 to 53 individuals per time point). These results reveal the
ability of CHIKV to efficiently disseminate from the midgut, as modelling for a larger dose range estimates
that all infected mosquitoes eventually disseminate even at the lowest dose considered (2 $\log_{10}$ FFU/mL).
CHIKV and ZIKV present a nearly identical $OID_{50\%}$ suggesting similar midgut infection potential in *Ae.*
albopictus. However, ZIKV dissemination is slower and, to a minor extend, reaches lower value compared
to CHIKV (Lequime et al., 2020). This discrepancy might be due to viral replication in the midgut as
dissemination rate correlates with midgut viral load (Houk et al., 1981; Bosio et al., 1998; Dickson et al.,
2014; Vazeille et al., 2019; Carpenter et al., 2021). CHIKV dissemination might arise from an efficient
replication in the midgut tissue. Recently, the CHIKV 3' untranslated region was recently shown to promote
dissemination through an increased viral replication in the mosquito midgut (Merwaiss et al., 2020).
Ultimately, arboviruses infect and replicate in mosquito salivary glands, this step being essential to
allow virus transmission to the host (Vega-Rúa et al., 2015; Raquin & Lambrechts, 2017). Virus prevalence
in the head is often used as a proxy for transmission potential but it is likely an overestimate due to salivary
glands barriers, notably for CHIKV (Sanchez-Vargas et al., 2021). Our study shows that ~60 % of the
mosquitoes with disseminated infection eventually become infectious, this being an underestimate of
mosquito-to-host transmission potential due to the use of forced salivation technique (Gloria-Soria et al.,
2022). Moreover, transmission rate strongly depends on the time post-exposure. Previously, the time to
reach 50 % of infectious mosquitoes in the population ($EIP_{50\%}$) was estimated by a meta-analysis at 7 days
(± 1 day), based on dissemination data and for mosquitoes exposed to relatively high virus doses

(Christofferson et al., 2014). Despite no overall effect of virus dose on the transmission rate (P_{dose} = 0.18),
CHIKV $EIP_{50\%}$ was 7.5 and 3.5 days in mosquitoes exposed to 5.68 or 8.06 $\log_{10}$ FFU/mL, respectively,
suggesting that virus dose might influence CHIKV transmission. Increasing sample size and/or testing an
intermediate virus dose (*e.g.* 6 $\log_{10}$ FFU/mL) could help to better capture the impact of virus dose on
transmission rate and resolve this discrepancy. Interestingly, when considering all CHIKV-positive salivas
(including the ones with only CHIKV RNA but found negative during infectious titration), the CHIKV RNA
load in the saliva depends on time post-exposure and virus dose but not in interaction. Overall, the CHIKV
load in saliva seems higher at high dose but decreases overtime, questioning arbovirus-salivary glands
interaction. In a previous study, individual ZIKV-disseminated *Ae. aegypti* mosquitoes were offered
successive non-infectious blood meals in an attempt to monitor expelled virus during feeding in a non-
sacrificial manner. Authors observed an on/off presence of ZIKV in the blood meal for a single individual
over time; however, whether this is due to biological or methodological causes is unclear (Mayton et al.,
2021). In *Ae. albopictus*, up to 10,000-fold difference of CHIKV RNA load in the saliva was found between
individuals at a given time point and dose, in accordance with previous results (Dubrulle et al., 2009; Bohers
et al., 2020; Robison et al., 2020). No correlation was found between CHIKV titer in the salivary glands and
in the saliva, that might be linked with such inter-individual variations (Sanchez-Vargas et al., 2019). Despite
several studies that identified histological and genetical factors modulating viral infection in this tissue, it
is still unclear how and in which amount infectious virions are produced in salivary glands and then
transferred into the saliva over time (Ciano et al., 2014; Modahl et al., 2019; Chowdhury et al., 2021;
Sanchez-Vargas et al., 2021). Notably, viral particles in the saliva might use specific viral factors and/or
mosquito saliva proteins to persist in the saliva and promote their transmission (Pompon et al., 2017;
Marin-Lopez et al., 2021). This is key as virus titer in the mosquito inoculum is associated with viraemia
level and symptoms severity in mice and macaques models (Labadie et al., 2010; Zhang et al., 2022).
531

Genetic and environmental factors impacting intra-vector dynamics
VComp is a composite phenotype that also depends on the interaction between virus genotype,
mosquito genotype and temperature (Zouache et al., 2014). From a virus perspective, some CHIKV
mutations that impact VComp were already described and could be useful for epidemiological monitoring,
even if data suggest that overall CHIKV swarm maintains an intermediate mutation frequency to avoid
fitness loss in the mosquito (Coffey et al., 2014). Transcriptomics and quantitative genetics studies led to
the identification of mosquito genetic loci that constitute interesting targets towards engineered vector
control approaches, as shown for *Flaviviridae* (Bosio et al., 2000; Raquin et al., 2017; Aubry et al., 2020;
Merkling et al., 2020; Williams et al., 2020; Dong et al., 2022). In addition, mosquitoes host a microbiota
composed of bacteria, viruses, fungi and protists that have a major impact on vector's biology (Guégan et
al., 2018). Some of these micro-organisms are associated with a decrease in arbovirus transmission and
constitute interesting vector control tools, like *Wolbachia* or *Delftia tsuruhatensis* bacteria blocking DENV
and *Plasmodium* infection, respectively or insect-specific viruses modulating *Ae. aegypti* vector
competence for DENV (Olmo et al., 2018; 2023; Huang et al., 2023). Despite an antiviral activity of
Wolbachia against CHIKV in *Ae. aegypti*, as well a in *Ae. albopictus* C6/36 cell line, no CHIKV blocking was
detected in *Ae. albopictus* mosquitoes (Mousson et al., 2012; Raquin et al., 2015; Aliota et al., 2016).
Moreover, *Ae. albopictus* infection by CHIKV impacts mosquito bacterial community composition while
several *Ae. albopictus* symbionts were associated with an increase of CHIKV infection (Zouache et al., 2012;
Monteiro et al., 2019). The reason for this lack of microbiota-mediated antiviral blocking against CHIKV in
Ae. albopictus remains obscure, but could depend on mosquito and virus genotype and/or temperature as
Ae. albopictus microbiota composition depends on the temperature (Bellone et al., 2023). This interaction
should be further studied as it could impact arbovirus transmission, as suggested by models estimating
that the release of *Wolbachia*-infected *Ae. aegypti* against DENV will be less efficient upon long heatwaves
due to loss of *Wolbachia* infection upon high temperatures (Vásquez et al., 2023). With the exception of
DENV genotype (Fontaine et al., 2018), the impact of aforementioned factors was not tested on VComp
dynamics and it will be interesting to determine if these factors, beyond modulating VComp at discrete
time, impact the proportion of infectious mosquitoes over time.
559

560 **Human viremia and infectiousness to mosquitoes**

A large majority of VComp studies exposed *Ae. albopictus* to a high (>7 $\log_{10}$ plaque-forming units (PFU)/mL) CHIKV titer in the blood meal (Coffey et al., 2014). According to our estimate, CHIKV viraemia in humans can reach >7 $\log_{10}$ PFU equivalent per mL of blood, although this corresponds to the highest value measured within a short time window (<2 days). Our CHIKV viraemia estimate from human longitudinal data lasts from 4 to 12 days with a mean maximum titer of 7.55 $\log_{10}$ PFU/mL, which is largely supported by previous human and nonhuman studies (Lanciotti et al., 2007; Panning et al., 2008; Labadie et al., 2010; Schwartz & Albert, 2010). The mean human viraemia is above 3.94 $\log_{10}$ FFU/mL during 5.5 days, implying that humans are infectious to mosquitoes during more than half of their viraemia. These data are key to improve sanitary guidelines for the management of CHIKV infections and spread. However, major differences in CHIKV viraemia magnitude and length are observed between individual hosts, which can be associated with host and/or virus genotypes as observed in dengue virus (DENV)-infected patients (Nguyet et al., 2013). In addition, if artificial mosquito blood feeding limits variation in blood composition and promotes reproducibility, it does not necessary represent the native infectiousness of human-derived arbovirus for mosquitoes. This is partly linked to host plasma factors level (IgM, IgG, low-density lipoproteins or gamma-aminobutyric acid) while asymptomatic DENV cases are more infectious to mosquitoes than symptomatic counterparts at a given dose suggesting a link between host immune response and vector transmission (Nguyet et al., 2013; Duong et al., 2015; Wagar et al., 2017; Zhu et al., 2017). Moreover, for CHIKV, ~85 % infections are symptomatic with a median blood titer about 100-fold higher compare to asymptomatic carriers, although this difference was not statistically significant (Appassakij et al., 2013). Thus, improving epidemiological models by implementing the time-dependant human host infectiousness to mosquitoes represents an interesting lead to better anticipate and prevent CHIKV outbreaks. This notably prompts the need for viraemia monitoring over time in large groups of patients infected by arboviruses, using standardized virus titration procedures to facilitate comparisons and calibrate dose-response experiments in mosquitoes. This also requires other improvements of modelling strategies to account for mosquito and human populations structure, sanitary measures as well as *Ae. albopictus* tendency to take several consecutive blood meals. Indeed, virus dissemination increases with gonotrophic cycles and that successive bloodmeals are associated with a shortened EIP (Delatte et al., 2010; Armstrong et al., 2019; Fikrig & Harrington, 2021; Mulatier et al., 2023). Although those limitations could modulate the explosiveness of the outbreak and could be improved, our data support that CHIKV transmission potential of local *Ae. albopictus* is not a limiting factor for local CHIKV emergence and spread. Finally, such a standardized experimental design can be used to investigate the impact of additional (a)biotic factors on VComp dynamics.

Beyond virus dose and time post virus exposure, mosquito and virus genotype, mosquito microbiota and temperature are currently identified as major VComp drivers. Therefore, locally-acquired VComp data from area at risk for arbovirus circulation are needed. This could be achieved by exposing autochthonous field-derived mosquito populations to virus strains currently circulating (or at risk of introduction) upon a range of virus dose and temperature spanning the human viremia and the mosquito season, respectively, while controlling experimentally mosquito microbiota. This is unlikely to be done within a single experiment but will require the incremental acquisition of data sets for each factor, underlying the importance of experimental procedure standardization. Interestingly, modelisation of VComp based on available data could help to target a range of the factor's values to be tested experimentally, as exemplified with temperature (Shocket et al., 2020). These factors act together in interaction making difficult to dissociate their real impact on VComp in the field. Deciphering the complex interplay between each factor on VComp is challenging but feasible (Audsley et al., 2017) and holistic interaction studies could help to address this issue experimentally (Brinker et al., 2019). Beyond experimental conditions, our work underlines that monitoring intra-vector dynamics rather than end-point VComp is key to accurately quantify VComp variations and better estimate VCap (Christofferson & Mores, 2011). Several studies estimated VCap through the lens of mosquito density upon environmental variations (*e.g.* temperature, micro-climate, land cover) but with limited ecological (mosquito survival rate, biting rate, density per host) and VComp data, notably regarding intra-vector dynamics (Murdock et al., 2017; Wimberly et al., 2020; Peña-García et al., 2023). This highlights the current need for additional VComp and VCap-related studies using field-derived material, as well as increasing efforts between vector biology and modeling fields towards an integrative VCap estimation notably regarding intra-vector arbovirus dynamics. Altogether, this will improve vector control strategies and case management by health authorities.

Appendices

FigS1 - CHIKV infectious titer is stable upon a one hour incubation at 37°C in human erythrocytes
suspension. Refers to raw data table "blood_titration_FFU.txt" (see data availability section).
FigS2 - Rescaled mosquito transmission dynamics. Refers to raw data table
"Data_titer_EIPdyna_final.txt" (see data availability section).
FigS3 - Time-course of CHIKV load in mosquito saliva. Refers to raw data table
"Data_CHIKV_RNA_load_saliva.txt" (see data availability section).
TabS1 - Proportion of infected, disseminated and infectious mosquitoes over time according to the
dose of CHIKV in the blood meal. Refers to raw data table "Raw_data_viginier_et_al2023" (see data
availability section).
626

Acknowledgements

This project was funded by the scientific breakthrough project Micro-Be-Have (Microbial impact on
insect behavior) of the Université de Lyon within the program Investissements d'Avenir (ANR-11-IDEX-
0007; ANR-16-IDEX-0005). We thank the Equipex InfectioTron program and its project manager Isabelle
Weiss. We also thank all the members of the Micro-Be-Have consortium for insightful discussions. We
thank Anna-Bella Failloux and Patrick Mavingui for the CHIKV 06.21 isolate. We acknowledge the
contribution of SFR Biosciences (UAR3444/CNRS, US8/Inserm, ENS de Lyon, Universite Claude Bernard Lyon
1) AniRa biosafety level 3 platform (Marie-Pierre Confort) and plateau Analyse Génétique et Cellulaire
(Bariza Blanquier) facilities in Lyon. We also thank 2 anonymous reviewers for helpful comments on a
previous version of this manuscript.
637

Data, scripts, code, and supplementary information availability

639   Data, R scripts, supplementary information and main figures in full size are available online:
640 https://doi.org/10.5281/zenodo.8033668

Conflict of interest disclosure

642   The authors declare that they comply with the PCI rule of having no financial conflicts of interest in
643 relation to the content of the article.
644   Sebastian Lequime is a recommender for PCI infections.

Author contributions

BV: Investigation; Data curation; Validation
LC: Investigation; Data curation; Validation
CG: Investigation; Data curation; Validation
EM: Resources
CM: Resources
CVM: Funding acquisition; Resources; Supervision; Writing – review and editing
GM: Funding acquisition; Writing – review and editing
AF: Data curation; Formal analysis; Software; Visualization; Methodology; Writing – review and editing
SL: Data curation; Formal analysis; Software; Visualization; Methodology; Writing – review and editing
MR: Funding acquisition; Supervision; Writing – review and editing
FA: Funding acquisition; Project administration; Supervision; Writing – review and editing

Funding

Micro-Be-Have (Microbial impact on insect behavior) of the Université de Lyon within the program
Investissements d'Avenir (ANR-11-IDEX-0007; ANR-16-IDEX-0005).

References

Aliota MT, Walker EC, Yepes AU, Velez ID, Christensen BM, Osorio JE (2016) The *w*Mel Strain of *Wolbachia*
Reduces Transmission of Chikungunya Virus in *Aedes aegypti*. *PLoS neglected tropical diseases*, **10**,
e0004677. https://doi.org/10.1371/journal.pntd.0004677

Appassakij H, Khuntikij P, Kemapunmanus M, Wutthanarungsan R, Silpapojakul K (2013) Viremic profiles
in CHIKV-infected cases. *Transfusion*, **53**, 2567–2574. https://doi.org/10.1111/j.1537-
2995.2012.03960.x

Armstrong PM, Ehrlich HY, Magalhaes T, Miller MR, Conway PJ, Bransfield A, Misencik MJ, Gloria-Soria A,
Warren JL, Andreadis TG, Shepard JJ, Foy BD, Pitzer VE, Brackney DE (2019) Successive blood meals
enhance virus dissemination within mosquitoes and increase transmission potential. *Nature*
microbiology, 1–9. https://doi.org/10.1038/s41564-019-0619-y

Aubry F, Dabo S, Manet C, Filipović I, Rose NH, Miot EF, Martynow D, Baidaliuk A, Merkling SH, Dickson
LB, Crist AB, Anyango VO, Romero-Vivas CM, Vega-Rúa A, Dusfour I, Jiolle D, Paupy C, Mayanja MN,
Lutwama JJ, Kohl A, Duong V, Ponlawat A, Sylla M, Akorli J, Otoo S, Lutomiah J, Sang R, Mutebi J-P,
Cao-Lormeau V-M, Jarman RG, Diagne CT, Faye O, Faye O, Sall AA, McBride CS, Montagutelli X, Rašić
G, Lambrechts L (2020) Enhanced Zika virus susceptibility of globally invasive *Aedes aegypti*
populations. *Science*, **370**, 991–996. https://doi.org/10.1126/science.abd3663

Audsley MD, Ye YH, McGraw EA (2017) The microbiome composition of *Aedes aegypti* is not critical for
Wolbachia-mediated inhibition of dengue virus. *PLoS Neglected Tropical Diseases*, **11**, e0005426.
https://doi.org/10.1371/journal.pntd.0005426

Bellone R, Lechat P, Mousson L, Gilbart V, Piorkowski G, Bohers C, Merits A, Kornobis E, Reveillaud J,
Paupy C, Vazeille M, Martinet J-P, Madec Y, Lamballerie XD, Dauga C, Failloux A-B (2023) Climate
change and vector-borne diseases: a multi-omics approach of temperature-induced changes in the
mosquito. *Journal of Travel Medicine*, **30**. https://doi.org/10.1093/jtm/taad062

Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, Moyes CL, Drake JM, Brownstein JS, Hoen AG,
Sankoh O, Myers MF, George DB, Jaenisch T, Wint GRW, Simmons CP, Scott TW, Farrar JJ, Hay SI
(2013) The global distribution and burden of dengue. *Nature*, **496**, 504–507.
https://doi.org/10.1038/nature12060

Bohers C, Mousson L, Madec Y, Vazeille M, Rhim A, M'ghirbi Y, Bouattour A, Failloux A-B (2020) The
recently introduced *Aedes albopictus* in Tunisia has the potential to transmit chikungunya, dengue
and Zika viruses. *PLOS Neglected Tropical Diseases*, **14**, e0008475.
https://doi.org/10.1371/journal.pntd.0008475

Bonilauri P, Bellini R, Calzolari M, Angelini R, Venturi L, Fallacara F, Cordioli P, Angelini P, Venturelli C,
Merialdi G, Dottori M (2008) Chikungunya Virus in *Aedes albopictus*, Italy. *Emerging Infectious*
Diseases, **14**, 852–854. https://doi.org/10.3201/eid1405.071144

Bosio CF, Beaty BJ, Black WC (1998) Quantitative genetics of vector competence for dengue-2 virus in
Aedes aegypti. *The American Journal of Tropical Medicine and Hygiene*, **59**, 965–970.
https://doi.org/10.4269/ajtmh.1998.59.965

Bosio CF, Fulton RE, Salasek ML, Beaty BJ, Black WC (2000) Quantitative trait loci that control vector
competence for dengue-2 virus in the mosquito *Aedes aegypti*. *Genetics*, **156**, 687–698.
https://doi.org/10.1093/genetics/156.2.687

Brinker P, Fontaine MC, Beukeboom LW, Salles JF (2019) Host, Symbionts, and the Microbiome: The
Missing Tripartite Interaction. *Trends in Microbiology*, **27**, 480–488.
https://doi.org/10.1016/j.tim.2019.02.002

Carpenter A, Bryant WB, Santos SR, Clem RJ (2021) Infection of *Aedes aegypti* Mosquitoes with Midgut-
Attenuated Sindbis Virus Reduces, but Does Not Eliminate, Disseminated Infection. *Journal of*
Virology, **95**, e00136-21. https://doi.org/10.1128/jvi.00136-21

Chowdhury A, Modahl CM, Missé D, Kini RM, Pompon J (2021) High resolution proteomics of *Aedes*
aegypti salivary glands infected with either dengue, Zika or chikungunya viruses identify new virus
specific and broad antiviral factors. *Scientific Reports*, **11**, 23696. https://doi.org/10.1038/s41598-
021-03211-0

Christofferson RC, Chisenhall DM, Wearing HJ, Mores CN (2014) Chikungunya Viral Fitness Measures
within the Vector and Subsequent Transmission Potential. *PLoS ONE*, **9**, e110538.
https://doi.org/10.1371/journal.pone.0110538

Christofferson RC, Mores CN (2011) Estimating the Magnitude and Direction of Altered Arbovirus
Transmission Due to Viral Phenotype. *PLoS ONE*, **6**, e16298.
https://doi.org/10.1371/journal.pone.0016298

Ciano K, Saredy J, Bowers D (2014) Heparan Sulfate Proteoglycan: An Arbovirus Attachment Factor
Integral to Mosquito Salivary Gland Ducts. *Viruses*, **6**, 5182–5197. https://doi.org/10.3390/v6125182

Coffey LL, Failloux A-B, Weaver SC (2014) Chikungunya Virus–Vector Interactions. *Viruses*, **6**, 4628–4663.
https://doi.org/10.3390/v6114628

Delatte H, Desvars A, Bouétard A, Bord S, Gimonneau G, Vourc'h G, Fontenille D (2010) Blood-Feeding
Behavior of *Aedes albopictus*, a Vector of Chikungunya on La Réunion. *Vector-Borne and Zoonotic*
Diseases, **10**, 249–258. https://doi.org/10.1089/vbz.2009.0026

Delisle E, Rousseau C, Broche B, Leparc-Goffart I, L'Ambert G, Cochet A, Prat C, Foulongne V, Ferré JB,
Catelinois O, Flusin O, Tchernonog E, Moussion IE, Wiegandt A, Septfons A, Mendy A, Moyano MB,
Laporte L, Maurel J, Jourdain F, Reynes J, Paty MC, Golliot F (2015) Chikungunya outbreak in
Montpellier, France, September to October 2014. *Eurosurveillance*, **20**. https://doi.org/10.2807/1560-
7917.es2015.20.17.21108

Dickson LB, Sanchez-Vargas I, Sylla M, Fleming K, Black WC (2014) Vector competence in West African
Aedes aegypti Is Flavivirus species and genotype dependent. *PLoS neglected tropical diseases*, **8**,
e3153. https://doi.org/10.1371/journal.pntd.0003153

Dong Y, Dong S, Dizaji NB, Rutkowski N, Pohlenz T, Myles K, Dimopoulos G (2022) The *Aedes aegypti*
siRNA pathway mediates broad-spectrum defense against human pathogenic viruses and modulates
antibacterial and antifungal defenses. *PLoS Biology*, **20**, e3001668.
https://doi.org/10.1371/journal.pbio.3001668

Dubrulle M, Mousson L, Moutailler S, Vazeille M, Failloux A-B (2009) Chikungunya virus and *Aedes*
mosquitoes: saliva is infectious as soon as two days after oral infection. *PloS One*, **4**, e5895.
https://doi.org/10.1371/journal.pone.0005895

Duong V, Lambrechts L, Paul RE, Ly S, Lay RS, Long KC, Huy R, Tarantola A, Scott TW, Sakuntabhai A,
Buchy P (2015) Asymptomatic humans transmit dengue virus to mosquitoes. *Proceedings of the*
National Academy of Sciences, **112**, 14688–14693. https://doi.org/10.1073/pnas.1508114112

Favier C, Schmit D, Mller-Graf CDM, Cazelles B, Degallier N, Mondet B, Dubois MA (2005) Influence of
spatial heterogeneity on an emerging infectious disease: the case of dengue epidemics. *Proceedings*
of the Royal Society B: Biological Sciences, **272**, 1171–1177. https://doi.org/10.1098/rspb.2004.3020

Fikrig K, Harrington LC (2021) Understanding and interpreting mosquito blood feeding studies: the case of
Aedes albopictus. *Trends in Parasitology*, **37**, 959–975. https://doi.org/10.1016/j.pt.2021.07.013

Fontaine A, Lequime S, Moltini-Conclois I, Jiolle D, Leparc-Goffart I, Reiner RC, Lambrechts L (2018)
Epidemiological significance of dengue virus genetic variation in mosquito infection dynamics. *PLOS*
Pathogens, **14**, e1007187. https://doi.org/10.1371/journal.ppat.1007187

Franke F, Giron S, Cochet A, Jeannin C, Leparc-Goffart I, Valk H de, Jourdain F, Lamballerie X de, L'Ambert
G, Paty MC (2019) Autochthonous chikungunya and dengue fever outbreak in Mainland France, 2010-
2018. *European Journal of Public Health*, **29**. https://doi.org/10.1093/eurpub/ckz186.628

Franz AWE, Kantor AM, Passarelli AL, Clem RJ (2015) Tissue Barriers to Arbovirus Infection in Mosquitoes.
Viruses, **7**, 3741–3767. https://doi.org/10.3390/v7072795

Gloria-Soria A, Brackney DE, Armstrong PM (2022) Saliva collection via capillary method may
underestimate arboviral transmission by mosquitoes. *Parasites & Vectors*, **15**, 103.
https://doi.org/10.1186/s13071-022-05198-7

Gloria-Soria A, Payne AF, Bialosuknia SM, Stout J, Mathias N, Eastwood G, Ciota AT, Kramer LD,
Armstrong PM (2020) Vector Competence of *Aedes albopictus* Populations from the Northeastern
United States for Chikungunya, Dengue, and Zika Viruses. *The American Journal of Tropical Medicine*
and Hygiene. https://doi.org/10.4269/ajtmh.20-0874

Gratz NG (2004) Critical review of the vector status of *Aedes albopictus*. *Medical and Veterinary*
Entomology, **18**, 215–227. https://doi.org/10.1111/j.0269-283x.2004.00513.x

Greiser-Wilke I, Moennig V, Kaaden O-R, Figueiredo LTM (1989) Most Alphaviruses Share a Conserved
Epitopic Region on Their Nucleocapsid Protein. *Journal of General Virology*, **70**, 743–748.
https://doi.org/10.1099/0022-1317-70-3-743

Guégan M, Zouache K, Démichel C, Minard G, Van VT, Potier P, Mavingui P, Moro CV (2018) The mosquito
holobiont: fresh insight into mosquito-microbiota interactions. *Microbiome*, **6**, 49.
https://doi.org/10.1186/s40168-018-0435-2

Heitmann A, Jansen S, Lühken R, Leggewie M, Schmidt-Chanasit J, Tannich E (2018) Forced Salivation As a
Method to Analyze Vector Competence of Mosquitoes. *Journal of Visualized Experiments : JoVE*,
57980. https://doi.org/10.3791/57980

Houk EJ, Hardy JL, Presser SB, Kramer LD (1981) Dissemination Barriers for Western Equine
Encephalomyelitis Virus in *Culex tarsalis* Infected after Ingestion of Low Viral Doses. *The American
Journal of Tropical Medicine and Hygiene*, **30**, 190–197. https://doi.org/10.4269/ajtmh.1981.30.190

Huang W, Rodrigues J, Bilgo E, Tormo JR, Challenger JD, Cozar-Gallardo CD, Pérez-Victoria I, Reyes F,
Castañeda-Casado P, Gnambani EJ, Hien DF de S, Konkobo M, Urones B, Coppens I, Mendoza-Losana
A, Ballell L, Diabate A, Churcher TS, Jacobs-Lorena M (2023) *Delftia tsuruhatensis* TC1 symbiont
suppresses malaria transmission by anopheline mosquitoes. *Science*, **381**, 533–540.
https://doi.org/10.1126/science.adf8141

Hurk AF van den, Hall-Mendelin S, Pyke AT, Smith GA, Mackenzie JS (2010) Vector Competence of
Australian Mosquitoes for Chikungunya Virus. *Vector-Borne and Zoonotic Diseases*, **10**, 489–495.
https://doi.org/10.1089/vbz.2009.0106

Islam ZU, Bishop SC, Savill NJ, Rowland RRR, Lunney JK, Trible B, Doeschl-Wilson AB (2013) Quantitative
Analysis of Porcine Reproductive and Respiratory Syndrome (PRRS) Viremia Profiles from
Experimental Infection: A Statistical Modelling Approach. *PLoS ONE*, **8**, e83567.
https://doi.org/10.1371/journal.pone.0083567

Labadie K, Larcher T, Joubert C, Mannioui A, Delache B, Brochard P, Guigand L, Dubreil L, Lebon P, Verrier
B, Lamballerie X de, Suhrbier A, Cherel Y, Grand RL, Roques P (2010) Chikungunya disease in
nonhuman primates involves long-term viral persistence in macrophages. *Journal of Clinical
Investigation*, **120**, 894–906. https://doi.org/10.1172/jci40104

Labeaud AD, Bashir F, King CH (2011) Measuring the burden of arboviral diseases: the spectrum of
morbidity and mortality from four prevalent infections. *Population Health Metrics*, **9**, 1.
https://doi.org/10.1186/1478-7954-9-1

Lanciotti RS, Kosoy OL, Laven JJ, Panella AJ, Velez JO, Lambert AJ, Campbell GL (2007) Chikungunya virus
in US travelers returning from India, 2006. *Emerging Infectious Diseases*, **13**, 764–767.
https://doi.org/10.3201/eid1305.070015

Lequime S, Bastide P, Dellicour S, Lemey P, Baele G (2020) nosoi: A stochastic agent-based transmission
chain simulation framework in r. *Methods in Ecology and Evolution*, **11**, 1002–1007.
https://doi.org/10.1111/2041-210x.13422

Lequime S, Dehecq J-S, Matheus S, Laval F de, Almeras L, Briolant S, Fontaine A (2020) Modeling intra-
mosquito dynamics of Zika virus and its dose-dependence confirms the low epidemic potential of
Aedes albopictus. *PLOS Pathogens*, **16**, e1009068. https://doi.org/10.1371/journal.ppat.1009068

Mariconti M, Obadia T, Mousson L, Malacrida A, Gasperi G, Failloux A-B, Yen P-S (2019) Estimating the
risk of arbovirus transmission in Southern Europe using vector competence data. *Scientific Reports*, **9**,
17852. https://doi.org/10.1038/s41598-019-54395-5

Marin-Lopez A, Jiang J, Wang Y, Cao Y, MacNeil T, Hastings AK, Fikrig E (2021) *Aedes aegypti* SNAP and a
calcium transporter ATPase influence dengue virus dissemination. *PLoS Neglected Tropical Diseases*,
15, e0009442. https://doi.org/10.1371/journal.pntd.0009442

Mayton EH, Hernandez HM, Vitek CJ, Christofferson RC (2021) A Method for Repeated, Longitudinal
Sampling of Individual *Aedes aegypti* for Transmission Potential of Arboviruses. *Insects*, **12**, 292.
https://doi.org/10.3390/insects12040292

Merkling SH, Raquin V, Dabo S, Henrion-Lacritick A, Blanc H, Moltini-Conclois I, Frangeul L, Varet H, Saleh
M-C, Lambrechts L (2020) Tudor-SN Promotes Early Replication of Dengue Virus in the *Aedes aegypti*
Midgut. *iScience*, **23**, 100870. https://doi.org/10.1016/j.isci.2020.100870

Merwaiss F, Filomatori CV, Susuki Y, Bardossy ES, Alvarez DE, Saleh M-C (2020) Chikungunya virus
replication rate determines the capacity of crossing tissue barriers in mosquitoes. *Journal of Virology*,
95. https://doi.org/10.1128/jvi.01956-20

Modahl C, Chowdhury A, Oliveira F, Kini RM, Pompon J (2019) Salivary gland RNA-seq from arbovirus-
infected *Aedes aegypti* and *Aedes albopictus* provides insights into virus transmission. *Access*
Microbiology, **1**. https://doi.org/10.1099/acmi.imav2019.po0039

Moloney RM, Kmush B, Rudolph KE, Cummings DAT, Lessler J (2014) Incubation Periods of Mosquito-
Borne Viral Infections: A Systematic Review. *The American Journal of Tropical Medicine and Hygiene*,
90, 882–891. https://doi.org/10.4269/ajtmh.13-0403

Mombouli J-V, Bitsindou P, Elion DOA, Grolla A, Feldmann H, Niama FR, Parra H-J, Munster VJ (2013)
Chikungunya Virus Infection, Brazzaville, Republic of Congo, 2011. *Emerging Infectious Diseases*, **19**,
1542–1543. https://doi.org/10.3201/eid1909.130451

Monteiro VVS, Navegantes-Lima KC, Lemos AB de, Silva GL da, Gomes R de S, Reis JF, Junior LCR, Silva OS
da, Romão PRT, Monteiro MC (2019) *Aedes*–Chikungunya Virus Interaction: Key Role of Vector
Midguts Microbiota and Its Saliva in the Host Infection. *Frontiers in Microbiology*, **10**, 492.
https://doi.org/10.3389/fmicb.2019.00492

Mousson L, Zouache K, Arias-Goeta C, Raquin V, Mavingui P, Failloux A-B (2012) The native *Wolbachia*
symbionts limit transmission of dengue virus in *Aedes albopictus*. *PLoS neglected tropical diseases*, **6**,
e1989. https://doi.org/10.1371/journal.pntd.0001989

Moutailler S, Barré H, Vazeille M, Failloux A (2009) Recently introduced *Aedes albopictus* in Corsica is
competent to Chikungunya virus and in a lesser extent to dengue virus. *Tropical Medicine &*
International Health, **14**, 1105–1109. https://doi.org/10.1111/j.1365-3156.2009.02320.x

Mulatier M, Boullis A, Dollin C, Cebrián-Torrejón G, Vega-Rúa A (2023) Chikungunya Virus Infection and
Gonotrophic Cycle Shape *Aedes aegypti* Oviposition Behavior and Preferences. *Viruses*, **15**, 1043.
https://doi.org/10.3390/v15051043

Murdock CC, Evans MV, McClanahan TD, Miazgowicz KL, Tesla B (2017) Fine-scale variation in
microclimate across an urban landscape shapes variation in mosquito population dynamics and the
potential of *Aedes albopictus* to transmit arboviral disease. *PLoS Neglected Tropical Diseases*, **11**,
e0005640. https://doi.org/10.1371/journal.pntd.0005640

Nguyet MN, Duong THK, Trung VT, Nguyen THQ, Tran CNB, Long VT, Dui LT, Nguyen HL, Farrar JJ, Holmes
EC, Rabaa MA, Bryant JE, Nguyen TT, Nguyen HTC, Nguyen LTH, Pham MP, Nguyen HT, Luong TTH,
Wills B, Nguyen CVV, Wolbers M, Simmons CP (2013) Host and viral features of human dengue cases
shape the population of infected and infectious *Aedes aegypti* mosquitoes. *Proceedings of the*
National Academy of Sciences of the United States of America, **110**, 9072–9077.
https://doi.org/10.1073/pnas.1303395110

Olmo RP, Ferreira AGA, Izidoro-Toledo TC, Aguiar ERGR, Faria IJS de, Souza KPR de, Osório KP, Kuhn L,
Hammann P, Andrade EG de, Todjro YM, Rocha MN, Leite THJF, Amadou SCG, Armache JN, Paro S,
Oliveira CD de, Carvalho FD, Moreira LA, Marois E, Imler J-L, Marques JT (2018) Control of dengue
virus in the midgut of *Aedes aegypti* by ectopic expression of the dsRNA-binding protein Loqs2.
Nature Microbiology, **3**, 1385–1393. https://doi.org/10.1038/s41564-018-0268-6

Olmo RP, Todjro YMH, Aguiar ERGR, Almeida JPP de, Ferreira FV, Armache JN, Faria IJS de, Ferreira AGA,
Amadou SCG, Silva ATS, Souza KPR de, Vilela APP, Babarit A, Tan CH, Diallo M, Gaye A, Paupy C,
Obame-Nkoghe J, Visser TM, Koenraadt CJM, Wongsokarijo MA, Cruz ALC, Prieto MT, Parra MCP,
Nogueira ML, Avelino-Silva V, Mota RN, Borges MAZ, Drumond BP, Kroon EG, Recker M, Sedda L,
Marois E, Imler J-L, Marques JT (2023) Mosquito vector competence for dengue is modulated by
insect-specific viruses. *Nature Microbiology*, **8**, 135–149. https://doi.org/10.1038/s41564-022-01289-
4

Pagès F, Peyrefitte CN, Mve MT, Jarjaval F, Brisse S, Iteman I, Gravier P, Tolou H, Nkoghe D, Grandadam
M (2009) Aedes albopictus Mosquito: The Main Vector of the 2007 Chikungunya Outbreak in Gabon.
PLoS ONE, **4**, e4691. https://doi.org/10.1371/journal.pone.0004691

Panning M, Grywna K, Esbroeck M van, Emmerich P, Drosten C (2008) Chikungunya fever in travelers
returning to Europe from the Indian Ocean region, 2006. *Emerging Infectious Diseases*, **14**, 416–422.
https://doi.org/10.3201/eid1403.070906

Paupy C, Delatte H, Bagny L, Corbel V, Fontenille D (2009) *Aedes albopictus*, an arbovirus vector: from the
darkness to the light. *Microbes and Infection / Institut Pasteur*, **11**, 1177–1185.
https://doi.org/10.1016/j.micinf.2009.05.005

Paupy C, Kassa FK, Caron M, Nkoghé D, Leroy EM (2012) A Chikungunya Outbreak Associated with the
Vector *Aedes albopictus* in Remote Villages of Gabon. *Vector-Borne and Zoonotic Diseases*, **12**, 167–
169. https://doi.org/10.1089/vbz.2011.0736

Peña-García VH, Luvall JC, Christofferson RC (2023) Arbovirus Transmission Predictions Are Affected by
Both Temperature Data Source and Modeling Methodologies across Cities in Colombia.
Microorganisms, **11**, 1249. https://doi.org/10.3390/microorganisms11051249

Pesko K, Westbrook CJ, Mores CN, Lounibos LP, Reiskind MH (2009) Effects of Infectious Virus Dose and
Bloodmeal Delivery Method on Susceptibility of *Aedes aegypti* and *Aedes albopictus* to Chikungunya
Virus. *Journal of Medical Entomology*, **46**, 395–399. https://doi.org/10.1603/033.046.0228

Pompon J, Manuel M, Ng GK, Wong B, Shan C, Manokaran G, Soto-Acosta R, Bradrick SS, Ooi EE, Missé D,
Shi P-Y, Garcia-Blanco MA (2017) Dengue subgenomic flaviviral RNA disrupts immunity in mosquito
salivary glands to increase virus transmission. *PLOS Pathogens*, **13**, e1006535.
https://doi.org/10.1371/journal.ppat.1006535

Raquin V, Lambrechts L (2017) Dengue virus replicates and accumulates in *Aedes aegypti* salivary glands.
Virology, **507**, 75–81. https://doi.org/10.1016/j.virol.2017.04.009

Raquin V, Merkling SH, Gausson V, Moltini-Conclois I, Frangeul L, Varet H, Dillies M-A, Saleh M-C,
Lambrechts L (2017) Individual co-variation between viral RNA load and gene expression reveals novel
host factors during early dengue virus infection of the *Aedes aegypti* midgut. *PLoS Neglected Tropical*
Diseases, **11**, e0006152. https://doi.org/10.1371/journal.pntd.0006152

Raquin V, Moro CV, Saucereau Y, Tran F-H, Potier P, Mavingui P (2015) Native *Wolbachia* from *Aedes*
albopictus Blocks Chikungunya Virus Infection *In Cellulo*. *PLOS ONE*, **10**, e0125066.
https://doi.org/10.1371/journal.pone.0125066

RCoreTeam (2022) R Core Team (2021) R: A Language and Environment for Statistical Computing.

Riswari SF, Ma'roef CN, Djauhari H, Kosasih H, Perkasa A, Yudhaputri FA, Artika IM, Williams M, Ven A van
der, Myint KS, Alisjahbana B, Ledermann JP, Powers AM, Jaya UA (2015) Study of viremic profile in
febrile specimens of chikungunya in Bandung, Indonesia. *Journal of clinical virology : the official*
publication of the Pan American Society for Clinical Virology, **74**, 61–5.
https://doi.org/10.1016/j.jcv.2015.11.017

Robison A, Young MC, Byas AD, Rückert C, Ebel GD (2020) Comparison of Chikungunya Virus and Zika
Virus Replication and Transmission Dynamics in *Aedes aegypti* Mosquitoes. *The American journal of*
tropical medicine and hygiene. https://doi.org/10.4269/ajtmh.20-0143

Sanchez-Vargas I, Harrington LC, Black WC, Olson KE (2019) Analysis of Salivary Glands and Saliva from
Aedes albopictus and *Aedes aegypti* Infected with Chikungunya Viruses. *Insects*, **10**, 39.
https://doi.org/10.3390/insects10020039

Sanchez-Vargas I, Olson KE, Black WC (2021) The Genetic Basis for Salivary Gland Barriers to Arboviral
Transmission. *Insects*, **12**, 73. https://doi.org/10.3390/insects12010073

Schuffenecker I, Iteman I, Michault A, Murri S, Frangeul L, Vaney M-C, Lavenir R, Pardigon N, Reynes J-M,
Pettinelli F, Biscornet L, Diancourt L, Michel S, Duquerroy S, Guigon G, Frenkiel M-P, Bréhin A-C,
Cubito N, Desprès P, Kunst F, Rey FA, Zeller H, Brisse S (2006) Genome Microevolution of Chikungunya
Viruses Causing the Indian Ocean Outbreak. *PLoS Medicine*, **3**, e263.
https://doi.org/10.1371/journal.pmed.0030263

Schwartz O, Albert ML (2010) Biology and pathogenesis of chikungunya virus. *Nature Reviews*
Microbiology, **8**, 491–500. https://doi.org/10.1038/nrmicro2368

Shocket MS, Verwillow AB, Numazu MG, Slamani H, Cohen JM, Moustaid FE, Rohr J, Johnson LR,
Mordecai EA (2020) Transmission of West Nile and five other temperate mosquito-borne viruses
peaks at temperatures between 23°C and 26°C. *eLife*, **9**, e58511. https://doi.org/10.7554/elife.58511

Smith DL, Battle KE, Hay SI, Barker CM, Scott TW, McKenzie FE (2012) Ross, Macdonald, and a Theory for
the Dynamics and Control of Mosquito-Transmitted Pathogens. *PLoS Pathogens*, **8**, e1002588.
https://doi.org/10.1371/journal.ppat.1002588

Tsetsarkin KA, Vanlandingham DL, McGee CE, Higgs S (2007) A single mutation in chikungunya virus
affects vector specificity and epidemic potential. *PLoS pathogens*, **3**, e201.
https://doi.org/10.1371/journal.ppat.0030201

Vásquez VN, Kueppers LM, Rašić G, Marshall JM (2023) *w*Mel replacement of dengue-competent
mosquitoes is robust to near-term climate change. *Nature Climate Change*, **13**, 848–855.
https://doi.org/10.1038/s41558-023-01746-w

Vazeille M, Madec Y, Mousson L, Bellone R, Barré-Cardi H, Sousa CA, Jiolle D, Yébakima A, Lamballerie X
de, Failloux A-B (2019) Zika virus threshold determines transmission by European *Aedes albopictus*
mosquitoes. *Emerging Microbes & Infections*, **8**, 1668–1678.
https://doi.org/10.1080/22221751.2019.1689797

Vazeille M, Moutailler S, Coudrier D, Rousseaux C, Khun H, Huerre M, Thiria J, Dehecq J-S, Fontenille D,
Schuffenecker I, Despres P, Failloux A-B (2007) Two Chikungunya Isolates from the Outbreak of La
Reunion (Indian Ocean) Exhibit Different Patterns of Infection in the Mosquito, *Aedes albopictus*. *PLoS*
ONE, **2**, e1168. https://doi.org/10.1371/journal.pone.0001168

Vega-Rúa A, Lourenço-de-Oliveira R, Mousson L, Vazeille M, Fuchs S, Yébakima A, Gustave J, Girod R,
Dusfour I, Leparc-Goffart I, Vanlandingham DL, Huang Y-JS, Lounibos LP, Ali SM, Nougairede A,
Lamballerie X de, Failloux A-B (2015) Chikungunya Virus Transmission Potential by Local *Aedes*
Mosquitoes in the Americas and Europe. *PLOS Neglected Tropical Diseases*, **9**, e0003780.
https://doi.org/10.1371/journal.pntd.0003780

Vega-Rúa A, Marconcini M, Madec Y, Manni M, Carraretto D, Gomulski LM, Gasperi G, Failloux A-B,
Malacrida AR (2020) Vector competence of *Aedes albopictus* populations for chikungunya virus is
shaped by their demographic history. *Communications Biology*, **3**, 326.
https://doi.org/10.1038/s42003-020-1046-6

Vega-Rúa A, Schmitt C, Bonne I, Locker JK, Failloux A-B (2015) Chikungunya Virus Replication in Salivary
Glands of the Mosquito *Aedes albopictus*. *Viruses*, **7**, 5902–5907. https://doi.org/10.3390/v7112917

Vega-Rua A, Zouache K, Caro V, Diancourt L, Delaunay P, Grandadam M, Failloux A-B (2013) High
Efficiency of Temperate *Aedes albopictus* to Transmit Chikungunya and Dengue Viruses in the
Southeast of France. *PLoS ONE*, **8**, e59716. https://doi.org/10.1371/journal.pone.0059716

Venturi G, Luca MD, Fortuna C, Remoli ME, Riccardo F, Severini F, Toma L, Manso MD, Benedetti E,
Caporali MG, Amendola A, Fiorentini C, Liberato CD, Giammattei R, Romi R, Pezzotti P, Rezza G, Rizzo
C (2017) Detection of a chikungunya outbreak in Central Italy, August to September 2017.
Eurosurveillance, **22**, 17–00646. https://doi.org/10.2807/1560-7917.es.2017.22.39.17-00646

Viglietta M, Bellone R, Blisnick AA, Failloux A-B (2021) Vector Specificity of Arbovirus Transmission.
Frontiers in Microbiology, **12**, 773211. https://doi.org/10.3389/fmicb.2021.773211

Wagar ZL, Tree MO, Mpoy MC, Conway MJ (2017) Low density lipopolyprotein inhibits *flavivirus*
acquisition in *Aedes aegypti*. *Insect Molecular Biology*, **26**, 734–742.
https://doi.org/10.1111/imb.12334

Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, Grolemund G, Hayes A, Henry L,
Hester J, Kuhn M, Pedersen T, Miller E, Bache S, Müller K, Ooms J, Robinson D, Seidel D, Spinu V,
Takahashi K, Vaughan D, Wilke C, Woo K, Yutani H (2019) Welcome to the Tidyverse. *Journal of Open*
Source Software, **4**, 1686. https://doi.org/10.21105/joss.01686

Williams AE, Franz AWE, Reid WR, Olson KE (2020) Antiviral Effectors and Gene Drive Strategies for
Mosquito Population Suppression or Replacement to Mitigate Arbovirus Transmission by *Aedes*
aegypti. *Insects*, **11**, 52. https://doi.org/10.3390/insects11010052

Wimberly MC, Davis JK, Evans MV, Hess A, Newberry PM, Solano-Asamoah N, Murdock CC (2020) Land
cover affects microclimate and temperature suitability for arbovirus transmission in an urban
landscape. *PLoS Neglected Tropical Diseases*, **14**, e0008614.
https://doi.org/10.1371/journal.pntd.0008614

Zhang Y, Yan H, Li X, Zhou D, Zhong M, Yang J, Zhao B, Fan X, Fan J, Shu J, Lu M, Jin X, Zhang E, Yan H
(2022) A high-dose inoculum size results in persistent viral infection and arthritis in mice infected with
chikungunya virus. *PLoS Neglected Tropical Diseases*, **16**, e0010149.
https://doi.org/10.1371/journal.pntd.0010149

Zhu Y, Zhang R, Zhang B, Zhao T, Wang P, Liang G, Cheng G (2017) Blood meal acquisition enhances
arbovirus replication in mosquitoes through activation of the GABAergic system. *Nature*
Communications, **8**, 1262. https://doi.org/10.1038/s41467-017-01244-6

Zouache K, Fontaine A, Vega-Rua A, Mousson L, Thiberge J-M, Lourenco-De-Oliveira R, Caro V,
Lambrechts L, Failloux A-B (2014) Three-way interactions between mosquito population, viral strain
and temperature underlying chikungunya virus transmission potential. *Proceedings of the Royal*
Society B: Biological Sciences, **281**, 20141078. https://doi.org/10.1098/rspb.2014.1078

Zouache K, Michelland RJ, Failloux A-B, Grundmann GL, Mavingui P (2012) Chikungunya virus impacts the
diversity of symbiotic bacteria in mosquito vector. *Molecular Ecology*, **21**, 2297–2309.
https://doi.org/10.1111/j.1365-294x.2012.05526.x

986