Title: Estimating the seroincidence of scrub typhus using antibody dynamics following infection

Authors: Kristen Aiemjoy[1,2], Nishan Kutawal[3], Krista Vaidya[1], Sony Shrestha[3], Melina Thapa[3], Peter Teunis[4], Isaac I. Bogoch[5], Paul Trowbridge[6], Pacharee Kantipong[7], Stuart D. Blacksell[8,9], George M Varghese[10], Richard Maude[8,9,11,12], Dipesh Tamrakar[3], Jason R. Andrews[13]

Affiliations:
1. Department of Public Health Sciences, University of California Davis School of Medicine, Davis, CA, USA
2. Department of Microbiology and Immunology, Faculty of Tropical Medicine, Mahidol University, Bangkok Thailand
3. Dhulikhel Hospital, Kathmandu University Hospital, Dhulikhel, Nepal
4. Center for Global Safe Water, Sanitation and Hygiene, Hubert Department of Global Health, Rollins School of Public Health, Emory University, Atlanta, GA, USA
5. Department of Medicine, University of Toronto, Toronto, ON, Canada
6. Michigan State University School of Human Medicine, Grand Rapids, MI, USA
7. Department of Internal Medicine, Chiangrai Prachanukroh Hospital, Chiang Rai, Thailand
8. Mahidol-Oxford Tropical Medicine Research Unit, Faculty of Tropical Medicine, Mahidol University, Bangkok Thailand
9. Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine, University of Oxford, Oxford, UK
10. Department of Infectious Diseases, Christian Medical College, Vellore, India
11. The Open University, Milton Keynes, UK
12. School of Public Health, Li Ka Shing Faculty of Medicine, University of Hong Kong, Hong Kong Department of Medicine, Stanford University School of Medicine, Stanford, CA, USA
13. Division of Infectious Diseases and Geographic Medicine, Stanford University School of Medicine, Stanford, CA, USA

Corresponding Author:
Kristen Aiemjoy
Assistant Professor
Division of Epidemiology
Department of Public Health Sciences,
University of California Davis School of Medicine, Davis, CA, USA
kaiemjoy@ucdavis.edu

Word Count: 2620

Abstract

Introduction: Scrub typhus is an acute febrile illness caused by the bacterium *Orientia tsutsugamushi.* Characterizing the population-level burden of scrub typhus is challenging due to the lack of accessible and accurate diagnostics. We describe a new approach using information about seroresponse after infection to generate population-level scrub typhus seroincidence estimates.

Methods: We use data from two clinical studies of scrub typhus patients enrolled in Chiang Rai, Thailand, and Vellore, India, and representative population data from two serosurveys in and around the Kathmandu valley, Nepal, and Vellore, India. The samples were tested for IgM and IgG responses to *Orientia tsutsugamushi*-derived recombinant 56-kDa antigen using commercial ELISA kits. We used Bayesian hierarchical models to fit two-phase models to the antibody responses from scrub typhus cases and used the joint distributions of the peak antibody titers and decay rates to estimate population-level incidence rates in the cross-sectional serosurveys. We compared this new method to a traditional cut-off-based approach for estimating seroincidence.

Results: Among 18 to 29-year-olds, the seroincidence of scrub typhus was 886 (95% CI 432-1817) per 100,000 person-years in India and 945 (95% CI: 616-1449) per 100,000 in Nepal. Seroincidence rose with age, reaching a rate of 3231 (95% CI: 2630-3969) per 100,000 among 50 to 89-year-olds in Vellore, India. The seroincidence rates estimated using a cutoff were half the rates we estimated using antibody dynamics.

Conclusion: The approach described here can be deployed prospectively, coupled with existing serosurveys, or leverage banked samples to rapidly characterize scrub typhus burden and generate scrub typhus seroincidence estimates that are comparable across populations, regions, and time.

Keywords: *Orientia tsutsugamushi, Scrub Typhus, Antibody kinetics, Seroepidemiology, incidence*

Introduction

Scrub typhus, an acute febrile illness caused by the bacterium *Orientia tsutsugamushi (OT),* is an important, under-recognized etiology of fever (1). Once thought to be restricted to the "tsutsugamushi triangle", a region spanning from Russia to Pakistan, Australia, and Japan, recent studies have identified scrub typhus transmission in South America, Africa, and the Middle East (2–4). Infections occur when trombiculid mite chiggers (larvae) enter a host's skin through hair follicles and feed on lysed skin tissue. The mites, both vectors and reservoirs of OT, feed on various mammals, including humans and rodents (5). In humans, symptoms are non-specific and include fever, myalgia, headache, gastrointestinal symptoms, and rash. An eschar is found at the inoculation site in a variable proportion of cases and is frequently missed on clinical examination. Case fatality rates are estimated to be 1-2% among treated patients and 6% among untreated patients (6).

Determining where scrub typhus transmission occurs is critical to inform public health interventions and research priorities. Clinical incidence underestimates the true underlying burden of disease due to non-specific symptoms and the lack of accessible and accurate diagnostics (7). Periodic serosurveillance studies across endemic countries have demonstrated significant heterogeneity in seroprevalence within and between countries(8). However, directly comparing seroprevalence is not straightforward because of differences in the age distributions of each sampled population and uncertainty in antibody-waning patterns.

Here, we apply a novel analytic approach to estimate scrub typhus seroincidence using antibody decay information from confirmed cases. The decay of antibody concentrations defines a timescale for inferring when an infection occurred. This approach does not depend on classification using cutoffs, which have been difficult to derive for scrub typhus across locations with varying forces of

infection(9). We first model longitudinal IgG and IgM antibody responses to OT-derived antigens

among confirmed scrub typhus cases in Thailand and India and then use these parameters to

estimate scrub typhus seroincidence from cross-sectional population serosurveys in Nepal and

India.

Methods

<u>Study populations and enrollment</u>

Longitudinal scrub typhus cases: We used antibody responses measured from confirmed scrub

typhus patients in Thailand and India. In Thailand, hospitalized patients >15 years old with acute

fever duration <2 weeks were recruited from a fever surveillance study in Prachanukhao Hospital

in Chiang Rai, Thailand (10,11). Enrollment occurred from August 2007 to August 2008. Patients

confirmed to be infected with scrub typhus were defined by meeting at least one of the following

criteria: (1) *In vitro* isolation of *O. tsutsugamushi*, (2) a ≥4-fold rise in IgM titer in paired serum

samples when tested by the indirect immunofluorescence assay, (3) a positive result in at least two

out of the three PCR assays described in Paris et al., 2011(11). Serum samples were collected at

admission and stored at −80°C until testing. In India, serum samples were collected from

individuals infected with scrub typhus who were > 18 years old and sought care at Christian

Medical College Teaching hospital in Vellore, India, between December 2011 and March 2015.

Confirmed patients with scrub typhus were defined by a positive IgM ELISA (optical density (OD)

>0.8) and a positive PCR for *O. tsutsugamushi*. The serum samples were collected cross-

sectionally from retrospective cases with diagnoses between 2 months and 3.5 years prior. Serum

samples were collected in the patient's household, stored at $^-80^0$C, and processed within two

weeks of collection.

Population samples: In Nepal, we enrolled a geographically random, population-based cross-

sectional sample of individuals aged 0 to 25 years from the catchment areas of Kathmandu

University Teaching Hospital in Kathmandu (urban), and Dhulikhel Hospital in Kavrepalanchok,

Nepal (periurban and rural) (12). Within catchment areas, we randomly selected geographically

defined grid clusters and enumerated all households in each cluster. From this census, we

randomly selected individuals and sought consent. We collected capillary blood samples from

consenting participants onto TropBioTM filter papers (Cellabs Pty Ltd., Brookvale, New South

Wales, Australia). The samples were air-dried for at least two hours at room temperature, then

stored with desiccant in individual plastic bags at -20℃ until processing. Study participants were

enrolled between February 2019 and Jan 2021. In India, we utilized a previously-conducted cross-

sectional serosurvey for scrub typhus conducted from September 2014 to December 2014(13).

The study enrolled adults ≥18 years of age in the Vellore District of Tamil Nadu in South India. A

two-stage clustered sample design was used to randomly select communities and individuals(13).

Venous blood was collected from consenting participants, transported on ice, and stored at -70°C

until tested.

<u>Laboratory methods</u>

All samples were tested using IgM and IgG responses to *O. tsutsugamushi*-derived recombinant

56-kDa antigen using the Scrub Typhus Detect ELISA kit (InBios International, Inc., Seattle, WA,

USA)(10) performed as per the manufacturer's instructions. All serum samples were tested at a

1:100 dilution, and the results were read at 450 nm using a microplate reader (Thermo Scientific

Multiskan FC) to generate a final optical density result (optical density (OD) at 450 nm). To prepare

the dried capillary blood samples used in Nepal, we cut two filter paper protrusions and submerged

them in 133 μL of 1XPBS 0.05% Tween buffer overnight at 4 C, then centrifuged to recover the

eluates. The eluate was assumed to be equivalent to a 1:10 dilution of plasma.

<u>Statistical methods</u>

We estimated seroincidence in two ways: 1) by deriving it from the age-dependent seroprevalence

and 2) as a function of the longitudinal antibody dynamics after infection. For method 1, we used

finite mixture models to determine the IgG seropositivity cutoffs using the mean plus three standard

deviations of the first mixture component (14). We calculated IgG seroprevalence as the proportion

of individuals classified as IgG seropositive. We then used an exponential survival model to derive

seroincidence (equations 1 & 2), where $\pi(a)$ equals the seroprevalence at age (a) and λ equals

the seroincidence rate.

165
166   $$eq\ 1. \qquad \pi(a) = 1 - e^{-\lambda a}$$
167
168   $$eq\ 2. \qquad \lambda = \frac{(-log_{10}\ (1 - \pi(a))}{a}$$
169

We fit a generalized linear model to the binomial seropositivity outcome conditional on age with a

complementary log-log link and estimated seroincidence from the model's intercept term (15,16).

This approach assumes there is no antibody waning over time and that seroincidence is constant

over age. To evaluate how seropositivity changes over age, we fit generalized additive models (17)

with a cubic spline for age and simultaneous confidence intervals using a parametric bootstrap of

the variance-covariance matrix of the fitted model parameters (18).

For method 2, we used information about antibody decay from confirmed cases to estimate

seroincidence. First, we modeled longitudinal antibody dynamics after scrub typhus infection using

two-phase models with an infection episode characterized by an exponential rise in antibody

response and a non-exponential power function decay (19–21). We model antibody kinetics from

onset of fever ($t = 0$) with initial antibody levels of y_0 reaching a peak antibody response of y_1. We

used a Bayesian hierarchical framework to estimate the above models for IgG and IgM (19,20) and

generate joint distributions using Markov chain Monte Carlo (MCMC) sampling, allowing for

individual variation. We implemented the models in R version 4.1.3 using JAGS(22).

We used the distributions of peak antibody response and the decay rate and shape to estimate

seroincidence in the cross-sectional population samples(23). We created a likelihood function for the

observed cross-sectional population data based on the longitudinal kinetics following scrub typhus

infection with the assumption that incident infections occur as a Poisson process with rate lambda

(λ)(24). We generated maximum likelihood profiles for λ using each isotype separately and jointly by

combining their likelihood functions. We accounted for two sources of noise in the observed serologic

responses: measurement noise of the assay and biologic noise as detailed in Teunis et. al. (23).

<u>Ethics statement</u>

Institutional Review Boards in India (Christian Medical College, Vellore), Thailand (Chiang Rai

Hospital, the Faculty of Tropical Medicine, Mahidol University, and the Thai Ministry of Public

Health), The United States (Stanford University Institutional Review Board), and Nepal (Nepal

Health Research Council Ethical Review Board)) approved the study forms and protocols.

Results

For the longitudinal cases, ELISA antibody responses were measured from 253 patients with scrub

typhus; 84 in Thailand and 211 in India(25). Only one serum sample was available per participant,

between 1 day and 1250 days after the date of symptom onset (median 440 days, inter-quartile

range (IQR): 140-840). The samples from Thailand were collected between 1 and 11 days after

symptom onset (median 5.5) and the samples in India were collected between 65 and 1250 days

after symptom onset (median 599). Both cohorts measured IgM responses, but IgG responses

were only available from the India cohort (Figure 1). The median ages of participants were 46

years (IQR 35-56); 47 in India (IQR 35-57), and 43 in Thailand (IQR 35-51) (Table 1).

Table 1: Summary of longitudinal scrub typhus patient data

	India (N=211)	Thailand (N=42)	Overall (N=253)
Age (years)			
Median (IQR)	47 (35 - 57)	43 (35 - 51)	46 (35 - 56)
Missing	0 (0%)	1 (2.4%)	1 (0.4%)
Days since symptom onset			
Median (IQR)	520 (340 - 880)	5.5 (4.0 - 7.0)	440 (140 - 840)

212
213

Median IgM responses were elevated above the kit cut-off for 6.65 months after symptom onset

and above the mixture model cut-off for 9.4 months after symptom onset. Median IgG responses

were elevated above the kit cut-off for 31.5 months after symptom onset and above the mixture

model cut-off for 29.7 months after symptom onset (Figure 2A). As expected, IgM responses had a

faster decay rate than IgG responses. Peak IgM responses were higher than IgG responses. The

time-to-peak response was similar between IgG and IgM. The model-predicted baseline IgG

responses were higher than baseline IgM (Figure 2B).

For the population data, ELISA antibody responses were measured from 721 participants in

Vellore, India, and 1105 participants in Kathmandu and Kavre districts, Nepal. The median age of

participants was 49 in India (IQR 40-62) and 11 in Nepal (IQR 5.5-17). In India, 63.2% (456/721) of

participants were female, compared to 48.7% (538/1105) in Nepal (Table 2).

Table 2: Population data summary

	India (N=721)	Nepal (N=1105)
Age, in years		
Median (IQR)	49 (40 - 62)	11 (5.5 - 17)
Age, in years, Categorical		
0-17	0 (0%)	876 (79.3%)
18-29	59 (8.2%)	229 (20.7%)
30-49	302 (41.9%)	0 (0%)
50-89	360 (49.9%)	0 (0%)
Gender		
Female	456 (63.2%)	538 (48.7%)
Male	265 (36.8%)	567 (51.3%)
Residence zone		
Rural	177 (24.5%)	316 (28.6%)
Periurban	284 (39.4%)	319 (28.9%)
Urban	190 (26.4%)	470 (42.5%)
Missing	70 (9.7%)	0 (0%)
Occupation		
Agriculture	85 (11.8%)	0 (0%)
Labor & Manufacturing	181 (25.1%)	14 (1.3%)
Minor, <=18 years	1 (0.1%)	880 (79.6%)
None	343 (47.6%)	152 (13.8%)
Other	21 (2.9%)	21 (1.9%)
Professional	44 (6.1%)	26 (2.4%)
Service	46 (6.4%)	12 (1.1%)
Household size		
Median (IQR)	4.0 (3.0 - 5.0)	4.0 (4.0 - 6.0)
Missing	3 (0.4%)	1 (0.1%)

In both India and Nepal, the seroprevalence of scrub typhus infection increased with age (Figure 3). In Vellore, India the overall seroprevalence was 29.7% (95% CI 26.3-33.3): growing from 10.2% (95% CI 2.4-17.9) among 18-29-year-olds to 37.8% (95% CI 32.8-42.8) among 50 to 89-year-olds. In Kathmandu and Kavre, Nepal, the overall seroprevalence was 1.1% (95% CI 0.5-1.7), rising from 0.1% (95%CI 0-.3) among 0 to 17-year-olds to 4.8% (95%CI 2-7.6%) among 18-29-year-olds (Table 3).

Table 3: Age-specific seroprevalence and seroincidence in Nepal and India

Age	N	N IgG seropositive	IgG seroprevalence (95% CI)	Seroincidence per 100,000 person-years (95% CIs)	
				IgG antibody kinetics	Derivative of age-dependent IgG seroprevalence
India					
0-17	0	0	–	–	–
18-29	59	6	10.2% (2.4-17.9)	886 (432-1817)	431 (194-958)
30-49	302	72	23.8% (19.0-28.7)	1602 (1254-2046)	675 (535-851)
50-89	360	136	37.8% (32.8-42.8)	3231 (2630-3969)	757 (639-897)
Overall	721	214	29.7% (26.3-33.0)	2199 (1893-2555)	712 (622-815)
Nepal					
0-17	876	1	0.1% (-0-0.3)	212 (120-374)	13 (2-90)
18-29	229	11	4.8% (2.0-7.6)	945 (616-1449)	232 (128-419)
30-49	0	0	–	–	–
50-89	0	0	–	–	–
Overall	1105	12	1.1% (0.5-1.7)	425 (303-596)	95 (54-168)

In Vellore, India, the overall scrub typhus seroincidence rate was 2199 per 100,000 person-years

(95% CI: 1893-2555) using IgG antibody dynamics. Seroincidence rose with age; growing from 886

(95% CI 432-1817) to 3231 (95% CI 2630-3969) among 50-89-year-olds. In the Kathmandu valley

of Nepal, the overall seroincidence was 425 per 100,000 person-years, also rising with age from

212 (95% CI 120-374) per 100,000 person-years among 0 to 17-year-olds to 945 (95%CI 616-

1449) among 18 to 29-year-olds. In both Nepal and India, the overall seroincidence using antibody

dynamics was more than double the seroincidence derived from the age-dependent IgG-

seroprevalence (Table 3). In both sites, scrub typhus seroincidence varied by resident zones. In

Vellore, India, seroincidence was highest in periurban settings, followed by rural and then urban,

whereas in Nepal, seroincidence was higher in urban settings (Figure 4). In India, females had

elevated seroincidence rates compared to males across all age groups (Figure 4).

Discussion

We describe a new approach for generating population-level scrub typhus seroincidence estimates

from cross-sectional serosurveys. Seroincidence, the number of new infections in a population per

year, characterizes transmission intensity and is highly informative for determining where and

among whom infection burden is highest. While classic approaches for estimating seroincidence

require cutoff points and ignore antibody waning overtime, in this approach we use the antibody

decay trajectories from confirmed cases to estimate seroincidence. We show here that ignoring

antibody waning and using a cutoff to calculate scrub typhus seroincidence underestimates the

rate by at least half.

Understanding the post-infection antibody kinetics of scrub typhus is also important for guiding the

clinical interpretation of ELISA antibody response data. Using data from patients infected with

scrub typhus in Thailand and India, we found that median IgM responses remained elevated above

the ELISA kit threshold for six and a half months after symptom onset. IgG responses remained

elevated above the ELISA kit threshold for over 24 months after symptom onset. These findings

are relevant to clinicians who should therefore interpret elevated IgM and IgG responses from a

single sample with caution given the many local causes of fever unrelated to scrub typhus.

Most scrub typhus seroepidemiologic studies dichotomize IgG responses using a cutoff threshold

and then calculate the proportion of the population whose values fall above that threshold

(seroprevalence). Cutoff thresholds are notoriously challenging to calculate and can vary across

regions and studies (9), making it challenging to compare seroprevalence results across regions

and time. The approach we describe in this paper has the advantage of not requiring a set cutoff

point. Additional strengths of the approach are that it incorporates information about measurement

noise, biologic noise, and age to generate more accurate uncertainty intervals around

seroincidence estimates.

We show here that seroprevalence is highly age-dependent; therefore, the age of the patients will

influence the overall seroprevalence estimate. In our study populations, the overall seroprevalence

was much higher in India than in Nepal; however, in India, the median age of patients was 49

years compared to the median age of 11 years in Nepal. Despite differences in overall

seroprevalence rates, the seroincidence rate among 18 to 29-year-olds in India and Thailand were

similar, although there may be micro-geographic differences in infection risk based on elevation,

vegetation, climate, and other factors

The results of this study should be interpreted within the context of several limitations. First, only

one sample was available from each patient with scrub typhus, limiting our precision in estimating

individual-level heterogeneity in decay rates. Moreover, with just one sample per individual, we

were unable to identify and remove potential reinfections. We hypothesize that individuals who are

re-exposed to scrub typhus will have elevated IgG responses; when these observations remain in

the data, our antibody decay rates will be biased towards slower decay. An additional limitation of

the available case data is that we had no early data on IgG responses and no responses from

children less that 15 years old. It is possible that antibody dynamics among young children are

different from those among older individuals, and these dynamics would influence the accuracy of

seroincidence estimates among younger ages. Future studies that enroll patients infected with

scrub typhus across ages and follow the same individuals over time are needed to refine the

longitudinal antibody decay rates, accurately define peak IgG responses, explore the influence of

reinfections of antibody kinetics and investigate the potential influence of age on antibody kinetics.

Scrub typhus remains an important but underrecognized etiology of acute fever with endemicity

expanding globally. There is a critical need for low-cost, accurate tools to quantify the burden of

scrub typhus infections to inform public health decision-making. We describe a sero-surveillance

approach that can efficiently generate population-level scrub typhus seroincidence estimates. This

approach can be deployed prospectively, coupled with existing serosurveys, or leverage banked

samples to rapidly characterize scrub typhus burden and generate estimates that are comparable

across populations, regions, and time.

310
311
312 **Acknowledgments**
313
314 We gratefully acknowledge the study participants for their valuable time and interest in participating

315 in these studies.

Funding

This work was supported by the Fogarty International Center at National Institutes of Health [K01

TW012177-01A1]

Figures

Figure 1: Quantitative antibody responses among (A) Scrub Typhus Cases and (B,C) population serosurveys in Nepal and India. Anti-OT IgA and IgG responses are measured using kinetic enzyme-linked immunosorbent assays (ELISAs) and depicted on the y-axis of all plots. The horizontal dashed line denotes the kit cutoff (light grey) and the mixture model cutoff (dark grey). In panel A, responses are compared over time from fever onset among confirmed cases from Thailand and India. Each dot is an individual case, the overlaying boxplots indicate the median and the inter-quartile range. Panel B shows the distribution of responses among the population-based samples in India and Nepal. Panel C shows the antibody among the population-based samples in India and Nepal as a function of age.

346
Figure 2: (A) Kinetics of IgM responses among Scrub Typhus cases and (B) Distributions of
model-predicted peak antibody response, decay rate and shape, time to peak antibody
response, and baseline antibody response. Longitudinal antibody dynamics were modeled from
ELISA-measured antibody responses using Bayesian hierarchical models. In panel A, the points
are the observed individual antibody concentrations; each point indicates one patient. The dark
solid line indicates the median, and dotted lines indicate 95% credible intervals for the model-fitted
antibody decay concentrations. The horizontal dashed line denotes the kit cutoff (light grey) and
the mixture model cutoff (dark grey). Panel B shows distributions and boxplots for the model-
predicted decay rate, decay shape, time to peak, baseline response, and peak antibody responses
for both IgM and IgG.
357

369
Figure 3: Age-dependent seroprevalence
Seroprevalence is depicted as a function of age. Seroprevalence was determined using the
mixture-model derived cutoff for IgG responses. The age-dependent seroprevalence was modeled
using a generalized additive model with with a cubic spline for age and simultaneous confidence
intervals using a parametric bootstrap of the variance-covariance matrix of the fitted model
parameters.
376

Figure 4: Seroincidence by age plus (A) resident zones and (B) gender
Seroincidence rates across are shown across age strata, residence zone (A), and gender (B)
comparing the India and Nepal population samples. Seroincidence is estimated using IgG antibody
kinetics. The point represents the incidence estimate per 100,000 person-years, and the bar
indicates the 95% confidence interval.

(A)

(B)

References

1. Paris DH, Shelite TR, Day NP, Walker DH. Unresolved Problems Related to Scrub Typhus: A Seriously Neglected Life-Threatening Disease. The American Journal of Tropical Medicine and Hygiene. 2013 Aug 7;89(2):301–7.

2. Weitzel T, Dittrich S, López J, Phuklia W, Martinez-Valdebenito C, Velásquez K, et al. Endemic Scrub Typhus in South America. N Engl J Med. 2016 Sep 8;375(10):954–61.

3. Izzard L, Fuller A, Blacksell SD, Paris DH, Richards AL, Aukkanit N, et al. Isolation of a Novel Orientia Species (O. chuto sp. nov.) from a Patient Infected in Dubai. Journal of Clinical Microbiology. 2010 Dec;48(12):4404–9.

4. Thiga JW, Mutai BK, Eyako WK, Ng'ang'a Z, Jiang J, Richards AL, et al. High Seroprevalence of Antibodies against Spotted Fever and Scrub Typhus Bacteria in Patients with Febrile Illness, Kenya. Emerg Infect Dis. 2015 Apr;21(4):688–91.

5. Luce-Fedrow A, Lehman ML, Kelly DJ, Mullins K, Maina AN, Stewart RL, et al. A Review of Scrub Typhus (Orientia tsutsugamushi and Related Organisms): Then, Now, and Tomorrow. Trop Med Infect Dis. 2018 Jan 17;3(1):8.

6. Bonell A, Lubell Y, Newton PN, Crump JA, Paris DH. Estimating the burden of scrub typhus: A systematic review. PLOS Neglected Tropical Diseases. 2017 Sep 25;11(9):e0005838.

7. John R, Varghese GM. Scrub typhus: a reemerging infection. Current Opinion in Infectious Diseases. 2020 Oct;33(5):365–71.

8. Xu G, Walker DH, Jupiter D, Melby PC, Arcari CM. A review of the global epidemiology of scrub typhus. PLOS Neglected Tropical Diseases. 2017 Nov 3;11(11):e0006062.

9. Saraswati K, Phanichkrivalkosil M, Day NPJ, Blacksell SD. The validity of diagnostic cut-offs for commercial and in-house scrub typhus IgM and IgG ELISAs: A review of the evidence. PLOS Neglected Tropical Diseases. 2019 Feb 4;13(2):e0007158.

10. Blacksell SD, Tanganuchitcharnchai A, Nawtaisong P, Kantipong P, Laongnualpanich A, Day NPJ, et al. Diagnostic Accuracy of the InBios Scrub Typhus Detect Enzyme-Linked Immunoassay for the Detection of IgM Antibodies in Northern Thailand. Clinical and Vaccine Immunology [Internet]. 2015 Dec 9 [cited 2022 Feb 20]; Available from: https://journals.asm.org/doi/abs/10.1128/CVI.00553-15

11. Paris DH, Blacksell SD, Nawtaisong P, Jenjaroen K, Teeraratkul A, Chierakul W, et al. Diagnostic Accuracy of a Loop-Mediated Isothermal PCR Assay for Detection of Orientia tsutsugamushi during Acute Scrub Typhus Infection. PLOS Neglected Tropical Diseases. 2011 Sep 13;5(9):e1307.

12. Aiemjoy K, Seidman JC, Saha S, Munira SJ, Sajib MSI, Sium SMA, et al. Estimating typhoid incidence from community-based serosurveys: a multicohort study. The Lancet Microbe [Internet]. 2022 Jun 21 [cited 2022 Jun 23];0(0). Available from: https://www.thelancet.com/journals/lanmic/article/PIIS2666-5247(22)00114-8/fulltext

13. Trowbridge P, P. D, Premkumar PS, Varghese GM. Prevalence and risk factors for scrub
typhus in South India. Tropical Medicine & International Health. 2017;22(5):576–82.

14. Bickel PJ, Klaassen CA, Bickel PJ, Ritov Y, Klaassen J, Wellner JA, et al. Efficient and
adaptive estimation for semiparametric models. Vol. 2. Springer New York; 1998.

15. Arnold BF, Martin DL, Juma J, Mkocha H, Ochieng JB, Cooley GM, et al. Enteropathogen
antibody dynamics and force of infection among children in low-resource settings. Ferguson
NM, Jit M, White M, Leung DT, Azman A, editors. eLife. 2019 Aug 19;8:e45594.

16. Jewell NP, van der Laan M. Generalizations of current status data with applications. Lifetime
Data Anal. 1995 Mar 1;1(1):101–9.

17. Wood SN. Generalized additive models: an introduction with R. Chapman and Hall/CRC; 2006.

18. Marra G, Wood SN. Coverage properties of confidence intervals for generalized additive model
components. Scandinavian Journal of Statistics. 2012;39(1):53–74.

19. de Graaf WF, Kretzschmar MEE, Teunis PFM, Diekmann O. A two-phase within-host model for
immune response and its application to serological profiles of pertussis. Epidemics. 2014
Dec;9:1–7.

20. Teunis PFM, van Eijkeren JCH, de Graaf WF, Marinović AB, Kretzschmar MEE. Linking the
seroresponse to infection to within-host heterogeneity in antibody production. Epidemics. 2016
Sep;16:33–9.

21. Teunis PFM, van der Heijden OG, de Melker HE, Schellekens JFP, Versteegh FGA,
Kretzschmar MEE. Kinetics of the IgG antibody response to pertussis toxin after infection with
B. pertussis. Epidemiol Infect. 2002 Dec;129(3):479–89.

22. Hornik K, Leisch F, Zeileis A. JAGS: A program for analysis of Bayesian graphical models
using Gibbs sampling. In: Proceedings of DSC. 2003.

23. Teunis PFM, van Eijkeren JCH. Estimation of seroconversion rates for infectious diseases:
Effects of age and noise. Stat Med. 2020 Sep 20;39(21):2799–814.

24. Teunis PFM, van Eijkeren JCH, Ang CW, van Duynhoven YTHP, Simonsen JB, Strid MA, et al.
Biomarker dynamics: estimating infection rates from serological data. Stat Med. 2012 Sep
10;31(20):2240–8.

25. Varghese GM, Rajagopal VM, Trowbridge P, Purushothaman D, Martin SJ. Kinetics of IgM and
IgG antibodies after scrub typhus infection and the clinical implications. International Journal of
Infectious Diseases. 2018 Jun 1;71:53–5.