1   **Title: Estimating the seroincidence of scrub typhus using antibody dynamics following**
2   **infection**
3
4
5   **Authors:** Kristen Aiemjoy[1,2], Nishan Katuwal[3], Krista Vaidya[1], Sony Shrestha[3], Melina Thapa[3],
6   Peter Teunis[4], Isaac I. Bogoch[5], Paul Trowbridge[6], Pacharee Kantipong[7], Stuart D. Blacksell[8,9], Tri
7   Wangrangsimakul[8], George M Varghese[10], Richard Maude[8,9,11,12], Dipesh Tamrakar[3], Jason R.
8   Andrews[13]
9
10
Affiliations:
1. Department of Public Health Sciences, University of California Davis School of Medicine, Davis,
CA, USA
2. Department of Microbiology and Immunology, Faculty of Tropical Medicine, Mahidol University,
Bangkok Thailand
3. Dhulikhel Hospital, Kathmandu University Hospital, Dhulikhel, Nepal
4. Center for Global Safe Water, Sanitation and Hygiene, Hubert Department of Global Health,
Rollins School of Public Health, Emory University, Atlanta, GA, USA
5. Department of Medicine, University of Toronto, Toronto, ON, Canada
6. Michigan State University School of Human Medicine, Grand Rapids, MI, USA
7. Department of Internal Medicine, Chiangrai Prachanukroh Hospital, Chiang Rai, Thailand
8. Mahidol-Oxford Tropical Medicine Research Unit, Faculty of Tropical Medicine, Mahidol
University, Bangkok Thailand
9. Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine, University of
Oxford, Oxford, UK
10. Department of Infectious Diseases, Christian Medical College, Vellore, India
11. The Open University, Milton Keynes, UK
12. School of Public Health, Li Ka Shing Faculty of Medicine, University of Hong Kong, Hong Kong
Department of Medicine, Stanford University School of Medicine, Stanford, CA, USA
13. Division of Infectious Diseases and Geographic Medicine, Stanford University School of
Medicine, Stanford, CA, USA
32
33
Corresponding Author:
Kristen Aiemjoy
Assistant Professor
Division of Epidemiology
Department of Public Health Sciences,
University of California Davis School of Medicine, Davis, CA, USA
Department of Microbiology and Immunology, Faculty of Tropical Medicine, Mahidol University,
Bangkok Thailand
kaiemjoy@ucdavis.edu
43
44
45
46   **Word Count: 3721**
47
48

Abstract

Introduction: Scrub typhus is an acute febrile illness caused by the bacterium *Orientia tsutsugamushi.* Characterizing the population-level burden of scrub typhus is challenging due to the lack of accessible and accurate diagnostics. In this study, we present a novel approach that utilizes information on antibody dynamics after infection to generate population-level scrub typhus seroincidence estimates from cross-sectional serosurveys.

Methods: We use data from three cohorts of scrub typhus patients enrolled in Chiang Rai, Thailand, and Vellore, India, and representative population data from two serosurveys in and around the Kathmandu valley, Nepal, and Vellore, India. The samples were tested for IgM and IgG responses to *Orientia tsutsugamushi*-derived recombinant 56-kDa antigen using commercial ELISA kits. We used Bayesian hierarchical models to fit two-phase models to the antibody responses from scrub typhus cases and used the joint distributions of the peak antibody titers and decay rates to estimate population-level incidence rates in the cross-sectional serosurveys. We compared this new method to a traditional cut-off-based approach for estimating seroincidence.

Results: Median IgG antibodies persisted above OD 1.7 for 22 months, while IgM displayed longer persistence than expected, with 50% of participants having an OD >1 for 5 months. We estimated an overall seroincidence of 18 per 1000 person-years (95% CI: 16-21) in India and 4 per 1000 person-years (95% CI: 3-6) in Nepal. Among 18 to 29-year-olds, the seroincidence was 8 per 1000 person-years (95% CI 4 -16) in India and 9 per 1000 person-years (95% CI: 6-14) in Nepal. In both India and Nepal, seroincidence was higher in urban and periurban settings compared to rural areas. Compared to our method, seroincidence estimates derived from age-dependent IgG-seroprevalence without accounting for antibody decay underestimated the disease burden by 50%. By incorporating antibody dynamics, the approach described here provides more accurate age-specific infection risk estimates, emphasizing the importance of considering both IgG and IgM decay patterns in scrub typhus seroepidemiology.

Conclusion: The sero-surveillance approach developed in this study efficiently generates population-level scrub typhus seroincidence estimates from cross-sectional serosurveys. This methodology offers a valuable new tool for informing targeted prevention and control strategies, ultimately contributing to a more effective response to scrub typhus in endemic regions worldwide.

Keywords: *Orientia tsutsugamushi, Scrub Typhus, Antibody kinetics, Seroepidemiology, incidence*

Introduction

Scrub typhus, an acute febrile illness caused by the bacterium *Orientia tsutsugamushi (OT),* is an important, under-recognized etiology of fever (1). Once thought to be restricted to the "tsutsugamushi triangle", a region spanning from Russia to Pakistan, Australia, and Japan, recent studies have identified scrub typhus transmission in South America, Africa, and the Middle East (2–4). Infections occur when trombiculid mite chiggers (larvae) enter a host's skin through hair follicles and feed on lysed skin tissue. The mites, both vectors and reservoirs of OT, feed on various mammals, including humans and rodents (5). In humans, symptoms are non-specific and include fever, myalgia, headache, gastrointestinal symptoms, and rash. An eschar is found at the inoculation site in a variable proportion of cases and is frequently missed on clinical examination. Case fatality rates are estimated to be 1-2% among treated patients and 6% among untreated patients (6).

Determining where scrub typhus transmission occurs is critical to inform public health interventions and research priorities. Clinical incidence underestimates the true underlying burden of disease due to non-specific symptoms and the lack of accessible and accurate diagnostics (7). Periodic serosurveillance studies across endemic countries have demonstrated significant heterogeneity in seroprevalence within and between countries (8). However, directly comparing seroprevalence is not straightforward because of differences in the age distributions of each sampled population and uncertainty in antibody-waning patterns.

Here, we apply a novel analytic approach to estimate scrub typhus seroincidence using antibody decay information from confirmed cases. The decay of antibody concentrations defines a timescale for inferring when an infection occurred. This approach does not depend on classification using cutoffs, which have been difficult to derive for scrub typhus across locations with varying forces of

infection (9). We first model longitudinal IgG and IgM antibody responses to OT-derived antigens

among confirmed scrub typhus cases in Thailand and India and then use these parameters to

estimate scrub typhus seroincidence from cross-sectional population serosurveys in Nepal and

India.

Methods

<u>Study populations and enrollment</u>

Scrub typhus cases: We used antibody responses measured from confirmed scrub typhus patients

enrolled in three studies: two in Thailand and one in India. The first study in Thailand enrolled

children less than 18 years old who were admitted to Chiang Rai Prachanukroh Hospital with fever

(temperature higher than 37.5°C) or a history of fever within the last 14 days(10). Enrollment

occurred between July 2015 to August 2016. Confirmed scrub typhus infections were defined by

meeting at least one of the following criteria: (1) a positive PCR or culture result from either a blood

or eschar sample, (2) a fourfold rise in the IgM titer to ≥1:3200 in paired serum or plasma samples,

or (3) a single IgM titer of ≥1:3200 in an acute-phase serum or plasma sample(11,12). Blood

samples were collected at baseline and at 2-, 12-, and 52-weeks after enrollment, stored at −80°C,

and transported to Bangkok for diagnostic processing.

The second study In Thailand enrolled hospitalized patients older than 15 years from a fever

surveillance study also at the Chiang Rai Prachanukhao Hospital (13,14). Enrollment occurred

from August 2007 to August 2008. Patients confirmed to be infected with scrub typhus were

defined by meeting at least one of the following criteria: (1) *In vitro* isolation of *O. tsutsugamushi*,

(2) a ≥4-fold rise in IgM titer in paired serum samples when tested by the indirect

immunofluorescence assay, (3) a positive result in at least two out of the three PCR assays

described in Paris et al., 2011(14). Serum samples were collected at admission and stored at

−80°C until testing.

In India, serum samples were collected from individuals infected with scrub typhus who were > 18

years old and sought care at Christian Medical College Teaching Hospital in Vellore, India,

between December 2011 and March 2015 (15). Confirmed patients with scrub typhus were defined

by a positive IgM ELISA (optical density (OD) >0.8) and/or a positive PCR for *O. tsutsugamushi*.

The serum samples were collected cross-sectionally from retrospective cases with diagnoses

between 2 months and 3.5 years prior. Serum samples were collected in the patient's household,

stored at $^{-}80^{0}$C, and processed within two weeks of collection.

Population samples: In Nepal, we enrolled a geographically random, population-based cross-

sectional sample of individuals aged 0 to 25 years from the catchment areas of Kathmandu

University Teaching Hospital in Kathmandu (urban), and Dhulikhel Hospital in Kavrepalanchok,

Nepal (periurban and rural) (16). Within catchment areas, we randomly selected geographically

defined grid clusters and enumerated all households in each cluster. From this census, we

randomly selected individuals and sought consent. We collected capillary blood samples from

consenting participants onto TropBioTM filter papers (Cellabs Pty Ltd., Brookvale, New South

Wales, Australia). The samples were air-dried for at least two hours at room temperature, then

stored with desiccant in individual plastic bags at -20℃ until processing. Study participants were

enrolled between February 2019 and Jan 2021. In India, we utilized a previously-conducted cross-

sectional serosurvey for scrub typhus conducted from September 2014 to December 2014(17).

The study enrolled adults ≥18 years of age in the Vellore District of Tamil Nadu in South India. A

two-stage clustered sample design was used to randomly select communities and individuals(17).

Venous blood was collected from consenting participants, transported on ice, and stored at -70°C

until tested.

<u>Laboratory methods</u>

All samples were tested using IgM and IgG responses to *O. tsutsugamushi*-derived recombinant

56-kDa antigen using the Scrub Typhus Detect ELISA kit (InBios International, Inc., Seattle, WA,

USA)(13) performed as per the manufacturer's instructions. All serum samples were tested at a

1:100 dilution, and the results were read at 450 nm using a microplate reader (Thermo Scientific

Multiskan FC) to generate a final optical density result (optical density (OD) at 450 nm). To prepare

the dried capillary blood samples used in Nepal, we cut two filter paper protrusions and submerged

them in 133 μL of 1XPBS 0.05% Tween buffer overnight at 4 C, then centrifuged to recover the

eluates. The eluate was assumed to be equivalent to a 1:10 dilution of plasma.

<u>Statistical methods</u>

We estimated seroincidence in two ways: 1) by deriving it from the age-dependent seroprevalence

and 2) as a function of the longitudinal antibody dynamics after infection. For method 1, we used

finite mixture models to determine the IgG seropositivity cutoffs using the mean plus two standard

deviations of the first mixture component (18). We calculated IgG seroprevalence as the proportion

of individuals classified as IgG seropositive. We then used an exponential survival model to derive

seroincidence (equations 1 & 2), where $\pi(a)$ equals the seroprevalence at age (a) and λ equals

the seroincidence rate.

184
185 $$eq\ 1. \qquad \pi(a) = 1 - e^{-\lambda a}$$
186
187 $$eq\ 2. \qquad \lambda = \frac{(-log_{10}\ (1 - \pi(a)))}{a}$$
188

We fit a generalized linear model to the binomial seropositivity outcome conditional on age with a

complementary log-log link and estimated seroincidence from the model's intercept term (19,20).

This approach assumes there is no antibody waning over time and that seroincidence is constant

over age. To evaluate how seropositivity changes over age, we fit generalized additive models (21)

with a cubic spline for age and simultaneous confidence intervals using a parametric bootstrap of

the variance-covariance matrix of the fitted model parameters (22).

For method 2, we used information about antibody decay from confirmed cases to estimate

seroincidence. First, we modeled longitudinal antibody dynamics after scrub typhus infection using

two-phase models with an infection episode characterized by an exponential rise in antibody

response and a non-exponential power function decay (23–25). We model antibody kinetics from

onset of fever ($t = 0$) with initial antibody levels of y_0 reaching a peak antibody response of y_1. We

used a Bayesian hierarchical framework to fit the above models for IgG and IgM (23,24) and

generate joint distributions using Markov chain Monte Carlo (MCMC) sampling, allowing for

individual variation. We implemented the models in R version 4.1.3 using JAGS(26).

We used the distributions of peak antibody response and the decay rate and shape to estimate

seroincidence in the cross-sectional population samples(27). We created a likelihood function for the

observed cross-sectional population data based on the longitudinal kinetics following scrub typhus

infection with the assumption that incident infections occur as a Poisson process with rate lambda

(λ(28). We generated maximum likelihood profiles for λ using each isotype separately and jointly by

combining their likelihood functions. We accounted for two sources of noise in the observed serologic

responses: measurement noise of the assay and biologic noise as detailed in Teunis et. al. (27).

<u>Ethics statement</u>

Institutional Review Boards in India (Christian Medical College, Vellore), Thailand (Chiang Rai

Hospital, the Faculty of Tropical Medicine, Mahidol University, and the Thai Ministry of Public

Health), The United States (Stanford University Institutional Review Board), and Nepal (Nepal

Health Research Council Ethical Review Board)) approved the study forms and protocols.

Results

ELISA antibody responses were measured from 288 patients with confirmed scrub typhus

infections: 77 in Thailand and 211 in India (15)**.** Samples were collected between 0 and 1300 days

after symptom onset. In the pediatric cohort in Thailand, 4 samples were collected between 0 400

days after symptom onset. In the adult cohorts in India and Thailand, just one sample was

collected between 1 and 11 days after symptom onset (median 5.5) in Thailand and between 65

and 1250 days after symptom onset (median 599) in India. The adult cohort in Thailand did not

measure IgG responses. The median ages of participants were 44 years (IQR 27-55, range 1.2 -

84); 47 in India (IQR 35-57), and 43 in the adult cohort in Thailand (IQR 35-51) and 6.6 (IQR 4 –

10) in the pediatric cohort in Thailand (Table 1).

238 **Table 1: Summary of longitudinal scrub typhus patient data**

	India: Adult (N=211)	Thailand: Pediatric (N=35)	Thailand: Adult (N=42)	Overall (N=288)
Age (years)				
Median (IQR)	47 (35 - 57)	6.6 (4.0 - 10)	43 (35 - 51)	44 (27 - 55)
Min/Max	19 - 79	1.2 - 13	15 - 84	1.2 - 84
Missing	0 (0%)	0 (0%)	1 (2.4%)	1 (0.3%)
Days from symptom onset to first visit				
Median (IQR)	520 (340 - 880)	10 (7.5 - 12)	5.5 (4.0 - 7.0)	380 (13 - 750)
Min/Max	65 - 1300	3.0 - 19	0.15 - 11	0.15 - 1300
Days from symptom onset to last visit				
Median (IQR)	-	350 (350 - 370)	-	350 (350 - 370)
Min/Max		10 - 400		10 - 400
Number of visits				
Median (IQR)	1.0 (1.0 - 1.0)	4.0 (4.0 - 4.0)	1.0 (1.0 - 1.0)	1.0 (1.0 - 1.0)
Min/Max	1.0 - 1.0	1.0 - 4.0	1.0 - 1.0	1.0 - 4.0

Median IgG responses were elevated above the mixture model cut-off for 22.3 months after

symptom onset (Figure 2A). An OD of 1 has been suggested as a reasonable cutoff to indicate a

recent infection (29,30). In our study population, median responses were elevated about an OD of

1 for 5.3 months after infection, 25% of the population had elevated for 12.5 months and 10% had

elevated responses for 26.5 months. IgM responses decayed at a rate of OD 1.7 (95% credible

interval (CrI) 0.77 – 3.77) per month compared to 0.11 (95% CrI 0.04 – 0.29) per month for IgG).

IgM responses peaked at OD 5.12 (95% CrI 4.00 – 7.03), 5.2 days (95% CrI 3.4 – 7.6) after

symptom onset. IgG responses peaked at OD 2.8 (95% CrI 2.1 – 3.73) 11 days (95% CrI 5.7 –

20.6) after symptom onset. (Supplemental Table 2).

For the population data, ELISA antibody responses were measured from 721 participants in

Vellore, India, and 1105 participants in Kathmandu and Kavre districts, Nepal. The median age of

participants was 49 in India (IQR 40-62) and 11 in Nepal (IQR 5.5-17). In India, 63.2% (456/721) of

participants were female, compared to 48.7% (538/1105) in Nepal (Table 2).

Table 2: Population data summary

	India (N=721)	Nepal (N=1105)
Age, in years		
Median (IQR)	49 (40 - 62)	11 (5.5 - 17)
Min/Max 18 - 87	0.90 - 27	
Age, in years, Categorical		
0-17	0 (0%)	876 (79.3%)
18-29	59 (8.2%)	229 (20.7%)
30-49	302 (41.9%)	0 (0%)
50-89	360 (49.9%)	0 (0%)
Gender		
Female	456 (63.2%)	538 (48.7%)
Male	265 (36.8%)	567 (51.3%)
Residence zone		
Rural	177 (24.5%)	316 (28.6%)
Periurban	284 (39.4%)	319 (28.9%)
Urban	190 (26.4%)	470 (42.5%)
Missing	70 (9.7%)	0 (0%)
Occupation		
Agriculture	85 (11.8%)	0 (0%)
Labor & Manufacturing	181 (25.1%)	14 (1.3%)
Minor, <=18 years	1 (0.1%)	880 (79.6%)
None	343 (47.6%)	152 (13.8%)
Other	21 (2.9%)	21 (1.9%)
Professional	44 (6.1%)	26 (2.4%)
Service	46 (6.4%)	12 (1.1%)
Household size		
Median (IQR)	4.0 (3.0 - 5.0)	4.0 (4.0 - 6.0)
Missing	3 (0.4%)	1 (0.1%)

In both India and Nepal, the seroprevalence of scrub typhus infection increased with age (Figure 3). In Vellore, India the overall seroprevalence was 29.8% (95% CI 26.5-33.2): growing from 10.2% (95% CI 2.4-17.9) among 18-29-year-olds to 38.1% (95% CI 33.0-43.1) among 50 to 89-year-olds. In Kathmandu and Kavre, Nepal, the overall seroprevalence was 1.2% (95% CI 0.5-1.8), rising from 0.1% (95%CI 0-0.3) among 0 to 17-year-olds to 5.2% (95%CI 2.3 – 8.1%) among 18-29-year-olds (Table 3).

Table 3: Age-specific seroprevalence and seroincidence in Nepal and India

| | | | | Seroincidence per 1000 person-years (95% CIs) | | |
Age	N	N IgG seropositive	IgG seroprevalence (95% CI)	IgG antibody kinetics	IgG and IgM antibody kinetics	Derivative of age-dependent IgG seroprevalence
India						
0-17	0	0	–	–	–	–
18-29	59	6	10.2% (2.4-17.9)	8 (4-16)	11 (6-18)	4 (2-10)
30-49	302	72	23.8% (19.0-28.7)	14 (11-18)	12 (10-15)	7 (5-9)
50-89	360	137	38.1% (33.0-43.1)	26 (21-32)	20 (17-24)	8 (6-9)
Overall	721	215	29.8% (26.5-33.2)	18 (16-21)	16 (14-18)	7 (6-8)
Nepal						
0-17	876	1	0.1% (0-0.3)	2 (1-4)	–	0 (0-1)
18-29	229	12	5.2% (2.3-8.1)	9 (6-14)	–	3 (1-4)
30-49	0	0	–	–	–	–
50-89	0	0	–	–	–	–
Overall	1105	13	1.2% (0.5-1.8)	4 (3-6)	–	1 (1-2)

In Vellore, India, the overall scrub typhus seroincidence rate was 18 per 1000 person-years (95% CI: 16 - 21) using IgG antibody dynamics. Seroincidence rose with age; growing from 8 (95% CI 4 -

16) among 18-29 year-olds to 26 (95% CI 21 - 32) among 50-89-year-olds. In the Kathmandu

valley of Nepal, the overall seroincidence was 4 (95% CI 3 – 6) per 1000 person-years, also rising

with age from 2 (95% CI 1 - 4) per 1000 person-years among 0 to 17-year-olds to 9 (95%CI 6 -14)

among 18 to 29-year-olds. In both Nepal and India, the overall seroincidence using antibody

dynamics was more than double the seroincidence derived from the age-dependent IgG-

seroprevalence, 2.6 times higher in India and 4 times higher in Nepal (Table 3). IgM responses

were only available from the study in India; there, the seroincidence using IgM was similar to IgG.

The confidence intervals using IgM and IgG together were more narrow than using IgG alone.

In both sites, scrub typhus seroincidence varied by resident zones. In Vellore, India, seroincidence

was highest in periurban settings across all age groups, followed by rural and then urban, whereas

in Nepal, seroincidence was higher in urban settings (Figure 4). In India, females had slightly

elevated seroincidence rates compared to males across all age groups, although the confidence

intervals overlapped (Figure 4).

Discussion

In this study, we developed and applied a novel analytic approach to estimate the seroincidence of

scrub typhus using antibody decay information from confirmed cases in Thailand and India and

cross-sectional population serosurveys in Nepal and India. Seroincidence, the number of new

infections in a population per year, characterizes transmission intensity and is highly informative for

determining where and among whom infection burden is highest. We found a high seroincidence

of scrub typhus in both study populations, underscoring the importance of addressing the burden of

scrub typhus infection in these regions. A key finding of our study involves delineating the

persistence of antibodies after infection, characterizing their decay rate and shape, and using this

information to interpret population-level responses. Our results demonstrate that employing the

cutoff method to estimate seroprevalence while disregarding antibody decay underestimates the

true burden of scrub typhus infection.

Most cross-sectional seroepidemiologic studies on scrub typhus rely on seroprevalence as a

marker of infection, employing a cutoff threshold to dichotomize IgG responses and calculating the

proportion of the population with values above the threshold (6,8,31). Antibody responses in a

population at any given time will be a function of exposure, the age of the population, and antibody

persistence. For example, in our study populations, the overall seroprevalence was much higher in

India than in Nepal; however, in India, the median age of patients was 49 years compared to the

median age of 11 years in Nepal. Despite differences in overall seroprevalence rates, the

discrepancies in age-specific seroprevalence rates between the two populations were less marked.

This underlines the importance of considering age-specific seroprevalence rates when comparing

seroprevalence between different populations, as the age distribution can have a considerable

impact on the observed seroprevalence. Our novel analytical approach, which accounts for age

and antibody decay, provides a more accurate representation of scrub typhus infection in these

regions, highlighting the need for targeted public health interventions and research.

In both India and Nepal, seroprevalence increased notably with age, a finding that has been

consistently observed in other scrub typhus serosurveys (17,32). An increase in seroprevalence

with age can result from gradual accumulating in IgG responses over time or from a genuinely

elevated risk of infection in older age groups. Both arguments have been put forth for scrub typhus

seroepidemiology, with the suggestion that elevated risk in older ages is attributable to changes in

skin health, immune system function, and physiological processes (29). We argue that reporting

seroincidence rather than seroprevalence is essential to disentangling the effects of IgG

accumulation over age versus age-specific risk. Seroincidence, which measures the number of

new infections in a population per year, characterizes transmission intensity and is highly

informative for determining where and at what ages the infection burden is highest. Incorporating

antibody dynamics into the calculation of seroincidence, as demonstrated in this paper, further

elucidates age-specific changes in infection risk by accounting for decaying antibody levels. In our

study populations, we observed a marked increase in seroprevalence with age, while the increase

in seroincidence at older ages was comparatively more gradual. For example, in India, individuals

aged 50-89 had 3.8 times higher IgG seroprevalence compared to 18-29 year-olds compared to

1.8 times higher seroincidence using IgG and IgM antibody dynamics. This highlights the

importance of using seroincidence to better understand the age-specific risk of infection and to

inform targeted public health interventions.

Defining antibody dynamics and persistence is crucial for interpreting population-level

seroresponses to Orientia tsutsugamushi and modeling seroincidence. In our study, we found that

median IgG responses remained elevated above a mixture-model cutoff of 1.7 OD for nearly two

years. These findings align with a recent study by Schmidt et al., which reported that half of the

cases had persistent IgG antibodies beyond two years (29). Interestingly, IgM responses persisted

longer than anticipated. We observed that half of the participants had an OD <1 after 5.3 months

after symptom onset, and 90% of participants had an OD <1 for 26.5 months. This extended

persistence contrasts with Schmidt et al's findings, where half of the participants had IgM OD <1

after 2.7 months and 90% had OD <1 after 7.7 months. A potential explanation for this discrepancy

could be the difference in the duration of sample collection between the two studies. In the Schmidt

study, samples were collected up to a year after infection, while our study included samples

collected up to 3.5 years after infection. Another explanation could be the differences in modeling

approaches. Our power-function model allows for the shape of the decay to be non-exponential,

permitting a slower decay at longer times since infection. In contrast, a restricted cubic spline

approach may require a constant rate and shape of decay. It would be ideal to test these

approaches with the same data in future research. The persistence of IgM revealed in our study

suggests that it would be a good complement to IgG to identify recent infections and improve the

precision around seroincidence estimates. Indeed, for the India data where IgM results were

available, the seroincidence estimates using both IgM and IgG were consistent, yielding similar

estimates for the seroincidence, thus strengthening the validity of our approach. The seroincidence

estimates using IgM and IgG had narrower confidence intervals than those using IgG alone.

One of the key strengths of our approach lies in its ability to account for the inherent heterogeneity

of antibody responses, as well as measurement and biologic noise. Traditional seroepidemiological

methods, which rely on a single cutoff value for seroprevalence and seroincidence, fail to

incorporate these sources of variability. By explicitly integrating the heterogeneity in antibody

responses, our method captures the distribution of responses among individuals across ages,

clinical severity and levels of immune function. Additionally, our approach takes into account the

measurement noise associated with the ELISA assay and the biologic noise that stems from non-

specific antibody binding. This more comprehensive incorporation of uncertainty leads to more

accurate and robust estimates of scrub typhus seroincidence, which in turn better informs public

health interventions and research priorities.

Our findings have several implications for public health. First, we reveal a substantial subclinical burden

of scrub typhus beyond what is captured by the reporting of clinical cases. In India, a recent

population-based fever surveillance study conducted reported a clinical incidence of 0.8 cases per 1000

person-years, defined as the incidence of scrub typhus among febrile individuals who sought care or

were hospitalized (32). By applying our seroincidence estimates derived from IgG and IgM antibody

dynamics, we estimate that there are around 14 subclinical infections for every clinical case in the

population surrounding CMC Hospital. This inflation factor is less than estimated by Devamani et al.,

who reported a seroincidence of 44 per 1000 person-years. This study was conducted in a population

that was purposefully selected for its elevated numbers of clinical scrub typhus cases, which could

explain the differences (30). Population-based clinical incidence estimates for scrub typhus are not

currently available for the study population in Nepal but would be valuable in distinguishing clinical

versus sub-clinical infections. While individual health implications of subclinical infections are currently

unclear, seroincidence studies like ours are useful for identifying areas with substantial scrub typhus

burden, thereby informing the allocation of resources for enhanced clinical diagnostic capacity and

targeted surveillance efforts. Furthermore, our study challenges the notion that scrub typhus is solely a

rural disease. We observed elevated incidence rates in both urban and peri-urban communities in

Nepal and India. However, we could not ascertain whether exposure occurred at the place of residence

or was linked to rural exposure, highlighting the need for further research targeting these populations

and investigating exposures. Expanding our understanding of the distribution and risk factors

associated with scrub typhus in various settings will help inform the development of more effective and

targeted prevention and control measures, ultimately reducing the burden of disease in affected

populations.

There are several limitations to our study that warrant consideration. First, the population

serosurveys in India and Nepal were conducted over different time periods, with the Indian survey

spanning a few months and the Nepalese survey taking place over a year. This discrepancy could

introduce variability in the observed seroprevalence and seroincidence rates between the two

populations due to seasonal differences in scrub transmission. Second, our study assumes that the

antibody decay patterns observed in the cohorts from Thailand and India are applicable to the

populations in Nepal and India. This may not necessarily hold true, as factors such as age,

geography, clinical severity, and population-specific immune characteristics could lead to variations

in antibody decay patterns. Third, we estimated antibody dynamics using cases from a wide range

of ages. Although our sample size was not large enough to investigate age-specific variability in

kinetics, it is possible that these dynamics may vary with immune function at different ages,

presenting an interesting avenue for future research. Additionally, our findings may be influenced

by the overrepresentation of higher-risk groups in our data, such as women and urban residents,

which could affect the population-level seroincidence estimates. Future studies may benefit from

adjusting for potential biases introduced by overrepresented demographic groups. Lastly, we do

not have sufficient data to investigate whether antibody decay rates vary by age, which could

influence our seroincidence estimates. Despite these limitations, our study offers valuable insights

into the seroepidemiology of scrub typhus and highlights the importance of accounting for antibody

dynamics and heterogeneity in such analyses.

Scrub typhus remains an important but underrecognized etiology of acute fever with endemicity

expanding globally. There is a critical need for low-cost, accurate tools to quantify the burden of

scrub typhus infections to inform public health decision-making. We describe a sero-surveillance

approach that can efficiently generate population-level scrub typhus seroincidence estimates from

cross-sectional serosurveys. This methodology offers a valuable new tool for informing targeted

prevention and control strategies, ultimately contributing to a more effective response to scrub

typhus in endemic regions worldwide.

Acknowledgments

We gratefully acknowledge the study participants for their valuable time and interest in participating in these studies.

Funding

This work was supported by the Fogarty International Center at National Institutes of Health [K01 TW012177-01A1]

466　**Figures**
467
Figure 1: Quantitative antibody responses among population serosurveys in Nepal and India.
Anti-OT IgM and IgG responses are measured using kinetic enzyme-linked immunosorbent assays
(ELISAs) and depicted on the y-axis of all plots. The horizontal dashed line denotes the mixture model
cutoff. Panel A shows the distribution of responses among the population-based samples in India and
Nepal. Panel B shows the antibody among the population-based samples in India and Nepal as a
function of age.
474

Figure 2: Kinetics of IgM responses among Scrub Typhus cases. Longitudinal antibody dynamics were modeled from ELISA-measured antibody responses using Bayesian hierarchical models. The points are the observed individual antibody concentrations; each point indicates one sample; a patient's samples are connected with light purple lines. The dark solid lines indicates the median, and the 95% credible intervals for the model-fitted antibody decay concentrations. The horizontal dashed line denotes the mixture model cutoff.

Figure 3: Age-dependent seroprevalence
Seroprevalence is depicted as a function of age. Seroprevalence was determined using the
mixture-model derived cutoff for IgG responses. The age-dependent seroprevalence was modeled
using a generalized additive model with a cubic spline for age and simultaneous confidence
intervals using a parametric bootstrap of the variance-covariance matrix of the fitted model
parameters.

Figure 4: Seroincidence by age plus (A) resident zones and (B) gender
541 Seroincidence rates across are shown across age strata, residence zone (A), and gender (B)
542 comparing the India and Nepal population samples. Seroincidence is estimated using IgG antibody
543 kinetics. The point represents the incidence estimate per 1000 person-years, and the bar indicates
544 the 95% confidence interval.

(A)

(B)

545
546

547
Supplemental material

549
550 Supplemental Figure 1: Sampling collection over time and geographic unit

552
553
554 **Supplemental Table 1: Modeled antibody kinetic parameters**

	Median (95% Credible interval)	
Paramenter	**IgG**	**IgM**
Decay rate (1/years)??	0.11 (0.04-0.29)	1.70 (0.77-3.77)
Decay shape	1.34 (1.17-1.68)	2.05 (1.76-2.45)
Time to peak (days)??	11.04 (5.66-20.55)	5.19 (3.36-7.95)
Baseline response (OD)??	0.13 (0.07-0.24)	0.06 (0.04-0.10)
Peak response (OD)??	2.80 (2.10-3.73)	5.12 (4.00-7.03)

556
557
558
559

References

1. Paris DH, Shelite TR, Day NP, Walker DH. Unresolved Problems Related to Scrub Typhus: A Seriously Neglected Life-Threatening Disease. The American Journal of Tropical Medicine and Hygiene. 2013 Aug 7;89(2):301–7.

2. Weitzel T, Dittrich S, López J, Phuklia W, Martinez-Valdebenito C, Velásquez K, et al. Endemic Scrub Typhus in South America. N Engl J Med. 2016 Sep 8;375(10):954–61.

3. Izzard L, Fuller A, Blacksell SD, Paris DH, Richards AL, Aukkanit N, et al. Isolation of a Novel Orientia Species (O. chuto sp. nov.) from a Patient Infected in Dubai. Journal of Clinical Microbiology. 2010 Dec;48(12):4404–9.

4. Thiga JW, Mutai BK, Eyako WK, Ng'ang'a Z, Jiang J, Richards AL, et al. High Seroprevalence of Antibodies against Spotted Fever and Scrub Typhus Bacteria in Patients with Febrile Illness, Kenya. Emerg Infect Dis. 2015 Apr;21(4):688–91.

5. Luce-Fedrow A, Lehman ML, Kelly DJ, Mullins K, Maina AN, Stewart RL, et al. A Review of Scrub Typhus (Orientia tsutsugamushi and Related Organisms): Then, Now, and Tomorrow. Trop Med Infect Dis. 2018 Jan 17;3(1):8.

6. Bonell A, Lubell Y, Newton PN, Crump JA, Paris DH. Estimating the burden of scrub typhus: A systematic review. PLOS Neglected Tropical Diseases. 2017 Sep 25;11(9):e0005838.

7. John R, Varghese GM. Scrub typhus: a reemerging infection. Current Opinion in Infectious Diseases. 2020 Oct;33(5):365–71.

8. Xu G, Walker DH, Jupiter D, Melby PC, Arcari CM. A review of the global epidemiology of scrub typhus. PLOS Neglected Tropical Diseases. 2017 Nov 3;11(11):e0006062.

9. Saraswati K, Phanichkrivalkosil M, Day NPJ, Blacksell SD. The validity of diagnostic cut-offs for commercial and in-house scrub typhus IgM and IgG ELISAs: A review of the evidence. PLOS Neglected Tropical Diseases. 2019 Feb 4;13(2):e0007158.

10. Wangrangsimakul T, Greer RC, Chanta C, Nedsuwan S, Blacksell SD, Day NPJ, et al. Clinical Characteristics and Outcome of Children Hospitalized With Scrub Typhus in an Area of Endemicity. Journal of the Pediatric Infectious Diseases Society. 2020 Apr 30;9(2):202–9.

11. Lim C, Blacksell SD, Laongnualpanich A, Kantipong P, Day NPJ, Paris DH, et al. Optimal Cutoff Titers for Indirect Immunofluorescence Assay for Diagnosis of Scrub Typhus. Journal of Clinical Microbiology. 2015 Nov;53(11):3663–6.

12. Jiang J, Chan TC, Temenak JJ, Dasch GA, Ching WM, Richards AL. DEVELOPMENT OF A QUANTITATIVE REAL-TIME POLYMERASE CHAIN REACTION ASSAY SPECIFIC FOR ORIENTIA TSUTSUGAMUSHI. The American Journal of Tropical Medicine and Hygiene. 2004 Apr 1;70(4):351–6.

13. Blacksell SD, Tanganuchitcharnchai A, Nawtaisong P, Kantipong P, Laongnualpanich A, Day NPJ, et al. Diagnostic Accuracy of the InBios Scrub Typhus Detect Enzyme-Linked

Immunoassay for the Detection of IgM Antibodies in Northern Thailand. Clinical and Vaccine
Immunology [Internet]. 2015 Dec 9 [cited 2022 Feb 20]; Available from:
https://journals.asm.org/doi/abs/10.1128/CVI.00553-15

14. Paris DH, Blacksell SD, Nawtaisong P, Jenjaroen K, Teeraratkul A, Chierakul W, et al.
Diagnostic Accuracy of a Loop-Mediated Isothermal PCR Assay for Detection of Orientia
tsutsugamushi during Acute Scrub Typhus Infection. PLOS Neglected Tropical Diseases. 2011
Sep 13;5(9):e1307.

15. Varghese GM, Rajagopal VM, Trowbridge P, Purushothaman D, Martin SJ. Kinetics of IgM and
IgG antibodies after scrub typhus infection and the clinical implications. International Journal of
Infectious Diseases. 2018 Jun 1;71:53–5.

16. Aiemjoy K, Seidman JC, Saha S, Munira SJ, Sajib MSI, Sium SMA, et al. Estimating typhoid
incidence from community-based serosurveys: a multicohort study. The Lancet Microbe
[Internet]. 2022 Jun 21 [cited 2022 Jun 23];0(0). Available from:
https://www.thelancet.com/journals/lanmic/article/PIIS2666-5247(22)00114-8/fulltext

17. Trowbridge P, P. D, Premkumar PS, Varghese GM. Prevalence and risk factors for scrub
typhus in South India. Tropical Medicine & International Health. 2017;22(5):576–82.

18. Bickel PJ, Klaassen CA, Bickel PJ, Ritov Y, Klaassen J, Wellner JA, et al. Efficient and
adaptive estimation for semiparametric models. Vol. 2. Springer New York; 1998.

19. Arnold BF, Martin DL, Juma J, Mkocha H, Ochieng JB, Cooley GM, et al. Enteropathogen
antibody dynamics and force of infection among children in low-resource settings. Ferguson
NM, Jit M, White M, Leung DT, Azman A, editors. eLife. 2019 Aug 19;8:e45594.

20. Jewell NP, van der Laan M. Generalizations of current status data with applications. Lifetime
Data Anal. 1995 Mar 1;1(1):101–9.

21. Wood SN. Generalized additive models: an introduction with R. Chapman and Hall/CRC; 2006.

22. Marra G, Wood SN. Coverage properties of confidence intervals for generalized additive model
components. Scandinavian Journal of Statistics. 2012;39(1):53–74.

23. de Graaf WF, Kretzschmar MEE, Teunis PFM, Diekmann O. A two-phase within-host model for
immune response and its application to serological profiles of pertussis. Epidemics. 2014
Dec;9:1–7.

24. Teunis PFM, van Eijkeren JCH, de Graaf WF, Marinović AB, Kretzschmar MEE. Linking the
seroresponse to infection to within-host heterogeneity in antibody production. Epidemics. 2016
Sep;16:33–9.

25. Teunis PFM, van der Heijden OG, de Melker HE, Schellekens JFP, Versteegh FGA,
Kretzschmar MEE. Kinetics of the IgG antibody response to pertussis toxin after infection with
B. pertussis. Epidemiol Infect. 2002 Dec;129(3):479–89.

26. Hornik K, Leisch F, Zeileis A. JAGS: A program for analysis of Bayesian graphical models
using Gibbs sampling. In: Proceedings of DSC. 2003.

27. Teunis PFM, van Eijkeren JCH. Estimation of seroconversion rates for infectious diseases:
Effects of age and noise. Stat Med. 2020 Sep 20;39(21):2799–814.

28. Teunis PFM, van Eijkeren JCH, Ang CW, van Duynhoven YTHP, Simonsen JB, Strid MA, et al.
Biomarker dynamics: estimating infection rates from serological data. Stat Med. 2012 Sep
10;31(20):2240–8.

29. Schmidt WP, Devamani CS, Rose W, Alexander N, Prakash JAJ. Antibody response following
scrub typhus infection: clinical cohort study. Tropical Medicine & International Health.
2019;24(12):1455–64.

30. Devamani CS, Prakash JAJ, Alexander N, Stenos J, Schmidt WP. The incidence of Orientia
tsutsugamushi infection in rural South India. Epidemiol Infect. 2022 Jun 29;150:e132.

31. Chanyasanha C, Kaeburong K, Chenchittikul M, Sujirarat D. Seroprevalence of scrub typhus
infection in patients with pyrexia at some malaria clinics in three western provinces of Thailand.
Asian Pac J Allergy Immunol. 1998;16(2–3):119–25.

32. Devamani CS, Prakash JAJ, Alexander N, Suzuki M, Schmidt WP. Hospitalisations and
outpatient visits for undifferentiated fever attributable to scrub typhus in rural South India:
Retrospective cohort and nested case-control study. PLoS Negl Trop Dis. 2019 Feb
25;13(2):e0007160.