Assessing the accuracy of California county level COVID-19
hospitalization forecasts to inform public policy decision making

Lauren A. White*[1], Ryan McCorvie[1], David Crow[1], Seema Jain[1], Tomás M. León[1]

[1]Division of Communicable Disease Control, California Department of Public Health

Corresponding Author: *Lauren A. White

Email: lauren.white@cdph.ca.gov

Author Contributions: LW and TL designed research and wrote the paper. LW performed the
research and analyzed data. RM and DC contributed new analytic tools. RM, DC, and SJ
revised/edited the manuscript. SJ supervised the project.

Competing Interest Statement: The findings and conclusions in this article are those of the
author(s) and do not necessarily represent the views or opinions of the California Department of
Public Health or the California Health and Human Services Agency.

Keywords: infectious disease modeling, forecasting, model evaluation, COVID-19, public health

Abstract

The COVID-19 pandemic has highlighted the role of infectious disease forecasting in informing public policy. However, significant barriers remain for effectively linking infectious disease forecasts to public health decision making, including a lack of model validation. Forecasting model performance and accuracy should be evaluated retrospectively to understand under which conditions models were reliable and could be improved in the future. Using archived forecasts from the California Department of Public Health's California COVID Assessment Tool (https://calcat.covid19.ca.gov/cacovidmodels/), we compared how well different forecasting models predicted COVID-19 hospitalization census across California counties and regions during periods of Alpha, Delta, and Omicron variant predominance. Based on mean absolute error estimates, forecasting models had variable performance across counties and through time. When accounting for model availability across counties and dates, some individual models performed consistently better than the ensemble model, but model rankings still differed across counties. Local transmission trends, variant prevalence, and county population size were informative predictors for determining which model performed best for a given county based on a random forest classification analysis. Overall, the ensemble model performed worse in less populous counties, in part because of fewer model contributors in these locations. Ensemble model predictions could be improved by incorporating geographic heterogeneity in model coverage and performance. Consistency in model reporting and improved model validation can strengthen the role of infectious disease forecasting in real-time public health decision making.

Significance Statement

Spurred by the COVID-19 pandemic, infectious disease forecasting has gained prominence as a source of public health intelligence that ultimately may shape public health policy. Importantly, validation of forecast results is a critical, but often missing step to refine evidence-based decision making. We conducted a retrospective analysis of forecasts from the California Department of Public Health's California COVID Assessment Tool. Model performance was variable across counties, and the best performing model could be predicted by local transmission dynamics, variant prevalence, and county population size. Less populous counties had fewer model contributors and generally had higher ensemble model error. Ensemble model performance could be improved by capturing these county-level differences and by incentivizing model coverage in less populous and underserved regions.

Main Text

Introduction

In public health, forecasting has been used to predict infectious disease dynamics for a variety of diseases including influenza, dengue fever, Ebola virus disease, Zika fever, and most recently COVID-19, which has highlighted the importance of infectious disease modeling to help inform public health decision making (1). Nevertheless, significant barriers remain for effectively linking infectious disease forecasts with public health decision making including a lack of model standardization and validation, and difficulty in successfully communicating model complexity and uncertainty (2). Moreover, public health practitioners may need different outcomes or indicators than what forecast models provide (2, 3).

In June 2020, as part of the COVID-19 response, the California Department of Public Health's (CDPH) COVID Modeling Team launched the [California COVID-19 Assessment Tool (CalCAT)](#) to compile available COVID-19 models, mostly from academic groups, to inform policy and public health action (4). CalCAT provides nowcasts (R-effective estimates), forecasts (short-term predictions for hospitalizations, ICU admissions, and deaths), and longer-range scenario models for a variety of COVID indicators at the state, regional, and county scales. Some contributors are national, forecasting for all states, while others focus only on California and may not be publicly available elsewhere (Table 1).

During the COVID-19 pandemic response, many California local health jurisdictions communicated the importance of forecasts focused on the relevant scale of decision making (e.g., county- vs. state-level forecasts) because there was significant geographic heterogeneity in COVID-19 outcomes at regional and local levels (5). A better understanding of how forecasting models have captured these geographical heterogeneities could help inform local public health decision making during future COVID-19 waves and enable local health jurisdictions to employ models judiciously given proven past performance. Lessons learned from COVID-19 forecasting efforts can also be applied to future modeling for other diseases including influenza.

We retrospectively evaluated archived forecasting predictions from CalCAT for models that consistently provided county-level hospitalization census predictions across a year long period from February 1, 2021 to February 1, 2022 (Table 1). To explore the effects of COVID-19 variants on model performance within that period, we also compared forecasting model accuracy during three phases of the COVID-19 pandemic at the county and regional level in California (Figure 1 A-C) with different variant predominance: Alpha (April 22- June 1, 2021), Delta (June 21 - September 1, 2021), and Omicron (December 21, 2021 - February 1, 2022). These periods also differed in their hospitalization burden (Figure 1A) and epidemic growth rates (Figure 1C).

Multiple methods exist for measuring epidemic forecast accuracy including metrics that evaluate specific point estimates and uncertainty (6). When full predictive estimates are available, metrics like the logarithmic score or continuous ranked probability score provide context for probabilistic models' predictions and uncertainty. When forecasts are provided in quantile/interval formats, the weighted interval score is a potential alternative (6). Since not all models incorporated into CalCAT provided full predictive or interval estimates, we focused on the median point estimates (50th percentile) from forecasting models for hospital census at the county scale. In addition to these models, we retroactively created a baseline California forecast that projected forward the 7-day rolling mean from the prior week. Each forecast has the following properties: (1) model (m): the organization or group issuing the forecast (Table 1); (2) location (j): the geographic location for which a forecast is issued (in this case, at the county-level): (3) publication date (i): the date that the forecast was displayed on CalCAT; and (4) target end date (k): the future forecast horizon date for which the prediction is made.

We utilized mean absolute error (MAE) and relative error at the 7-, 14-, and 21-day
forecast horizons to evaluate the accuracy of these point estimates. To better compare across
counties with different population sizes, we normalized both error types by the median hospital
capacity of each county (14-day horizon results are highlighted in the main text; the remaining
forecast horizons are provided in the Supporting Information). From the MAE scores, we
computed a standardized ranking score for every forecasted observation relative to other models
issuing a prediction for that same publication date and location (7). In addition, we also conducted
pairwise tournaments of model performance to control for the frequency of model participation.
Finally, using a classification regression approach, we explored which county-level
epidemiological and socio-economic covariates could help explain the "winning" model for a given
location and date based on the MAE scores for a given forecast horizon.

County results are grouped by health officer regions, which are contiguous groupings of
58 counties used for health mandates in California (Figure 1D): Association of Bay Area Health
Officers (ABAHO); Greater Sacramento Region Health Officers (GSRHO); Rural Association of
Northern California Health Officers (RANCHO); San Joaquin Valley Consortium (SJVC); and
Southern California (SCAL). Some counties do not have major hospitals and therefore lack
forecasting predictions, actual numbers of hospitalizations, or both. For this reason, Alpine,
Sierra, and Sutter Counties are not included in the analyses that follow.

Results

Model performance varied across locations and under different periods of variant
predominance

Model performance was heterogeneous across counties and during different periods of
variant predominance (Figure 2A, 3A, 4A), in part reflecting that the number of models available
for a given publication date and location varied through time; fewer models were available during
the Omicron variant period and for less populous health officer regions such as RANCHO (SI
Figures S3, S11, S15). For example, in Trinity County – one of California's least populous
counties – the Simple Growth model had the lowest 14-day normalized MAE for most forecast
publication dates during the Alpha and Omicron predominant periods (Figure 2A, 4A), whereas
the Columbia model had the lowest 14-day normalized MAE during the Delta period (Figure 3A).
In San Diego County, California's second most populous county, the LEMMA model had the
lowest 14-day normalized MAE during the Alpha period (Figure 2A), and the COVID NearTerm
model had the lowest 14-day MAE for the most days during Delta and Omicron periods (Figure
3A & 4A). Overall, the Simple Growth model performed particularly well in the RANCHO region
during the Omicron period as demonstrated by a lower 14-day MAE for many counties in that
region (Figure 4A). The LEMMA model had the lowest 14-day MAE across many regions during
the Omicron period on or after January 13, 2022 (Figure 4A). In general, the magnitude of relative
error increased with longer time horizons and during the Omicron period (SI Figures1-2). During
the Omicron period, most models had median relative error values less than zero, indicating a
tendency for underprediction (SI Figure 1).

The sum of the standardized rank score ($\Sigma sr_{m,i,j}$) in each county, j, rewards both
performance (model accuracy) and frequent participation (model coverage). During the Alpha
period, the LEMMA model had the highest score in 20/55 counties with the Ensemble model a
close second with the highest score in 17/55 counties (Figure 2B). During the Delta period, the
Ensemble model had the highest score in 21/55 counties (Figure 3B). During the Omicron period,
the Ensemble model had the highest score in 22/55 counties, and the Simple Growth model was
a close second with 20/55 counties (Figure 4B).

The density distributions of standardized rank ($sr_{m,i,j}$) allow for comparison of model
performance while controlling for frequency of model participation (Figures 2C, 3C, 4C). Although

COVID NearTerm did not have the highest sum of the standardized rank score in any counties, it
had the highest median standardized rank score during the Alpha and Delta periods (Figures 2C,
3C). The LEMMA model had the highest median standardized. rank score during the Omicron
period (Figure 4C). The same pattern of ranking was present for 7-day MAE (SI Figures 4-6). For
21-day MAE, the COVID NearTerm model had the highest median standardized rank score
during the Alpha and Omicron periods (SI Figures 16 & 18), while the LEMMA model had the
highest median rank scores during the Delta period (SI Figures 17).

When controlling for participation, some models outperformed the ensemble, but pairwise
model rankings varied across counties

When matching across all locations and all observation dates, two models– COVID
NearTerm and LEMMA– performed better in pairwise comparisons relative to the Ensemble
model for 14-day MAE (Figure 5A & B). However, pairwise rankings were quite variable when
disaggregated by county and also highlighted the differences in coverage and availability across
locations for different models (Figure 5C). For example, although the Simple Growth model came
fourth in the overall pairwise ranking (Figure 5A), it came first in twelve individual counties.
Similarly, the Columbia model came last in the overall pairwise ranking (Figure 5A) and generally
performed worse than average ($\theta_m > 1$), but performed better than average ($\theta_m < 1$) in Plumas
and Inyo counties (Figure 5C).

Overall pairwise rankings were robust to forecast horizon length for the complete analysis
period (SI Figures 5A & 17A). However, overall pairwise rankings were more unstable during
specific periods of variant predominance, particularly for shorter forecast horizons (SI Figures 6-8,
10-12, 18-20) and for county-specific rankings (SI Figures 7C-9C, 12C-14C, 19C-21C).

Epidemiological traits, county population size, and variant traits best predicted forecast
"winners"

For the entire analysis period (February 1, 2021-February 1, 2022), time-varying vaccine
coverage at the county-level, local transmission dynamics (R-effective and 7-day change in R-
effective), county population size, and regional proportion of variants, were most important in
predicting which model had the lowest MAE for a given county on a given publication date (SI
Figure 23). Other static socio-economic variables like income, percent unemployment, percent of
residents with a university degree, and percent of residents in poverty were less important for
predicting model outcomes. These variable importance rankings were robust to the forecast
horizon used for MAE calculations (SI Figure 23).

Less populated counties have ensemble predictions with higher median MAE and more
variable MAE

When comparing 14-day MAE normalized by hospital capacity, counties with smaller
population sizes typically had a higher median MAE score and more variable MAE distributions
compared to more populous counties (SI Figure 24B). Based on a linear regression, the
logarithmic of the normalized MAE score was negatively correlated with county population size
(coefficient estimate: $1.7 \cdot 10^{-7}$; p-value $< 2 \cdot 10^{-16}$). This relationship held true regardless of the
forecast horizon used for MAE calculations (SI Figure 24 A & C).

186
187

Discussion

Ensemble model could be improved by incorporating geographic heterogeneity in model coverage and performance

Echoing other analyses of COVID-19 forecast performance that have described a large variation in model accuracy by location (7, 8), forecasting models performed differentially across California counties and regions and for different periods of variant predominance during the COVID pandemic (Figures 2-4, 5C, SI Figures 4-6, 16-18). Moreover, location-specific features like local transmission dynamics or county population size helped explain model performance (SI Figure 24). This geographic variation in model performance points to the importance of location-specific model evaluation in order for local health jurisdictions to best employ forecasts for public health decision making.

In general, combining multiple models into ensembles allows for better performance (7, 9–12). However, in this case, COVID NearTerm and LEMMA consistently outperformed the Ensemble model when controlling for frequency of participation (Figures 2-4C, Figure 5), although pairwise ranking scores remained variable at the county level (Figure 5C). The higher performance of individual models over the Ensemble model combined with the variability in performance at the county-level suggests that the Ensemble model does not have to be applied uniformly across all locations; public health decision making could benefit from model selection and ensemble weighting that reflects location-specific past performance as well as local transmission trends (13).

Lower forecast coverage in less populated counties weakens evidence-based decision making

One interesting question from a public health decision making context is whether model coverage (i.e., frequent issuing of forecasts across all potential locations) and model accuracy should be weighed equally when establishing the criteria for a "winning" forecast. In this analysis, there was typically a mismatch between raw model performance based on availability as measured by the sum of standardized ranking (Figures 2-4) and model performance when controlling for participation via pairwise tournaments (Figure 5). In part, this disagreement reflects that not all models provided estimates for all counties, especially for less populous regions/counties (SI Figures 1, 9, 13). For example, the COVID NearTerm model ranked first in the pairwise ranking but provided no coverage for any counties in the less populous RANCHO region (Figure 5C). In contrast, the Ensemble model came first in the majority of counties during the Delta and Omicron periods as measured by sum of the standardized rank score for that period ($\Sigma sr_{m,i,j}$) (Figures 3B, 4B), but was generally outperformed in pairwise ranking evaluations both overall and for individual counties (Figure 5). Further highlighting the potential consequences of fewer contributors, the Ensemble model in less populous counties exhibited a higher median normalized MAE and a more variable normalized MAE regardless of the forecast horizon (SI Figure 22).

While maximizing model accuracy is important, a forecast cannot add value if it is not available for decision making. As county-level contributors are lost to attrition, ensemble estimates may further decrease in accuracy or may not be possible in these less populous counties. Policy and public health decision makers should evaluate what investments or innovations in modeling are needed to improve results for underserved counties with lower forecast coverage. In addition, decision makers could seek to incentivize the best-performing models to serve smaller counties that neither have the resources to do this work in-house nor have academic partners readily available.

The lack of coverage in smaller counties also points to the inherent complexities of
interpreting in hospitalization burden—since hospitalizations are typically recorded via hospital
location rather than patient residency (14). As others have suggested, forecasting at the
geographic unit of hospital referral networks could be another solution to low model coverage in
less populous counties (14).

Continuity of contributors, forecast structure, and documentation helps real-time public
health decision making and post-hoc analysis

The overall continuity of forecasting contributions has proved challenging for post-hoc
evaluation. Although CalCAT has had roughly ten unique forecast contributors through time,
many of these groups have ceased contributing as the COVID landscape has increased in
complexity (e.g., emerging variants, prior immunity, boosters). Although less relevant to
forecasting hospitalizations, changes in case ascertainment and testing practices make
retrospective analyses more challenging. Interruptions to forecast continuity can also limit post-
hoc evaluation. For example, some modeling groups paused forecasts in order to reset or
recalibrate for new variants like Omicron. While initiatives like the COVID-19 Forecast Hub have
worked to standardize forecast output and reporting (7), changing repository structures, file
nomenclature, and data formatting can also disrupt the archiving process necessary for ensemble
generation and subsequent post-hoc review. This analysis is a snapshot of what was available on
CalCAT—and therefore to the general public and public health decision makers—and may not
entirely reflect what model contributors would intend to be their contributing forecast at all times.
The CalCAT team updated data and results iteratively as often as possible, but not all model
changes were announced. As the COVID-19 pandemic necessitated rapid changes in data
reporting and data infrastructure, other information technology issues may have introduced
unintended errors.

The current classification regression analysis in this manuscript does not include model-
specific traits. In order to truly evaluate whether underlying model traits and assumptions help to
explain performance for specific locations through time, it would be necessary to have a larger
number of forecasting contributors and consistent metadata on both the changes in model
construction and the timing of those changes. Therefore, another potential area of documentation
might include not just existing model assumptions and structure but how those characteristics
have changed over time. This analysis may be easier to do at a state or national scale, where
more model contributors are available, and reporting is better standardized through initiatives like
the Forecast and Scenario Hubs.

Reporting and communicating infectious disease forecasting results, with all their
inherent uncertainty and complexity, remain areas for improvement and growth for public health
departments and their academic and industry collaborators to support evidence-based public
health policy planning and decision making. Importantly, forecasting models may also serve as a
communication tool to influence behavior change by the general public. One phenomenon not
explored in this analysis is the potential for forecasts to alter human behavior, and subsequently
model accuracy.

Major progress in infectious disease forecasting has been made during the COVID-19
pandemic, while ongoing challenges, such as those around data and communication, have
persisted. We retrospectively investigated hospitalization census model performance at the
county level in the state of California. Model performance and ranking varied through space and
time and by metric, highlighting the difficulty of making blanket recommendations for which
models to use for individual counties, including an ensemble approach. Calibrating based on past
model performance may help improve ensemble forecast generation, and counties may benefit
by considering which individual model contributors have historically served them the best. Going
forward, closer collaboration between forecasters, researchers, and policymakers may create

positive feedback loops that inform the ongoing COVID-19 response and other future public
health action.

285
Materials and Methods

Mean absolute error

The raw mean absolute error (MAE) for each publication date (i) with associated target
end dates k is calculated as: $\frac{1}{N}\sum_{k=i}^{i+N}|x_{i,k} - \hat{x}_k|$ where N is the number of days into the future that
the forecast is made, $x_{i,k}$ is the prediction made on publication date i for target end date k and
$\hat{x}_k$ is the actual observed value for a given target end date (7). We then standardized the MAE by
h, the median non-surge hospital capacity of a given county: $\frac{MAE}{h} \cdot 100$

Note that not all model forecasts were available for all counties or all dates. A model only
receives an MAE score for a given publication date if it has predictions available for the target end
dates of interest (e.g., to receive a 7-day MAE score, a model must have made predictions for 1-7
days ahead of the publication date). Here we used CA-state specific data (15) for post-hoc
evaluation, whereas many model teams may be relying on other data sources (e.g., Health and
Human Services) for fitting or calibration.

Relative error

The relative error for each publication date (i) with associated target end dates k is
calculated as: $\frac{1}{N}\sum_{k=i}^{i+N}(x_{i,k} - \hat{x}_k)$ where N is the number of days into the future that the forecast is
made, $x_{i,k}$ is the prediction made on publication date i for target end date k and $\hat{x}_k$ is the actual
observed value for a given target end date (7). We then standardized the relative error by h, the
median non-surge hospital capacity of a given county: $\frac{relative\ error}{h} \cdot 100$

Therefore, a positive relative error corresponds to a model overestimating the hospital census,
while a negative relative error corresponds to an underestimation.

Standardized ranking score

For each publication date i and location j, we calculated a standardized rank for every
available model m based on its associated MAE: $sr_{m,i,j} = 1 - \frac{r_{m,i,j}-1}{n_{i,j}}$ where $r_{m,i,j}$ is the ranking of
the MAE of model m out of the n other models that made predictions for publication date i and
location j (adapted slightly from (7)). Thus, for a given publication date i and location j, the
highest possible standardized ranking score for any given model is 1 and the lowest is the inverse
of the lowest possible ranking ($1/n_{i,j}$). Models not participating for a given publication date i and
location j receive a zero.

Pairwise tournament

To conduct a pairwise tournament, we calculated a relative MAE for each pair of models
m and m': $\theta_{m,m'} = median\left\{\frac{MAE(m;\ i,j,k)}{MAE(m';\ i,j,k)}\right\}$ where $\theta_{m,m'}$ is the median of the ratio of the
simultaneously available MAE scores or model m to model m' with shared publication dates i,
target end dates k, and locations j (16). Importantly, the common locations, publication dates,

and observation dates may differ for each pair of models *m* and *m'*. This approach varies slightly
from some previous examples, as the order of operations is scale than aggregate rather than
aggregate than scale (7, 17).

An overall performance score of a given model, *m* is then calculated as the geometric
mean of all relative MAE scores: $\theta_m = \left(\prod_{m'=1}^{M} \theta_{m,m'}\right)^{1/M}$ where *M* is the total number of all
models available for comparison. At the county level for counties with smaller hospital capacities,
there was a non-trivial probability of certain models achieving an MAE of zero, which leads to
relative MAE scores of zero or infinity depending on the order of comparison. To eliminate these
irregularities in the pairwise comparisons, we excluded counties with median non-surge hospital
capacities ≤ 25 (i.e., Calaveras, Lassen, Mariposa, Modoc, Mono, San Benito, and Trinity).

Random forest classification analysis

To explore whether the model with the lowest MAE score for a given location and
observation date could be explained by county-specific epidemiological or socioeconomic factors,
we conducted a random forest classification analysis. We explored the factors of: progressive
vaccination coverage at the county level, county-level R-effective, 7-day change in R-effective,
variant prevalence at the health officer region level (18), county population size, percent of county
residents in poverty (2019), percent unemployment (2020), median income (2019), five-year
average percentage completing college (university degree) (2015-2019), and 2013 Rural-Urban
Continuum code. All socioeconomic variables were taken from U.S. Department of Agriculture
Economic Research Service county-level data sets (19). For pre-processing, data were centered
and scaled using the caret package (20). For model training and tuning, 70% of original data was
used with K-fold validation (four-fold, repeated four times).

Data and code availability

The forecasts and R-effective values analyzed in this paper are available from CalCAT (4).
California-specific hospitalization data is available on the California Open Data Portal (15).
Because of reporting delays and backfilling, datasets used in the analysis may represent a
snapshot of what was available at that point in time. All data and code used for analysis and
figure generation is available in the public repository:
https://github.com/whit1951/CA_COVID_Forecasting_Accuracy. Analyses were performed in R (v
3.6.0) (22).
350
351 **Acknowledgments**
352

353 The authors thank the CMU Delphi Group including Ryan Tibshurani and Daniel J. McDonald,
354 MIDAS members, and the COVID-19 Forecasting Hub for discussion and feedback. The authors
355 also thank Californian local health jurisdictions and members of the CDPH Modeling and
356 Advanced Analytics team including Chris Hoover, Mugdha Thakur, Natalie Linton, Phoebe Lu,
357 Sindhu Ravuri, and Sophie Zhu for conversations and insights that improved these analyses.

358
359
360

References

1. A. L. Bertozzi, E. Franco, G. Mohler, M. B. Short, D. Sledge, The challenges of modeling and forecasting the spread of COVID-19. *Proc. Natl. Acad. Sci.* **117**, 16732–16738 (2020).

2. C. S. Lutz, *et al.*, Applying infectious disease forecasting to public health: a path forward using influenza forecasting examples. *BMC Public Health* **19**, 1659 (2019).

3. C. Doms, S. C. Kramer, J. Shaman, Assessing the Use of Influenza Forecasts and Epidemiological Modeling in Public Health Decision Making in the United States. *Sci. Rep.* **8**, 12406 (2018).

4. California Department of Public Health, California COVID Assessment Tool (2022). https://calcat.covid19.gov/cacovidmodels/

5. California State Government, California's commitment to health equity. *Calif. All* (2022) (May 3, 2022). https://covid19.ca.gov/equity/

6. J. Bracher, E. L. Ray, T. Gneiting, N. G. Reich, Evaluating epidemic forecasts in an interval format. *PLOS Comput. Biol.* **17**, e1008618 (2021).

7. E. Y. Cramer, *et al.*, Evaluation of individual and ensemble probabilistic forecasts of COVID-19 mortality in the United States. *Proc. Natl. Acad. Sci. U. S. A.* **119**, e2113561119 (2022).

8. N. G. Reich, R. J. Tibshirani, E. L. Ray, R. Rosenfeld, On the predictability of COVID-19 (2021) (June 3, 2022).

9. N. G. Reich, *et al.*, Accuracy of real-time multi-model ensemble forecasts for seasonal influenza in the U.S. *PLOS Comput. Biol.* **15**, e1007486 (2019).

10. M. A. Johansson, *et al.*, An open challenge to advance probabilistic forecasting for dengue epidemics. *Proc. Natl. Acad. Sci.* **116**, 24268–24274 (2019).

11. C. Viboud, *et al.*, The RAPIDD ebola forecasting challenge: Synthesis and lessons learnt. *Epidemics* **22**, 13–21 (2018).

12. The Influenza Forecasting Working Group, *et al.*, Collaborative efforts to forecast seasonal influenza in the United States, 2015–2016. *Sci. Rep.* **9**, 683 (2019).

13. E. L. Ray, N. G. Reich, Prediction of infectious disease epidemics via weighted density ensembles. *PLOS Comput. Biol.* **14**, e1005910 (2018).

14. R. Rosenfeld, R. J. Tibshirani, Epidemic tracking and forecasting: Lessons learned from a tumultuous year. *Proc. Natl. Acad. Sci. U. S. A.* **118**, e2111456118 (2021).

15. California Department of Public Health, COVID-19 Hospital Data. *Calif. Open Data Portal*. https://data.ca.gov/group/covid-19

16. A. Green, *et al.*, CMU Delphi Covid-19 Forecasts (2021). https://github.com/cmu-delphi/covid-19-forecast/tree/develop#delphi-forecasting-efforts

17.   J. Bracher, Evaluating probabilistic COVID19 forecasts under partial missingness: A
pairwise comparison approach (2020). https://covid19forecasthub.org/talks/2020-10-27-
Bracher_Pairwise_Comparisons.pdf

18.   California Department of Public Health, Variants. *https://covid19.ca.gov/variants/*.

19.   U.S. Department of Agriculture Economic Research Service, County-level Data Sets. *Data*
Prod. (2021) (March 23, 2022).

20.   M. Kuhn, caret: Classification and Regression Training (2021).
https://github.com/topepo/caret/

21.   S. Pei, J. Shaman, Initial Simulation of SARS-CoV2 Spread and Intervention Effects in the
Continental US (2020) https:/doi.org/10.1101/2020.03.21.20040303 (March 22, 2022).

22.   A. B. Olshen, *et al.*, "COVIDNearTerm: A Simple Method to Forecast COVID-19
Hospitalizations" (Infectious Diseases (except HIV/AIDS), 2021)
https:/doi.org/10.1101/2021.10.08.21264785 (March 22, 2022).

23.   J. Schwab, M. Peterson, Local Epidemic Modeling for Management and Action (LEMMA)
(2021). https://localepi.github.io/LEMMA/index.html

411
412
413

 Figures and Tables

Figure 1. Time courses of **(A)** California COVID-19 hospitalization census, **(B)** variant
prevalence, **(C)** statewide R-effective estimate, and **(D)** California health officer regions. The
period displayed for panels A-C corresponds to the complete period of analysis February 1, 2021-
February 1, 2022 used for pairwise tournament and random forest analysis. Shaded regions for
panels A:C correspond to the dates of analysis for the three variant predominant periods: Alpha,
Delta, and Omicron.

426
Figure 2. Forecasting accuracy results at the county level during the Alpha wave in California as
measured by mean absolute error (MAE). **(A)** Heat map of the best daily performing model for a
given prediction date as measured by 14-day MAE. Each cell in the heat map corresponds to a
normalized mean absolute error calculated for the day that a model forecast was published.
Counties are grouped into panels by California health officer regions. **(B)** A summary map of
California where the color of the county corresponds to the model with the highest sum of the
standardized rank score for that period ($\Sigma sr_{m,i,j}$). Note that by using the summation of the
standardized ranking score, models are penalized for lack of participation. **(C)** A density
distribution of the standardized rank score ($sr_{m,i,j}$) that depicts the median (dashed) and mean
(solid) as vertical lines for each model distribution.

Figure 3. Forecasting accuracy results at the county level during the Delta wave in California as measured by mean absolute error (MAE). **(A)** Heat map of the best daily performing model for a given prediction date as measured by 14-day MAE. Each cell in the heat map corresponds to a standardized mean absolute error calculated for the day that a model forecast was published. Counties are grouped into panels by California health officer regions. **(B)** A summary map of California where the color of the county corresponds to the model with the highest sum of the standardized rank score for that period ($\Sigma sr_{m,i,j}$). Note that by using the summation of the standardized ranking score models are penalized for lack of participation. **(C)** A density distribution of the standardized rank score ($sr_{m,i,j}$) that depicts the median (dashed) and mean (solid) as vertical lines for each model distribution.

Figure 4. Forecasting accuracy results at the county level during the Omicron wave in California as measured by mean absolute error (MAE). (**A)** Heat map of the best daily performing model for a given prediction date as measured by 14-day MAE. Each cell in the heat map corresponds to a standardized mean absolute error calculated for the day that a model forecast was published. Counties are grouped into panels by California health officer regions. (**B)** A summary map of California where the color of the county corresponds to the model with the highest sum of the standardized rank score for that period ($\Sigma sr_{m,i,j}$). Note that by using the summation of the standardized ranking score models are penalized for lack of participation. (**C)** A density distribution of the standardized rank score $(sr_{m,i,j})$ that depicts the median (dashed) and mean (solid) as vertical lines for each model distribution.

Figure 5. Pairwise tournament median rankings of models for the whole analysis period for 14-day MAE. **(A)** Overall median rankings (θ_m) across all locations and observation dates. **(B)** Median pairwise rankings $(\theta_{m,m'})$ comparing each model m relative to every other model m'. The grid is symmetrical, so the ratio of model m: model m' is the inverse score of the ratio of model m': model m. **(C)** Overall median rankings for all available observation dates disaggregated by county.

470 **Table 1.** Constituent models providing county-level hospitalization census predictions that are
471 archived on CalCAT and included in the analysis.
472

Model	Forecast update frequency	Methods/Approach	Documentation
Columbia	Weekly	County level metapopulation model	(21)
UCSF, COVID NearTerm	Daily	Bootstrap-based method based on an autoregressive model	(22)
UCB LEMMA	Daily	SEIR compartmental model with parameters fit using case series data of COVID-19 hospital and ICU census, hospital admissions, deaths, cases and seroprevalence	(23)
CDPH Simple Growth	Daily	Assumes new cases grow exponentially according to the rate given by the latest ensemble R-effective. Assumes a fixed severity and average length of stay to generate hospitalizations	(4)
CalCAT Ensemble	Daily	The ensemble forecast takes the median of all the forecasts available on a given date and fits a smoothed spline to the trend.	(4)
CA Baseline	Daily	Retroactive 7-day rolling average mean of past hospitalization values	Methods