Title: Community carriage of ESBL-producing *Escherichia coli* and *Klebsiella*

***pneumoniae*: A cross-sectional study of risk factors and comparative genomics of**

carriage and clinical isolates

Authors: Niclas Raffelsberger (ORCID 0000-0002-9463-8915)[a,b]*, Dorota Julia Buczek

(ORCID 0000-0002-2282-6298)[b], Kristian Svendsen (ORCID 0000-0003-3481-3539)[c], Lars

Småbrekke (ORCID 0000-0001-6475-3368)[c], Anna Kaarina Pöntinen (ORCID 0000-0002-

3160-2042)[d,e], Iren H. Löhr (ORCID 0000-0002-8847-0044)[f], Lotte Leonore Eivindsdatter

Andreassen[a], Gunnar Skov Simonsen (ORCID 0000-0003-0043-7045)[a,b], Norwegian *E. coli*

ESBL Study Group[g], Arnfinn Sundsfjord (ORCID 0000-0002-3728-2270)[b,d], Kirsten

Gravningen (ORCID 0000-0002-8194-3208)[a,h,i#], Ørjan Samuelsen (ORCID 0000-0002-

5525-2614)[c,d#]

[a]*Department of Microbiology and Infection Control, University Hospital of North Norway,*

Tromsø, Norway

[b]*Department of Medical Biology, Faculty of Health Sciences, UiT The Arctic University of*

Norway, Tromsø, Norway

[c]*Department of Pharmacy, Faculty of Health Sciences, UiT The Arctic University of Norway,*

Tromsø, Norway

[d]*Norwegian National Advisory Unit on Detection of Antimicrobial Resistance, Department of*

Microbiology and Infection Control, University Hospital of North Norway, Tromsø, Norway

[e]*Department of Biostatistics, University of Oslo, Oslo, Norway*

[f]*Department of Medical Microbiology, Stavanger University Hospital, Stavanger, Norway*

[g]*Collaborators forming the Norwegian E. coli ESBL Study Group are listed in the*

Acknowledgement.

*h*Department of Microbiology and Infection Control, Akershus University Hospital,

Nordbyhagen, Norway

*i*Division of Medicine and Laboratory Sciences, Institute of Clinical Medicine, University of

Oslo, Oslo, Norway

[#] equal contribution

*Corresponding author: Niclas Raffelsberger, Correspondence address: Department of

Microbiology and Infection Control, University Hospital of North Norway, N-9038 Tromsø,

Norway E-mail: niclas.raffelsberger@unn.no

Abstract

The global prevalence of infections caused by ESBL-producing Enterobacterales (ESBL-E) is

increasing and for *Escherichia coli* observations indicate that this is partly driven by

community-onset cases. The ESBL-E population structure in the community is scarcely

described and data on risk factors for carriage are conflicting. Here, we report the prevalence

and population structure of fecal ESBL-producing *E. coli* and *Klebsiella pneumoniae* (ESBL-

Ec/Kp) in a general adult population, examine risk factors, and compare carriage isolates with

contemporary clinical isolates.

Fecal samples obtained from 4999 participants (54% women) ≥40 years in the seventh survey

of the population-based Tromsø Study, Norway (2015-2016) were screened for ESBL-Ec/Kp.

In addition, we included 118 ESBL-Ec clinical isolates from the Norwegian surveillance

program in 2014. All isolates were whole-genome sequenced. Risk factors associated with

carriage were analyzed using multivariable logistic regression.

ESBL-Ec gastrointestinal carriage prevalence was 3.3% (95% CI 2.8-3.9%, no sex difference)

and 0.08% (0.02-0.20%) for ESBL-Kp. For ESBL-Ec, travel to Asia was the only

independent risk factor (AOR 3.47, 95% CI 2.18-5.51). *E. coli* ST131 was most prevalent in

both collections. However, the ST131 proportion was significantly lower in carriage (24%)

vs. clinical isolates (58%, p<0.001). Carriage isolates were genetically more diverse with a

higher proportion of phylogroup A (26% vs. 5%, p<0.001), indicating that ESBL gene

acquisition occurs in a variety of *E. coli* lineages colonizing the gut. STs commonly related to

extra-intestinal infections were more frequent in clinical isolates also carrying a higher

prevalence of antimicrobial resistance, which could indicate clone associated pathogenicity.

Keywords: *Escherichia coli*, *Klebsiella pneumoniae*, Extended-spectrum β-lactamase,

antimicrobial resistance, carriage, risk factors, general population, bacterial genomics

Importance

ESBL-producing *E. coli* (ESBL-Ec) and *K. pneumoniae* (ESBL-Kp) are major pathogens in

the global burden of antimicrobial resistance. However, there is a gap in knowledge

concerning the bacterial population structure of human ESBL-Ec/Kp carriage isolates in the

community. We have examined ESBL-Ec/Kp isolates from a population-based study and

compared these to contemporary clinical isolates. The large genetic diversity of carriage

isolates indicates frequent ESBL gene acquisition, while those causing invasive infections are

more clone dependent and associated with a higher prevalence of antibiotic resistance. The

knowledge of factors associated with ESBL carriage helps to identify patients at risk to

combat the spread of resistant bacteria within the healthcare system. Particularly, previous

travel to Asia stands out as a major risk factor for carriage and should be considered in

selecting empirical antibiotic treatment in critically ill patients.

Introduction

Extended spectrum β-lactamase (ESBL) producing *Escherichia coli* (ESBL-Ec) and *Klebsiella pneumoniae* (ESBL-Kp) are major contributors to the global burden of disease due to antibiotic resistant bacteria.[1] The prevalence of infections caused by ESBL-Ec or ESBL-Kp is increasing, even in countries with low antibiotic use and availability on prescription only, like Norway.[2] The mortality of invasive infections with ESBL-producing Enterobacterales (ESBL-E) ranges from 10-35%, depending on bacterial species, host factors, severity of disease and initial antibiotic therapy.[3,4] Consequently, ESBL-Ec/Kp are high-priority pathogens when developing new drugs to combat the threat from antibiotic resistant bacteria.[5]

ESBL-Ec/Kp colonization precedes invasive infections[6-8] and recent reviews report an increasing global prevalence of human ESBL-E carriage in the community, with an 8-fold rise over the past two decades.[9,10] The highest prevalence is detected in Asian and African regions (20-70%), the lowest in Europe and the Americas (<10%).[9,10] A study from the USA shows that the increased incidence of ESBL-Ec infections was driven by an increase in community-onset cases[11], and human-to-human transmission was the attributable source to 60% of community-acquired gastrointestinal ESBL-Ec carriage in the Netherlands [12].

Several risk factors for ESBL-E gastrointestinal carriage have been described. However, these have, with a few exceptions[13,14], mainly been investigated in small and/or selected study populations, such as international travelers[15], patients with gastroenteritis[16], patients recruited by general practitioners[17], persons with recent healthcare contact[18], pregnant women[19], children[20] or persons living in a livestock-dense area[21]. Many studies have identified international travel as a risk factor for ESBL acquisition, whilst the significance of sex, age, antibiotic or proton pump inhibitor use, hospitalization and diet are conflicting.[13-15,17,18,22-24]

We have comprehensive knowledge of the prevalence and population structure of clinical

ESBL-E isolates showing a dominance of CTX-M-group ESBL enzymes and the association

with specific extraintestinal pathogenic *E. coli* (ExPEC) and multidrug-resistant *K.*

pneumoniae high-risk clones.[25-28] Large-scale genomic studies have identified specific

subclades of *E. coli* sequence type (ST) 131 and *K. pneumoniae* ST307 as major contributors

to the increasing prevalence of ESBL infections.[25,27,29,30] Community-based studies from the

Netherlands and Sweden also observed a predominance of ST131, but a high genetic diversity

within the ESBL-Ec population.[13,14]

Improved knowledge of risk factors for ESBL-E carriage and their population structure in the

general human population may provide information for risk stratification and targeted

infection control measures. It is important to consider ESBL-E carriage in critically ill septic

patients with respect to the choice of empirical antibiotic treatment. The aims of this study

were to examine the prevalence of, and risk factors associated with, ESBL-Ec/Kp

gastrointestinal carriage in a general adult population in Norway, and to compare the ESBL-

Ec population structure with a national collection of clinical isolates.

Results

We detected gastrointestinal carriage of putative ESBL-Ec/Kp in 188 of 4,999 randomly

selected participants who provided fecal samples in the seventh survey of the population-

based Tromsø study (Tromsø7)[31] (Suppl. Table 1, Figure 1). Overall, 87% of participants

receiving a sampling kit returned a fecal sample. In total, 180 Ec and nine Kp putative ESBL-

positive isolates were isolated from the 188 ESBL screening-positive fecal samples. Both

ESBL-Ec and -Kp were detected in one sample.

Figure 1. Flowchart and results of fecal sample screening for gastrointestinal carriage of ESBL-producing *E. coli* and *K. pneumoniae* in 4,999 participants in Tromsø7, 2015-2016.

Pheno- and genotypic analyses showed that 14 putative ESBL-Ec isolates were either pheno- and/or genotypically ESBL-negative. Of these, one isolate harbored plasmid-mediated AmpC (bla_{CMY-2}). After exclusion of the ESBL-negative isolates, prevalence of ESBL-Ec gastrointestinal carriage was 3.3% (95% confidence interval (CI) 2.8-3.9%, 166 of 4,999 participants); 3.1% (2.5-3.9%) in women and 3.5% (2.8-4.4%) in men.

Among the nine screening ESBL-positive Kp isolates, one harbored bla_{CMY-2} and four did not express an ESBL phenotype. Consequently, four samples were considered positive for ESBL-producing Kp, corresponding to a prevalence of 0.08% (0.02-0.20%), all among male participants.

Factors associated with gastrointestinal carriage of ESBL-E. coli

We analyzed data from 4,996 participants, excluding those three who were positive for

ESBL-Kp only (Suppl. Table 1). Median age was 65 years (interquartile range 58-70 years,

no sex difference). In multivariable logistic regression analyses adjusted for all explanatory

variables, only travel to Asia the past 12 months was associated with ESBL-Ec

gastrointestinal carriage with an adjusted odds ratio (AOR) of 3.47 (2.18-5.51) (Table 1).

Among participants reporting hospitalization in the past year, or recent use of antibiotics or

acid suppressive medication, we observed a non-significant increase in prevalence of ESBL-

Ec.

Table 1. ESBL-producing *E. coli* gastrointestinal carriage and associated factors among 4,996
participants in Tromsø7.

Characteristics	% (ESBL-*E. coli*)	n (ESBL-*E. coli*)	N	AOR	95% CI	p-value
Age (years)						0.236
40-49	3.6	22	605	1.00		
50-59	4.1	33	814	1.15	0.66-2.01	
60-69	3.2	68	2,127	0.81	0.49-1.33	
70-84	3.0	43	1,450	0.73	0.42-1.26	
Hospitalization past 12 months						0.128
No	3.3	142	4,341	1.00		
Yes	4.0	24	593	1.43	0.90-2.27	
Antibiotic use past 14 days[a]						0.258
No	3.3	158	4,831	1.00		
Yes	5.2	8	153	1.58	0.72-3.47	
Acid suppressive medication past 4 weeks						0.951
Not used	3.3	123	3,760	1.00		
≤weekly	3.2	13	402	1.02	0.57-1.83	
Every week, but not daily	3.5	9	260	1.10	0.55-2.20	
Daily	3.7	11	297	1.20	0.63-2.26	
Travel abroad past 12 months[b]						<0.001
No	2.5	53	2,145	1.00		
Other regions (excl. Asia)	3.2	70	2,212	1.36	0.93-1.98	
Asia exclusively or Asia + other regions	8.2	38	461	3.47	2.18-5.51	
Traveler`s diarrhea past 12 months[c]						0.972
No	3.3	156	4,721	1.00		
Yes	4.8	8	166	1.01	0.48-2.16	

ESBL, extended spectrum β-lactamase; N, denominator; AOR, adjusted odds ratio; CI, confidence interval; AOR adjusted for
age, hospitalization past 12 months, antibiotic use past 14 days, acid suppressive medication past 4 weeks, travel abroad past
12 months and traveler`s diarrhea past 12 months.
The multivariable model includes 4,488 participants with complete information on all variables.
[a] Have you taken any antibiotics (tablets or oral suspensions, nasal ointments, eye drops or eye ointment) during the past 14
days?
[b] Traveled outside the Nordic countries >1 week duration in the past 12 months.
[c] For each travel abroad the past 12 months, the participants were asked if they had experienced diarrhea in connection with
the travel.
159
160

Both in the literature and in our previous study[32], we identified several factors associated with

Kp gastrointestinal carriage, overlapping with those associated with ESBL-Ec

carriage.[15,22,33,34] Thus, in 2,972 participants (of the total 4,996 participants in this current

study) previously screened for Kp, we performed a multivariable logistic regression analysis

including Kp carriage as an additional explanatory variable (Suppl. Table 2). Prevalence of

ESBL-Ec gastrointestinal carriage was 2.7% among participants without Kp carriage and

4.8% (p=0.016) among those with Kp carriage. In the full model, AOR was 1.66 (0.99-2.79,

p=0.055), indicating a possible association between Kp and ESBL-Ec carriage.

Comparative analysis of ESBL-E. coli carriage and clinical isolates

To explore the population structure and genomic characteristics of ESBL-Ec in community

carriage we whole-genome sequenced (WGS) all 166 isolates. Furthermore, we sequenced a

contemporary national collection of 118 clinical ESBL-Ec isolates (NORM 2014) for

comparative analysis (Figure 2, Suppl. Table 3).

Tree scale: 0.1 ⊢━━━┥

Figure 2. Maximum-likelihood phylogenetic tree based on core genome alignment of the genomes of ESBL-*E. coli* carriage isolates from Tromsø7 (labeled grey, n=166) and clinical isolates from NORM 2014 (blood isolates labeled with red and urine isolates with yellow, n=118). The innermost ring illustrates phylogroups, followed by a ring with sequence types (STs), a ring with ST131 subclades, and a ring with *fimH* types (ND, not detected). The heatmap shows presence (blue color) or absence (white) of ESBL gene variants.

At the phylogroup level, 52.4% of the ESBL-Ec carriage isolates belonged to either

phylogroup A (26.5%) or D (25.9%), and 33.7% belonged to phylogroup B2 (Figure 2 and

Figure 3, Suppl. Table 3). This contrasts with the clinical isolates where 64.4% (p<0.001) of

the isolates belonged to phylogroup B2 and only 20.4% (p<0.001) belonged to phylogroup A

(5.1%) and D (15.3%) combined.

Figure 3. Phylogroup (labeled on the top) and sequence type (ST) distribution of ESBL-*E.*

coli carriage isolates from Tromsø7 (n=166) and clinical isolates from NORM 2014 (n=118).

Carriage isolates had higher ST diversity (Figure 2 and 3, Suppl. Table 3) with Simpson's

diversity index 92.4% compared to clinical isolates (65.9%, p<0.001). We identified 58

different STs among the carriage isolates while the clinical isolates included 27 STs (Figure 2

and 3, Suppl. Table 3). ST131 (phylogroup B2) was the dominant ST in both collections,

however, more prevalent among clinical isolates (57.6%, n=68) than carriage isolates (24.1%,

n=40, p<0.001). Comparison of prevalence of ESBL-producing ST131 colonization vs.

infection in our study results in a crude odds ratio (OR) for infection of 4.3 (2.57-7.13,

p<0.001).

Within ST131, the multidrug resistant subclades C1 (alternatively referred to as *H*30-R) and

C2 (*H*30-Rx) accounted for 67.5% in carriage and 73.5% clinical isolates (Figure 2, Suppl.

Figure 1, Suppl. Table 3). The subclade C1 (47.5%, n=19) was the most prevalent among

carriage isolates and C2 (42.6%, n=29) among clinical isolates. The difference in the

proportion of C2 between clinical and carriage isolates was statistically significant (20.0%,

n=8, p=0.017). For the overall ESBL-Ec population, crude OR for infection among C2 was

6.44 (2.82-14.68, p<0.001) (Suppl. Table 4).

Although non-significant, a higher proportion of subclades A, B and B0, less associated with

antibiotic resistance, was observed among ST131 carriage isolates (32.5%) than clinical

isolates (26.5%, p=0.508) (Suppl. Figure 1). However, with regard to the overall ESBL-Ec

population, subclade A was associated with infection (carriage 6.0% vs. clinical 13.6%, OR

2.45, 1.07-5.60, p=0.034) (Suppl. Table 4).

The most prevalent STs in carriage isolates following ST131 were ST10 (7.2%, n=12,

phylogroup A), ST69 (7.2%, n=12) and ST38 (6.6%, n=11, both phylogroup D) compared to

ST405 (7.6%, n=9), ST38 (5.9%, n=7, both phylogroup D), and ST648 (5.1%, n=6,

phylogroup F) in clinical isolates (Figure 2 and 3, Suppl. Table 3). We did not identify the

common ExPEC lineage ST73[25,29] among the carriage isolates but found one ST95[25,29] (0.6%)

and five isolates of the emerging ST1193[35,36] (3.0%).

Using a single nucleotide polymorphism (SNP) cut-off of ≤17[37], we detected two putative

clusters (Suppl. Table 5) among five of 284 isolates. All five were carriage isolates. One

ST357 cluster (6-9 SNP differences) consisted of three isolates and the other cluster of two

ST131 isolates (4 SNP difference). We detected no clusters among the clinical isolates.

Antimicrobial resistance and plasmid replicon content

CTX-M enzymes accounted for at least 97.0% of the ESBL phenotypes in both collections

(Figure 2 and Figure 4, Suppl. Table 3). The most prevalent ESBL genes among both carriage

and clinical isolates were $bla_{CTX-M-15}$ (40.4%, n=67 vs. 61.0%, n=72, p=0.001), $bla_{CTX-M-14}$

(20.5%, n=34 vs. 14.4%, n=17, p=0.188), and $bla_{CTX-M-27}$ (19.9%, n=33 vs. 15.3%, n=18,

p=0.321). One clinical isolate of ST38 harbored both $bla_{CTX-M-15}$ and $bla_{CTX-M-14}$. $Bla_{CTX-M-3}$,

$bla_{CTX-M-8}$, $bla_{CTX-M-32}$ and $bla_{CTX-M-101}$ were exclusively detected in carriage isolates, whereas

$bla_{CTX-M-24}$ and $bla_{CTX-M-104}$ were present only in clinical isolates. The remaining ESBL

producers contained bla_{SHV-12} in 3.0% of the carriage and 1.7% of the clinical isolates. Bla_{TEM}

ESBL genes were not detected.

230
Figure 4. ESBL gene prevalence in ESBL-*E. coli* carriage isolates from the Tromsø7 study

(n=166) and clinical isolates from NORM 2014 (n=118).

The clinical isolates showed an overall higher proportion of phenotypic resistance compared

to the carriage isolates (Table 2, Suppl. Table 3). Only 2.4% of carriage isolates were

phenotypically resistant to piperacillin-tazobactam compared to 31.8% (p<0.001) of clinical

isolates. We also detected a high prevalence of phenotypic co-resistance to non-β-lactam

antibiotics such as gentamicin, ciprofloxacin and trimethoprim-sulfamethoxazole both in

clinical and carriage isolates. Isolates co-resistant to all these three antibiotic classes,

accounted for 10.2% (17/166) of carriage and 33.1% (39/118, p<0.001) of clinical isolates.

Table 2. Susceptibility profile of ESBL-*E. coli* carriage isolates from Tromsø7 (n=166) and

clinical isolates from NORM 2014 (n=118).

	Carriage			Clinical		
	%S	%I	%R	%S	%I	%R
Amoxicillin-clavulanic acid i.v.	57.2	-	42.8	25.4	-	74.6
Amoxicillin-clavulanic acid*	90.4	-	9.6	54.2	-	45.8
Piperacillin-tazobactam	97.6	-	2.4	68.2†	-	31.8†
Cefuroxime*	0.6	-	99.4	1.7	-	98.3
Ceftazidime	20.5	32.5	47.0	10.2	11.0	78.8
Cefotaxime	0.0	2.4	97.6	0.8	0.8	98.4
Cefepime	7.8	54.8	37.3	4.7†	5.9†	89.4†
Ceftazidime-avibactam	100.0	-	0.0	NA	NA	NA
Ertapenem	100.0	-	0.0	NA	NA	NA
Meropenem	100.0	0.0	0.0	99.2	0.8	0.0
Aztreonam	5.4	23.5	71.1	NA	NA	NA
Amikacin	100.0	-	0.0	NA	NA	NA
Gentamicin	80.1	-	19.9	48.3	-	51.7
Tobramycin	76.5	-	23.5	NA	NA	NA
Ciprofloxacin	49.4	13.9	36.7	14.4	8.5	77.1
Trimethoprim-sulfamethoxazole	48.8	0.6	50.6	27.1	0.0	72.9
Nitrofurantoin	99.4	-	0.6	93.9‡	-	6.1‡
Fosfomycin	99.4	-	0.6	NA	NA	NA
Fosfomycin*	96.4	-	3.6	NA	NA	NA
Colistin	98.8	-	1.2	NA	NA	NA
Tigecycline	100.0	-	0.0	100.0†	-	0.0†

S, susceptible; I, susceptible, increased exposure; R, Resistant; i.v. intravenous; NA, not available; *Breakpoints
for uncomplicated urinary tract infections; †available for blood culture isolates only (n=85); ‡available for urine
culture isolates only (n=33)

Phenotypic ciprofloxacin resistance was different for ST131 (70.0%) vs. non-ST131 (26.2%,

p<0.001) among carriage isolates (Suppl. Table 6) in contrast to the clinical isolates (Suppl.

Table 7). Two carriage isolates of ST484 and ST1324 (both phylogroup A) were

phenotypically resistant to colistin and harbored *mcr*-1.1 and *mcr*-3.5, respectively. No

isolates expressed clinical resistance against carbapenems. However, two clinical isolates of

ST95 (phylogroup B2) and ST410 (phylogroup C) harbored bla_{IMP-26} and $bla_{OXA-181}$,

respectively. All isolates were susceptible to tigecycline.

Despite the overall higher proportion of phenotypic resistance among clinical isolates, the

average number of different plasmid replicon-types per isolate (both 3.1/isolate) did not differ

between the carriage and clinical collections. We identified 43 and 39 different plasmid

replicon types among 97.1% (162/166) carriage and 95.0% (112/118) clinical isolates (Suppl.

Table 3, Suppl. Figure 2). The most prevalent were IncFIB(AP001918) (carriage 22.7% vs.

clinical 22.3%) followed by Col156 (carriage 11.1% vs. clinical 11.5%) and IncFIA (carriage

10.9% vs. clinical 17.7%, p=0.101). Replicon type Col156 and IncFIA were mainly detected

in globally disseminated ExPEC clones (ST131, ST1193, ST648, ST69, ST405), whereas

IncFIB(AP001918) was additionally frequently spread among other STs (Suppl. Table 3).

ESBL-K. pneumoniae carriage isolates

We detected only four ESBL-Kp, each of a different ST (ST29, ST211, ST261 and ST2459)

harboring $bla_{CTX-M-15}$ (n=2), $bla_{CTX-M-14}$ (n=1), or bla_{SHV-12} (n=1). None of the isolates were

genomically assigned as hypervirulent.

Discussion

Our study contributes to the knowledge of prevalence of, and factors associated with, ESBL-

Ec/Kp gastrointestinal carriage in a general adult population, and the bacterial population

structure of carriage isolates. The comparison to a contemporary collection of clinical ESBL-

Ec isolates revealed differences in the population structure and the prevalence of phenotypic

resistance between carriage and clinical isolates. Travel to Asia was identified as a major risk

for ESBL-Ec gastrointestinal carriage.

An ESBL-Ec carriage prevalence of 3.3% (2.8-3.9) is lower but comparable to previous

community-based data from Europe including Sweden 4.4% (3.5-5.3, n=2,134, data collected

2012-2013)[13], the Netherlands 4.5% (3.9-5.1, n=4,177, 2014-2016)[14] and a Norwegian study

4.9% (2.7-8.1, n=284, 2014-2016)[17] using similar screening approaches.

We identified significant differences in the ESBL-Ec population structure between the

community and clinical isolates. The globally disseminated phylogroup B2 clone ST131 has

been identified as a key contributor to the increase in ESBL prevalence[38] and in a longitudinal

study of *E. coli* bloodstream isolates we identified ST131 to be the single largest contributor

to the increase in prevalence of ESBL-Ec in Norway.[25] We also observed the predominance

of ST131 in both our collections. However, the proportion is significantly lower in carriage

isolates due to lower numbers of the multidrug resistant subclade C2. Moreover, the carriage

population had a higher proportion of phylogroup A, associated with asymptomatic intestinal

carriage in humans[39,40] and a significantly greater ST diversity overall, compared to the

clinical isolates. These observations indicate that acquisition of ESBL genes frequently occur

in a variety of *E. coli* lineages colonizing the gut. However, there are differences in the

colonization potential of *E. coli* lineages and the risk of invasive infection by ESBL-Ec which

seems to be clone dependent.[38,41] The higher odds for infection that we detected for ST131 is

similar to that of the Swedish study (AOR 3.4, 1.8-6.4) indicating a higher pathogenicity

potential of ST131 compared to commensal *E. coli* lineages of phylogroup A, such as ST10.[13]

Moreover, we found that ST131 subclade A, previously reported with less resistance, and the

multidrug resistant subclade C2 had higher odds for infection, and this may contribute to the

sustained establishment of these subclades among bloodstream infections in Norway.[25]

Assuming a 100% colonization rate of *E. coli*, the large proportion of STs notorious as

common causes of extra-intestinal clinical infections (e.g. ST131, ST405, ST38 and

ST648)[25,27,28] could at least partly explain the higher prevalence of ESBL among *E. coli*

causing bloodstream infections in Norway (5.8% in 2016)[42] compared to the carriage

prevalence of 3.3% identified here. We also observed emerging clones such as ST1193 which

appears to have disseminated rapidly worldwide over the last decade.[35,36] The low prevalence

of ESBL-Kp gastrointestinal carriage (0.08%) is consistent with previous community based

reports.[9,14,17]

The identification of clusters with closely related isolates could indicate putative clonal

spread. This could include within-household and social network transmission or nosocomial

spread. However, we did not have access to epidemiological data to examine this further.

In line with other studies, we found a strong association between ESBL-Ec carriage and travel

to Asian regions.[13-15,17,22] This supports the current patient screening recommendations for

ESBL-producing gram-negative bacteria after hospital stay abroad in the past year before

hospital admission in Norway.[43] In contrast to a Swedish and a Dutch study,[15,44] we did not

identify travelers' diarrhea as a risk factor for ESBL-Ec carriage. However, our study was not

designed to specifically investigate international travelers but rather focused on risk factors in

the general adult population.

We found no association between ESBL-Ec gut carriage and factors such as hospitalization,

antibiotic use, and acid suppressive medication, and conflicting results have been detected in

previous studies.[22,45] Hospitalization as a risk factor has mainly been reported from studies

investigating patients with ESBL-E infections.[23,46] In line with most studies that assessed risk

factors regarding ESBL-E carriage in individuals in the community, we did not identify

hospitalization as an independent risk factor.[13,22,44] This may be due to the increased ESBL

prevalence not only in hospitals, but also in the community over the last decades and implies

that boundaries have become blurred between those two settings.[9,22,47,48]

The nonsignificant effect of antibiotic use is mainly due to limitations of the drug variable in

our study which is based on self-reported data, including topical antibiotics and covers only

the last two weeks before self-sampling. As antibiotic use has been found as a risk factor for

resistance in many other studies,[14,15,22] we cannot rule out that antibiotic use plays a role in

ESBL-E carriage. There are reports identifying an association between the use of gastric acid

suppressive medication and intestinal colonization or infections with ESBL-E.[18,21,23,33,34,46]

However, a Dutch study comparable to ours did not find an association between proton pump

inhibitor use and ESBL-E carriage in the overall analysis.[14]

Interestingly, we found a possible association between Kp and ESBL-Ec carriage. An

association between ESBL-E carriage and vancomycin-resistant enterococci (VRE) has been

described previously[7,49], and also that VRE colonization is significantly associated with Kp

colonization among intensive care unit patients.[50] These associations warrant further

investigations to assess if a common set of risk factors for carriage of different clinically

important pathogens can be identified.

An important strength of our study is the non-selective recruitment from the official

population-registry, and the high participation (87%) compared to 18.3% and 18.8% in

comprehensive studies from the Netherlands[14] and Sweden[13], respectively. It is a limitation

that we only captured the general population 40 years and older. However, other studies have

not found an association between age and ESBL-E carriage.[13,14] Moreover, more extensive

data on drug use would have strengthened the analyses. Additionally, the genomic diversity of

carriage isolates is likely to be underestimated due to the isolation and sequencing of only one

colony per fecal sample. Although, we compared local carriage isolates with a national

collection of clinical ESBL-Ec, there was no discernible spatiotemporal spread or

phylogenetic structure within Norway according to a nationwide genomic study on *E. coli*

causing bloodstream infections.[25]

Conclusions

The prevalence of ESBL-Ec carriage in a general adult urban Norwegian population was low

reflecting the relatively low prevalence of ESBL-Ec in clinical isolates. Travel to Asia was

the only independent risk factor for ESBL-Ec carriage and should be considered in terms of

screening recommendations before hospital admission. The differences in ESBL-Ec

populations between carriage and clinical isolates indicating a higher risk of infection

dependent on the ESBL-Ec clone warrants further investigations to elucidate if risk

stratification should include determination of the bacterial genetic background.

Subjects and Methods

Study population and design

Our study sample was drawn from Tromsø7, the last of seven cross-sectional health surveys

conducted between 1974 and 2016 in Tromsø municipality, Norway

(https://uit.no/research/tromsostudy). Tromsø is representative of a Northern European, urban

population.[31] Tromsø7 (March 2015- October 2016) included questionnaires and two clinical

visits (https://uit.no/research/tromsostudy/project?pid=708909). Unique national identity

numbers from the official population-registry were used to invite all citizens $\geq$40 years

(n=32,591). Sixty-five percent (n=21,083, 11,074 women) attended the first clinical visit in

the study (Figure 5).

Figure 5. Flow diagram of the study population, Tromsø7, 2015-2016. *Study population

screened for *K. pneumoniae* carriage is described in.[32]

A selection of 9,320 persons attending the first visit, were invited for a second visit, also

including 3,154 former Tromsø Study participants not already included in the random

selection process, which was required for other clinical research purposes. From March 2015

to March 2016, 5,800 participants at the first visit were consecutively provided a fecal self-

sampling kit. Participants collected fecal material using nylon-flocked ESwab 490CE.A

(Copan, Brescia, Italy). In total, 87% (n=5,042) returned a sample either at the second visit, or

by mail to the laboratory.

All 5,042 fecal samples were screened for the presence of ESBL-producing Ec and Kp via

selective culture (see below). All participants completed two self-administered structured

questionnaires on socio-demographics, smoking, alcohol use, hospitalization, drug use, and

travel abroad. We excluded 13 participants with wrong or missing sample identification

number, two retracting consent to medical research, and 28 with incomplete questionnaires

for a final study population of 4,999 participants (Figure 5). We analyzed the association

between ESBL-Ec gastrointestinal carriage and different risk factors in 4,996 participants

(Figure 5 and Table 1). Next, we investigated the association between ESBL-Ec carriage and

Kp gastrointestinal carriage among 2,972 participants additionally screened for Kp in our

previous study,[32] irrespective of resistance (Figure 5, Supplementary Table 2).

Isolation of ESBL-producing E. coli and K. pneumoniae

We added 200 µl 85% glycerol to the ESwab tubes upon arrival at the local microbiological

laboratory, and stored the samples at -80°C. From the thawed media, 100 µl were plated onto

CHROMagar[TM] ESBL (CHROMagar, Paris, France) and incubated for 48 hours at 37°C.

Pink, purple and blue colonies suspected of being ESBL-producing Ec or *Klebsiella* spp. were

identified using mass spectrometry (MALDI-TOF, Bruker Daltonics, Bremen, Germany). The

first colony identified as either Ec, *K. pneumoniae* or *K. variicola* from each sample was kept

and further analyzed. All samples were plated on cysteine lactose electrolyte deficient agar

(MAST Group, Bootle, UK) to assess growth of fecal flora and validity of the samples.

K. pneumoniae isolation

The screening strategy and isolation procedure for Kp, is described in detail elsewhere[32].

Briefly, we plated and screened the fecal samples onto the selective SCAI (Simons citrate

agar with inositol; both Sigma-Aldrich, Darmstadt, Germany) medium and identified

suspected colonies using MALDI-TOF.

Antimicrobial susceptibility testing and phenotypic ESBL-identification

Susceptibility testing was performed according to the EUCAST broth microdilution method

for carriage isolates, disc diffusion method[51] for the clinical isolates and both interpreted

using the EUCAST 2023 breakpoint table (https://eucast.org/). For the confirmation of ESBL-

producing Ec and Kp, we followed the EUCAST algorithm for phenotypic detection of

ESBLs using the BD BBL[TM] combination disk test (Becton Dickinson and Company, Sparks,

USA).

Genomic sequencing and bioinformatic analysis

Genomic DNA from 166 Ec of the Tromsø7 collection was extracted with the MagNA Pure

96 system (Roche Applied Science, Mannheim, Germany) and sequencing libraries were

prepared according to the Nextera Flex sample preparation protocol (Illumina, San Diego,

CA, USA). Samples were sequenced on the Illumina MiSeq platform to generate 300 bp

paired-end reads. All reads were trimmed with TrimGalore v0.6.4 and assembled with

Unicycler v0.4.8 including SPAdes v3.13.0.[52-54] STs were assigned using the multi locus

sequence type (MLST) software *E.coli* scheme v2.19.0 and Enterobase.[55-57] AMRFinderPlus

v3.10.16 was used to determine resistance genes among the Ec isolates.[58] Plasmid replicons

were identified using Abricate v1.0.1 including the PlasmidFinder 2021-Mar-27 database.[59,60]

Phylogroup assignment was based on ClermonTyping v20.03 (Mar. 2020).[61] The *fimH* type

was identified using FimTyper v1.0 and BLAST+ v2.12.0.[62,63] Regarding the *Klebsiella*

isolates, Kleborate v2.0.0 was used to determine species identification, ST and acquired genes

encoding virulence or antibiotic resistance.[64,65]

The clinical ESBL-producing Ec included 118 isolates out of 123 representing all ESBL-Ec

isolates collected in 2014[66], as part of the yearly surveillance program in NORM. Genome

sequencing of five isolates were unsuccessful. Before sequencing, the NORM 2014 isolates

were stored at -80°C and then sent to GATC Biotech AG (part of Eurofins

Genomics/Eurofins Scientific) in Germany for DNA isolation and WGS. Raw reads were

trimmed using Trimmomatic v0.39 and assembled with SPAdes v3.15.0.[67,68] Contigs shorter

than 200 bp were discarded. The genomic data were analyzed as described above.

Phylogenetic and population structure analysis of ESBL-E. coli

We used Prokka v.1.14.6[69] to annotate genomes and snippy v.4.6.0[70] to map the sequence

reads to the ST131 Ec EC958 chromosome [HG941718.1]

(https://www.ncbi.nlm.nih.gov/nuccore/HG941718.1) to create the core genome alignment.

We used snp-dists v.0.8.2, (https://github.com/tseemann/snp-dists/) to create the SNP distance

matrix from the core genome alignment. We used a 17 SNP cut-off to define similarity

between two genomes and to identify probable ESBL-Ec transmission events among cases.[37]

To assess the phylogenetic relatedness, we used the core SNP alignment to infer a maximum-

likelihood tree using RAxML v.8.2.8 with the GTR + Gamma rate model and 100 rapid

bootstraps visualized in iTol (v6.5.2).[71,72] ST131 subclades were determined based on

subclade specific SNPs and *fimH* alleles, subclade membership was corrected when

assignment based on SNP profile of sporadic isolates did not fit with the phylogenetic

distribution of clades.[25,30,62,73]

Statistical analysis

Our primary analyses were multivariable logistic regression models, performed in two

overlapping study populations with outcome variable ESBL-Ec gastrointestinal carriage using

SPSS v.26.0 (SPSS, Inc., Chicago, IL, USA). First, we analyzed factors associated with

ESBL-Ec gastrointestinal carriage among 4,996 participants (Table 1). Next, in 2,972

participants also screened for Kp,[32] we performed a multivariable logistic regression analysis

including Kp carriage as an additional explanatory variable (Suppl. Table 2). Explanatory

variables were selected with the help of a directed acyclic graph constructed using DAGitty

v3.0 (Suppl. Figure 3 and 4).[74] All explanatory variables were kept in the fully adjusted

models. Multicollinearity between the entered variables was assessed calculating the variance

inflation factor (VIF) and tolerance statistic. Multicollinearity was not a problem with VIF>10

and tolerance statistic <0.2.[75] The strength of the associations was examined by calculating

AORs with 95% CI. Two-sided p-values <0.05 were considered statistically significant. The

prevalence of ST131 among carrier and clinical isolates was compared calculating the OR

with 95% CI using logistic regression in SPSS. The comparison of proportions was assessed

using χ^2-test.

Ethics

The study was approved by the Regional Committee for Medical and Health Research Ethics,

North Norway (REC North reference: 2016/1788 and 2014/940) and the Data Protection

Officer at University Hospital of North Norway (reference: 2019/4264). The study complied

with the Declaration of Helsinki. All participants in Tromsø7 signed an informed consent

form prior to participation.

Acknowledgements

We are grateful for technical assistance from Bjørg Haldorsen, Bettina Aasnæs and Ellen

Josefsen in organizing the collection of fecal samples in the laboratory, Eva Bernhoff and

Ragna-Johanne Bakksjø for performing WGS, Marit Andrea Klokkhammer Hetland for initial

sequence analysis of carriage isolates. Rod Wolstenholme for figure editing. We thank

Sigma2, the national provider of e-infrastructure, for access to High-Performance Computing

and large-scale data storage (Project ID: NN9794K).

Author contributions:

ØS conceptualized and acquired funding for the study in collaboration with AS, KG and IHL.

ØS was responsible for organizing the collection of fecal samples. LLEA did the screening of

fecal samples and phenotypic testing. NR, KS, LS and KG conceptualized and conducted the

epidemiological analysis of risk factors. IHL organized WGS of carriage isolates. GSS and

the Norwegian *E. coli* ESBL Study Group provided data and clinical isolates. DJB, AKP, NR

and ØS analyzed the WGS data. NR, ØS, KS and KG prepared first manuscript draft. All

authors contributed to review and editing of the manuscript and approved the final version.

Declaration of interests:

All authors report no conflicts of interest.

Role of funding source

This study was supported by grants from the Northern Norway Regional Health Authority

(HNF1415-18) and the Trond Mohn Foundation (TMF2019TMT03). The funders of the study

played no role in study design, data collection, data analysis, data interpretation, or writing of

the report.

Data Availability

Bacterial genome data (raw Illumina reads) are publicly available in NCBI under BioProject

PRJEB53319 (NORM 2014 collection) and PRJEB57251 (Tromsø7 collection). This study is

based on data owned by a third party (The Tromsø Study, Department of Community

Medicine, UiT The Arctic University of Norway). Confidentiality requirements according to

Norwegian law prevents sharing of individual patient level data in public repositories.

Application of legal basis and exemption from professional secrecy requirements for the use

of personal health data in research may be sent to a regional committee for medical and health

research ethics (https://rekportalen.no/). The authors gained access to the data through the

Tromsø Study's application process. Guidelines on how to access the data are available at the

website https://uit.no/research/tromsostudy. All enquiries about the Tromsø Study should be

sent by e-mail to tromsous@ism.uit.no. All the questionnaire variables are published in the

NESSTAR program system and results can be viewed online:

http://tromsoundersokelsen.uit.no/tromso/.

References

1. Antimicrobial Resistance C. Global burden of bacterial antimicrobial resistance in 2019: a
systematic analysis. *Lancet.* 2022 Feb 12;399(10325):629-655.
2. NORM/NORM-VET 2021. Usage of antimicrobial agents and occurrence of antimicrobial
resistance in Norway. Tromsø/Oslo/Ås; 2022. ISSN: 1502-2307 (print)/1890- 9965
(electronic).
3. Schwaber MJ, Carmeli Y. Mortality and delay in effective therapy associated with
extended-spectrum beta-lactamase production in Enterobacteriaceae bacteraemia: a
systematic review and meta-analysis. *J Antimicrob Chemother.* 2007 Nov;60(5):913-20.

4. Scheuerman O, Schechner V, Carmeli Y, et al. Comparison of predictors and mortality
between bloodstream infections caused by ESBL-producing *Escherichia coli* and ESBL-
producing *Klebsiella pneumoniae. Infect Control Hosp Epidemiol.* 2018 Jun;39(6):660-667.
5. Tacconelli E, Carrara E, Savoldi A, et al. Discovery, research, and development of new
antibiotics: the WHO priority list of antibiotic-resistant bacteria and tuberculosis. *Lancet*
Infect Dis. 2018 Mar;18(3):318-327.
6. Prevel R, Boyer A, M'Zali F, et al. Extended spectrum beta-lactamase producing
Enterobacterales faecal carriage in a medical intensive care unit: low rates of cross-
transmission and infection. *Antimicrob Resist Infect Control.* 2019 Jul 10;8:112.
7. Wyres KL, Hawkey J, Mirceta M, et al. Genomic surveillance of antimicrobial resistant
bacterial colonisation and infection in intensive care patients. *BMC Infect Dis.* 2021 Jul
14;21(1):683.
8. Denkel LA, Maechler F, Schwab F, et al. Infections caused by extended-spectrum beta-
lactamase-producing Enterobacterales after rectal colonization with ESBL-producing
Escherichia coli or *Klebsiella pneumoniae. Clin Microbiol Infect.* 2020 Aug;26(8):1046-
1051.
9. Woerther PL, Burdet C, Chachaty E, Andremont A. Trends in human fecal carriage of
extended-spectrum beta-lactamases in the community: toward the globalization of CTX-M.
Clin Microbiol Rev. 2013 Oct;26(4):744-58.
10. Bezabih YM, Sabiiti W, Alamneh E, et al. The global prevalence and trend of human
intestinal carriage of ESBL-producing *Escherichia coli* in the community. *J Antimicrob*
Chemother. 2021 Jan 1;76(1):22-29.
11. Jernigan JA, Hatfield KM, Wolford H, et al. Multidrug-Resistant Bacterial Infections in
U.S. Hospitalized Patients, 2012-2017. *N Engl J Med.* 2020 Apr 2;382(14):1309-1319.
12. Mughini-Gras L, Dorado-Garcia A, van Duijkeren E, et al. Attributable sources of
community-acquired carriage of *Escherichia coli* containing beta-lactam antibiotic resistance
genes: a population-based modelling study. *Lancet Planet Health.* 2019 Aug;3(8):e357-e369.
13. Ny S, Lofmark S, Borjesson S, et al. Community carriage of ESBL-producing
Escherichia coli is associated with strains of low pathogenicity: a Swedish nationwide study.
J Antimicrob Chemother. 2017 Feb;72(2):582-588.
14. van den Bunt G, van Pelt W, Hidalgo L, et al. Prevalence, risk factors and genetic
characterisation of extended-spectrum beta-lactamase and carbapenemase-producing
Enterobacteriaceae (ESBL-E and CPE): a community-based cross-sectional study, the
Netherlands, 2014 to 2016. *Euro Surveill.* 2019 Oct;24(41):1800594.
15. Arcilla MS, van Hattem JM, Haverkate MR, et al. Import and spread of extended-
spectrum beta-lactamase-producing Enterobacteriaceae by international travellers (COMBAT
study): a prospective, multicentre cohort study. *Lancet Infect Dis.* 2017 Jan;17(1):78-85.
16. Jorgensen SB, Samuelsen O, Sundsfjord A, et al. High prevalence of faecal carriage of
ESBL-producing Enterobacteriaceae in Norwegian patients with gastroenteritis. *Scand J*
Infect Dis. 2014 Jun;46(6):462-5.
17. Ulstad CR, Solheim M, Berg S, Lindbaek M, Dahle UR, Wester AL. Carriage of
ESBL/AmpC-producing or ciprofloxacin non-susceptible *Escherichia coli* and *Klebsiella* spp.
in healthy people in Norway. *Antimicrob Resist Infect Control.* 2016 Dec 15;5:57.
18. Kohler P, Seiffert SN, Kessler S, et al. Molecular Epidemiology and Risk Factors for
Extended-Spectrum beta-Lactamase-Producing Enterobacterales in Long-Term Care
Residents. *J Am Med Dir Assoc.* 2022 Mar;23(3):475-481.e5.
19. Rettedal S, Lohr IH, Bernhoff E, Natas OB, Sundsfjord A, Oymar K. Extended-spectrum
beta-lactamase-producing Enterobacteriaceae among pregnant women in Norway: prevalence
and maternal-neonatal transmission. *J Perinatol.* 2015 Nov;35(11):907-12.

20. Birgy A, Cohen R, Levy C, et al. Community faecal carriage of extended-spectrum beta-
lactamase-producing Enterobacteriaceae in French children. *BMC Infect Dis.* 2012 Nov
21;12:315.
21. Wielders CCH, van Hoek A, Hengeveld PD, et al. Extended-spectrum β-lactamase- and
pAmpC-producing Enterobacteriaceae among the general population in a livestock-dense
area. *Clin Microbiol Infect.* 2017 Feb;23(2):120.e1-120.e8.
22. Karanika S, Karantanos T, Arvanitis M, Grigoras C, Mylonakis E. Fecal colonization with
extended-spectrum beta-lactamase-producing Enterobacteriaceae and risk factors among
healthy individuals: A systematic review and metaanalysis. *Clin Infect Dis.* 2016 Aug
1;63(3):310-8.
23. Richelsen R, Smit J, Laxsen Anru P, Schønheyder HC, Nielsen H. Risk factors of
community-onset extended-spectrum β-lactamase *Escherichia coli* and *Klebsiella*
pneumoniae bacteraemia: an 11-year population-based case-control-control study in
Denmark. *Clin Microbiol Infect.* 2021 Jun;27(6):871-877.
24. Leistner R, Meyer E, Gastmeier P, et al. Risk factors associated with the community-
acquired colonization of extended-spectrum beta-lactamase (ESBL) positive *Escherichia coli.*
an exploratory case-control study. *PLoS One.* 2013 Sep 11;8(9):e74323.
25. Gladstone RA, McNally A, Pontinen AK, et al. Emergence and dissemination of
antimicrobial resistance in *Escherichia coli* causing bloodstream infections in Norway in
2002-17: a nationwide, longitudinal, microbial population genomic study. *Lancet Microbe.*
2021 Jul;2(7):e331-e341.
26. Fostervold A, Hetland MAK, Bakksjø R, et al. A nationwide genomic study of clinical
Klebsiella pneumoniae in Norway 2001-2015: Introduction and spread of ESBL facilitated by
CG15 and CG307. *J Antimicrob Chemother.* 2022 Feb 23;77(3):665-674.
27. Dunn SJ, Connor C, McNally A. The evolution and transmission of multi-drug resistant
Escherichia coli and *Klebsiella pneumoniae*: the complexity of clones and plasmids. *Curr*
Opin Microbiol. 2019 Oct;51:51-56.
28. Manges AR, Geum HM, Guo A, Edens TJ, Fibke CD, Pitout JDD. Global extraintestinal
pathogenic *Escherichia coli* (ExPEC) lineages. *Clin Microbiol Rev.* 2019 Jun
12;32(3):e00135-18.
29. Kallonen T, Brodrick HJ, Harris SR, et al. Systematic longitudinal survey of invasive
Escherichia coli in England demonstrates a stable population structure only transiently
disturbed by the emergence of ST131. *Genome Res.* 2017 Jul 18;27(8):1437-1449.
30. Ben Zakour NL, Alsheikh-Hussain AS, Ashcroft MM, et al. Sequential acquisition of
virulence and fluoroquinolone resistance has shaped the evolution of *Escherichia coli* ST131.
mBio. 2016 Apr 26;7(2):e00347-16.
31. Hopstock LA, Grimsgaard S, Johansen H, Kanstad K, Wilsgaard T, Eggen AE. The
seventh survey of the Tromsø Study (Tromsø7) 2015-2016: study design, data collection,
attendance, and prevalence of risk factors and disease in a multipurpose population-based
health survey. *Scand J Public Health.* 2022 Nov;50(7):919-929.
32. Raffelsberger N, Hetland MAK, Svendsen K, et al. Gastrointestinal carriage of *Klebsiella*
pneumoniae in a general adult population: a cross-sectional study of risk factors and bacterial
genomic diversity. *Gut Microbes.* 2021 Jan-Dec;13(1):1939599.
33. Willems RPJ, van Dijk K, Ket JCF, Vandenbroucke-Grauls C. Evaluation of the
association between gastric acid suppression and risk of intestinal colonization with
multidrug-resistant microorganisms: A systematic review and meta-analysis. *JAMA Intern*
Med. 2020 Apr 1;180(4):561-571.
34. Reuland EA, Al Naiemi N, Kaiser AM, et al. Prevalence and risk factors for carriage of
ESBL-producing Enterobacteriaceae in Amsterdam. *J Antimicrob Chemother.* 2016
Apr;71(4):1076-82.

35. Johnson JR, Johnston BD, Porter SB, et al. Rapid emergence, subsidence, and molecular
detection of *Escherichia coli* sequence type 1193-fimH64, a new disseminated multidrug-
resistant commensal and extraintestinal pathogen. *J Clin Microbiol.* 2019 Apr
26;57(5):e01664-18.
36. Tchesnokova V, Radey M, Chattopadhyay S, et al. Pandemic fluoroquinolone resistant
Escherichia coli clone ST1193 emerged via simultaneous homologous recombinations in 11
gene loci. *Proc Natl Acad Sci U S A.* 2019 Jul 16;116(29):14740-14748.
37. Ludden C, Coll F, Gouliouris T, et al. Defining nosocomial transmission of *Escherichia*
coli and antimicrobial resistance genes: a genomic surveillance study. *Lancet Microbe.* 2021
Sep;2(9):e472-e480.
38. Nicolas-Chanoine MH, Bertrand X, Madec JY. *Escherichia coli* ST131, an intriguing
clonal group. *Clin Microbiol Rev.* 2014 Jul;27(3):543-74.
39. Tenaillon O, Skurnik D, Picard B, Denamur E. The population genetics of commensal
Escherichia coli. *Nat Rev Microbiol.* 2010 Mar;8(3):207-17.
40. Denamur E, Clermont O, Bonacorsi S, Gordon D. The population genetics of pathogenic
Escherichia coli. *Nat Rev Microbiol.* 2021 Jan;19(1):37-54.
41. Marin J, Clermont O, Royer G, et al. The population genomics of increased virulence and
antibiotic resistance in human commensal *Escherichia coli* over 30 years in France. *Appl*
Environ Microbiol. 2022 Aug 9;88(15):e0066422.
42. NORM/NORM-VET 2016. Usage of antimicrobial agents and occurrence of antimicrobial
resistance in Norway. Tromsø/Oslo; 2017. ISSN: 1502-2307 (print)/1890- 9965 (electronic).
43.Health NIoP. ESBL-holdige gramnegative stavbakterier - smitteverntiltak i
helseinstitusjoner. 2015. https://www.fhi.no/sv/forebygging-i-helsetjenesten/smittevern-i-
institusjoner/tiltak/esbl-holdige-gramnegative-stavbakte/.
44. Ostholm-Balkhed A, Tarnberg M, Nilsson M, et al. Travel-associated faecal colonization
with ESBL-producing Enterobacteriaceae: incidence and risk factors. *J Antimicrob*
Chemother
. 2013 Sep;68(9):2144-53.
45. Nicolas-Chanoine MH, Gruson C, Bialek-Davenet S, et al. 10-Fold increase (2006-11) in
the rate of healthy subjects with extended-spectrum β-lactamase-producing *Escherichia coli*
faecal carriage in a Parisian check-up centre. *J Antimicrob Chemother.* 2013 Mar;68(3):562-8.
46. Søgaard M, Heide-Jørgensen U, Vandenbroucke JP, Schønheyder HC, Vandenbroucke-
Grauls C. Risk factors for extended-spectrum β-lactamase-producing *Escherichia coli* urinary
tract infection in the community in Denmark: a case-control study. *Clin Microbiol Infect.*
2017 Dec;23(12):952-960.
47. Vihta K-D, Stoesser N, Llewelyn MJ, et al. Trends over time in *Escherichia coli*
bloodstream infections, urinary tract infections, and antibiotic susceptibilities in Oxfordshire,
UK, 1998–2016: a study of electronic health records. *Lancet Infect Dis.* 2018
Oct;18(10):1138-1149.
48. Thaden JT, Fowler VG, Sexton DJ, Anderson DJ. Increasing incidence of extended-
spectrum β-lactamase-producing *Escherichia coli* in community hospitals throughout the
Southeastern United States. *Infect Control Hosp Epidemiol.* 2016 Jan;37(1):49-54.
49. Reddy P, Malczynski M, Obias A, et al. Screening for extended-spectrum β-lactamase-
producing Enterobacteriaceae among high-risk patients and rates of subsequent bacteremia.
Clin Infect Dis. 2007 Oct 1;45(7):846-52.
50. Collingwood A, Blostein F, Seekatz AM, et al. Epidemiological and microbiome
associations between *Klebsiella pneumoniae* and vancomycin-resistant *Enterococcus*
colonization in intensive care unit patients. *Open Forum Infect Dis.* 2020 Jan 12;7(1):ofaa012.

51. Matuschek E, Brown DF, Kahlmeter G. Development of the EUCAST disk diffusion antimicrobial susceptibility testing method and its implementation in routine microbiology laboratories. *Clin Microbiol Infect.* 2014 Apr;20(4):O255-66.

52. Krueger F. TrimGalore. 2019. https://github.com/FelixKrueger/TrimGalore (accessed August 2019.

53. Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: Resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol.* 2017 Jun 8;13(6):e1005595.

54. Bankevich A, Nurk S, Antipov D, et al. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol.* 2012 May;19(5):455-77.

55. Jolley KA, Bray JE, Maiden MCJ. Open-access bacterial population genomics: BIGSdb software, the PubMLST.org website and their applications. *Wellcome Open Res.* 2018 Sep 24;3:124.

56. Seemann T. mlst Github. 2021. https://github.com/tseemann/mlst2021).

57. Zhou Z, Alikhan NF, Mohamed K, Fan Y, Achtman M. The EnteroBase user's guide, with case studies on *Salmonella* transmissions, *Yersinia pestis* phylogeny, and *Escherichia* core genomic diversity. *Genome Res.* 2020 Jan;30(1):138-152.

58. Feldgarden M, Brover V, Gonzalez-Escalona N, et al. AMRFinderPlus and the reference gene catalog facilitate examination of the genomic links among antimicrobial resistance, stress response, and virulence. *Sci Rep.* 2021 Jun 16;11(1):12728.

59. Carattoli A, Zankari E, Garcia-Fernandez A, et al. In silico detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. *Antimicrob Agents Chemother.* 2014 Jul;58(7):3895-903.

60. Seemann T. *Abricate*, Github. 2019. https://github.com/tseemann/abricate.

61. Beghain J, Bridier-Nahmias A, Le Nagard H, Denamur E, Clermont O. ClermonTyping: an easy-to-use and accurate in silico method for *Escherichia* genus strain phylotyping. *Microb Genom.* 2018 Jul;4(7):e000192.

62. Roer L, Tchesnokova V, Allesøe R, et al. Development of a web tool for *Escherichia coli* subtyping based on *fim*H alleles. *J Clin Microbiol.* 2017 Aug;55(8):2538-2543.

63. Madden T. The BLAST sequence analysis tool. in: McEntyre J OJ, editor. http://www.ncbi.nlm.nih.gov/books/NBK21097/: The NCBI Handbook. Bethesda (MD): National Center for Biotechnology Information (US); 2002-. Chapter 16.; 2002.

64. Lam MM, Wick RR, Wyres KL, Holt KE. A genomic surveillance framework and genotyping tool for *Klebsiella pneumoniae* and its related species complex *Nat Commun.* 2021 Jul 7;12(1):4188.

65. Lam MMC, Wyres KL, Judd LM, et al. Tracking key virulence loci encoding aerobactin and salmochelin siderophore synthesis in *Klebsiella pneumoniae*. *Genome Med.* 2018 Oct 29;10(1):77.

66. NORM/NORM-VET 2014. Usage of antimicrobial agents and occurrence of antimicrobial resistance in Norway. Tromsø/Oslo; 2015. ISSN: 1502-2307 (print)/1890- 9965 (electronic).

67. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics.* 2014 Aug 1;30(15):2114-20.

68. Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. Using SPAdes *de novo* assembler. *Curr Protoc Bioinformatics.* 2020 Jun;70(1):e102.

69. Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics.* 2014 Jul 15;30(14):2068-9.

70. Seemann T. snippy: fast bacterial variant calling from NGS reads. 2015. https://github.com/tseemann/snippy.

71. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 2014 May 1;**30**(9):1312-3.

72. Letunic I, Bork P. Interactive tree of life (iTOL) v5: an online tool for phylogenetic tree
display and annotation. *Nucleic Acids Res.* 2021 Jul 2;49(W1):W293-W296.
73. Price LB, Johnson JR, Aziz M, et al. The epidemic of extended-spectrum-beta-lactamase-
producing *Escherichia coli* ST131 is driven by a single highly pathogenic subclone, *H30*-Rx.
mBio. 2013 Dec 17;4(6):e00377-13.
74. Textor J, van der Zander B, Gilthorpe MS, Liskiewicz M, Ellison GT. Robust causal
inference using directed acyclic graphs: the R package 'dagitty'. *Int J Epidemiol.* 2016 Dec
1;45(6):1887-1894.
75. Field A. Discovering statistics using IBM SPSS statistics: SAGE Publications Ltd; 2017.

734