Excess Mortality and Years of Potential Life Lost Among Black People in the US from 1999 to 2020

César Caraballo, MD;[1,2] Daisy S. Massey, BA;[3] Chima D. Ndumele, PhD;[4] Trent Haywood, MD, JD;[5] Shayaan Kaleem;[6] Terris King, BS, MS, ScD;[7] Yuntian Liu, MPH;[1] Yuan Lu, ScD;[1,2] Marcella Nunez-Smith, MD, MHS;[8] Herman A. Taylor, MD, MPH;[9] Karol E. Watson, MD, PhD;[10] Jeph Herrin, PhD;[2] Clyde W. Yancy, MD, MSc;[11] Jeremy S. Faust, MD, MS;[12] Harlan M. Krumholz, MD, SM;[1,2,4]

[1]Center for Outcomes Research and Evaluation, Yale New Haven Hospital, New Haven, Connecticut, United States of America

[2]Section of Cardiovascular Medicine, Department of Internal Medicine, Yale School of Medicine, New Haven, Connecticut, United States of America

[3]University of Massachusetts Chan Medical School, Worcester, Massachusetts, United States of America

[4]Department of Health Policy and Management, Yale School of Public Health, New Haven, Connecticut, United States of America

[5]Zing Health, Chicago, Illinois, United States of America

[6]Department of Human Biology, University of Toronto, Toronto, Canada

[7]Lukan Group, Randallstown, Maryland, United States of America

[8]Equity Research and Innovation Center, Section of General Internal Medicine, Yale School of Medicine, New Haven, Connecticut, United States of America

[9]Cardiovascular Research Institute, Morehouse School of Medicine, Atlanta, Georgia, United States of America

[10]Division of Cardiology, Department of Medicine, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, California, United States of America

[11]Division of Cardiology, Feinberg School of Medicine, Northwestern University, Chicago, Illinois, United States of America

[12]Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, United States of America

Corresponding author:

Harlan M. Krumholz, MD, SM

195 Church Street, Fifth Floor, New Haven, CT 06510

203-497-1246; harlan.krumholz@yale.edu

Word count: 2813; references: 32; tables/figures: 5

ABSTRACT

Background: There is scarce information on recent trends in excess deaths and years of potential life lost among Black populations compared with White populations in the United States.

Methods: We used death certificates data from the Centers for Disease Control and Prevention WONDER from 1999-2020 and evaluated trends in excess age-adjusted mortality rate (deaths per 100,000 individuals) and years of potential life lost (YPLL) among Black people compared with White people. We also evaluated these metrics by cause of death and age.

Results: From 1999 to 2020, the total number of excess deaths was 658,356 and 1,154,108 among Black females and males, representing 34,938,070 and 47,005,048 excess YPLL among Black females among males, respectively. The excess deaths and YPLL were the greatest in deaths due to heart disease and among infants and non-elderly adults. From 1999, the age-adjusted excess mortality rate among Black people declined at an annual average of -9.0 (95% CI: -10.0, -8.0; P<0.001) among females and -16.3 (95% CI: -20.9, -11.6; P<0.001) among males. It plateaued between 2015-2019 among females and 2012-2019 among males. In 2020, rates increased to 395 and 192 excess deaths per 100,000 males and females, respectively, reaching rates close to those of 1999. The trends in rates of excess YPLL followed a similar pattern.

Conclusions: Over a recent 22-year period, Black people in the US lost more than 80 million years of life when compared with White people. After a period of some progress, improvements stalled, and most gains were eliminated in 2020.

Word count: 250

INTRODUCTION

In 1985, the U.S. Department of Health and Human Services (HHS) Secretary Margaret M.

Heckler issued the Report of the Secretary's Task Force on Black and Minority Health, also

known as the Heckler Report.[1] The report found that Black People had strikingly higher

mortality rates than White people, resulting in almost 60,000 excess deaths a year relative to

White People. Almost four decades later, the U.S. continues to have persistently higher mortality

rates in Black People compared with White People with little progress in eliminating disparities

in health and health care access despite continuing national calls for addressing health

inequities.[2-4] Race is a social construct so there is no intrinsic biological reason for those

categorized as Black individuals to have worse outcomes than White individuals. The most likely

reasons for the differences are the influence of social determinants of health, limitations in access

to care and structural barriers indicative of bias (i.e., structural racism).[5-8]

 The Heckler report focused on excess mortality; a metric that conveys the number of

additional lives lost by Black populations compared with what would have occurred if the rates

were the same as those in White populations. Quantifying excess years of potential life lost

(YPLL) provides a deeper measure of the differential burden of excess mortality on Black People

by giving more weight to deaths at younger ages. Publicly reporting excess deaths and YPLL

among Black people could focus attention on the magnitude of the disparities and the urgent

need for progress,[9] but there is relatively little recent information on the trends in YPLL and

excess deaths among Black populations compared with White populations. Addressing this

knowledge gap could motivate national reporting and tracking of excess deaths in the United

States.

Accordingly, we sought to construct contemporary metrics to assess any progress in reducing excess mortality among Black People. We calculated, over the most recent 22-year period for which data are available, the difference in age-adjusted excess mortality and YPLL for Black people compared with White people in the U.S. We also investigated these disparities by causes of death, and how they varied by age. We focus on the period before the COVID-19 pandemic, but also report results for 2020, which provide an early perspective on differences attributable to the pandemic.

METHODS

Data source

We used national U.S. death certificates data from the Centers for Disease Control and Prevention Wide-ranging Online Data for Epidemiologic Research (CDC WONDER) for years 1999 to 2020.[10] For this period, we obtained the annual life expectancy by 5-year age groups from the National Center for Health Statistics (NCHS) life tables.[11] The Yale Institutional Review Boards waived this study from review.

Study population

We included data from non-Hispanic Black and non-Hispanic White people aged <85 years old. From CDC WONDER, for each race group we obtained annual number of deaths, population size (N), and age-adjusted mortality rate (per 100,000 individuals) overall and by sex (female, male), age (years), and cause of death (Supplementary Methods and Table S1)

Race and ethnicity were determined by the information recorded in the death certificates, which is reported by the funeral director as provided by an informant (e.g., next of kin) or on the basis of observation if there is no informant available.[10]

Statistical analysis

All analyses described below were stratified by sex, and all rates were reported per 100,000 individuals.

For each year, we first calculated excess mortality rate by subtracting the year-specific age-adjusted mortality rate among White people from that of Black people. Then, we estimated annual YPLL (defined as the number of years that a person would have lived should they not died when they did) rate for each race by multiplying each 5-year age group crude mortality rate by their respective annual sex- and age-specific life expectancy in years and obtaining the mean rate across all age groups. For both racial groups, we used year-, age-, and sex-specific White life expectancy as the reference (Table S2).[11] We then estimated the Black-White age-adjusted mortality rate ratio, YPLL rate ratio, and the cumulative 22-year total number of excess deaths and excess YPLL among Black people (Supplementary Methods).

To assess the trends over the study period, we graphically assessed the relationship between each metric and study year. Based on this assessment, we used autoregressive integrated moving average models using a 1-year correlation and modelled time as a linear spline with knots that reflected the observed inflection points (Supplementary Methods). Separately, we calculated and visually evaluated annual rates in age-adjusted excess deaths and YPLL by the leading causes of death (Supplementary Methods and Table S1).

Finally, to estimate mortality differences by age, we combined all study years and used the average crude mortality rate for each 5-year age group and subtracted the White people's crude mortality from that of Black people of the same age group. Similarly, to estimate the YPLL rate differences by age, we estimated the annual crude mortality rate for each age group, multiplied it by their respective annual sex- and age-specific life expectancy in years, and then estimated the mean YPLL rate for each age group across the entire study period.

Cause of death heatmaps were produced using Python version 3.7.[12-16] All statistical tests performed were two-sided, with a level of significance of 0.05, using Stata SE version 17.0 (StataCorp, College Station TX).

RESULTS

Cumulative 22-year Excess Mortality and YPLL Among Black People

Overall, from 1999 to 2020, the mean excess mortality rate (deaths per 100,000 individuals) among Black females and males was 148.6 and 284.2, respectively, representing an estimated total of 658,356 excess deaths among Black females and 1,154,108 excess deaths among Black males over the study period. The mean 22-year excess YPLL rate (per 100,000 individuals) was 7932 among females and 11579 among males, representing an estimated total of 34,938,070 excess YPLL among Black females and 47,005,048 excess YPLL among Black males.

Temporal Trends in Racial Differences in Mortality and YPLL Rates

From 1999 to 2011, the age-adjusted excess mortality rate declined from 404 to 211 excess deaths per 100,000 individuals among Black males, at an annual average estimated decline of -16.3 excess deaths per 100,000 individuals (95% CI: -20.9, -11.6; P<0.001; Figure 1).

However, the rate plateaued from 2012 to 2019 (-1.4 per year; 95% CI: -9.6, +6.7; P=0.73), with a subsequent single-year increase to 395 excess deaths per 100,000 individuals in 2020; reaching rates close to those of 20 years before. Among Black females, the estimated age-adjusted excess mortality rate declined from 224 excess deaths (per 100,000 individuals) in 1999 to 87 in 2015, at an annual average estimated decline of -9.0 (95% CI: -10.0, -8.0; P<0.001) excess deaths. There was no significant trend from 2016 to 2019 (+0.9 per year; 95% CI: -3.7, +5.5; P=0.70), and rates increased to 192 excess deaths per 100,000 individuals in 2020; similar to rates of 2005 (Figure 1 and Figure S1). The age-adjusted mortality rate ratio between the two groups followed a similar pattern over the 22-year period, ranging from 1.13 to 1.31 among females and from 1.21 to 1.40 among males (Figure 1).

In 1999, the excess YPLL for Black males and Black females was 14,964 and 10,806 per 100,000 people, respectively. There was a continuous decline until 2011 among males, reaching 9348 excess YPLL per 100,000 individuals, and until 2012 among females, reaching 6210 excess YPLL per 100,000 individuals (P<0.001 for males and females; Figure 2). Then there was a plateau until 2019 for both groups (P=0.90 for males; P=0.89 for females), after which the excess YPLL among both Black males and Black females increased in 2020 to rates similar to those of 19 and 13 years before, respectively (Figure 2 and Figure S1). The annual YPLL rate ratio between the two groups followed a similar pattern over the 22-year period (Figure 2).

When analyzed by cause of death, the highest excess mortality rates among Black people through the study period were due to heart disease (Figure 3); in both sex groups there was a gradual decrease in mortality due to heart disease in the decade of 2010, metric that increased again in 2020. Among males, there was a gradual reduction in excess mortality rates due to cancer, diabetes, and HIV, whereas excess mortality due to assault and cerebrovascular diseases

were persistently high over the study period, with an increase in 2020. Among females, the

excess mortality rates due to cancer, diabetes, and cerebrovascular disease gradually decreased

from 1999 to 2020. There were consistent patterns in YPLL rates by cause of death for females

and males. In 2020, the excess age-adjusted mortality rates due to COVID-19 were 47 per

100,000 Black females and 80 per 100,000 Black males; similarly, the excess YPLL (per

100,000 individuals) due to COVID-19 were 1759 among Black females and 2572 among Black

males.

Racial Differences in Mortality Rate and YPLL Rate by Age Group

The relationship between excess mortality rates and age among females and males is

shown in Figure 4. Among Black females and Black males younger than 1 year, there were 654

and 776 excess deaths per 100,000 individuals, respectively. Such rates narrowed among those

aged 1 and 14 years. Among males, the excess mortality rate increased with age starting among

those 15 years and older, reaching 1302 excess deaths per 100,000 individuals among those

between 75-79 years. A similar pattern, although at a smaller magnitude, was observed among

Black females. The increase in mortality rate difference increased among Black females older

than 20 years and reached a maximum of 677 excess deaths per 100,000 individuals among those

between 75-79 years old. For both males and females, there was a slight decrease in excess

deaths among those aged 80-84 years. In relative terms, patterns for the mortality rate ratio

between Black and White individuals were similar (Figure 4).

The greatest difference in YPLL rate between Black people and White people was among

the youngest (Figure 5). Among those younger than 1 year old, excess YPLL rate was 59,232 per

100,000 Black males and 53,061 per 100,000 Black females, decreasing to its minimum among

those between 1 and 14 years old. Consistent with the excess mortality pattern, the excess YPLL

rate increased with age among males older than 15 years and among females older than 20 years,

peaking among those aged 65-69 years old and decreasing among those older. The YPLL rate

ratio across age groups followed a similar pattern (Figure 5).

DISCUSSION

In this study of US mortality data from 1999 to 2020, we estimate that there were 35

million excess YPLL among Black females and 47 million excess YPLL among Black males.

After a period of some progress, improvements stalled and even reversed up until the first year of

the pandemic, when the number of excess deaths was about the same as 1999, and the YPLL loss

was even greater. Moreover, even at the nadir of excess of deaths and YPLL over this 22-year

period, there were a massive annual number of excess deaths and more than a million years of

potential life lost among Black people compared with White people. This loss of life was most

prominent among infants, with Black-White mortality and YPLL rate ratios greater than 2.3

among those <1 year old. Importantly, heart disease in females and males and cancer in males

had the most excess deaths.

This study quantifies the numbers of years lost to Black populations in the

United States. This massive difference in deaths and YPLL are the result of how

social factors largely related skin color, a characteristic with little intrinsic biological

meaning but immense social consequences. In the absence of biological

determinants, the role of social construct-–including, social determinants of health,

access to care, structural barriers, and racism – are leading causal factors.[5-8]

Moreover, this study shows a brief period of progress that stalled and then was erased

with the inequities of harm associated with the early pandemic. The findings herald a need for new approaches and the metrics are useful for national accountability.

Prior studies have found persistently large numbers of excess deaths among Black people but have not explicitly quantified excess death or YPLL. For example, a study found that the mortality rate disparity between Black people and White people remained mostly stable from 1960 to 2000,[17] and another report estimated that 2,687,051 excess deaths occurred among Black people from 1970 to 2004.[18] From 1999 through 2014, we found that excess mortality declined and was followed by a period of stalling of progress, consistent with several other studies of Black and White mortality rates.[19,20] This observation could be at least partially explained by a disproportionally higher increase in mortality among White adults due to accidental drug overdoses in the late 2000s.[21,22] Notably, between 2010 and 2017, the largest relative increase in fatal drug overdoses occurred among Black adults,[21,23] which could also partially explain the plateau observed in the second decade of the study period. In addition, the health disparity in life expectancy, which had been narrowing in the 2000s, stagnated and may have begun to widen in 2015.[24,25] Our finding reveal that the disparity in mortality rates between Black and White people did not improve further up to 2019.

Another key observation is the marked increase in excess Black mortality and YPLL rates during the first year of the COVID-19 pandemic. Many studies have described disparities in the COVID-19 pandemic toll, but this study places it in context of excess mortality trends. The increase in excess mortality during the pandemic was so large that it eliminated the gains achieved in reducing mortality

differences with White people during the prior two decades. Such a dramatic reversal underscores the pernicious social factors disproportionally positioning Black people at a higher risk of adverse health outcomes, including barriers to health care, higher multimorbidity prevalence, and worse average health status.[2,3,26] It also indicates the fragility of recent gains.

The excess deaths and YPLL were elevated among most of the major causes of death, though heart disease was the most prominent. This is consistent with the findings from Kyalwazi and colleagues[27] who found that, despite a decline in the absolute difference, age-adjusted cardiovascular mortality rates were persistently higher among Black people when compared with White people from 1999 to 2019. It is likely that disparities in rates of hypertension control and other factors associated with heart disease are responsible for this excess loss of life.[28,29] Cancer was also an important source of the disparity. Importantly, heart disease and cancer have many modifiable risk factors and are importantly affected by social determinants of health. Nevertheless, the fact that there are excess deaths in most causes suggests that social determinants related to race are key to these conditions. Targeted and renewed efforts aimed at diseases that disproportionately drive differences are necessary to make sure clinical advancements are experienced uniformly across the population.

There is a marked disparity among infants, accounting for a markedly elevated number of excess deaths and an even more pronounced disparity in YPLL. After infancy, the excess deaths and YPLL among Black people became evident in early adulthood and generally increased with age. Among those older than age 65 years

these disparity metrics declined, as would be expected because of the persistently lower average lifespan among Black people.[25,30]

The implications of the study are profound. Black people in the US continue to die at much greater rates than White people, with a measurable excess loss of years of life. These metrics are suitable to be publicly reported and tracked frequently to drive strategies directed toward more in-depth root cause analysis and where qualified by evidence of efficacy, implementation of steps to improve these disparities. Moreover, this study identifies populations and conditions that may produce the greatest public health returns in addressing disparities, even as the overall finding suggests there needs to be global strategies directed against the embedded structural barriers to health and healthcare in all cohorts, not just by race.

Our study has limitations. First, there are known inaccuracies in death certificates,[31,32] from which CDC WONDER estimates are derived. However, such inaccuracies are less common for non- Hispanic Black and White decedent's death certificates.[33] Less importantly, cause of death in 2020 may have been unreliable and some deaths due to COVID-19 may be undercounted. However, these limitations are unlikely to have impacted the overall finding of an important, sustained difference in mortality between these two groups of people in the U.S.

In conclusion, in the US, massive numbers of excess deaths and YPLL among Black people persist and by scale warrant our urgent national attention and indicate the need for new strategies. Over a recent 22-year period, Black people in America lost more than 80 million years of life. With the pandemic, most gains were eliminated and YPLL were higher than they had ever been. The calls to action from

the Heckler Report remain just as relevant, or more so, today.

REFERENCES

1.   Heckler M. Report of the Secretary's task force on Black & minority health. US Department of Health and Human Services; 1985.

2.   Mahajan S, Caraballo C, Lu Y, et al. Trends in Differences in Health Status and Health Care Access and Affordability by Race and Ethnicity in the United States, 1999-2018. JAMA 2021;326(7):637-648. DOI: 10.1001/jama.2021.9907.

3.   Caraballo C, Herrin J, Mahajan S, et al. Temporal Trends in Racial and Ethnic Disparities in Multimorbidity Prevalence in the United States, 1999-2018. Am J Med 2022 DOI: 10.1016/j.amjmed.2022.04.010.

4.   Caraballo C, Ndumele CD, Roy B, et al. Trends in Racial and Ethnic Disparities in Barriers to Timely Medical Care Among Adults in the US, 1999 to 2018. JAMA Health Forum 2022;3(10):e223856-e223856. DOI: 10.1001/jamahealthforum.2022.3856.

5.   Churchwell K, Elkind MSV, Benjamin RM, et al. Call to Action: Structural Racism as a Fundamental Driver of Health Disparities: A Presidential Advisory From the American Heart Association. Circulation 2020;142(24):e454-e468. DOI: 10.1161/cir.0000000000000936.

6.   Bailey ZD, Krieger N, Agénor M, Graves J, Linos N, Bassett MT. Structural racism and health inequities in the USA: evidence and interventions. Lancet 2017;389(10077):1453-1463. DOI: 10.1016/s0140-6736(17)30569-x.

7.   Bailey ZD, Feldman JM, Bassett MT. How Structural Racism Works - Racist Policies as a Root Cause of U.S. Racial Health Inequities. N Engl J Med 2021;384(8):768-773. DOI: 10.1056/NEJMms2025396.

8.   Cogburn CD. Culture, Race, and Health: Implications for Racial Inequities and Population Health. Milbank Q 2019;97(3):736-761. DOI: 10.1111/1468-0009.12411.

9.   Krumholz HM, Massey DS, Dorsey KB. Racism as a leading cause of death in the United States. BMJ 2022;376:o213. DOI: 10.1136/bmj.o213.

10.   Centers for Disease Control and Prevention. CDC WONDER. March 14, 2022 (https://wonder.cdc.gov/).

11.   National Center for Health Statistics. Life Tables. April 25, 2022 (https://www.cdc.gov/nchs/products/life_tables.htm).

12.   Python Software Foundation. Python Language Reference, version 3.7. (http://www.python.org).

13.   Hunter JD. Matplotlib: A 2D Graphics Environment. Computing in Science & Engineering 2007;9(3):90-95. DOI: 10.1109/MCSE.2007.55.

14.   McKinney W. Data structures for statistical computing in python. Proceedings of the 9th Python in Science Conference: Austin, TX; 2010:51-56.

15.   Waskom ML. Seaborn: statistical data visualization. J Open Source Softw 2021;6(60):3021.

16.   Harris CR, Millman KJ, van der Walt SJ, et al. Array programming with NumPy. Nature 2020;585(7825):357-362. DOI: 10.1038/s41586-020-2649-2.

17.   Satcher D, Fryer GE, Jr., McCann J, Troutman A, Woolf SH, Rust G. What if we were equal? A comparison of the black-white mortality gap in 1960 and 2000. Health Aff (Millwood) 2005;24(2):459-64. DOI: 10.1377/hlthaff.24.2.459.

18. Rodriguez JM, Geronimus AT, Bound J, Dorling D. Black lives matter: Differential mortality and the racial composition of the U.S. electorate, 1970-2004. Soc Sci Med 2015;136-137:193-9. DOI: 10.1016/j.socscimed.2015.04.014.

19. Cunningham TJ, Croft JB, Liu Y, Lu H, Eke PI, Giles WH. Vital Signs: Racial Disparities in Age-Specific Mortality Among Blacks or African Americans - United States, 1999-2015. MMWR Morb Mortal Wkly Rep 2017;66(17):444-456. DOI: 10.15585/mmwr.mm6617e1.

20. Pathak EB. Mortality Among Black Men in the USA. J Racial Ethn Health Disparities 2018;5(1):50-61. DOI: 10.1007/s40615-017-0341-5.

21. Woolf SH, Schoomaker H. Life Expectancy and Mortality Rates in the United States, 1959-2017. JAMA 2019;322(20):1996-2016. DOI: 10.1001/jama.2019.16932.

22. Ruhm CJ. Drug Mortality and Lost Life Years Among U.S. Midlife Adults, 1999-2015. Am J Prev Med 2018;55(1):11-18. DOI: 10.1016/j.amepre.2018.03.014.

23. Scholl L, Seth P, Kariisa M, Wilson N, Baldwin G. Drug and Opioid-Involved Overdose Deaths - United States, 2013-2017. MMWR Morb Mortal Wkly Rep 2018;67(5152):1419-1427. DOI: 10.15585/mmwr.mm675152e1.

24. Bilal U, Diez-Roux AV. Troubling Trends in Health Disparities. N Engl J Med 2018;378(16):1557-1558. DOI: 10.1056/NEJMc1800328.

25. Life expectancy by county, race, and ethnicity in the USA, 2000-19: a systematic analysis of health disparities. Lancet 2022;400(10345):25-38. DOI: 10.1016/s0140-6736(22)00876-5.

26. Lavizzo-Mourey RJ, Besser RE, Williams DR. Understanding and Mitigating Health Inequities - Past, Current, and Future Directions. N Engl J Med 2021;384(18):1681-1684. DOI: 10.1056/NEJMp2008628.

27. Kyalwazi AN, Loccoh EC, Brewer LC, et al. Disparities in Cardiovascular Mortality Between Black and White Adults in the United States, 1999 to 2019. Circulation 2022;146(3):211-228. DOI: 10.1161/circulationaha.122.060199.

28. Shah NS, Lloyd-Jones DM, O'Flaherty M, et al. Trends in Cardiometabolic Mortality in the United States, 1999-2017. JAMA 2019;322(8):780-782. DOI: 10.1001/jama.2019.9161.

29. Lopez KN, Morris SA, Tejtel SKS, Espaillat A, Salemi JL. US Mortality Attributable to Congenital Heart Disease Across the Lifespan From 1999 Through 2017 Exposes Persistent Racial/Ethnic Disparities. Circulation 2020;142(12):1132-1147. DOI: doi:10.1161/CIRCULATIONAHA.120.046822.

30. Johnson CO, Boon-Dooley AS, DeCleene NK, et al. Life Expectancy for White, Black, and Hispanic Race/Ethnicity in U.S. States: Trends and Disparities, 1990 to 2019. Ann Intern Med 2022 DOI: 10.7326/m21-3956.

31. Arias E, Schauman WS, Eschbach K, Sorlie PD, Backlund E. The validity of race and Hispanic origin reporting on death certificates in the United States. Vital Health Stat 2 2008(148):1-23.

32. Hoffman RA, Venugopalan J, Qu L, Wu H, Wang MD. Improving Validity of Cause of Death on Death Certificates. Acm bcb 2018;2018:178-183. DOI: 10.1145/3233547.3233581.

33.   Arias E, Heron M, Hakes J. The Validity of Race and Hispanic-origin Reporting on

Death Certificates in the United States: An Update. Vital Health Stat 2 2016(172):1-21.

ACKNOWLEDGMENTS

Disclosures: In the past three years, **Harlan Krumholz** received expenses and/or personal fees from UnitedHealth, Element Science, Aetna, Reality Labs, Tesseract/4Catalyst, F-Prime, the Siegfried and Jensen Law Firm, Arnold and Porter Law Firm, and Martin/Baughman Law Firm. He is a co-founder of Refactor Health and HugoHealth, and is associated with contracts, through Yale New Haven Hospital, from the Centers for Medicare & Medicaid Services and through Yale University from Johnson & Johnson. **Herman A. Taylor** has worked as an advisor and consultant for United Health Group, Pfizer, and Novartis. The other authors had nothing to disclose.

Funding: This study did not receive external funding.

Data availability statement: All data used in this study is publicly available from https://wonder.cdc.gov/. All the code files used to analyze the data are publicly available from https://zenodo.org/record/7278473 Dr. Caraballo had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

FIGURES

Figure 1. Age-Adjusted Excess Mortality Rate (Left Panel) and Rate Ratio (Right Panel) Among Black People, 1999-2020

Legend: Left panel displays the annual excess age-adjusted mortality rates among Black people, defined as the difference in each year's mortality rate between Black people and White people. Right panel displays the annual age-adjusted mortality rate ratio, obtained by dividing the annual mortality rate among Black people by that of White people. An excess mortality rate of zero represents the absence of a difference, and a rate ratio of one represents equal rates between both groups. Rates were estimated per 100,000 individuals.

Figure 2. Excess Years of Potential Life Lost Rate (Left Panel) and Rate Ratio (Right Panel) Among Black People, 1999-2020

Legend: Left panel displays the annual excess YPLL rates among Black people, defined as the difference in each year's YPLL rate between Black people and White people. Right panel displays the annual YPLL rate ratio, obtained by dividing the annual YPLL rate among Black people by that of White people. An excess YPLL rate of zero represents the absence of a difference, and a rate ratio of one represents equal rates between both groups. Rates were estimated per 100,000 individuals.

Abbreviations: YPLL, years of potential life lost.

Figure 3. Black People Excess Mortality and Years of Potential Life Lost Rates by Cause of

Death, 1999-2020

    A) Females

B) Males

Legend: Top panels display annual excess age-adjusted mortality rates among Black females (A, in shades of blue) and Black males (B, in shades of red). Bottom panels display annual excess years of potential life lost among Black people. Excess mortality and YPLL rates were defined as the difference in each year's mortality rate (or YPLL rate) between Black people and White people. A higher number in both estimates represent a socially negative outcome; zero represents the absence of a difference. All rates are estimated per 100,000 individuals. Information related

to COVID-19 deaths are not displayed in the figure because data were only applicable to a single

year. Abbreviations: YPLL, years of potential life lost.

Figure 4. Excess Mortality Rate (Left Panel) and Rate Ratio (Right Panel) Among Black People by Age Group

Legend: Left panel displays the excess mortality rates among Black people by age group, defined as the difference in each age group mortality rate between Black people and White people. Right panel displays the mortality rate ratio by age group, obtained by dividing each age group mortality rate among Black people by that of White people. An excess mortality rate of zero represents the absence of a difference, and a rate ratio of one represents equal rates between both groups. Rates were estimated per 100,000 individuals.

Figure 5. Excess Years of Potential Life Lost Rate (Left Panel) and Rate Ratio (Right Panel) Among Black People by Age Group

Legend: Left panel displays the excess YPLL rates among Black people by age group, defined as the difference in each age group YPLL rate between Black people and White people. Right panel displays the YPLL rate ratio by age group, obtained by dividing each age group YPLL rate among Black people by that of White people. An excess YPLL rate of zero represents the absence of a difference, and a rate ratio of one represents equal rates between both groups. Rates were estimated per 100,000 individuals.

Abbreviations: YPLL, years of potential life lost.