TITLE

Foetal Allogeneic Intracerebroventricular Neural Stem Cell Transplantation in

People with Secondary Progressive Multiple Sclerosis: A phase I dose-

escalation clinical trial

AUTHORS

Leone MA[1*], Gelati M[1*], Profico DC[1], Conti C[2], Spera C[2], Muzi G[2], Grespi V[1], Bicchi I[1],

Ricciolini C[1], Ferrari D[4], Zarrelli M[1], Amoruso L[1], Placentino G[1], Crociani P[1], Apollo F[1], Di

Viesti P[1], Fogli D[1], Popolizio T[1], Colosimo C[2], Frondizi D[2], Stipa G[2], Tinella E[2], Ciampini A[2],

Sabatini S[2], Paci F[2], Silveri G[1], Gobbi C[3,6], Pravatà E[3,6,7], Zecca E[3,6], Balzano RF[7], Kuhle J[5],

Copetti M[1], Fontana A[1], Carella M[1], D'Aloisio G[1], Abate L[1], Pluchino S[8], Peruzzotti-Jametti L[8]

and Vescovi AL[1,4]

* The two Authors contributed equally to the manuscript

1. IRCCS Casa Sollievo della Sofferenza, Viale Cappuccini 1, 71013 San Giovanni Rotondo,

Foggia, Italy

2. AOSP Santa Maria, via Tristano di Joannuccio 1, 05100, Terni, Italy

3. Multiple Sclerosis Centre (MSC), Department of Neurology, Neurocentre of Southern

Switzerland, EOC, 6900 Lugano, Switzerland

4. BTBS Università Bicocca, piazza della Scienza 2, 20126, Milan, Italy

5. Department of Neurology, University Hospital Basel, and University of Basel/Switzerland

6. Faculty of biomedical Sciences, Università della Svizzera Italiana (USI), 6900 Lugano,

Switzerland

7. Department of Neuroradiology, Neurocentre of Southern Switzerland, EOC, 6900 Lugano,

Switzerland

8. Department of Clinical Neurosciences and NIHR Biomedical Research Centre, University of

Cambridge, CB2 0QQ Cambridge, United Kingdom.

Corresponding Author: Prof. Angelo Luigi Vescovi. Email vescovia@gmail.com

IRCCS Casa Sollievo della Sofferenza, Viale Cappuccini 1, 71013 San Giovanni Rotondo,

Foggia, ItalyTel +39 0882 416566

ABSTRACT

Background: Advanced cell therapeutics are emerging as potentially effective treatments

for chronic neurological diseases, including secondary progressive multiple sclerosis (SPMS).

Here we report the results of a phase I trial in which good manufacturing practice-grade

foetal allogeneic human neural stem cells (hNSCs) were implanted via

intracerebroventricular (ICV) injection in 15 individuals with active and non-active SPMS.

Methods: This is a phase I, open-label, multicentre, dose-escalation, international study. The

primary objective was to assess the feasibility, safety, and tolerability of ICV injections of

allogeneic hNSCs in patients affected by SPMS over a study follow up of 12 months. We also

evaluated the number and type of adverse events (AEs) leading to a maximum tolerated

dose, the general health status, and mortality. The secondary objectives were the

therapeutic benefit of allogeneic hNSCs using assessment scales, magnetic resonance

imaging (MRI), and laboratory and neurophysiologic parameters.

Findings: Fifteen unrelated SPMS patients were enrolled and treated between 2018 and

2020. The participants had a median age of 49.8 years. Their mean extended disability

status scale (EDSS) at enrolment was 7.6, the mean disease duration was 22 years, and

mean time from diagnosis to progression was 10.1 years. Neither treatment-related deaths

nor serious AEs were reported during the study (1 year follow up after treatment). All the

other AEs were classified as non-serious and were associated to non-study concomitant

therapy or other medical conditions not connected to the experimental treatment. During

the study, none of the participants worsened in the progression of their SPMS as shown by

the evaluation scales implemented to assess their progress. Laboratory and

neurophysiologic parameters showed no clinically significant variations. MRI follow-up

showed non-clinically significant type 1, 2, and 3 changes.

Interpretation: The intracerebroventricular injection of foetal allogeneic hNSCs in people

with SPMS is feasible, tolerated and safe. Study participants displayed a substantial clinical

stability during the 12-month follow-up. The absence of relevant adverse reactions (Ars)

arising from the transplantation of hNSCs indicates a short-term neutral balance between

benefits and risks and suggests a concrete, though perspective therapeutic possibility for

SPMS patients. Further studies are needed to confirm and extend the findings herein and

evaluate the actual therapeutic potential of advanced cell therapeutics for a condition

where the lack of effective disease modifying therapies is a major unmet clinical need.

INTRODUCTION

Multiple sclerosis (MS) is the most common cause of non-traumatic disability in young adults, with

over 2·5 million sufferers worldwide.[1] The last decade has seen a fast development of effective

disease modifying treatments (DMTs) for patients with a relapsing remitting form of MS (RRMS).

However, as the disease evolves into secondary progressive MS (SPMS), DMTs have limited efficacy.

Thus, SPMS patients currently have a major unmet need.[2,3] A solution to the current lack of effective

treatments for SPMS may come from emerging cell therapy approaches, which have shown

promising initial results in other central nervous system (CNS) diseases, such as amyotrophic lateral

sclerosis (ALS) and Parkinson's disease.[4]

In vivo studies in rodent and non-human primate models of MS have shown that human neural stem

cells (hNSCs)[5-8] are an effective and safe tool to induce CNS functional recovery due to their tissue

specificity and multiple mechanisms of action.[9,10] hNSCs not only repair the damaged CNS by

replacing cells lost to injury, but also exert immunomodulatory actions on both innate and adaptive

immune responses via secretion of trophic factors, cross-correction of missing enzymatic activities,

and metabolic reprogramming.[11-14] Additional data suggest that delivering these cells directly to the

CNS via a single intracerebroventricular (ICV) injection allows maximising the number of cells that

reach the CNS[15-17] and may be key to target the intense compartmentalised inflammation that drives

progression in SPMS.[18] Indeed, following ICV injection, transplanted hNSCs spread throughout the

ventricular and subarachnoid space,[16,19] enabling their inflammation-guided migration into the CNS,

where they may reach axons and myelinating cells directly without crossing the blood brain barrier.[20]

The ICV cell injection is based on a widely used, technically simple, rapid, and standardised

neurosurgical procedure with a minimal rate of complications (6·8%), mainly due to catheter

malposition, haemorrhage, and infection (Morgenstern et al., 2016) . ICV is also a relatively

standardised experimental procedure, since it has already been used for the injection of growth

factors in ALS patients[21] and for chemotherapeutic agents in anti-tumour therapy. [22]

Here, we report the results of the first phase I, open-label, multicentre, dose-escalation study in

which good manufacturing practice (GMP)-grade foetal allogeneic human neural stem cells (hNSCs)

were implanted via intracerebroventricular (ICV) injection in 15 individuals with SPMS. To the best of

our knowledge, this is also the largest phase I trial with hNSCs in people with SPMS conducted to

date.

METHODS

Study design

In this phase I, open-label, multicentre, international, dose-escalation study, the participating patients were affected by SPMS with progressive accumulation of disability after an initial relapsing course, with or without disease activity. Fifteen patients, between 18 and 60 years of age were enrolled according to a "standard" dose-escalation phase I design following a modified Fibonacci sequence (100%, 60% and 50% dose increments). After the initial screening, all screened patients entered a 3-month run-in phase. After that, the patients were prospectively enrolled into four cohorts receiving four different doses of allogenic hNSCs (5, 10, 16 and 24 million cells).

This study was performed in three participating centres located in Italy and Switzerland. The Italian centres, the "IRCCS Casa Sollievo della Sofferenza" Research Hospital (Site 1) and the "Santa Maria di Terni" Hospital (Site 2), recruited the patients. The MSC of the Neurocentre of Southern Switzerland performed the magnetic resonance imaging (MRI) analysis, while ICV treatment was performed at Site 2.

The study was approved by the Ethical Committee of the Istituto Tumori "Giovanni Paolo II" (Bari) from the "Fondazione IRCCS Casa Sollievo della Sofferenza" Research Hospital (01PU/2016–21-01-2016); the Ethical Committee of the "Aziende Sanitarie dell'Umbria" (2404/17); the Agenzia Italiana del Farmaco (AIFA); the Istituto Superiore di Sanità (3090(16)-PRE21-1408–06-04-2016). The trial was subsequently registered in the European Clinical Trials Database (EudraCT, 2015-004855-37), and in ClinicalTrials.gov (NCT03282760).

Participants

Eligible patients were adults of either sex with a diagnosis of SPMS, with or without disease activity (Lublin 2014) with an Expanded Disability Status Score (EDSS)[23] ≥ 6.5 and ≤ 8, showing a progressive

accumulation of disability after initial relapsing course over the 2 years before recruitment (≥ 1.0

point for patients with EDSS =6.5 at the time of inclusion, and ≥ 0.5 points for patients with EDSS >

6.5 at the time of inclusion), and ineligibility to other therapeutic alternatives (as assessed by the

treating neurologist). All patients signed a written informed consent to be enrolled in the study.

Exclusion criteria included: other neurologic conditions; psychiatric/personality disorders or severe

cognitive decline; history of significant systemic, infectious, oncologic, or metabolic disorders; other

autoimmune diseases; chronic infections (HBV, HCV, HIV, tuberculosis); inability to undergo MRI

scans; inability to provide informed consent, received immunomodulant/immunosuppressive

treatments <6 months before inclusion; participated in other research; any contra-indication to

lumbar puncture; and were pregnant or breast feeding.

At the end of the run-in period, and if no serious co-morbidity nor health status changes occurred,

patients were deemed eligible for the intervention. After eligibility assessment and provision of

informed consent, each eligible patient was registered in the "Database for Clinical Studies with

Gene and Somatic Therapy" of the "Istituto Superiore di Sanità" (ISS). Before registration, the ISS

verified patient eligibility to ensure criteria were respected. Upon registration, the database

assigned a unique identification number to each patient which was used for anonymisation

purposes.

To establish baseline clinical features, a 3-month run-in period was started after the screening

examination. All patients were evaluated at the onset and the end of the run-in period by

investigating and performing physical and neurological examination, vital signs, pregnancy test in

fertile women, haematological and urine tests, lumbar puncture for standard CSF examination and

JC Virus test, serum and CSF collection (stored at minus 80 degrees), motor, sensory and visual

evoked potentials, optical coherence tomography (OCT), EDSS, MFSC, RAO brief repeatable battery,

MS-QOL54 for the evaluation of quality of life (Solari et al. 1999), brain and spinal MRI. At the end of

the run-in period, and if no serious co-morbidity nor health status changes occurred, patients were

deemed eligible for the intervention. Around 50% of patients had at least one new or one enlarging

T2-visible lesion, while 43% of patients had at least one lesion with contrast enhancement during

the run-in period.

Study objectives and outcomes

The primary objective of the study was to assess the feasibility, safety, and tolerability of foetal

allogeneic hNSCs delivered via ICV injection in people with active and non-active SPMS patients by

evaluating the following outcomes: mortality, the number and type of adverse reactions (ARs) or

events (AEs) leading to a maximum tolerated dose.

The secondary objectives were to evaluate the functional effects of hNSC therapetics by monitoring

the following outcomes of disease progression: functional disability via EDSS and Multiple Sclerosis

Functional Composite (MSFC)[24] ; annualised relapse rate and time to confirmed relapse; cognitive

function by Rao's brief repeatable battery [BRB] of neuropsychological tests; visual, sensory and

motor functions via combination of electrophysiological measurements (visual, somato-sensory,

motor evoked potentials [EP]); and optical coherence tomography (OCT). Brain MRI evaluations

were performed to monitor structural changes related to both the intervention and disease activity.

MRIs were acquired at the two recruiting centres, using a Philips Ingenia scanner (Philips Medical

Systems, Best, The Netherlands) and a Siemens Verio scanner (Siemens, Erlangen, Germany).

Sequences type and parameters were harmonised between the two vendors to improve

reproducibility. The MRIs were performed at run-in onset (month -3), run-in end (month 0), and at

months +1, +2, +3, +4, +5, +6, +9 and +12.

Potential effects of the treatment on biomarkers of neuronal loss and inflammation was also

evaluated. Cerebrospinal fluid (CSF) and serum neurofilament (NfL) levels were used as markers of

neurodegeneration/neuronal damage, while CHI3L1 (also known as YKL-40) as indicator of reactive

astrocytes.[25] We also investigated serum and CSF levels of IL-17a, IL-2, IL-8, TNF-α, CCL2, CCL3,

CX3CL1, VEGF-a, OPN and GFAP as additional exploratory objectives. All the biomarkers were

evaluated in both CSF and serum collected before and after treatment (see table 1 for time points)

Clinical evaluation

After ICV injection, the participants were followed up for 12 months post-treatment (**Table 1**).

General health status and mortality evaluations were based on the occurrence of any serious co-

morbidities or changes in the general health of the participants post-intervention and the number of

deaths due to the treatment, or the procedure itself.

Table 1: Clinical trial study plan

	Screening	Run in START	Run in END	Surgery	M1	M2	M3	M4	M5	M6	M7	M8	M9	M10	M11	M12
Informed consent signature	x	x														
Medical history	x															
Neurological history	x															
Physical examination	x	x	x													
Infectious screening	x															
Vital Signs	x	x	x		x		x			x			x			x
Pregnancy test for fertile women	x	x	x													
Hematological tests	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
Electrocardiogram	x															
Chest X-ray	x															
standard urine test	x	x	x		x		x			x			x			x
Neurological examination	x	x	x		x	x	x	x	x	x	x	x	x	x	x	x
VAS		x	x		x	x	x	x	x	x	x	x	x	x	x	x
EDSS		x	x		x		x			x			x			x
MSFC		x	x		x		x			x			x			x
Evoked potential test		x	x		x		x			x			x			x
OCT		x	x		x		x			x			x			x
Brain and spinal MRI		x	x	x	x	x	x	x	x	x			x			x
RAO battery		x	x		x		x			x			x			x
MS-QOL54		x	x		x		x			x			x			x
Conselling with psycologist	x				x											
Serum collection for biomarkers		x	x		x		x			x			x			x
CSF collection for biomarkers		x	x		x					x						x
JC virus examination		x	x		x					x						x
AE record				x		x	x	x	x	x	x	x	x	x	x	x
Thin slice cranial CT or MRI				x	x											

All AEs were recorded and evaluated for their relationship with the hNSCs or injection procedure.

Each AE was classified and categorised according to the International Conference on Harmonisation

guidance for Clinical Safety Data Management's Definitions and Standards for Expedited Reporting

E2A and the Common Terminology Criteria for Adverse Events (CTCAE). Clinical relapses were

defined as the appearance of a new neurologic deficit, or worsening of previously stable or

improving pre-existing neurologic deficit, separated by at least 30 days from the onset of a preceding

clinical demyelinating event.[26] The neurologic deficit must have been present for at least 24 hours

and occurred in the absence of fever (<37·5°C) or known infection.

hNSCs dosage and production

hNSCs were produced following a previously described method[27] and in full compliance with the

conditions and practices required by GMP regulations. The hNSCs consisted of a highly enriched

population of cells extracted from a single female foetal human donor (spontaneous miscarriage

16weeks after conception) under Ethics Committee approval, with the donors' parents' informed

consent, and according to the Helsinki declaration. hNSCs were cultured for 10-17 passages *in*

vitro.[27], the last passage was performed 24-96 hours before formulation of the final drug product.

On the day of treatment, hNSCs were collected from culture flasks, centrifuged counted and

suspended in HBSS at a concentration of 50,000 cells/μl. After batch release, the cells were

maintained at 4 +/-2 °C for up to 1·5 hours prior to implantation. See Profico et al 2022 for complete

quality control strategy for batch release.

Intervention

Trial participants were admitted to the Neurosurgery Unit of the "Azienda Ospedaliera Santa Maria"

(Terni, Italy) for the ICV injection of hNSCs. Before the intervention, they underwent thin-slice cranial

CT or MRI scans to evaluate the ventricular system and plan the procedure. Image guidance was

provided by the frameless stereotactic AxiEM system (Stealth station AxiEM electromagnetic

tracking system, Medtronic navigation, Louisville, CO, USA) to perform the ventricular cannulation.

The correct placement of the catheter was verified based on the egress of CSF. Then, a Rickham

reservoir was connected to the ventricular catheter. The participants were then prospectively

assigned to receive 5 (n=3), 10 (n=3), 16 (n=3), and 24 (n=6) x 10^6 cells ICV. All the participants

underwent a post-operative CT scan within 24 hours to evaluate any complications.

All the participants received methylprednisolone orally 125 mg 2 hours pre-intervention, and

cefazolin 1g IV immediately before and after the hNSCs injection. The immunosuppressive treatment

also included oral prednisone with a 28-day taper (consisting of a dose change per week, from 60,

40, 20, to 10 mg q.d). The participants also received Tacrolimus (0·05 mg/kg, oral, b.i.d.), 12 hours

after the intervention and then every 12 hours, for 6 months. This drug was titrated to maintain

blood levels ranging 5–10 ng/ml.

Data management and study monitoring

All trial demographic and clinical data was collected by designated investigators at the screening,

run-in, and follow-up visits using ad-hoc case report forms (CRFs). A clinical trial monitor periodically

reviewed the CRFs with source data verification and requested corrections as needed. The data were

then entered into a eCRF and underwent systematic quality control of the study manager prior to

database locking. The database was periodically reviewed by an independent Data and Safety

Monitoring Board (DSMB) to ensure protocol adherence and to monitor possible AEs. The DSMB

made recommendations concerning the continuation, modification, or termination of the trial as

needed.

Statistical Methods

Demographic, clinical, and laboratory patients' characteristics were reported as median and

interquartile ranges (IQR) or as mean and standard deviation (SD), as appropriate, for continuous

variables and as frequency and percentage for categorical variables. Normal distribution was

checked using the Shapiro-Wilk test. Safety analyses were performed in all subjects receiving at least

one injection of hNSCs. All AEs and severe AEs (SAEs) were recorded at follow-up visits and at the

end of the study. Exploratory efficacy analyses were conducted fashion in all subjects receiving at

least one injection of hNSCs (here, FAS and ITT populations matched). Because of small sample size

and of almost all variables having non-normal distributions, pre-post differences (12-month visit vs

run-in end visit) were assessed via linear models using ranks. New or enlarging T2-visible lesions and

lesions with contrast enhancement were analysed using negative binomial models with follow-up

time as the offset and results reported as annualised rates.

A p-value <0.05 was considered statistically significant. All analyses have been performed using SAS

Release 9.4, SAS Institute, Cary, NC, USA.

RESULTS

Clinical Results

A total of 220 candidates applied to participate in this study between September 26, 2017, and

January 13, 2020. After undergoing the screening visit and completing the run-in period, fifteen

SPMS patients were enrolled in the study. The participants had a median age of 49.8 years (range:

37.8–56.6), eight of them were females and seven were males (1.14 : 1 ratio), and an almost equal

proportion was recruited at the two study sites. The mean EDSS was 7.6 (range 7–8), mean disease

duration was 22 years (range 16–29), and mean time from diagnosis to progression was 10.1 years

(range: 1–20). All the patients' detailed demographic and clinical characteristics are reported in

Table 2.

Table 2: Demographic data and group assignment.

DOSE	Gender	EDSS	AGE AT TREATMENT (range)
$5*10^6$	M	7,5	55-59
	M	7	50-54
	M	8	50-54
$10*10^6$	M	7,5	36-39
	F	7	50-54
	F	8	50-54
$16*10^6$	F	7	50-54
	F	8	50-54
	F	8	45-49
$24*10^6$	F	7,5	50-54
	F	8	45-49
	M	7	40-45
	F	7	55-59
	M	8	36-39
	F	8	50-54

262	No deaths nor serious ARs were reported during the study period. The AEs that occurred, including

263	their severity, relationship to study and non-study therapies, actions taken, and outcomes are

264	reported in **Table 3**.

266   Table 3: Adverse events

Patient	Description of event	expected?	Severity	Relationship to study therapy	Action taken	If other, please specify	New treatment/therapy given/taken?	Outcome (at the end of the study)
#1	Tremor	Yes	Mild	Unrelated	Tacrolimus dose changed		No	Recovered
	Flu like syndrome	No	Mild	Unrelated	None		Yes	Recovered
	Respiratory failure	Yes	Mild	Unrelated	Concomitant therapy changed		Yes	Recovered
	Cataract (left eye; surgical procedure)	No	Mild	Unrelated	None		No	Recovered
	Upper respiratory infection	No	Mild	Unrelated	Concomitant therapy changed		Yes	Recovered
#2	Hyperglycemia	Yes	Moderate	Unrelated	Concomitant therapy changed		No	Recovered
	Urinary retention	Yes	Moderate	Unrelated	Other, specify	Catheterization	Yes	Recovering (improving)
	Back pain	No	Mild	Unrelated	Other, specify	New therapy given	Yes	Recovered
#4	Seizure	Yes	Mild	Possible	Other, specify	Diazepam, levetiracetam	Yes	Recovered
#7	Psychosis	No	Moderate	Unrelated	Concomitant therapy changed		Yes	Recovered
#8	Fracture (right femur)	No	Moderate	Unrelated	Other, specify	Follow-up 10 omitted	No	Recovered with sequelae
#9	Leukoencephalopathy	No	Mild	Unrelated	Other, specify	MRI evaluations	No	Recovering (improving)
	Urinary tract infection	No	Mild	Unrelated	Other, specify	Drugs administration	Yes	Recovered
#12	Depression	No	Moderate	Unrelated	None		No	Unknown
#14	Urinary tract infection	No	Mild	Unrelated	Other, specify	Antibiotics administration	Yes	Not recovered

Among these, SAEs (defined as AEs requiring hospitalisation) occurred in 2 patients (2/15), but none

of them were related to the hNSC injection: of these two, one patient developed a steroid-induced

acute psychiatric disorder 1 month post-injection, but recovered completely within 1 month with the

administration of valproate, lorazepam, olanzapine, and psychotherapy; Another patient

experienced femur fracture during a physiotherapy session at 18 months post-injection.

Other AEs were classified as non-serious and only one was possibly related to the study. This was a

patient (1/15) who experienced a first-ever partial motor seizure at month 6 (expected, possibly

related to MS). As per the possible complications related to the long-term immunosuppressive

treatment, one patient (1/15) developed a tremor during treatment with Tacrolimus which

disappeared with dose-adjustment, one patient (1/15) developed respiratory failure, and infections

were detected in three patients (3/15) (1 upper respiratory tract, 2 urinary tract). All the other

reported AEs were related to non-study concomitant therapy or other medical conditions not

related to the experimental protocol.   The immunosuppressive treatment was successfully

completed by patients. Tacrolimus was well tolerated by all patients except for the one already

mentioned. In all patients, Tacrolimus blood levels were within the therapeutic target range (below

20 ng/mL).

As per the secondary objectives of the study, no changes were measured in EDSS and MSFC for the

whole length of the study (**Table 4**).

Table 4. EDSS and MSFC evaluation

	hNSC Dose		Run-in Phase Onset (N=15)	Run-in Phase End (N=15)	1 month (N=15)	3 months (N=15)	6 months (N=15)	9 months (N=15)	12 months (N=15)	p value
289 290	5 millions	MSFC (Total Score)								0.753
291		Median	-5.5	-5.5	-5.6	-5.4	-5.6	-5.6	-5.3	
292		Nobs	3	3	3	3	3	3	3	
293		EDSS								0.114
294		Median	7.5	7.5	7.5	7.5	7.5	7.5	7.0	
295		Nobs	3	3	3	3	3	3	3	
296 297	10 millions	MSFC (Total Score)								0.150
298		Median	-6.2	-6.0	-6.0	-6.0	-5.9	-5.9	-5.9	
299		Nobs	3	3	3	3	3	3	3	
300		EDSS								1.000
301		Median	7.5	7.5	7.5	7.5	7.5	7.5	7.5	
302		Nobs	3	3	3	3	3	3	3	
303 304	16 millions	MSFC (Total Score)								0.924
305		Median	-6.0	-5.7	-6.2	-6.1	-5.8	-5.8	-5.6	
306		Nobs	3	3	3	3	3	3	2	
307		EDSS								1.000
308		Median	8.0	8.0	8.0	8.0	8.0	8.0	8.0	
309		Nobs	3	3	3	3	3	3	3	
310 311	24 millions	MSFC (Total Score)								0.772
312		Median	-6.1	-6.1	-6.0	-6.2	-5.9	-6.1	-6.2	
313		Nobs	6	6	6	5	3	4	4	
314		EDSS								0.831
315		Median	7.8	7.8	7.8	7.8	7.8	7.8	7.8	
316		Nobs	6	6	6	6	6	6	6	

319 EDSS: Expanded Disability Status Scale; MSFC: Multiple Sclerosis Functional Composite; Std.dev: Standard Deviation; IQR: Interquartile Range (i.e. first, third quartiles)

Test for linear trend from Run-In End to the end of the study: p-values from linear mixed-effects models

The median EDSS score did not change from the end of the run-in phase to the end of the study. Two

patients (2/15) had a change in the functional systems score of >1 point, both in the pyramidal area:

one decreased from 4.5 to 3.0 and another one increased from 0.0 to 2.0. The MSFC scores also did

not significantly change from the end of the run-in phase to the end of the study.

None of the patients reported symptoms indicative of clinically active disease and cognitive

functions, as measured by Rao's BRB, did not show significant changes during the study period for

any test. Neurophysiological parameters were monitored with EPs. Linear model analyses on ranks

did not show any variation trends throughout the study for any visual, somatosensory, and motor

EPs. We did not observe any changes on the OCT, except for one patient that showed an increase in

retinal nerve fibre layer's (RNFL) thickness in both eyes at month 6, which was interpreted as an

artefact.

As part of the secondary objectives, of the 40 scheduled MRIs for the 15 patients, 25 were

completed according to the study protocol and 15 were not performed (11 due to patient refusal,

two due to patient inability, two due to the COVID-19 pandemic). One patient did not have a

baseline MRI and post-Gadolinium images were not available for 8 MRIs due to exam interruption.

Brain changes on MRI were classified as type 1, 2, and 3 to facilitate comparison between time

points.

Type 1 changes were likely related to the surgical procedure. They included a linear T2-

hyperintensity in the parenchyma beneath the surgical right frontal cranial hole, passing through the

right frontal lobe white matter to the homolateral ventricle and corpus callosum (**Figure 1**).x

Figure 1. Example of type 1 change, interpreted as parenchymal gliotic modifications following
ventricular cannulation. Coronal-oblique 3D-T2-FLAIR subtraction map in A (Month 12 minus Run-in
End) detected a linearly-shaped positive signal change, passing through the right frontal lobe to the
homolateral ventricle. This change corresponded to a subtle contrast-enhancing tract which
occurred at Month 1 (B, post-Gadolinium T1-weighted image) at the level of the surgical cranial hole
(circle in B). Month 6 (C) and Month 12 (D) T2-FLAIR images demonstrated the persisting chronic
changes (arrows).

This was variable across all participants, ranging from a very subtle to a manifest signal change, and

it was detectable in all patients on Month 1 images, together with a linear contrast enhancement on

the corresponding post-Gadolinium T1-weighted images in most of cases.

Type 2 changes were MS-related and were used to compare the pre-transplantation with the post-

transplantation period.

Type 3 changes were undetermined and detected in only one participant, where subtraction images

highlighted a triangle-shaped, subtle T2-hyperintense signal change in the left frontal lobe white

matter, without contrast enhancement or water diffusion restriction (**Figure 2**).

Figure 2– Type 3 changes. Coronal-oblique 3D-T2-FLAIR subtraction map (Month 6 minus Run-in-End) detects a triangle-shaped, positive signal change in the left frontal white matter (arrows). This corresponded to a subtle T2-FLAIR signal hyperintensity on the corresponding Month 6 source image (B, arrows). No contrast enhancement and/or water diffusivity restriction was noted (not shown).

This change remained stable at the subsequent follow-ups and was categorised as non-specific. This patient had additional MRI follow-up during which the lesion did not change.

Pre-transplantation (during the three-month run-in period), nine new-onset and two enlarging T2-visibile lesions were detected in 7/14 patients (50% of patients, with an average of 1.3 and 0.3 of new and enlarging T2 lesions per patient) (**Table 5**).

372 Table 5 Summary of pre- and post-transplantation new-onset and enlarging T2-visible lesions
373 obtained by using subtraction imaging (see main text for details). N/A = data not available *When
374 the Month12 time-point data was not available, the last previous available data was employed for
375 subtraction

Subject #		Pre-Transplantation		Post-Transplantation		
	Compared Time Points	Number of New (Enlarging) T2-visibile lesions	Monthly Lesion Activity Rate	Compared Time Points*	Number of New (Enlarging) T2-visibile lesions	Monthly Lesion Activity Rate
#1	Run In End – Run In Onset	0	0.00	Month12 – Run In End	8	0.67
#2		0	0.00	Month12 - Run In End	1	0.08
#3		0	0.00	Month12 - Run In End	0	0.00
#4		2	0.67	Month12 - Run In End	6	0.50
#5		1	0.33	Month12 - Run In End	0	0.00
#6		0	0.00	Month12 - Run In End	1	0.08
#7		0	0.00	Month12 - Run In End	0	0.00
#8		0	0.00	Month 6 - Run In End	0 (1)	0.17
#9		0 (2)	0.67	Month12 - Run In End	8	0.67
#10		3	1	Month12 - Run In End	26	2.17
#11		1	0.33	Month12 - Run In End	3	0.25
#12		1	0.33	Month 3 - Run In End	0	0.00
#13		0	0.00	Month 9 - Run In End	0	0.00
#14		N/A	N/A	Month12 - Run In End	6	0.50
#15		1	0.33	Month12 - Run In End	5 (2)	0.58
Total		**9 (2)**			**64 (3)**	

After transplantation (during the 12-month follow-up period), 64 new-onset and three enlarging T2-

visible lesions were detected in 10 out of 15 patients (66% of patients, with an average of 6.4 and 0.3

of new and enlarging T2 lesions per patient). The annualised pre-transplantation rate of new or

enlarging T2-visibile lesions was 3.30 (95% CI: 1·86–5·83), which was not statistically different (p-

value= 0·26) to the post-transplantation rate (4·72, 95% CI: 2·32–9·61). When the 12-month data

was not available, the last previously available data was used to compare).

Pre-transplantation, 10 lesions with contrast enhancement were detected in 6/14 patients (43%)

(**Table 6**).

Subject #	Number of lesions with contrast enhancement (Time point)	Number of lesions with contrast enhancement (Time point)
#1	0	1 (Month6)
#2	0	0
#3	0	0
#4	2 (Run in Onset)	3 (Month3), 2 (Month6)
#5	1 (Run in End)	0
#6	0	0
#7	0	0
#8	0	1 (Month3)
#9	0	1 (Month6)
#10	1 (Run in Onset), 1 (Run in End)	9 (Month3), 12 (Month 6), 4 (Month12)
#11	1 (Run in End)	1 (Month9)

387   Table 6 Summary of pre- and post-transplantation lesions showing contrast enhancement on T1-

388   weighted images, detected upon visual inspection. N/A = data not available.

#12	1 (Run in Onset), 1 (Run in End)	0
#13	0	0
#14	N/A	1 (Month12), 1(Month6)
#15	1 (Run in Onset), 1 (Run in End)	4 (Month3)
Total	**5 (Run in End), 5 (Run in Onset)**	**40**

After transplantation, 40 gadolinium enhancing lesions were detected in multiple scans from 8 out

of 15 patients. The annualised pre-transplantation rate of lesions with contrast enhancement was

2·86 (95% CI: 1·51–5·40) and it was not statistically different (p-value=0·98) to the post-

transplantation values (2·83, 95% CI: 0·92–8·73).

Of note, 40% of the new-onset T2 lesions and the majority of gadolinium enhancing lesions

happened in one single patient who had the highest disease activity in the run-in period.

Additionally, one patient who showed one new T2 lesion and two lesions with contrast

enhancement in the pre-transplantation period, had no new T2 or gadolinium enhancing lesions at

follow-up.

Among the laboratory exams performed during the study, the only clinically significant variation was

in the number of white blood cells that ranged from a mean of 7·4 x10^9/L at the end of the run-in

period to a mean of 8·3 x10^9/L at the 12-month follow-up. The vital signs of the participants (blood

pressure and temperature) were within the normal range and urine tests did not show clinically

significant abnormalities. Biomarker analyses of NfL and CHI3L1/YKL-40 showed a statistically

significant increase in the NfL concentration (pg/mL) at 1 month after intervention in both serum

and CSF, followed by a gradual return to pre-intervention values (**Table 7**).

Table 7 Estimated marginal means, along with 95% confidence interval, from repeated measurements ANOVA models of serum neurofilament light
concentrations (pg/mL) collected in NSC-SPMS patients at different groups and time points. P-values are referred to within-group and within-time
comparisons and were derived from statistical contrasts defined into each model

Biomarker	hNSC assigned dose groups	Run-In onset (RSTART)	Run-In end (REND)	1 month (T1)	3 months (T3)	6 months (T6)	9 months (T9)	12 months (T12)	Within-group comparisons (p-values)						
									RSTART vs. REND	T1 vs. REND	T3 vs. REND	T6 vs. REND	T9 vs. REND	T12 vs. REND	p-value for trend
NfL CRMs	5 millions	13.84 (7.85-24.39)	11.33 (6.43-19.97)	31.61 (17.94-55.70)	29.87 (16.95-52.63)	20.62 (11.70-36.33)	12.93 (7.34-22.78)	13.21 (7.50-23.28)	0.552	0.003	0.005	0.079	0.695	0.648	0.664
	10 millions	14.59 (8.28-25.71)	16.91 (9.60-29.80)	45.64 (25.90-80.43)	21.78 (12.36-38.38)	18.21 (10.33-32.08)	17.17 (9.74-30.26)	19.68 (11.17-34.68)	0.660	0.004	0.452	0.826	0.964	0.652	0.825
	16 millions	11.80 (6.70-20.79)	12.33 (7.00-21.73)	98.03 (50.35-190.86)	27.94 (15.85-49.23)	23.07 (13.09-40.65)	14.33 (8.13-25.25)	14.84 (7.62-28.90)	0.896	<0.001	0.018	0.066	0.654	0.626	0.491
	24 millions	18.70 (12.53-27.91)	14.90 (9.98-22.24)	41.41 (27.74-61.81)	26.05 (16.91-40.12)	19.49 (11.41-33.29)	16.18 (8.51-30.79)	14.32 (8.91-23.00)	0.340	<0.001	0.030	0.370	0.814	0.882	0.540
	Overall	14.53 (11.15-18.94)	13.70 (10.51-17.86)	49.19 (37.21-65.03)	26.23 (20.06-34.29)	20.27 (15.33-26.80)	15.06 (11.23-20.20)	15.33 (11.52-20.41)	0.708	<0.001	<0.001	0.019	0.576	0.499	0.078
	Within-time comparisons (p-values)	0.594	0.743	0.090	0.881	0.948	0.905	0.778							
GFAP	5 millions	77.40 (45.16-132.65)	64.20 (37.46-110.02)	79.41 (46.33-136.08)	96.66 (56.40-165.65)	83.13 (48.51-142.47)	71.45 (41.69-122.45)	81.26 (47.41-139.26)	0.259	0.200	0.016	0.121	0.516	0.156	0.846
	10 millions	169.04 (98.64-289.70)	144.04 (84.05-246.85)	184.13 (107.44-315.55)	144.11 (84.09-246.97)	173.45 (101.21-297.26)	191.89 (111.97-328.85)	187.90 (109.64-322.01)	0.333	0.140	0.998	0.262	0.086	0.111	0.469
	16 millions	148.40 (86.59-254.32)	179.56 (104.77-307.73)	222.80 (126.50-392.41)	204.55 (119.35-350.55)	176.34 (102.89-302.20)	197.01 (114.96-337.64)	180.73 (102.61-318.32)	0.250	0.252	0.430	0.912	0.574	0.972	0.957
	24 millions	92.62 (63.28-135.57)	97.47 (66.60-142.67)	110.93 (75.79-162.36)	124.85 (84.58-184.30)	117.19 (77.00-178.37)	110.82 (70.20-174.92)	112.42 (75.25-167.96)	0.661	0.269	0.049	0.215	0.461	0.287	0.301
	Overall	115.80 (90.01-148.99)	112.79 (87.67-145.11)	137.87 (106.77-178.04)	137.34 (106.66-176.83)	131.39 (101.73-169.68)	131.53 (101.45-170.54)	132.72 (102.57-171.71)	0.732	0.015	0.014	0.061	0.070	0.051	0.056
	Within-time comparisons (p-values)	0.155	0.077	0.063	0.303	0.200	0.051	0.131							

410 The same pattern was evident across the four dose groups. These changes are likely due to post-procedure tissue damage that was detectable in all the
411 participants on MRI imaging. On the contrary, GFAP concentration did not show any peculiar pattern and remained stable throughout the trial for all the
412 dose groups (Table 7).

DISCUSSION

The ICV injection of hNSCs was safe and did not lead to clinically relevant AEs post-intervention. The only SAEs that occurred were not related to hNSCs or the intervention, and no withdrawals or deaths were registered. Importantly, none of the patients has shown either a clinical relapse or a progression of their condition during the study period. The follow-up disability evaluation scales and cognitive testing documented clinical stability of all the participants. The neurophysiologic parameters and OCT also showed no significant variations. The MRI disease activity, assessed by annualised rate of new or enlarging T2-visible lesions and lesions with contrast enhancement, was not affected by the treatment. Finally, despite an initial fluctuation of the levels of NfL and CHI3L1/YKL-40, these markers later returned to their baseline values.

Currently, there is a lack of treatment options for SPMS. Although another study using a similar NSC preparation and method of administration is underway in another Italian centre (NCT03269071), to our knowledge this manuscript represents the first report of the use of hNSCs in SPMS patients. In addition, this study utilises a cell line that has been already shown safe in a prior clinical trial on ALS (Mazzini et al., 2015, 2019).

This study has limitations inherent to its early phase, non-randomised design, and small sample. Also, the fact that some MRIs could not be performed could have weakened the imaging results. Nevertheless, this research provides novel information on the safety and potential effectiveness of this treatment modality for SPMS. It also describes a system by which, owed to the peculiar expansion technique adopted for cGMP expansion (vescovi et al, 1997), the very same hNSCs from a single donor used here, can be used in a broad number of future clinical trials, thereby obviating to the current outstanding issues of inter-trial variability of cell drug products.

It is worth noting that, unlike other clinical trials, the cells used in this study are hNSCs that grow as stable, reproducible unmodified cell lines (**Table 8**)

Table 8: Comparison between primary culture (2014) and final product release test (2018 –
2020). Data for final product are expressed as mean value of the release tests conducted on
all the batches used for the clinical trial. See Profico et al 2022 for complete quality control
strategy. IP: intermediate product, FP: final product, CE: Clonal Efficiency; N: Neurons; O:
Oligodendrocytes; A: Astrocytes; GC: Growth Curve, expressed as the slope of the growth
curve..
445

	IP	FP
CE	1,70%	1,45%
N	27,90%	31,50%
A	36,10%	50,50%
O	17,70%	18,70%
GC	0,076	0,057

This is crucial in view of the emerging concept that one key issue in cell therapy is the inter-trial

standardization of the cell drug, which may be homogeneous within the same clinical protocol but

does vary significantly between different trials, due to the scarcity of the donor cells. The approach

used here provides the significant advantage that the very same cGMP hNSCs used here - with all

the patients receiving the same cell drug - will also be available for a number of future trials both on

SPMS and other disorders. To the best of our knowledge, this is the first report of such an approach

being possible.

In conclusion, NSC transplantation via ICV injection appears to be a safe procedure with neither

major nor short-term deleterious effects. The study participants experienced a substantial clinical

stability during 12 months of follow-up. The considerable absence of risks for the patients indicates a

short-term neutral balance between benefits and risks and a therapeutic possibility on the horizon

for SPMS patients. Further studies are needed to confirm and extend the findings herein and

evaluate the actual therapeutic potential of advanced cell therapeutics for a condition where the

lack of effective disease modifying therapies is a major unmet clinical need.

DECLARATION OF INTERESTS

All of the authors declare no conflict of interest

AUTHOR CONTRIBUTIONS

All the authors reviewed and approved the final manuscript

ML	Clinical coordinating investigator, Patients recruitment, data analysis, manuscript writing and approval
CS	Clinical investigator, Patients recruitment, data analysis
DCP	Clinical protocol design, regulatory affairs and administrative management of the study. Biological fluids collection and management, biomarkers analysis, manuscript writing.
MG	Clinical protocol design, regulatory affairs and administrative management of the study. Drug product release, data collection and analysis, manuscript writing.
CC, MZ, LA, GP, CP, FA, PDV, DF, GS, ET, AC, SS, FP, GS, RFB	Clinical data collection and analysis
CG, EP, EZ	Clinical protocol drafting, clinical data collection and analysis
SP, LP	Data analysis, manuscript writing
GM, VG, IB, CR,	Drug product production and release, data collection and evaluation
DF	Non clinical testing, manuscript drafting and revision
ALV	Study coordinator, clinical protocol concept and writing, regulatory affair, provided funding and administrative support, manuscript writing and final approval
MC, AF	Study design, data collection, statistical analysis, manuscript writing
MC	Study management, administrative and regulatory support
JK, GD, LA	Biomarker analysis, data collection and evaluation

DATA SHARING STATEMENT

All relevant data are included in this manuscript. The data that support the findings of this study are

available from the corresponding authors upon reasonable request.

ACKNOWLEDGEMENTS

Writing assistance in the preparation of this article was provided by Maria Carolina Rojido (Medical

Writing Consultant). Proofreading and submission assistance was provided by Laura C Collada Ali

(Medical Writing Consultant).

FUNDING

Support for writing, proofreading and submission assistance was funded by IRCCS Casa Sollievo della

Sofferenza Research Hospital, Italy.

REFERENCES

1.   Browne P., Chandraratna D., Angood C., et al. Atlas of Multiple Sclerosis 2013: A growing global
problem with widespread inequity. Neurology. 2014;83(11):1022-4.
2.   Comi G. Disease-modifying treatments for progressive multiple sclerosis. Mult Scler. 2013;19(11):1428-
36.
3.   Giovannetti A.M., Pietrolongo E., Borreani C., et al. Conversion to secondary progressive multiple
sclerosis: Multistakeholder experiences and needs in Italy. PLOS ONE. 2020;15(2):e0228587.
4.   Alessandrini M., Preynat-Seauve O., De Bruin K., Pepper M.S. Stem cell therapy for neurological
disorders. S Afr Med J. 2019;109(8b):70-7.
5.   Lindvall O., Kokaia Z. Stem cells in human neurodegenerative disorders--time for clinical translation? J
Clin Invest. 2010;120(1):29-40.
6.   Vescovi A.L., Parati E.A., Gritti A., et al. Isolation and cloning of multipotential stem cells from the
embryonic human CNS and establishment of transplantable human neural stem cell lines by epigenetic
stimulation. Exp Neurol. 1999;156(1):71-83.
7.   Ferrari D., Zalfa C., Nodari L.R., et al. Differential pathotropism of non-immortalized and immortalized
human neural stem cell lines in a focal demyelination model. Cell Mol Life Sci. 2012;69(7):1193-210.
8.   Rota Nodari L., Ferrari D., Giani F., et al. Long-term survival of human neural stem cells in the ischemic
rat brain upon transient immunosuppression. PLoS One. 2010;5(11):e14035.
9.   Pluchino S., Quattrini A., Brambilla E., et al. Injection of adult neurospheres induces recovery in a
chronic model of multiple sclerosis. Nature. 2003;422(6933):688-94.
10.   Pluchino S., Gritti A., Blezer E., et al. Human neural stem cells ameliorate autoimmune
encephalomyelitis in non-human primates. Ann Neurol. 2009;66(3):343-54.
11.   Cossetti C., Alfaro-Cervello C., Donegà M., Tyzack G., Pluchino S. New perspectives of tissue remodelling
with neural stem and progenitor cell-based therapies. Cell Tissue Res. 2012;349(1):321-9.
12.   Giusto E., Donegà M., Cossetti C., Pluchino S. Neuro-immune interactions of neural stem cell
transplants: from animal disease models to human trials. Exp Neurol. 2014;260:19-32.
13.   Pluchino S., Cossetti C. How stem cells speak with host immune cells in inflammatory brain diseases.
Glia. 2013;61(9):1379-401.
14.   Peruzzotti-Jametti L., Bernstock J.D., Vicario N., et al. Macrophage-Derived Extracellular Succinate
Licenses Neural Stem Cells to Suppress Chronic Neuroinflammation. Cell Stem Cell. 2018;22(3):355-68.e13.
15.   Zhang C., Cao J., Li X., et al. Treatment of multiple sclerosis by transplantation of neural stem cells
derived from induced pluripotent stem cells. Sci China Life Sci. 2016;59(9):950-7.
16.   Takahashi Y., Tsuji O., Kumagai G., et al. Comparative study of methods for administering neural
stem/progenitor cells to treat spinal cord injury in mice. Cell Transplant. 2011;20(5):727-39.
17.   Ben-Hur T., Fainstein N., Nishri Y. Cell-based reparative therapies for multiple sclerosis. Curr Neurol
Neurosci Rep. 2013;13(11):397.

18. Peruzzotti-Jametti L., Pluchino S. Therapy with mesenchymal stem cell transplantation in multiple
sclerosis ready for prime time: Commentary. Mult Scler. 2022;28(9):1328-9.
19. Jin K., Sun Y., Xie L., et al. Comparison of ischemia-directed migration of neural precursor cells after
intrastriatal, intraventricular, or intravenous transplantation in the rat. Neurobiol Dis. 2005;18(2):366-74.
20. Morgenstern P.F., Connors S., Reiner A.S., Greenfield J.P. Image Guidance for Placement of Ommaya
Reservoirs: Comparison of Fluoroscopy and Frameless Stereotactic Navigation in 145 Patients. World
Neurosurg. 2016;93:154-8.
21. Van Damme P., Robberecht W. Developments in treatments for amyotrophic lateral sclerosis via
intracerebroventricular or intrathecal delivery. Expert Opin Investig Drugs. 2014;23(7):955-63.
22. Xu X., Chen W., Zhu W., et al. Adeno-associated virus (AAV)-based gene therapy for glioblastoma.
Cancer Cell Int. 2021;21(1):76.
23. Kurtzke J.F. Rating neurologic impairment in multiple sclerosis: an expanded disability status scale
(EDSS). Neurology. 1983;33(11):1444-52.
24. Fisher D., Beutler L.E., Williams O.B. Making assessment relevant to treatment planning: the STS
Clinician Rating Form. Systemic Treatment Selection. J Clin Psychol. 1999;55(7):825-42.
25. Burman J., Raininko R., Blennow K., Zetterberg H., Axelsson M., Malmeström C. YKL-40 is a CSF
biomarker of intrathecal inflammation in secondary progressive multiple sclerosis. J Neuroimmunol.
2016;292:52-7.
26. McDonald W.I., Compston A., Edan G., et al. Recommended diagnostic criteria for multiple sclerosis:
guidelines from the International Panel on the diagnosis of multiple sclerosis. Ann Neurol. 2001;50(1):121-7.
27. Mazzini L., Gelati M., Profico D.C., et al. Human neural stem cell transplantation in ALS: initial results
from a phase I trial. J Transl Med. 2015;13:17.

Solari, A., G. Filippini, L. Mendozzi, A. Ghezzi, S. Cifani, E. Barbieri, S. Baldini, A. Salmaggi, L. L. Mantia,M.
Farinotti, D. Caputo and P. Mosconi (1999). "Validation of Italian multiple sclerosis quality of life 54
questionnaire." J Neurol Neurosurg Psychiatry 67(2): 158-162.

Profico, D.C.; Gelati, M.; Ferrari, D.; Sgaravizzi, G.; Ricciolini, C.; Projetti Pensi, M.; Muzi, G.; Cajola, L.; Copetti,
M.; Ciusani, E.; et al. Human Neural Stem Cell-Based Drug Product: Clinical and Nonclinical Characterization.
Int. J. Mol. Sci. 2022, 23, 13425. https://doi.org/10.3390/ijms232113425

Mazzini L, Gelati M, Profico DC, Sorarù G, Ferrari D, Copetti M, Muzi G, Ricciolini C, Carletti S, Giorgi C, Spera C,
Frondizi D, Masiero S, Stecco A, Cisari C, Bersano E, De Marchi F, Sarnelli MF, Querin G, Cantello R, Petruzzelli
F, Maglione A, Zalfa C, Binda E, Visioli A, Trombetta D, Torres B, Bernardini L, Gaiani A, Massara M, Paolucci S,
Boulis NM, Vescovi AL; ALS-NSCs Trial Study Group. Results from Phase I Clinical Trial with Intraspinal Injection
of Neural Stem Cells in Amyotrophic Lateral Sclerosis: A Long-Term Outcome. Stem Cells Transl Med. 2019
Sep;8(9):887-897. doi: 10.1002/sctm.18-0154. Epub 2019 May 18. PMID: 31104357; PMCID: PMC6708070.