

1 ***Factors influencing the use of multiple HIV prevention services among Transport workers in a***
2 ***City in Southwestern Uganda.***

3 Benjamin Betunga^a, Phionah Atuhaire^a, Catherine Nakasiita^a, Christa Kanyamuneza^a, Proscovia
4 Namiiro^a, Joseph Tugume^a, Matovu Hairat^a, Ahmed M Sarki^{b,g}, Benedicto Mugabi^c, Birungi Lilian^f,
5 Richard Mugisha^d, Edward Kumakech^e, John Baptist Asiimwe^b

6 ^aFaculty of Nursing and Health Sciences, Bishop Stuart University, P.O.BOX 09, Mbarara,
7 Uganda

8 ^bAga Khan University, Uganda Campus. P. O. Box 8842, Kampala, Uganda

9 ^cBaylor College of Medicine Children's Hospital, PO Box 72052, Kampala, Uganda

10 ^dMbarara University of Science and Technology P.O Box 1410, Mbarara, Uganda

11 ^eFaculty of Nursing and Midwifery Lira University, P.O. Box 1035 Lira, Uganda.

12 ^fCollege of Nursing, Sultan Qaboos University, P.O Box 50, Postal code 123, Muscat, Oman.

13 ^gFamily and Youth Health Initiative (FAYOHI), Dutse, Jigawa State, Nigeria.

14 **Correspondence to john.asiimwe@aku.edu**

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31

32 **Abstract**

33 **Introduction:** Over the last three decades, there has been tremendous progress in curbing the HIV
34 epidemic in Uganda, although the HIV prevalence remains high among transport workers. Using
35 multiple HIV prevention services has been found to decrease the risk of acquiring HIV when
36 tailored to individuals at risk of HIV exposure, including transport workers. Therefore, we assessed
37 the uptake of multiple HIV prevention services (≥ 2) and associated factors among transport
38 workers in a city in Southwestern Uganda.

39 **Methods:** A cross-sectional study was conducted between November 2021 and February 2022.
40 Transport workers comprising motorcycle taxi riders, and motor vehicle and truck drivers, aged
41 18 to 55 years were selected and responded to an interviewer-administered questionnaire. We
42 conducted descriptive statistical and logistic regression analysis to identify independent factors
43 associated with the uptake of multiple HIV prevention services.

44 **Results:** Out of 420 participants, 97.6% were male, with a median age of 28 years and the majority
45 were aged < 34 years (84.6%). Overall, less than half (45.3%) of the participants had used multiple
46 (≥ 2) HIV prevention services within a one-year period. Most of the participants had used condoms
47 (32.2%) followed by voluntary HIV counseling and testing (27.1%), and safe male circumcision
48 (17.3%). Most participants who tested for HIV had ever used condoms (16.2%), followed by those
49 who received safe male circumcision and had ever used condoms (15%), and those who tested for
50 HIV and had started on antiretroviral therapy (ART) (9.1%). In the adjusted model, factors that
51 were significantly associated with the use of multiple HIV prevention services included religion
52 (AOR=2.69, 95% CI=1.38-5.23), the number of concurrent sexual partners (AOR=2.52, 95%
53 CI=1.45-4.37), prior HIV testing and awareness of HIV serostatus (AOR=0.23, 95% CI=0.13-
54 0.41), awareness of HIV prevention services (AOR=5.45, 95% CI=1.82-16.33), and financial
55 payment to access HIV services (AOR=4.80, 95% CI=2.33-9.88).

56 **Conclusions**

57 The uptake of multiple HIV prevention services among transport workers remains suboptimal.
58 This study suggests that individual behavioral factors influence the use of multiple HIV services
59 compared with other factors. Therefore, differentiated strategies are needed to increase the
60 utilization of HIV prevention services to curb the new HIV infections among transport workers.

61

62 **Keywords:** HIV prevention, transport workers, factors, determinants, association

63 **Introduction**

64 Over the last three decades, there has been tremendous progress in curbing the HIV epidemic in
65 Uganda. HIV prevalence has declined from 18% in the 1990s to 6.2% in 2016 and 5.4% in 2020
66 among 15–49-year-olds[1-3]. Like in other sub-Saharan countries, the decline in the prevalence of
67 HIV is partly attributed to the use of HIV prevention services such as condom use, and
68 antiretroviral based prevention services [4].

69
70 One of the current drivers of the HIV epidemic that threatens to undermine the success story is the
71 high incidence of HIV among high-risk groups or key populations [2, 5]. Transport workers which
72 include motorcycle taxi riders, truck drivers, and motor vehicle taxi drivers as defined for this
73 study are among the high-risk groups for acquiring HIV [6-11]. HIV prevalence among transport
74 workers varies across countries. It ranged from 9.2 to 13.9% among truck drivers in Cambodia
75 [12]. In some sub-Saharan Africa countries, the prevalence exceeds the national average for HIV
76 prevalence[11]. As an example, the prevalence of HIV among truck drivers in South Africa and
77 motorcycle taxi riders in Mbarara city in south-western Uganda were 26% and 9.9%, much higher
78 than the national averages of 12% and 5.4%, respectively[13-15].

79
80 The high HIV prevalence among transport workers, particularly long-distance truck drivers' is
81 attributable to engaging in transactional sex with sex workers along their transit routes, risky sexual
82 behaviors such as having multiple concurrent sexual partners and alcoholism, lack of access to and
83 or poor utilization of HIV prevention services, low education, lack of knowledge, and awareness
84 of HIV prevention methods among other factors [10, 16, 17]. In fact, transport workers, especially

85 truck drivers make up to 70% of the clients of sex workers, and about 26% of the new HIV
86 infections in sub-Saharan Africa in 2021[15, 18, 19].

87

88 Given that HIV transmission is driven by multiple biomedical, behavioral, and structural factors,
89 the use of a single HIV prevention method by high-risk populations is unlikely to alter the HIV
90 epidemic trajectory, putting the attainment of the 2025 goal of less than 370, 000 annual new
91 infections worldwide in jeopardy[5, 18, 20, 21]. To maximize population-level effects and
92 potentially bring the HIV epidemic under control, countries (Uganda inclusive) have undertaken a
93 policy to provide HIV prevention services in series at health facilities and during community health
94 outreaches to enable multiple uptake of these services[2, 20, 22]. The outcomes of this policy shift
95 has been impressive so far. For instance, using multiple HIV prevention services was found to
96 decrease the risk of acquiring HIV when tailored to individuals at high risk of HIV exposure or
97 key populations. In Kenya, providing multiple HIV prevention services to key populations reduced
98 the incidence of new HIV infections by 44%, whereas in Thailand, Cambodia, and Vietnam, new
99 HIV infections reduced by over 60% between 2010 and 2020[18, 21]

100

101 Mechanisms behind the beneficial effect of the use of multiple HIV prevention services on HIV
102 infections are well known. The prompt use of multiple HIV prevention services such as voluntary
103 HIV counselling and testing (VCT) followed by antiretroviral therapy (ART) preferably within a
104 year of HIV diagnosis is critical in preventing delays in treatment initiation, reducing missed
105 opportunities for treatment, preventing HIV progression and ultimately reducing the HIV
106 transmission risk[23]. In fact, the prompt use of VCT and antiretroviral drugs (ARVs) for the
107 elimination of mother-to-child transmission of HIV (EMTCT) series of HIV services is critical in

108 the prevention of mother-to-child transmission of HIV (MTCT)[24]. Whereas the use of the other
109 series of HIV prevention services such as VCT and condom use or Preexposure prophylaxis (PrEP)
110 are critical in preventing new HIV infections among HIV sero-discordant partners[25]

111

112 Given the aforementioned significance, the WHO and ministries of health across sub-Saharan
113 Africa including Uganda have instituted policies that enable multiple uses of HIV prevention
114 services such as "test and treat" for HIV test-positive cases, "test, and condoms" or "test and PrEP"
115 for HIV sero-discordant couples, "test, and PEP (post-exposure prophylaxis)" for occupational or
116 accidental or forced exposure cases. There is also the policy of "test and ART" for EMTCT for
117 pregnant and breast-feeding women living with HIV[2, 20].

118

119 Despite the availability of policies and a multitude of HIV prevention services at various health
120 service delivery points in Uganda, the uptake of multiple HIV prevention services and factors
121 associated with its use, especially among transport workers, remains poorly documented. A few
122 previous studies such as the one conducted among men who have sex with men (MSM) in Brazil,
123 found low (2.8%) use of behavioral and biomedical HIV prevention services [26]. Additionally,
124 previous research findings from South Africa indicated suboptimal use of HIV prevention services
125 by transport workers, despite interventions by various stakeholders to increase transport workers'
126 access to the HIV services[7, 8, 11].

127

128 Individual factors such as sociodemographic characteristics, knowledge, and awareness about HIV
129 prevention, perceived risk of HIV infection, and HIV stigma among others, were reported to
130 predict the use of single HIV prevention services. In agreement, good knowledge about HIV

131 transmission, perceived high risk of acquiring HIV infection, awareness of HIV status, and having
132 a high level of education among others, were found to be associated with higher likelihood of using
133 either condoms or HIV testing services among truck drivers and motorcycle taxi riders in a multi-
134 country study which included Togo, Nigeria, and Uganda, respectively[17, 27, 28]. In this study,
135 we hypothesized these factors as determinants of the uptake of multiple HIV prevention services
136 among transport workers through similar mechanisms.

137 Therefore, we set out to investigate the influence of individual factors on the utilization of multiple
138 HIV prevention services among transport workers in southwestern Uganda. The findings of this
139 study will inform tailor-made interventions aimed at increasing access to and utilization of HIV
140 prevention services among transport workers in Uganda and other similar sub-Saharan Africa
141 settings.

142 **Materials and Methods**

143 **Study design and setting**

144 This study used a cross-sectional design. The study setting was the four slum areas of Mbarara city
145 in southwestern Uganda where the HIV prevalence was higher than the Uganda national average.
146 The slum areas (also known as hot spots) of Mbarara city included Kakoba, Kiyanja, Kizungu, and
147 Ruti. The area have many sex workers, motorcycle taxi riders, motor vehicle taxi drivers (who are
148 residents of the area and its neighborhood), and stopover stations for truck drivers on transit.
149 According to the 2020 estimates, Mbarara district has the third highest level of HIV prevalence
150 (13.1%) in Uganda when compared to the national average of 5.4% [2, 18]. Among a subset of
151 transport workers in this case motorcycle riders (also known as bodabodas) in a Mbarara city, a
152 previous study reported a 9.9% prevalence of HIV infections which was also above the national
153 average of 5.4%[13].

154

155 There are several health facilities in and around the four slum areas of Mbarara city that provide
156 all the recommended HIV prevention services including Kakoba Health Center, Nyamityobora
157 health center, Mbarara City Hospital, and Mbarara Regional Referral Hospital. Besides the health
158 facilities, there are other non-governmental organizations such as The AIDS Support Organization
159 (TASO) that also provide HIV prevention services at mobile community outreach events to people
160 living in Mbarara city's slum areas. Mbarara city is located in southwestern Uganda at a distance
161 of about 280 kilometers from Kampala the capital city of Uganda.

162 **Sample, sample size, and sampling method**

163 We included in the study male and female motorcycle taxi riders, and motor vehicle taxi and truck
164 drivers aged 18 and 55 years, who were residents and have operated their transport trade within
165 Mbarara city for at least one year preceding the study. We excluded transport workers who were
166 feeling sick or unwell and thus unable to withstand the study procedures. The initial study sample
167 size of 385 transport workers was obtained using the Cochrane (1963) formula[29]. For calculating
168 the sample size, the proportion of transport workers utilizing multiple HIV prevention services (p)
169 was assumed to be 50% ($p=0.5$), the Z score from standard normal curve corresponding with the
170 95% confidence interval, and the allowable error were set at 1.96 and 0.05 respectively. We
171 adjusted the initial sample size of 385 participants for 10% non-response rate to give the desired
172 sample size of 424 participants. From the register of transport workers obtained from their leaders,
173 a complete register was established. We performed simple random sampling to select the
174 participants for the study. Out of the 424 participants sampled and approached for informed
175 consent, 422 participants provided informed consent and participated in the study (a 99.5%

176 response rate). Only two participants declined to provide informed consent and participate in the
177 study because of their busy work schedules.

178 **Data collection procedure**

179 The data were collected between November 2021 and February 2022. We obtained the transport
180 workers' registers with their contact details from their leaders. Simple random sampling (the lottery
181 method with replacement) was used to select eligible participants. The leaders of the respective
182 transport workers then contacted the selected participants by phone calls and introduced the team
183 of researchers to them. This was followed by appointments to meet the study participants. When a
184 participant declined to participate in the study, two failed follow up attempts were made before a
185 replacement was sampled from the register to participate in the study. Prior to the commencement
186 of data collection, the principal investigator trained research assistants (who were Bachelor of
187 Nursing professionals) about the study protocol including the study tools. The research assistants
188 participated in pretesting of the questionnaires on 10 participants which included motorcycle taxi
189 riders, motor vehicle taxi, and truck drivers at another slum area known as Ruharo trading center
190 within Mbarara city. Adjustments to the study tools were made to assure validity before their use
191 in the main data collection.

192 At the respective stopovers or workplaces of transport workers, the research assistants identified
193 the selected participants, introduced, and informed them about the study, and requested their
194 participation in the study. Motorcycle riders who accepted to participate in study would then be
195 moved aside from the other members of the group for the informed consent, and data collection to
196 assure privacy and confidentiality. Tax drivers and truck drivers were interviewed in their vehicles
197 when they did not have customers to assure privacy and confidentiality. All data were collected in
198 the local languages (Runyakole-Rukiga) or English. Each data collection procedure took about 10-

199 15 minutes. After each data collection session, the research assistants checked the questionnaire
200 for completeness before leaving the participant and the field site. The two research assistants of
201 nursing educational background who worked in a pair collected all the data.

202 **Measurement of the use of multiple HIV prevention services**

203 Using multiple HIV prevention services was the primary outcome of interest for the study.
204 Participants were asked to mention the HIV prevention services they had ever used within the
205 previous 12 months. Using multiple HIV prevention services was indicated when a participant
206 used two or more of the recommended HIV prevention services within a 12 months period. The
207 HIV prevention services comprised evidence-informed high-impact biomedical and behavioral
208 HIV prevention services and methods such as ARVs for the EMTCT or ART, condoms, VCT,
209 health education on HIV, safe male circumcision (SMC), PrEP and PEP. Participants were also
210 asked to mention the source where the services were obtained from and how often they used the
211 HIV services within the 12 months period.

212 **Measurements of factors of interest**

213 We categorized the factors which were potential predictor variables into three namely the
214 participant's socio-demographic characteristics, knowledge, awareness and perceptions (KAP)
215 and behavioral factors among the study participants:

216 **1. Sociodemographic characteristics**

217 The participants were asked to state their status on nine items which were considered as measures
218 of the sociodemographic characteristics of the participants namely age, gender, marital status,
219 education level, income level, occupation, length of stay in the study area, distance to a nearby
220 health facility, and religion.

221 **2. Knowledge, awareness and perception factors among the participants'**

222 *a). Awareness of HIV prevention services*

223 Participants were asked to mention any biomedical or behavioral HIV prevention services they
224 were aware of and their source of information. Awareness was categorized into awareness of a
225 single (1) service and awareness of multiple HIV prevention services (2+) as measures of
226 awareness.

227 *b) Knowledge of HIV prevention*

228 Participant's knowledge about HIV prevention was assessed using a validated questionnaire
229 adapted from a previous study conducted in South Africa and Cambodia[30, 31]. The questionnaire
230 has twelve knowledge items about HIV prevention with answer options of "Yes", "No", or "Don't
231 Know". A score of 1 was awarded to participants who correctly answered a question, and 0 for
232 incorrect or don't know responses. The maximum attainable score was twelve (12) which was
233 further grouped into poor knowledge level (0-6 score), and good knowledge level (7-12 score).

234

235 *c) Perceived risk of HIV infection*

236 The perceived risk of HIV infection was measured using a validated perceived risk of HIV
237 infection questionnaire adapted from Napper et al., [32]. The questionnaire measures the cognitive
238 and affective aspects of the perceived risk of HIV infection namely how an individual
239 conceptualizes the risk of HIV infection, feels about it, and how often they think about it. The tool
240 has ten (10) risk statements with strongly agree to strongly disagree measured on a 5-point likert
241 scale. Participants were asked to provide their level of agreement with each of the statements
242 regarding the perceived risk of contracting HIV infection. The maximum attainable score was 43

243 points. The scores were further grouped into the low perceived risk of HIV infection (0-22.5) and
244 high perceived risk of HIV infection (22.6-43.0).

245 *d) Perceived HIV stigma*

246 Perceived HIV stigma was measured using a validated 11-itemed questionnaire which was
247 adapted from the questionnaire used to assess perceived personal and community HIV stigma in
248 South Africa [33]. On a 5-point Likert scale (strongly agree to strongly disagree), participants
249 were asked to provide their level of agreement with the 11 statements measuring stigma. The
250 maximum attainable score was forty-four (44) which was further grouped into low perceived HIV
251 stigma (0-22), and high perceived HIV stigma (23-44).

252 **3. Behavioral factors**

253 The participants were asked to state their status regarding the three (3) items which were
254 considered as measures of behaviors likely to influence the access and or utilization of HIV
255 prevention services. These behaviors included sexual behavior in particular the number of
256 concurrent sexual partners, HIV self-care behavior in particular prior HIV testing (as well as
257 knowing own HIV status) and willingness to pay or having ever paid to access or obtain HIV
258 prevention services.

259 **Reliability analyses**

260 Reliability analyses were conducted for knowledge of HIV prevention, perceived risk of HIV
261 infection, and HIV stigma subscales. Before reliability analyses, all negatively worded items plus
262 their respective scores within the perceived risk of HIV infection and perceived HIV stigma
263 subscales were reversed. Internal consistency of the subscales was evaluated using the Cronbach
264 alpha (α) coefficient. The Cronbach alpha (α) coefficient values of 0.7 or higher were considered
265 acceptable [34], see Table 1). Overall, the Cronbach alpha coefficient for the knowledge of HIV

266 prevention, perceived risk of HIV infection, and perceived HIV stigma subscales were 0.707, 0.88,
267 and 0.91, respectively, demonstrated all the three subscales were reliable at measuring the
268 construct they were meant to measure. Notably, both the mean inter-item correlation (MIIC) and

Table 1. Internal Consistency and Measurement Properties of Items

Subscales	Items	MIIC	Cronbach's α	NCITC>0.3
Knowledge of HIV prevention	12	0.162	0.707	7
Perceived risk of HIV*	8	0.497	0.88*	8
Perceived HIV Stigma	11	0.483	0.91	11

269 the number of items with corrected item-total correlation (NCITC, >0.3) were within acceptable
270 limits.

271
272 *Note: The coefficient * in the perceived risk of HIV subscale means two items were eliminated*
273 *from the scale to compute the coefficient as recommended by the authors of the tool (Napper et al.*
274 *[32]).*

275

276 Data Analysis

277 The data were entered and cleaned in EpiData 3.1 software and exported to the Statistical Package
278 for Social Sciences (SPSS) version 20 for analysis. The primary outcome variable (with response
279 score options of 1 or 0) was created for the purpose of logistic regression analysis. Descriptive
280 analyses were conducted for categorical and continuous variables. Categorical variables are
281 reported as frequency counts and percentages whereas continuous variables were summarized as
282 mean (when normally distributed) and median (when not normally distributed). Chi-square

283 statistics were used to establish the relationship between categorical factors and the use of multiple
284 HIV prevention services. Variables found statistically significantly associated with the use of
285 multiple HIV prevention services ($p < 0.05$) were entered into the binary logistic regression models
286 including demographic characteristics, KAP factors, and behavioral factors to obtain their
287 predictive and explanatory effects. The level of significance and adjusted odds ratios were reported
288 at 95% confidence intervals.

289

290 **Ethical consideration**

291 The Mbarara University of Science and Technology research ethics committee (MUREC
292 1/7/15/12-20) approved the study. Administrative permission to conduct the research was obtained
293 from the district health officer of the Mbarara district and the leaders of the transport workers.
294 Informed consent was obtained from all study participants after introducing the study to them using
295 the informed consent form. Using either a thumbprint (illiterate) or a signature confirmed consent.
296 For confidentiality, participants were interviewed away from the other transport workers in private
297 spaces. The consenting and interview processes were conducted in the local languages
298 (Runyakole-Rukiga) or English. Interviews were stopped, and participants were dropped and
299 replaced if they received a customer for their transportation trade during the study and this
300 happened with 26 participants. Research assistants and participants wore masks and disinfected
301 their hands with alcohol gel to avoid cross-infection with COVID-19. We paid time compensation
302 with an amount equivalent to USD 2 to every participant upon completion of the data collection
303 session. Copies of the completed questionnaires were kept in a lockable cabinet in the first author's
304 office whereas the electronic databases were password protected and only accessible to the
305 research team.

306 **Results**

307 **Demographic characteristics of the study participants**

308 The demographic characteristics of the study participants are summarized in Table 2. Out of the
309 420 participants interviewed (99.5% response rate), the majority were male (97.6%) and middle-
310 aged adults aged below 34 years, with a median age of 28 years (54.4%). About half of the study
311 participants had lived, temporarily stayed, or worked in Mbarara city for over 5 years (51.0%) with
312 a median of 6 years, in the same city. The majority were single (41.2%), Christians by religion
313 (78.6%), and had obtained a primary (38.5%) or ordinary-level certificate of education (39.5%).
314 Most participants were motorcycle taxi riders (boda-boda riders, 69.7%), followed by motor
315 vehicle taxi and truck drivers (25.9%). Most earned a monthly income of 400,001-500,000/= (~111
316 to 137 USD, 59.5%). Many of the study participants knew their current HIV serostatus (54.1%);
317 most were seronegative (43.9%). However, a large proportion of participants had multiple
318 concurrent sex partners (52.4%), with a median of 2 partners per participant. Last, most
319 participants lived, stayed, or worked within 1-3 kilometers (km) of the nearby health facility
320 (86.6%), with a median of 2 km.

321

322

323

324

325

326

327

328

329

330 **Table 2: Demographic Characteristic of the Participants**

Variable	n (%)
Gender	
Male	400 (97.6)
Female	10 (2.5)
Age in years	28 (18-48)*
18-24 yrs.	134 (30.2)
25-34 yrs.	223 (54.4)
≥ 35 yrs.	63 (15.4)
Length of stay in the area (yrs.)	6 (1-28)*
<5years	201 (49.0)
≥ 5year	209 (51.0)
Marital status	
Single	169 (41.2)
Married	154 (37.6)
Cohabiting	22 (5.4)
Separated/ divorced/ widowed	65 (15.9)
Religion	
Anglican	131 (32.0)
Catholic	118 (28.8)
Pentecost	73 (17.8)
Moslem	84 (20.5)
Others	4 (1.0)
Education level	
None	23 (5.6)
Primary	158 (38.5)
Ordinary level	162 (39.5)
Advanced level	56 (13.7)
Tertiary	11 (2.7)
Occupation	
Motorcycle rider (Boda Boda)	286 (69.7)
Motor vehicle driver/conductor	106 (25.9)
Truck driver/conductor	18 (4.4)
Monthly income	
82-110 USD	48 (11.7)
111-137USD	244 (59.5)
>137 USD	118 (28.8)
Serostatus (self-report)	
Positive	42 (10.2)
Negative	180 (43.9)
Unknown	188 (45.9)
Sex partners	2 (0-11)*
Single	195 (47.6)
Multiple	215 (52.4)
Distance to health facility/HIV care service provider	2 (1-10)*
1-3km	355 (86.6)
+3km	55 (13.4)

331 Key: *Median (Minimum-Maximum)

332 **The use of multiple HIV prevention services**

333 When participants were asked to disclose the number of HIV prevention services they had ever
334 used within the past 12 months, results indicate that less than half of the participants (45.3%), had
335 ever used two or more services (Table 3). Approximately one-third of the participants had used
336 condoms (32.2%) followed by voluntary counselling and testing (27.1%), and safe male
337 circumcision (17.3%).

338

339 Regarding multiple use of HIV prevention services, the findings show that most of the participants
340 had tested for HIV and then used condoms (16.2%), followed by those who had received safe male
341 circumcision and then used condoms (15%), and those who had tested for HIV and then began on
342 ART (9.1%). Most of the HIV prevention services were obtained from the public health facilities
343 and not for profit civil society organizations that support patients with HIV (83.3%) and were free
344 (82.8%).

345

346

347

348

349

350

351

352

353

354

355 **Table 3. Use of HIV prevention services among transport workers in Mbarara City,**
 356 **Southwestern Uganda (N=420)**

Variables	n (%)
Number of Used HIV prevention services	
1 or None	223 (54.7)
≥ 2 services	185 (45.3)
Use of a single HIV prevention service.	
VCT	171 (27.1)
Condom Use	203 (32.2)
SMC	109 (17.3)
PREP	10 (1.6)
EMTCT/ART	66 (10.5)
Others	43 (6.8)
Common combinations used	
VCT + ART/EMTCT	38 (9.1)
VCT + PrEP	06 (1.4)
VCT + condoms	68 (16.2)
VCT + SMC	36 (8.6)
SMC+ condoms	63 (15)
Sources of HIV prevention services	
Government	197 (56.8)
CSO and others	92 (26.5)
Private health facility	58 (16.7)
Pay for services	
Yes	58 (16.9)
No	280 (82.8)

357 **Note:** VCT is voluntary testing and counselling, SMC is safe male circumcision, PrEP is pre-
 358 exposure prophylaxis, EMTCT is the elimination of mother-to-child transmission of HIV, ART
 359 is Anti-retroviral therapy, and CSO is civil society organizations.

360 **Factors associated with the use of multiple HIV prevention services**

361 The results of the bivariate shown in table 4 indicate that participant’s sociodemographic
 362 characteristics namely age, education level, marital status, religion and years of residence in the
 363 study area were significantly associated with the utilization of multiple HIV prevention services.
 364 Similarly as shown in Table 4, the results indicate that awareness, knowledge, perceptions and
 365 behavioral factors were significantly associated with the utilization of multiple HIV prevention
 366 services.

367 The results of the univariate logistic regression table shown in table 4, participants who were older
368 (≥ 25 years) compared with the younger participants (< 25 years); participants who had multiple
369 concurrent sex partners compared with those with a single sex partner; participants who were in a
370 marital relationship compared with those not in a marital relationship; participants who subscribed
371 to Muslim or other religious faith compared with those who belonged to the Christian faith, and
372 participants who had attained the secondary and above level of education compared with those
373 who attained primary or no education were 2.278, 1.487, 1.637, 2.157, and 1.599 times more likely
374 to use multiple HIV prevention services, respectively.

375
376 In addition, participants who had worked for a longer period (> 5 years) around the town compared
377 with those who had worked for a short period (< 5 years), and participants who were aware of two
378 or more HIV prevention services compared with those who were aware of one, participants who
379 had good knowledge compared with those with poor knowledge about HIV prevention,
380 participants who perceived the risk of acquiring HIV to be high compared with those who
381 perceived it to be low, and participants who did not pay to access HIV prevention services
382 compared to those who paid, were 1.529, 8.600, 2.019, 3.179, and 4.365 times more likely to use
383 multiple HIV prevention services, respectively.

384
385 Contrariwise, participants who were not aware of their HIV status compared to those who knew it
386 and participants who had perceived high level of HIV stigma compared with a low perceived HIV
387 stigma were 0.196 and 0.494 times less likely to use multiple HIV prevention services,
388 respectively.

389

390 **Table 4. Factors associated with the Use of multiple HIV prevention services**

Variable	Use of multiple HIV Prevention services		Pearson chi-square	P-value	Crude odds ratio
	2 and + (n, %)	1 or None n (%)	Value (d.f)	Asymp. sig. (2-sided)	cOR (95% CI)
Overall	185 (45.3)	223 (54.7)			
Gender			0.90 (1)	0.765	1.00
Male	180 (44.1)	218 (53.4)			
Female	5 (1.2)	5 (1.2)			1.211 (0.345-4.249)
Occupation			0.999 (2)	0.607	1.00
Motorcycle taxi rider (Boda Boda)	125 (30.6)	160 (39.2)			
Motor vehicle Driver/conductor	52 (12.7)	53 (13.0)		0.319	1.256 (0.802-1.966)
Truck Driver/conductor	8 (2.0)	10 (2.5)		0.961	1.024 (0.393-2.671)
Age			15.349 (2)	0.000*	
18-24	38 (9.3)	85 (20.8)			1.00
25-34	112 (27.5)	110 (27.0)		0.001*	2.278 (1.432-3.623)
≥35	35 (8.6)	28 (6.9)		0.001*	2.796 (1.494-5.235)
Number of Sexual partners			3.938 (1)	0.047*	1.00
Single	78 (19.1)	116 (28.4)			
Multiple	107 (26.2)	107 (26.2)			1.487 (1.004-2.202)
Marital status			5.997 (1)	0.014*	1.00
Not in relationship	93 (22.8)	139 (4.1)			
In relationship	92 (22.5)	84 (20.6)			1.637 (1.102-2.431)
Religion			10.030 (1)	0.002*	1.00
Christians	132 (32.4)	188 (46.1)			
Moslem and Others	53 (13.0)	35 (6.1)			2.157 (1.333-3.490)
Income level			0.184 (1)	0.688	1.00
≤500,000/= (below average)	130 (31.9)	161 (39.5)			
≥500, 000/=	55 (13.5)	62 (15.2)			1.099 (0.714-1.690)
Education level			5.414 (1)	0.020*	1.00
≤ Primary	70 (17.2)	110 (27.0)			
≥Secondary	115 (28.2)	113 (27.7)			1.599 (1.076-2.377)
HIV status			58.601 (1)	0.000*	1.00
Known	139 (34.1)	83 (20.3)			
Unknown	46 (11.3)	140 (34.3)			0.196 (0.128-0.302)
Distance to facility			1.315(1)	0.251	1.00
1-3km	164 (40.2)	189 (46.3)			
>3km	21 (5.1)	34 (8.3)			0.712 (0.397-1.275)
Years of work in the area			4.519 (1)	0.034*	1.00
<5years	80 (19.6)	120 (29.4)			
>5year	105 (25.7)	103 (25.2)			1.529 (1.033-2.264)
HIV Services Awareness			26.776 (1)	0.000*	1.00
1 service	5 (1.2)	43 (10.5)			
2+ services	180 (44.1)	180 (44.1)			8.600 (3.330-22.211)
Knowledge about HIV prevention			10.028 (1)	0.002*	1.00
Poor	42 (10.3)	83 (20.3)			
Good	143 (35.0)	140 (34.3)			2.019 (1.302-3.129)
Perceived risk of HIV			21.290 (1)	0.000*	1.00
Low	25 (6.1)	74 (18.1)			
High	160 (39.2)	149 (36.5)			3.179 (1.918-5.269)
Perceived stigma			11.733 (1)	0.001*	1.00
Low	87 (21.3)	68 (16.7)			
High	98 (24.0)	155 (38.0)			0.494 (0.329-0.741)
Pay for HIV services			23.050 (1)	0.000*	1
Yes	43 (12.7)	15 (4.4)			
NO	111 (32.8)	169 (50)			4.365 (2.314-8.233)

391 *Note:* D.f is degree of freedom, COR is crude odds ratio

392

393 **Factors that independently Predict the use of multiple HIV** 394 **prevention services**

395 The factors that were significantly associated with the use of multiple HIV services ($p < 0.1$) from
396 the binary logistic regression were entered into three separate multivariate logistic regression
397 models (Table 5). The first model comprised sociodemographic factors such as age, education,
398 religion, marital status, and length of stay in the area. In this model, results showed that age
399 ($p = 0.019$), and religion ($p = 0.01$) independently predicted the use of multiple HIV prevention
400 services. Participants aged between 24-25 years were almost twice more likely to use multiple HIV
401 prevention services than those aged 18-24 years after controlling for other sociodemographic
402 factors. In addition, participants who were Muslim or other faith were also twice more likely to
403 use multiple HIV prevention services than those who subscribed to the Christian faith after
404 controlling for other sociodemographic factors.

405
406 In model 2, knowledge of HIV prevention, perceived risk of HIV, perceived HIV stigma, and
407 awareness of HIV prevention services were then added to model 1. The results from model 2
408 multivariate regression model in table 5 indicate that religion, perceived risk of HIV, and
409 awareness of HIV services independently predicted the use of multiple HIV prevention services.
410 Whereas age ($p = 0.086$) became statistically non-significant. Again, participants who subscribed
411 to the Muslim or other faith were almost twice more likely to use multiple HIV prevention services
412 than those who subscribed to the Christian faith. Participants who perceived the high risk of
413 acquiring HIV were twice more likely to use multiple HIV prevention services than those who
414 perceived low risk of HIV. Participants who were aware of two or more HIV prevention services

415 were almost seven times more likely to use multiple HIV prevention services than those who knew
416 about only one HIV prevention service.

417 In model 3, behavioral factors in this case the number of concurrent sex partners, HIV testing, or
418 awareness of HIV status, and paying for HIV prevention services were added into model 2. In
419 model 3, religion, awareness of HIV status, awareness of HIV prevention services, paying for HIV
420 prevention services, and the number of concurrent sexual partners independently predicted the use
421 of multiple HIV prevention services. In this model 3, participants who subscribed to the Muslim
422 or other faith were almost thrice more likely to use multiple HIV prevention services than those
423 who subscribed to the Christian faith. Participants who had two or more sexual partners were
424 almost thrice more likely to use multiple HIV prevention services than those with single or no
425 partners. Additinally, Participants who were aware of two or more HIV prevention services were
426 five times more likely to use multiple HIV prevention services than those who knew only one HIV
427 prevention service. Participants who never paid to access the HIV prevention services were almost
428 five times more likely to use HIV prevention services than those who paid to access the HIV
429 prevention services. However, participants who did not know their HIV serostatus were 0.228
430 times less likely to use HIV prevention services than those who knew their HIV serostatus.

431

432

433

434

435

436

437

438 **Table 5. Odds (and confidence interval) from binary logistic regressions of individual factors**
 439 **that predict the use of multiple HIV prevention services.**

Variable	aOR Exp (B) (95% CI)		
	Model 1	Model 2	Model 3
Age (years)			
18-24yrs (Ref.)	1.00	1.00	1.00
25-34yrs	1.853(1.106-3.105)*	1.618(0.786-3.332)	1.468(0.768-2.808)
>35YRS	1.802(0.842-3.860)	1.774(0.609-5.167)	1.365(0.523-3.563)
Length of stay			
5yrs &< (Ref.)	1.00	1.00	1.00
> 5 years	1.314(0.854-2.020)	1.2976(0.721-2.364)	1.226(0.717-2.099)
Religion			
Christian (Ref)	1.00	1.00	1.00
Muslim & others	1.99(1.171-3.147)*	1.924(1.969-3.819) *	2.685(1.379-5.226)*
Education			
Primary & none	1.00	1.00	1.00
Secondary & above	1.274(0.828-1.959)	0.665(0.481-1.683)	0.737(0.423-1.284)
Marital status			
Not in relationship	1.00	1.00	1.00
In relationship	1.125(0.709-1.783)	0.943(0.498-1.786)	0.832(0.459-1.507)
Awareness of HIV services			
Services	-	1.00	1.00
2 services and above	-	6.988(1.928-25.333)*	5.452(1.821-16.328)*
Knowledge about HIV			
Poor	-	1.00	1.00
Good	-	1.186(0.593-2.370)	1.148(0.614-2.145)
Perceived risk of HIV			
Low (Ref)	-	1.00	1.00
High	-	2.258(1.060-4.810)*	0.911(0.446-1.861)
Perceived stigma			
Low (Ref.)	-	1.00	1.00
High	-	0.688(0.375-1.262)	0.737(0.435-1.251)
Number of sexual partners			
Single or none (Ref)	-	-	1.00
2 & above	-	-	2.518(1.453-4.365)*
Multiple	-	-	-
HIV status			
Known (Ref)	-	-	1.00
Unknown	-	-	0.228(0.128-0.406)*
Pay for HIV services			
Yes (Ref.)	-	-	1.00
No	-	-	4.801(2.333-9.879)*

440 *Note: *Significant at 95% (P<0.05) and 99% CI (P<0.001), N/A=Not applicable. - not evaluated*
 441 *in that model, and aOR=adjusted odds ratio.*

442

443

444

445 **Discussion**

446 Curbing the HIV pandemic among transport workers partly requires sustainable utilization of
447 multiple HIV services either singly or in combination [22]. Prior to this study, evidence about the
448 provision of HIV prevention services in combination by various stakeholders and the efficacy of
449 using them existed. However, literature was scanty about the utilization of multiple HIV
450 prevention services by transport workers. We found that less than half of the study participants
451 (43%) utilized over one HIV prevention service in the past year preceding the study. Similarly, in
452 a study among men who have sex with men (MSM) in Brazil, the utilization of behavioral and
453 biomedical HIV prevention services was found to be as low as 2.8% [26]. In both contexts, the
454 utilization were low, given that the success of HIV prevention services requires a high sustained
455 uptake of these services.

456
457 Voluntary counselling and HIV testing is the first and entry point into other HIV prevention
458 services such as ART, PREP, and EMTCT and these have been found to reduce risky sexual
459 behaviors[5]. On the other hand, safe male circumcision must precede the use of condoms for
460 effective HIV control. We found that most participants had tested for HIV and used condoms
461 (16.2%), followed by those who had received safe male circumcision also used condoms (15%),
462 and those who had tested for HIV and began on ART (9.1%). Figures reported in this study are
463 still much lower than those reported by health facility-based studies conducted in Uganda among
464 the general population. For example, a retrospective conducted in eastern Uganda found that
465 92.4% of the participants who had tested and were diagnosed with HIV were immediately started
466 on ART[23]. The low utilization of the multiple HIV prevention services in this study may be
467 related to the COVID-19 pandemic restrictions that led to the suspension of community outreach

468 services and camps, particularly for HIV testing (VCT), safe male circumcision (SMC), condom
469 awareness and demand creation[2, 18]. In addition, the low integration of ARV-based prevention
470 services such as PREP in many health facilities in Uganda may also be responsible for the low
471 uptake of multiple HIV prevention services in this study[2].

472

473 Nevertheless, our analysis indicates that across single use, most study participants used condoms
474 (32.2%) followed by voluntary counseling and testing (27.1%), and safe male circumcision (SMC,
475 17.3%). Similarly, the literature indicates that the utilization of HIV testing services among
476 transport workers is much higher than the utilization of other HIV prevention services in other
477 countries. In Zambia, the uptake of HIV counselling and testing (VCT) was reported at 83%,
478 compared with 39% who used safe male circumcision (SMC), and 34.8 to 65% who used condoms
479 in Nigeria[17, 35, 36]. Although previous studies conducted in Uganda among motorcycle taxi
480 riders and truck drivers found the use of VCT and condom to be as high as 87%, our study indicates
481 a much lower figure, particularly in a city where the HIV rates are high, which is deeply concerning
482 and requires shift actions to raise the demand for use of HIV prevention services [27, 37, 38]

483

484 Across demographic characteristics, we found that age, religion, marital status, number of
485 concurrent sex partners, awareness of HIV status, and length of stay in the area were statistically
486 significantly associated with the utilization of multiple HIV prevention services. This partly
487 resonates with similar results in Zambia where multiple sex partners and marital status were found
488 to be associated with the uptake of HIV testing services among long-distance truck drivers [35].
489 Whereas age, serostatus, stigma, and HIV knowledge were associated with the use of VCT services
490 among motorcycle taxi riders in Uganda[27]. Additionally, knowledge of HIV prevention,

491 perceived risk of HIV acquisition, and higher education levels were statistically significantly
492 associated with the utilization of condoms among truck drivers in Togo and Nigeria[17, 28]

493
494 However, from our multivariate analysis only religion, the number of sex partners, awareness of
495 HIV prevention services, awareness of one's serostatus, and paying for HIV services significantly
496 predicted the use of multiple HIV prevention services. We found that being a Muslim or a person
497 of another faith was associated with higher odds of using multiple HIV prevention services
498 compared with participants who subscribe to the Christian faith. Our possible explanation for this
499 finding may lay in the fact that Uganda is a conservative community, with certain religious beliefs
500 that prevent the use of certain specific HIV prevention services that act as artificial family planning
501 methods, such as condoms among Catholics. Thus, strategies that aim to increase the use of
502 multiple HIV services need to involve the religious community, most especially of the Christian
503 faith. Similar to a study conducted in Zambia about condom use among long-distance truck drivers,
504 we found that participants who had two or more concurrent sex partners compared with those who
505 had single, or no partners had higher odds of the use of multiple HIV prevention services[35].
506 Those with multiple partners may understand the risky behaviors they are engaged in and take
507 serious precautions to avoid acquiring HIV, but it may also indicate the effectiveness of the
508 messages and current campaigns by various government and non-government players to combat
509 the HIV scourge.

510
511 This study found that awareness of two or more HIV prevention services was associated with
512 higher odds of using multiple HIV prevention services compared with the awareness of only one
513 HIV prevention service. This finding calls for mention of available HIV prevention services each

514 time health workers have an interface with the transport workers. However, because of the limited
515 time transport workers have, their interface with health workers may be minimal. Thus, other
516 platforms need to be explored such as through their associations, leadership, or social media. Using
517 mainstream media (radio and television), other than other targeted approaches mentioned above,
518 in creating a demand for HIV prevention services, may not be an effective strategy, particularly in
519 mobile workers, such as transport workers, who always have limited time and access to such
520 services at their workplaces.

521

522 Knowing one's HIV status has been found to reduce the risk of HIV, partly through reducing risky
523 sexual behaviors. Also, it increases the awareness and utilization of HIV services, especially those
524 linked to an HIV diagnosis such as PEP, PREP, ART, and Emtct [5]. Similarly, we found that
525 participants who did not know their Serostatus were less likely to use HIV prevention services than
526 those who knew their HIV Serostatus, as earlier reported in Zambia, and Uganda among truck
527 drivers and motorcycle taxi riders using HIV testing services[27, 35].

528

529 The findings of this study indicate that paying for HIV prevention services is a barrier whereas
530 obtaining free services may be a facilitator. Specifically, we found that participants who did not
531 pay to access the HIV prevention services were 4.801 times more likely to use HIV prevention
532 services than those who paid to access the HIV prevention services. Given the current tough
533 economic times brought by the COVID-19 pandemic, stakeholders must make additional
534 investments in the providing free HIV prevention services, if we are to overcome the current HIV
535 epidemic in sub-Saharan Africa. However, the free services need to be provided innovatively
536 targeting the transport workers given that they have limited time. For example, transport workers

537 may access services such as condom use through their leadership, offices, commercial stages, and
538 workshops. In summary, this study suggests that behavioral factors influence the use of multiple
539 HIV prevention services compared with other individual factors.

540 **Limitation of the study.**

541 This study has limitations that may influence the conclusions made in this paper. First, we also did
542 not assess the health system factors related to the use of HIV prevention services, which could
543 have enriched the study. The cross-sectional study design does not also allow for causal
544 attributions. Lastly, the administration of the questionnaire by interviewers carried the risk of
545 social desirability bias.

546 **Conclusion**

547 Our study conducted in Mbarara city-southwestern Uganda found suboptimal uptake of multiple
548 HIV prevention services among transport workers. The study also found that factors that influence
549 the use of the multiple HIV prevention services included religion, awareness of HIV prevention
550 services, knowledge of HIV status, number of sexual partners and experience of paying for HIV
551 prevention services. Therefore, differentiated strategies are needed to increase the utilization of
552 HIV prevention services among the transport workers to curb the HIV epidemic. We recommend
553 interventions aimed at increasing awareness about HIV prevention services, awareness about HIV
554 prevention service-related costs to be borne by the patients, encouraging HIV testing among
555 persons with multiple sex partners and Christians. Future qualitative research to explore attitudes
556 and perceptions regarding HIV prevention services are recommended.

557

558

559 **Funding**

560 The research reported in this publications was partly supported by the Fogarty International Center
561 (U.S. Department of state's Office of the US Global Aids Coordinator and Health Diplomacy
562 (S/GAC) and the president's Emergency Plan for AIDs Relief (PEPFAR) of the national institutes
563 of Health under the award number R25TW011210. The content is solely the responsibility of the
564 authors and does not necessarily represent the official views of the National Institutes of Health.

565 **Conflict of Interest**

566 Author declares no conflict of interest

567 **Acknowledgment**

568 Special thanks go to the funding body, transport workers and their leadership, the city clerk, and
569 the director of health services Mbarara districts where the study was conducted.

570 **Author contributions**

571 **Concept and proposal development:** Asiimwe John Baptist, Benjamin Betunga, Phionah
572 Atuhaire, Catherine Nakasiita, Christa Kanyamuneza, Proscovia Namiiro, Joseph Tugume,
573 Matovu Hairat

574 **Data collection, and analysis:** Asiimwe John Baptist, Benjamin Betunga, Phionah Atuhaire,
575 Catherine Nakasiita, Christa Kanyamuneza, Proscovia Namiiro, Joseph Tugume, Matovu Hairat

576 **Supervision:** Asiimwe John Baptist, Ahmed M Sarki,, Richard Mugisha, Edward Kumakeche

577 **Writing the original draft:** Asimwe John Baptist, Benjamin Betunga, Phionah Atuhaire,
578 Catherine Nakasiita, Christa Kanyamuneza, Proscovia Namiiro, Joseph Tugume, Matovu Hairat, Benedicto
579 Mugabi, Birungi Lilian, Ahmed M Sarki,, Richard Mugisha, Edward Kumakeche

580 **Writing- review, and editing:** Asimwe John Baptist, Benjamin Betunga, Phionah Atuhaire,
581 Catherine Nakasiita, Christa Kanyamuneza, Proscovia Namiiro, Joseph Tugume, Matovu Hairat, Benedicto
582 Mugabi, Birungi Lilian, Ahmed M Sarki,, Richard Mugisha, Edward Kumakeche

583 **References**

- 584 1. Commission UA. Uganda population based HIV impact assessment. UAC: Kampala, Uganda.
585 2017:62-5.
- 586 2. UAC. ANNUAL JOINT AIDS REVIEW REPORT 2020/21. 2021.
- 587 3. UNAIDS. Country factsheets Uganda. 2019.
- 588 4. Hull MW, Montaner JS. HIV treatment as prevention: the key to an AIDS-free generation. *Journal*
589 *of food and drug analysis*. 2013;21(4):S95-S101.
- 590 5. Baxter C, Abdool Karim S. Combination HIV prevention options for young women in Africa. *Afr J*
591 *AIDS Res*. 2016;15(2):109-21. Epub 2016/07/12. doi: 10.2989/16085906.2016.1196224. PubMed PMID:
592 27399041.
- 593 6. Strauss M, George G, Lansdell E, Mantell JE, Govender K, Romo M, et al. HIV testing preferences
594 among long distance truck drivers in Kenya: a discrete choice experiment. *AIDS Care*. 2018;30(1):72-80.
595 Epub 2017/08/30. doi: 10.1080/09540121.2017.1367086. PubMed PMID: 28847156; PubMed Central
596 PMCID: PMC5903847.
- 597 7. Kelvin EA, George G, Mwai E, Nyaga EN, Mantell JE, Romo ML, et al. Offering Self-administered
598 Oral HIV Testing as a Choice to Truck Drivers in Kenya: Predictors of Uptake and Need for Guidance While
599 Self-testing. *AIDS Behav*. 2018;22(2):580-92. Epub 2017/05/26. doi: 10.1007/s10461-017-1783-9.
600 PubMed PMID: 28540563; PubMed Central PMCID: PMC5818565.
- 601 8. Kelvin EA, George G, Mwai E, Nyaga E, Mantell JE, Romo ML, et al. Offering self-administered oral
602 HIV testing to truck drivers in Kenya to increase testing: a randomized controlled trial. *AIDS Care*.
603 2018;30(1):47-55. Epub 2017/08/23. doi: 10.1080/09540121.2017.1360997. PubMed PMID: 28826229;
604 PubMed Central PMCID: PMC5901679.
- 605 9. Strauss M, George G, Mantell JE, Romo ML, Mwai E, Nyaga EN, et al. Stated and revealed
606 preferences for HIV testing: can oral self-testing help to increase uptake amongst truck drivers in Kenya?
607 *BMC Public Health*. 2018;18(1):1231. Epub 2018/11/08. doi: 10.1186/s12889-018-6122-1. PubMed PMID:
608 30400898; PubMed Central PMCID: PMC6219162.
- 609 10. Ibe Sally N, Nwoke E, Emerole C, Onyeocha F, Nwawume I, Nwaokoro JC. Knowledge, attitude and
610 perception of road transport workers to HIV/AIDS, STIs and condom use in southeast, Nigeria. *Technology*.
611 2014;1(2):50-4.
- 612 11. Lalla-Edward ST, Ncube S, Matthew P, Hankins CA, Venter WDF, Gomez GB. Uptake of health
613 services among truck drivers in South Africa: analysis of routine data from nine roadside wellness centres.

- 614 BMC Health Serv Res. 2017;17(1):649. Epub 2017/09/15. doi: 10.1186/s12913-017-2595-3. PubMed
615 PMID: 28903727; PubMed Central PMCID: PMC5598062.
- 616 12. NAA. Monitoring Progress towards the Targets of the 2011 UN Political Declaration on HIV and
617 AIDS: Cambodia Country Progress Report. 2014.
- 618 13. Mathias T, Emmanuel OO, Kakwezi MR, John M. Prevalence and predisposing factors of human
619 immunodeficiency virus infection among the boda-boda riders in Mbarara municipality-Uganda. 2020.
- 620 14. Delany-Moretlwe S, Bello B, Kinross P, Oliff M, Chersich M, Kleinschmidt I, et al. HIV prevalence
621 and risk in long-distance truck drivers in South Africa: a national cross-sectional survey. *Int J STD AIDS*.
622 2014;25(6):428-38. Epub 2013/12/20. doi: 10.1177/0956462413512803. PubMed PMID: 24352131.
- 623 15. Awosan K, Ibrahim MO, Arisegi S, Erhiano E. Knowledge of HIV/AIDS, risk perception, sexual
624 lifestyle and condom use among drivers in Sokoto, Nigeria. *Journal of Infectious Diseases and Immunity*.
625 2014;6(3):19-25.
- 626 16. Atilola GO, Akpa OM, Komolafe IO. HIV/AIDS and the long-distance truck drivers in south-west
627 Nigeria: a cross-sectional survey on the knowledge, attitude, risk behaviour and beliefs of truckers. *J Infect*
628 *Public Health*. 2010;3(4):166-78. Epub 2010/12/04. doi: 10.1016/j.jiph.2010.08.002. PubMed PMID:
629 21126721.
- 630 17. Yaya I, Landoh DE, Saka B, Vignikin K, Aboubakari AS, N'Dri K M, et al. Consistent Condom Use
631 during Casual Sex among Long-Truck Drivers in Togo. *PLoS One*. 2016;11(4):e0153264. Epub 2016/04/14.
632 doi: 10.1371/journal.pone.0153264. PubMed PMID: 27071059; PubMed Central PMCID:
633 PMC4829217.
- 634 18. UNAIDS. EN DANGER. 2022.
- 635 19. Malta M, Bastos F, Pereira-Koller E, Cunha M, Marques C, Strathdee SA. A qualitative assessment
636 of long distance truck drivers' vulnerability to HIV/AIDS in Itajai, southern Brazil. *AIDS care*.
637 2006;18(5):489-96.
- 638 20. Health Mo. Consolidated guidelines for the prevention and treatment of HIV and AIDS in Uganda.
639 2020.
- 640 21. UNAIDS. Creating HIV prevention cascades. 2021.
- 641 22. Chang LW, Serwadda D, Quinn TC, Wawer MJ, Gray RH, Reynolds SJ. Combination implementation
642 for HIV prevention: moving from clinical trial evidence to population-level effects. *The Lancet infectious*
643 *diseases*. 2013;13(1):65-76.
- 644 23. Opito R, Mpagi J, Bwayo D, Okello F, Mugisha K, Napyo A. Treatment outcome of the
645 implementation of HIV test and treat policy at The AIDS Support Organization (TASO) Tororo clinic, Eastern
646 Uganda: A retrospective cohort study. *Plos one*. 2020;15(9):e0239087.
- 647 24. Larsson EC, Ekström AM, Pariyo G, Tomson G, Sarowar M, Baluka R, et al. Prevention of mother-
648 to-child transmission of HIV in rural Uganda: Modelling effectiveness and impact of scaling-up PMTCT
649 services. *Global health action*. 2015;8(1):26308.
- 650 25. Ying R, Sharma M, Heffron R, Celum CL, Baeten JM, Katabira E, et al. Cost-effectiveness of pre-
651 exposure prophylaxis targeted to high-risk serodiscordant couples as a bridge to sustained ART use in
652 Kampala, Uganda. *J Int AIDS Soc*. 2015;18(4 Suppl 3):20013. Epub 2015/07/23. doi:
653 10.7448/IAS.18.4.20013. PubMed PMID: 26198348; PubMed Central PMCID: PMC4509901.
- 654 26. Firestone R, Rivas J, Lungo S, Cabrera A, Ruether S, Wheeler J, et al. Effectiveness of a combination
655 prevention strategy for HIV risk reduction with men who have sex with men in Central America: a mid-
656 term evaluation. *BMC Public Health*. 2014;14(1):1-15.
- 657 27. Ssekankya V, Githaiga SK, Aleko T, Munguciada EF, Nabakka VP, Kyalisiima JJ, et al. Factors
658 Influencing Utilization of HIV Testing Services among Boda-Boda Riders in Kabarole District, Southwestern
659 Uganda: A Cross-Sectional Study. *Biomed Res Int*. 2021;2021:8877402. Epub 2021/04/20. doi:
660 10.1155/2021/8877402. PubMed PMID: 33869636; PubMed Central PMCID: PMC8035009
661 publication of this article.

- 662 28. Sunmola A. Sexual practices, barriers to condom use and its consistent use among long distance
663 truck drivers in Nigeria. *AIDS care*. 2005;17(2):208-21.
- 664 29. Israel GD. Determining sample size. 1992.
- 665 30. Kalichman SC, Simbayi LC. HIV testing attitudes, AIDS stigma, and voluntary HIV counselling and
666 testing in a black township in Cape Town, South Africa. *Sexually transmitted infections*. 2003;79(6):442-7.
- 667 31. Yi S, Tuot S, Chhoun P, Pal K, Chhim K, Ngin C, et al. Sexual behaviors, HIV knowledge, HIV testing
668 attitudes and recent HIV testing among female entertainment workers in Cambodia: A cross-sectional
669 study. *PLoS One*. 2018;13(7):e0198095. Epub 2018/07/03. doi: 10.1371/journal.pone.0198095. PubMed
670 PMID: 29965968; PubMed Central PMCID: PMC6028079.
- 671 32. Napper LE, Fisher DG, Reynolds GL. Development of the perceived risk of HIV scale. *AIDS Behav*.
672 2012;16(4):1075-83. Epub 2011/07/26. doi: 10.1007/s10461-011-0003-2. PubMed PMID: 21785873;
673 PubMed Central PMCID: PMC3338003.
- 674 33. Visser MJ, Makin JD, Vandormael A, Sikkema KJ, Forsyth BW. HIV/AIDS stigma in a South African
675 community. *AIDS Care*. 2009;21(2):197-206. Epub 2009/02/21. doi: 10.1080/09540120801932157.
676 PubMed PMID: 19229689; PubMed Central PMCID: PMC4238924.
- 677 34. Asiimwe JB, Muwema M, Drake K. Factors that influence transition to advanced roles by RN to
678 BSN nurses, in three selected hospitals of Central-Uganda. *BMC nursing*. 2019;18(1):1-13.
- 679 35. Mutale LS, Chola M, Chongwe G, Kasongo W, Mwakazanga DK, Owiny M, et al. Uptake of HIV
680 testing and its associated factors among long-distance truck drivers in Zambia, 2015. *Journal of*
681 *Interventional Epidemiology and Public Health*. 2018;1. doi: 10.37432/jieph.2018.1.1.10.
- 682 36. Ekanem E, Afolabi B, Nuga A, Adebajo S. Sexual behaviour, HIV-related knowledge and condom
683 use by intra-city commercial bus drivers and motor park attendants in Lagos, Nigeria. *African Journal of*
684 *Reproductive Health*. 2005:78-87.
- 685 37. Matovu JK, Ssebadduka BN. Sexual risk behaviours, condom use and sexually transmitted
686 infection treatment-seeking behaviours among female sex workers and truck drivers in Uganda. *Int J STD*
687 *AIDS*. 2012;23(4):267-73. Epub 2012/05/15. doi: 10.1258/ijsa.2011.011313. PubMed PMID: 22581951.
- 688 38. Matovu JK, Ssebadduka NB. Knowledge, attitudes & barriers to condom use among female sex
689 workers and truck drivers in Uganda: a mixed-methods study. *Afr Health Sci*. 2013;13(4):1027-33. Epub
690 2014/06/19. doi: 10.4314/ahs.v13i4.24. PubMed PMID: 24940328; PubMed Central PMCID:
691 PMC4056494.

692