Title: Understanding the effectiveness of water, sanitation, and hygiene interventions: a counterfactual

simulation approach to generalizing the outcomes of intervention trials

Authors: Andrew F. Brouwer PhD[1], Mondal H. Zahid PhD[1], Marisa C. Eisenberg PhD[1], Benjamin F.

Arnold PhD[2,3], Sania Ashraf PhD[4], Jade Benjamin-Chung PhD[5,6], John M. Colford, Jr MD[7], Ayse

Ercumen PhD[8], Stephen P. Luby MD[9], Amy J. Pickering PhD[10], Mahbubur Rahman MBBS[4], Alicia N.M.

Kraay PhD[11], Joseph N.S. Eisenberg PhD[1,†], Matthew C. Freeman PhD[12,†]

[†]: these authors contributed equally

Affiliations

1. Department of Epidemiology, University of Michigan, Michigan, USA

2. Francis I. Proctor Foundation, University of California, San Francisco, San Francisco, CA, USA

3. Department of Ophthalmology, University of California, San Francisco, San Francisco, CA, USA

4. Environmental Interventions Unit, Infectious Disease Division, The International Centre for

Diarrhoeal Disease Research, Bangladesh, Dhaka, Bangladesh

5. Department of Epidemiology and Population Health, Stanford University, CA, USA

6. Chan Zuckerberg Biohub, San Francisco, CA, USA

7. School of Public Health, University of California Berkeley, Berkeley, CA, USA

8. Department of Forestry and Environmental Resources, North Carolina State University, Raleigh,

NC, USA

9. Division of Infectious Diseases and Geographic Medicine, Stanford University, CA, USA

10. Department of Civil and Environmental Engineering, University of California Berkeley,

Berkeley, CA, USA

11. Institute for Disease Modeling, a program within the Global Health Division of the Bill &

Melinda Gates Foundation, Seattle, Washington, USA

12. Rollins School of Public Health, Emory University, Atlanta, GA, USA

Corresponding author: Andrew F Brouwer; Department of Epidemiology, University of Michigan, 1415

Washington Heights, Ann Arbor, MI 48109; brouweaf@umich.edu; 734-764-7373

Declaration of conflicts of interest: ANMK's contributions were directly funded by the Bill & Melinda

Gates Foundation and not as part of the foundation grant to the authors. All other authors declare no

conflicts of interest.

Abstract

Background: While water, sanitation, and hygiene (WASH) interventions can reduce diarrheal disease, many large-scale trials have not found the expected health gains for young children in low-resource settings. Evidence-based guidance is needed to inform interventions and future studies.

Objectives: We aimed to estimate how sensitive the intervention effectiveness found in the WASH Benefits Bangladesh randomized controlled trial was to underlying WASH contextual and intervention factors (e.g.., baseline disease prevalence, compliance, community coverage, efficacy) and to generalize the results of the trial other contexts or scenarios.

Methods: We developed a disease transmission model to account for transmission across multiple environmental pathways, multiple interventions (water (W), sanitation (S), hygiene (H), nutrition (N)) applied individually and in combination, adherence to interventions, and the impact of individuals not enrolled in the study. Leveraging a set of mechanistic parameter combinations fit to the WASH Benefits Bangladesh trial (n=17,187) using a Bayesian sampling approach, we simulated trial outcomes under counterfactual scenarios to estimate how changes in intervention completeness, coverage, compliance, and efficacy, as well as preexisting WASH conditions and baseline disease burden, impacted intervention effectiveness.

Results: Increasing community coverage had the greatest impact on intervention effectiveness (e.g., median increases in effectiveness of 34.0 and 45.5% points in the WSH and WSHN intervention arms when increasing coverage to 20%). The effect of community coverage on effectiveness depended on how much transmission was along pathways not modified by the interventions. Intervention effectiveness was reduced by lower levels of preexisting WASH conditions or increased baseline disease burden. Individual interventions had complementary but not synergistic effects when combined.

Discussion: To realize the expected health gains, future WASH interventions must address community coverage and transmission along pathways not traditionally covered by WASH. The effectiveness of individual-level WASH improvements will be blunted the further the community is from the high community coverage needed to achieve herd protection.

Introduction

Enteric diseases, primarily spread through contact with fecally contaminated environments (e.g., water, surfaces, food), are one of the leading causes of morbidity and mortality in young children.[1] An estimated nearly 500,000 children under five years of age die from diarrheal disease globally each year[2,3], and it is hypothesized that repeated sub-clinical infections may lead to growth shortfalls.[4] Much of this burden is in low- and middle-income countries (LMICs).[5] Enteric pathogens are transmitted by myriad pathways, including fluids, fomites, food, flies, and fauna, as summarized in the classic "F-diagram".[6] Studying and preventing diarrheal disease is complicated because a diverse array of pathogens can cause similar symptoms,[7,8] pathogens can exploit multiple transmission pathways, and asymptomatic infections can contribute to the community pathogen burden.[9]

Diarrheal disease is greatly reduced in communities with robust water, sanitation, and hygiene (WASH) infrastructure, with mutually reinforcing levels of community and individual protection. Household-level WASH improvements can result in considerable reductions in diarrheal disease burden in LMICs, and many WASH interventions—such as improved latrines and handwashing with soap—have demonstrable efficacy to reduce fecal exposure.[10] A recent meta-analysis by Wolf et al. of WASH intervention randomized controlled trials (RCTs) highlighted that WASH interventions reduce diarrhea in children in low-resource settings.[10] However, there is substantial heterogeneity in the effect estimates across the studies, and many large-scale trials reported less-than-expected or null results.[11–17] In particular, the results of the WASH Benefits (WASH-B) Bangladesh and Kenya trials and the SHINE trial, all of which found no impacts of WASH on linear growth but some mixed effects on diarrhea, were particular subjects of substantial discussion in the literature.[18–22] The sub-optimal performance of the interventions in these trials is likely due to a combination of multiple factors, including incomplete blocking of all transmission pathways (low intervenable fraction, also called completeness), inadequate community coverage of the intervention, or a lack of intervention compliance or efficacy.[19,23] Additionally, a community's preexisting WASH conditions and baseline disease burden can also impact the real-world intervention effectiveness.[19,23] Assessing which factors are the largest barriers to diarrheal disease reduction will aid policy-makers, practitioners, and researchers in deciding how best to invest in WASH programs and design the next generation of programs and trials.[18–24]

RCTs are considered the gold-standard for estimating causal relationships, and they are rigorous assessments of a particular intervention within a particular context at a particular point in time. But, their findings do not necessarily generalize to other contexts or conditions—e.g., different populations, disease burdens, pathogens, transmission pathways, intervention fidelity and adherence—when there are effect modifiers that vary across field settings and intervention implementations.[20,23] Mechanistic infectious

disease transmission models, unlike meta-analyses, have the potential to generalize findings by directly

accounting for these location-specific contexts and conditions, exploring counterfactual questions through

simulation of alternate scenarios, and developing location-specific programmatic targets. This approach is

used extensively in other contexts to assess public health interventions or counterfactual conditions.[25–27] A

mechanistic, counterfactual approach could lead to better-targeted public health WASH interventions,

policy recommendations, and field trials.[28,29]

The aims of this work are to evaluate hypotheses about what led to the sub-optimal reductions in diarrhea

among intervention households in an RCT (WASH-B Bangladesh) using a compartmental transmission

model and to provide a framework to support planning of WASH interventions and context-specific

WASH programming. We previously developed this model framework accounting for multiple

environmental transmission pathways, shared environments, pre-existing WASH conditions, and

adherence to multiple interventions and applied it to the empirical trial data.[30] Our approach generates

thousands of combinations of coverage, intervention efficacy, and transmission pathway strengths that

could reasonably underlie the trial results. In this analysis, we leveraged those parameter combinations to

simulate how intervention effectiveness would have been different under alternate scenarios. These

counterfactual simulations provide evidence for policy recommendations, programmatic targets, and an

evaluation framework for next-generation WASH interventions.[20,21,24]

Methods

Summary of approach. In prior work, we developed a compartmental transmission model framework to

explain the outcomes of a RCT.[30] Then, we explored how effectiveness of a single intervention depends

on six WASH factors:[31]

- *Preexisting WASH conditions:* We account for the fact that a fraction of the population already
has WASH infrastructure comparable to that provided by the intervention trial.
- *Disease transmission potential:* We account for the baseline disease prevalence through the basic
reproduction number R_0, a summary measure of the disease transmission potential. Note that,
given the values of the other WASH factors, there is a one-to-one correspondence between the
disease prevalence and the basic reproduction number. In this analysis, we vary the baseline
disease prevalence rather than R_0, as it what intervention trials measure.
- *Intervention compliance*: We account for both intervention fidelity (whether the intervention was
delivered) and adherence (whether participants used the intervention), defining compliance as the
fraction of participants assigned to an intervention who are using it.

- *Intervenable fraction of transmission*: Any individual intervention targets some, but often not all, of the transmission pathways that pathogens exploit. We account for how much transmission is along pathways that the intervention attenuates (even if imperfectly) and how much even a perfect intervention would not affect.
- *Intervention efficacy*: Because interventions do not result in perfect reduction of transmission or shedding, we account for the actual reduction.
- *Community coverage fraction*: Intervention trials typically only provide the intervention to a subset of the population. We account for the fraction of the population is enrolled in the trial.

In this analysis, we use a multi-intervention version of the model to investigate outcomes in the WASH-B Bangladesh randomized controlled trial specifically. In prior work, we demonstrated how to find mechanistic parameter sets that were consistent with individual-level diarrheal outcomes.[30] Here, we use these mechanistic parameter sets to simulate what intervention effectiveness would have been in each of the WASH-B Bangladesh trial arms under each of six counterfactuals corresponding to the six WASH factors above, accounting for uncertainty in the parameters underlying the real data (original scenario). By simulating what the intervention effectiveness would have been in the trial under alternative circumstances, we evaluated the extent to which each factor may have contributed to the observed outcomes.

Data. The WASH-B Bangladesh trial was a cluster-randomized trial of the efficacy of water, sanitation, hygiene, and nutrition interventions, alone and in combination, on diarrhea prevalence and linear growth.[14] The investigators measured (child-guardian-reported, past-seven-day, all-cause) diarrheal prevalence in children at three time points approximately one year apart. Households in the study area are typically organized into compounds in which a patrilineal family shares a common space and resources, such as a water source and latrine. A total of 5551 compounds were enrolled, contingent on having a pregnant woman in her second trimester during the enrollment period. The study followed one or more target children born after baseline, as well as any other children in the compound who were under age 3 at baseline. These compounds were grouped into 720 clusters. Each cluster was assigned to one of seven arms testing combinations of four interventions: water chlorination (W), a double-pit, pour-flush improved latrine (S), handwashing with soap and water (H), and supplementary nutrition sachets (N). The control arm (C) consisted of 180 compounds, while 90 were assigned to each of the water (W), sanitation (S), handwashing (H), nutrition (N), combined water, sanitation, and handwashing (WSH), and all interventions (WSH-N) arms. Specific details on trial design, interventions, and results are published elsewhere[14,32]. We assessed whether any individual was using an intervention or a substantively

equivalent preexisting WASH condition through four indicators defined and assessed by the investigators:

detection of free chlorine in water (W), latrine with a functional water seal (S), handwashing station with

soap and water (H), at least 50% of nutrition sachets consumed (N). The W and H interventions were

intended for the households of the target children, but we were not able to determine whether other

children in the compound were in that household or not. For this analysis, we assumed that non-target

children were covered by the interventions; any misspecification will attenuate the estimated efficacy of

the W and H interventions.[14] We removed individuals with negative reported ages (n=2), missing reported

diarrhea (n=2,745), or missing in any of the four use indicators (n=2,660), which left 17,187 individual

observations (76% of the original sample) over the three surveys.

Ethics. This secondary analysis of deidentified data was not regulated as human-subjects research.

Model. Our compartmental transmission model, denoted SISE-RCT, is a susceptible-infectious-

susceptible (SIS) model with transmission through environmental (E) compartments and simulated to

steady state to approximate an RCT.[30] The SISE-RCT model accounts for the six key mechanistic factors

underlying WASH RCT outcomes described above. We previously developed the single-intervention

model,[31] and we include a description of it in the supplemental material for convenience.

As discussed in the Data section above, the WASH-B Bangladesh trial included 720 clusters of

households each assigned to one of seven arms (control, W, S, H, N, WSH, WSHN). We extended the

single-intervention SISE-RCT model to a multi-intervention model by accounting for transmission across

three environmental pathways (water, fomites & hands, and all others combined), four interventions

applied individually (W, S, H, N) and in combination (WSH, WSHN), and individual-level compliance

with interventions or preexisting conditions. In brief, we modeled each of the 720 clusters with

susceptible and infectious compartments for each of 2^4=16 combinations of interventions/conditions

depending on household adherence, i.e., in every cluster, we modeled the infection prevalence for each

combination of having or not having each intervention or equivalent preexisting WASH condition. For

example, for a cluster in the WSH arm, we estimated how many people were not using any interventions,

how many were using W only, how many were using S only, etc., and what the infection prevalence was

among each group given their collective interaction through the shared environments.

Specifically, for a given cluster, we denote the fraction of the population that is susceptible to infection

and is using intervention(s) or preexisting WASH condition(s) i in $\{0, W, S, H, N, WS, \dots, WSHN\}$ as S_i,

where 0 indicates the use of no intervention or preexisting WASH condition. Analogously, we denote the

fraction of the population that is infected analogously by I_i. The intervention and control arms are

simulated separately. The populations with regular and attenuated exposure are modeled in every cluster

simulation, accounting for the fraction of the population 1) enrolled in the study (the community

coverage, ω), 2) with preexisting WASH conditions (ρ_0), and 3) complying with the intervention (ρ).

Note that ρ_0 and ρ are vectors of length 16 that each sum to 1; i.e., everyone is categorized into one of the

16 exposure groups. For each cluster, the overall fraction of the population in each exposure group is

given by the vector

$$N = (1 - \omega)\rho_0 + \omega\rho. \#(1)$$

That is, the fraction of the population enrolled in the study (ω) follows the intervention compliance

distribution of exposure groups (ρ), and the fraction of the population not enrolled ($1 - \omega$) follows the

preexisting WASH condition distribution of exposure groups (ρ_0). In the control arm, $\rho = \rho_0$.

The environment is partitioned into the three environmental pathways: water (E_w), fomites & hands (E_f),

and all other pathways (E_o). We assume that chlorination reduces transmission along the water pathway,

sanitation reduces shedding into the water pathway, handwashing reduces transmission along the fomite

pathway, and nutrition reduces susceptibility (transmission) for all three pathways. For each of the three

pathways j, an environmental compartment E_j is characterized by the shedding into the environment (α_j),

the decay of pathogens in the environment (ξ_j), and the transmission of pathogens from the environment

to susceptible individuals (β_j). The relative magnitude of shedding into E_j and relative transmission from

E_j for the attenuated compared to the exposed populations are given by ϕ_{α_j} and ϕ_{β_j}, respectively. We

also accounted for the possibility that the preexisting conditions were less efficacious than the RCT

intervention. Once infected, individuals clear the infection at rate γ.

The model diagram of the multi-intervention SISE-RCT is given in Figure 1, although only two of the

sixteen different exposure populations are shown. The full equations are given below (Eqs 2). A

transmission term $\beta_j E_j$ denotes transmission from the environmental pathway j. The transmission term

$\beta_j E_j$ is attenuated by ϕ_{β_j}only for people in an attenuated exposure group (S_i) using an intervention or

preexisting condition that reduces transmission from pathway j, and contamination of that environmental

pathway is attenuated by ϕ_{α_j} only for infectious people in an exposure group (I_i) using an intervention or

preexisting condition that reduces shedding into pathway j. In the following equations, the subscripts *w, f,*

and *o* represent the water, fomite and other pathways. The subscripts *0, W, S, H, N,* and any combinations

represent the exposure groups as defined above. Parameters $\omega, \rho,$ and ρ_0 do not show up in these

equations but are accounted for in the constraints, as discussed below. For brevity, we omit the $\frac{dS_i}{dt}$

equations, each of which is given by $\frac{dS_i}{dt} = -\frac{dI_i}{dt}$ for the corresponding subpopulation.

$$\frac{dI_0}{dt} = (\beta_w E_w + \beta_f E_f + \beta_o E_o)S_0 - \gamma I_0 \#(2)$$

$$\frac{dI_W}{dt} = (\varphi_{\beta_w,W} \cdot \beta_w E_w + \beta_f E_f + \beta_o E_o)S_W - \gamma I_W$$

$$\frac{dI_S}{dt} = (\beta_w E_w + \beta_f E_f + \beta_o E_o)S_S - \gamma I_S$$

$$\frac{dI_H}{dt} = (\beta_w E_w + \varphi_{\beta_f,H} \cdot \beta_f E_f + \beta_o E_o)S_H - \gamma I_H$$

$$\frac{dI_N}{dt} = \varphi_{\beta,N} \cdot (\beta_w E_w + \beta_f E_f + \beta_o E_o)S_H - \gamma I_H$$

$$\frac{dI_{WS}}{dt} = (\varphi_{\beta_w,W} \cdot \beta_w E_w + \beta_f E_f + \beta_o E_o)S_{WS} - \gamma I_{WS}$$

$$\frac{dI_{WH}}{dt} = (\varphi_{\beta_w,W} \cdot \beta_w E_w + \varphi_{\beta_f,H} \cdot \beta_f E_f + \beta_o E_o)S_{WH} - \gamma I_{WH}$$

$$\frac{dI_{WN}}{dt} = \varphi_{\beta,N} \cdot (\varphi_{\beta_w,W} \cdot \beta_w E_w + \beta_f E_f + \beta_o E_o)S_{WN} - \gamma I_{WN}$$

$$\frac{dI_{SH}}{dt} = (\beta_w E_w + \varphi_{\beta_f,H} \cdot \beta_f E_f + \beta_o E_o)S_{SH} - \gamma I_{SH}$$

$$\frac{dI_{SN}}{dt} = \varphi_{\beta,N} \cdot (\beta_w E_w + \beta_f E_f + \beta_o E_o)S_{SN} - \gamma I_{SN}$$

$$\frac{dI_{HN}}{dt} = \varphi_{\beta,N} \cdot (\beta_w E_w + \varphi_{\beta_f,H} \cdot \beta_f E_f + \beta_o E_o)S_{HN} - \gamma I_{HN}$$

$$\frac{dI_{WSH}}{dt} = (\varphi_{\beta_w,W} \cdot \beta_w E_w + \varphi_{\beta_f,H} \cdot \beta_f E_f + \beta_o E_o)S_{WSH} - \gamma I_{WSH}$$

$$\frac{dI_{WSN}}{dt} = \varphi_{\beta,N} \cdot (\varphi_{\beta_w,W} \cdot \beta_w E_w + \beta_f E_f + \beta_o E_o)S_{WSN} - \gamma I_{WSN}$$

$$\frac{dI_{WHN}}{dt} = \varphi_{\beta,N} \cdot (\varphi_{\beta_w,W} \cdot \beta_w E_w + \varphi_{\beta_f,H} \cdot \beta_f E_f + \beta_o E_o)S_{WHN} - \gamma I_{WHN}$$

$$\frac{dI_{SHN}}{dt} = \varphi_{\beta,N} \cdot (\beta_w E_w + \varphi_{\beta_f,H} \cdot \beta_f E_f + \beta_o E_o)S_{SHN} - \gamma I_{SHN}$$

$$\frac{dI_{WSHN}}{dt} = \varphi_{\beta,N} \cdot (\varphi_{\beta_w,W} \cdot \beta_w E_w + \varphi_{\beta_f,H} \cdot \beta_f E_f + \beta_o E_o)S_{WSH} - \gamma I_{WSH}$$

$$\frac{dE_w}{dt} = \alpha_w \left(\sum_{S \text{ not in } i} I_i + \varphi_{\alpha,S} \sum_{S \text{ in } i} I_i \right) - \xi_w E_w$$

$$\frac{dE_f}{dt} = \alpha_f \left(\sum_i I_i \right) - \xi_f E_f$$

$$\frac{dE_o}{dt} = \alpha_o \left(\sum_i I_i \right) - \xi_o E_o$$

215   To find the steady state values (denoted by *) for the human compartments in the intervention arm, we set
216   the above equations to 0 and simplify:

$$0 = \left(R_{0,w} \bar{E}_w + R_{0,f} \bar{E}_f + R_{0,o} \bar{E}_o \right) S_0^* - I_0^*, \#(2)$$

$$0 = \left(\varphi_{\beta_w,W} \cdot R_{0,w} \bar{E}_w + R_{0,f} \bar{E}_f + R_{0,o} \bar{E}_o \right) S_W^* - I_W^*,$$

$$0 = \left(R_{0,w} \bar{E}_w + R_{0,f} \bar{E}_f + R_{0,o} \bar{E}_o \right) S_S^* - I_S^*,$$

$$0 = \left(R_{0,w} \bar{E}_w + \varphi_{\beta_f,H} \cdot R_{0,f} \bar{E}_f + R_{0,o} \bar{E}_o \right) S_H^* - I_H^*,$$

$$0 = \varphi_{\beta,N} \cdot \left(R_{0,w} \bar{E}_w + R_{0,f} \bar{E}_f + R_{0,o} \bar{E}_o \right) S_N^* - I_N^*,$$

$$0 = \left(\varphi_{\beta_w,W} \cdot R_{0,w} \bar{E}_w + R_{0,f} \bar{E}_f + R_{0,o} \bar{E}_o \right) S_{WS}^* - I_{WS}^*,$$

$$0 = \left(\varphi_{\beta_w,W} \cdot R_{0,w} \bar{E}_w + \varphi_{\beta_f,H} \cdot R_{0,f} \bar{E}_f + R_{0,o} \bar{E}_o \right) S_{WH}^* - I_{WH}^*,$$

$$0 = \varphi_{\beta,N} \cdot \left(\varphi_{\beta_w,W} \cdot R_{0,w} \bar{E}_w + R_{0,f} \bar{E}_f + R_{0,o} \bar{E}_o \right) S_{WN}^* - I_{WN}^*,$$

$$0 = \left(R_{0,w} \bar{E}_w + \varphi_{\beta_f,H} \cdot R_{0,f} \bar{E}_f + R_{0,o} \bar{E}_o \right) S_{SH}^* - I_{SH}^*,$$

$$0 = \varphi_{\beta,N} \cdot \left(R_{0,w} \bar{E}_w + R_{0,f} \bar{E}_f + R_{0,o} \bar{E}_o \right) S_{SN}^* - I_{SN}^*,$$

$$0 = \varphi_{\beta,N} \cdot \left(R_{0,w} \bar{E}_w + \varphi_{\beta_f,H} \cdot R_{0,f} \bar{E}_f + R_{0,o} \bar{E}_o \right) S_{HN}^* - I_{HN}^*,$$

$$0 = \left(\varphi_{\beta_w,W} \cdot R_{0,w} \bar{E}_w + \varphi_{\beta_f,H} \cdot R_{0,f} \bar{E}_f + R_{0,o} \bar{E}_o \right) S_{WSH}^* - I_{WSH}^*,$$

$$0 = \varphi_{\beta,N} \cdot \left(\varphi_{\beta_w,W} \cdot R_{0,w} \bar{E}_w + R_{0,f} \bar{E}_f + R_{0,o} \bar{E}_o \right) S_{WSN}^* - I_{WSN}^*,$$

$$0 = \varphi_{\beta,N} \cdot \left(\varphi_{\beta_w,W} \cdot R_{0,w} \bar{E}_w + \varphi_{\beta_f,H} \cdot R_{0,f} \bar{E}_f + R_{0,o} \bar{E}_o \right) S_{WHN}^* - I_{WHN}^*,$$

$$0 = \varphi_{\beta,N} \cdot \left(R_{0,w} \bar{E}_w + \varphi_{\beta_f,H} \cdot R_{0,f} \bar{E}_f + R_{0,o} \bar{E}_o \right) S_{SHN}^* - I_{SHN}^*,$$

$$0 = \varphi_{\beta,N} \cdot \left(\varphi_{\beta_w,W} \cdot R_{0,w} \bar{E}_w + \varphi_{\beta_f,H} \cdot R_{0,f} \bar{E}_f + R_{0,o} \bar{E}_o \right) S_{WSHN}^* - S_{WSHN}^*,$$

$$\bar{E}_w = \sum_{S \text{ not in } i} I_i^* + \varphi_{\alpha,S} \sum_{S \text{ in } i} I_i^*,$$

$$\bar{E}_f = \bar{E}_o = \sum_i I_i.$$

Here, $R_{0,j} = {\alpha_j \beta_j}\big/{\xi_j \gamma}$ is the pathway-specific reproduction number for transmission through environment

E_i. The variables $\bar{E}_j = \xi_j E_j^*/\alpha_j$ are conveniently scaled environmental steady states. For this model, the

overall basic reproduction number is $R_0 = R_{0,w} + R_{0,f} + R_{0,o}$, denoting the sum of the transmission

potential through each pathway.

To solve for the steady states solutions for our 32 state variables (16 exposure groups times two

susceptible/infection states), we solve the nonlinear system of equations (Eqs (3)) subject to the

population constraint given in Eqs (1). The prevalence of disease in the population is denoted $\pi^* = \sum I_i^*$.

The prevalence in an intervention arm π^* is compared to the prevalence π_c^* in the control arm, and the

intervention effectiveness for that arm is defined as $\varepsilon = (\pi_c^* - \pi^*)/\pi_c^*$, namely the fractional reduction in

prevalence in the intervention arm relative to the control arm.

In summary, the model includes 18 parameters: i) the overall basic reproduction number R_0, which

defines the transmission potential in the control arm at baseline, ii) two parameters partitioning R_0 into

the strengths of the drinking water $R_{0,w}$, fomite & hands $R_{0,f}$, and all other transmission pathways $R_{0,o}$,

iii) eight relative reproduction numbers accounting for systematic differences in disease pressure over the

trial time periods (baseline, midline, and endline) and across arms independently, iv) the community

coverage ω, and v) efficacy parameters defining the effect of each intervention (four) or preexisting

WASH condition (two) on the transmission pathways (the W intervention (chlorination) reduces

transmission via the water pathway $\varphi_{\beta_w,W}$; the S intervention (latrine with water seal) reduces shedding

into the shared water environment with different efficacy for preexisting conditions $\bar{\varphi}_{\alpha_w,S}$ and the trial

intervention $\varphi_{\alpha_w,S}$; the H (handwashing with soap and water) reduces transmission via the fomite

pathway with different efficacy for preexisting conditions $\bar{\varphi}_{\beta_f,H}$ and trial intervention $\varphi_{\beta_f,H}$; and the N

intervention (nutrition supplementation) reduces susceptibility to all transmission $\varphi_{\beta,N}$). In the Eqs (3),

the basic reproduction number parameters are adjusted by the time and arm-specific and relative basic

reproduction numbers corresponding to the cluster being modeled, and the intervention efficacy

parameters $\varphi_{\alpha_w,S}$ and $\varphi_{\beta_f,H}$ are replaced by $\bar{\varphi}_{\alpha_w,S}$ and $\bar{\varphi}_{\beta_f,H}$ in clusters without the S and H

interventions, respectively, and at baseline.

When solving for the steady state of these differential equations for a given cluster in a given time period,

we use the distribution of interventions and preexisting WASH conditions ρ recorded in the data for those

participants and assume that participants not in the study have the same distribution of preexisting

conditions as the control arm participants ρ_0. We solved this system using the nleqslv package in R.

To fit the model to the trial data, we employed a hybrid sampling-importance resampling and estimation

framework to obtain 50,000 parameter combinations that represented a good fit to the diarrheal outcomes

of each participant using a Bernoulli statistical likelihood.[33] We resampled, with replacement, from our

initial 50,000 parameter combinations, based on their goodness of fit; 3,774 unique parameter

combinations were included in the final sample, with varying frequency. These parameters sets are similar

to but not exactly the same as in Brouwer et al,[30] as they include a small code correction and use the

computational improvement described previously.[31] The fit to the data is given in Figure S1, and the

distributions of parameters are given in Figures S2-5.

WASH-B Bangladesh counterfactual analysis. We conducted two types of counterfactual analysis.

First, we estimated the counterfactual intervention effectiveness in each arm across a range of each of the

WASH factors starting from the scenario based on the median value of each parameter (which resulted in

a fit close to the best-fit and was more representative of the parameter distributions than the specific best-

fit parameter set). Because we are using the actual preexisting WASH conditions and intervention

compliance recorded for each individual in the trial, there is not a well-defined way to continuously scale

these two factors. So, we only compare the actual simulation to a "no preexisting conditions" and "full

compliance" counterfactual, respectively.

Second, to account for the uncertainty in the parameters underlying the actual intervention outcomes, for

each of the 50,000 parameter sets k identified by fitting the model to WASH-B Bangladesh, we defined

the corresponding original scenario matching the WASH-B Bangladesh trial outcomes and the

corresponding intervention effectiveness ε_0^k. Because we are now accounting for uncertainty across these

50,000 parameters sets, it is not possible to succinctly capture changes as we continuously vary the

factors. Thus, we considered six specific counterfactual scenarios, detailed in Table 2. Any parameter sets

that eliminated disease in the control arm in a counterfactual simulation were censored from the results as

they did not provide information on intervention effectiveness.

The main outcome of a counterfactual simulation was the (absolute) change intervention effectiveness

compared to the original scenario, namely $\varepsilon_*^k - \varepsilon_0^k$, where ε_*^k is the intervention effectiveness in the

given counterfactual scenario for the jth parameter set. We used absolute change rather than percentage

change because absolute change, unlike percent change, is bounded (between -100 and 100 percentage

points). The intervention is more effective (i.e., a greater reduction in diarrheal prevalence in the

intervention arm compared to the control arm) in the counterfactual scenario than the original scenario

when the change is positive. To assess whether the intervention factors modified the intervention

effectiveness in the counterfactual scenarios, for a subset of counterfactuals, we assessed how the effect

of the counterfactual depended on quantiles of the other parameter values.

The counterfactuals scenarios are not intended to be "plausible" for some specific, real-world changes.

Indeed, changing the contextual factors of the preexisting WASH conditions and baseline disease

prevalence is not possible. Instead, we can imagine these counterfactuals representing running the same

trial in a different location to assess what the results would have been. The intervenable fraction is also

not changeable for a given intervention (and underlying set of pathogens) but could be changed by adding

additional intervention aspects to reduce transmission along other pathways. More broadly, we believe

that investigating a broad range of counterfactual scenarios improves our understanding of the disease–

intervention system, so that more effective interventions may be designed in the future.

Results

WASH-B Bangladesh counterfactual analysis from the median parameters

At the median parameter values (Figs S2-S5), intervention effectivenesses were 8.5% in the W arm,

40.8% in the S arm, 37.2% in the H arm, 32.1% in the WSH arm, 35.0% in N arm, and 34.6% in the

WSHN arm, reflecting the results of the WASH-B Bangladesh trial. We estimated that removing all

preexisting sanitation and hygiene infrastructure would have resulted in a modest reduction in

intervention efficacy in all arms except for W, where the reduction was negligible (Figure 2a). Increasing

baseline disease prevalence (by changing increasing the R_0 parameter) nonlinearly decreased intervention

effectiveness in all arms, with decreasing reductions in effectiveness as baseline prevalence increased

(Figure 2b). We estimated that there would be negligible-to-modest improvements in intervention

effectiveness if there was full compliance (Figure 2c). Increasing the intervenable fraction (by reducing

the strength of the "other" pathway and proportionally increasing the strength of the water and fomite &

hands pathways while keeping the overall R_0 constant), increasingly improved the intervention

effectiveness (Figure 2d), with the WSH and WSHN interventions nearly achieving disease elimination if

transmission were 100% intervenable. Increasing community coverage increased effectiveness

approximately linearly, with each intervention achieving disease elimination at a different level of
community coverage (Figure 2e; W: 75%, S: 90%, H: 75%, WSH: 35%, N: 55%, WSHN: 30%).
Increasing intervention efficacy approximately linearly increased intervention effectiveness in the
corresponding arms (Figure 2f-i). Increasing efficacy of the sanitation and hygiene interventions to 100%
resulted in approximate disease elimination in the corresponding arms, but elimination would not have
been achieved by increasing the efficacy of the W and N interventions.

WASH-B Bangladesh counterfactual analysis accounting for parameter uncertainty

Because we do not know that the median values of the parameters in the scenario investigated in the first
counterfactual analysis are accurate, we also considered the distribution of changes in intervention
effectiveness when each counterfactual scenario (Table 2) was applied to the distribution of samples that
fit the original data well. The median baseline disease prevalence in the original scenario was 7.1% (range
5.9–8.2%), decreasing to 5.7% (range 5.2–6.3%) for midline/endline (Figure S1). The median
intervention effectivenesses were 8.2% in the W arm, 36.4% in the S arm, 33.1% in the H arm, 30.3% in
the WSH arm, 34.0% in N arm, and 34.6% in the WSHN arm (Table 3). The percentage point change
varied across arms in each counterfactual scenario. Figure 3 shows the distribution of percentage point
change in intervention effectiveness over the 50,000 parameter sets for each arm and counterfactual
scenario, and Table 3 gives the median values.

Eliminate preexisting WASH Conditions. We found that implementing the interventions in a community
with no handwashing stations with soap and water or latrines with water seals would have likely resulted
in less effective interventions compared to the actual community's higher baseline WASH conditions
(e.g., 9.5 percentage points less in the WSH arm; Figure 3a). The W arm is the exception because it had
lower effectiveness in the original scenario. The uncertainty in change in intervention effectiveness in
each arm was largely driven by uncertainty in what the baseline disease prevalence would have been in
the counterfactual scenario (median 8.9%, range 6.4–23.1%).

Double baseline disease prevalence. A higher transmission potential corresponding to a doubling of the
baseline diarrheal disease prevalence (median 14.2% vs median 7.1%) would also have resulted in less-
effective interventions compared to the true baseline diarrheal disease prevalence (e.g., 11.9 percentage
points less in the WSH arm; Figure 3b). As above, the W arm is the exception because it had lower
effectiveness in the original scenario.

Full compliance. The impact of increasing intervention adherence was negligible-to-modest (e.g., 4.4
percentage points more in the WSH arm; Figure 3c). Note that intervention compliance, as defined by the
investigators, was already high.[14,34]

Half of the other pathway transmission can be intervened on. We found that intervention effectiveness

could have been greater if more of the total disease transmission was via the water and fomite pathways

rather than through pathways that were not intervened on (e.g., 20.1 percentage points more in the WSH

arm; Figure 3d). There was potential for a substantial increase in intervention effectiveness, as indicated

by the distribution of the individual simulation outcomes, but the median impact was modest, with a less

than 25 percentage point increase in effectiveness in the multi-intervention arms and a less than 15

percentage point increase in the single-intervention arms (Table 3). The uncertainty in the potential

impact was largely driven by uncertainty in how much of the disease transmission was through other

pathways in the original scenario.

Double intervention efficacy. We assessed the impact of increasing efficacy—defined as increasing the

reduction of transmission along the relevant pathway(s)—of the four interventions. We found that in each

of these increased efficacy scenarios, substantial increases in intervention efficacy could have improved

intervention effectiveness in the corresponding arms (Figure 3e–h), with median improvements between

5% and 20% points.

Increase community coverage. The median estimated community coverage in the trial was 5.4%, but this

estimate was highly uncertain, ranging from nearly 0% to 20% (Figure S5). For our main coverage

counterfactual, we increased the community coverage in each simulation to 20.0%, chosen as a

substantial but not unreasonable increase in coverage.[35] This counterfactual scenario was associated with

the greatest median increase in intervention effectiveness (among all households now covered by the

intervention) of any of the considered counterfactual scenarios (e.g., 34.0 and 45.5 percentage points

more in the WSH and WSHN arms; Figure 4a). Following the results of the single-intervention model

that highlighted that the effect of coverage depended on the other WASH factors, we plotted the

intervention effectiveness distributions for this community coverage counterfactual by quintiles of the

values of the other WASH factors. Effect modification is present if the effect of increased coverage

depends on the quintile of the WASH factor. Note that when looking at quintiles of one factor, the values

of the other factors may not be evenly distributed across the quintiles if values of the factors are

correlated. We found that the increase intervention effectiveness with increased community coverage in

the W, S, WSH, and WSHN intervention arms depended partly on the strength of transmission via the

water pathway (Figure 4b). The increases in intervention effectiveness in these arms could only reach

their full potential if the strength of the water pathway were high. A similar, but more modest effect was

seen for the H arm and the strength of the fomite pathway (Figure 4c). The greatest overall effect modifier

of the impact of increased coverage on intervention effectiveness is the strength of the other pathways

(i.e., the intervenable fraction, Figure 4d). When the strength of other pathways was high, increasing

coverage had less of an impact. Intervention efficacy also modified the impact of increased coverage but

only in the intervention arms with those interventions (Figure 4e–h).

To further understand the joint impact of community coverage and the intervenable fraction (i.e., the

strength of the other transmission pathway), we simulated the intervention effectiveness as a function of

increased coverage for the highest and lowest quintiles of intervention completeness (Figure 5). The

impact of increased coverage on intervention effectiveness depended on the intervenable fraction most

strongly for the W, WSH, and WSHN arms, moderately for the S arm, and little for the H and N arms.

For example, in the W arm, increasing community coverage to 50% resulted in a median increase of 80%

points for samples with the highest intervenable fractions but only 23 percentage points for samples with

the lowest intervenable fractions. In contrast, in the H arm, increasing community coverage to 50%

resulted in a median increase of 31 percentage points for samples with the highest intervenable fractions

compared only 22 percentage points for samples with the lowest intervenable fractions. (Note that the fact

that intervenable fraction was relevant for the N arm at all is a result of the correlations between the

intervenable fraction and the other parameters in the original parameter sets.)

Discussion

Our model-based analysis used counterfactual simulations to generalize the results of a WASH

intervention trial and develop guidance for policymakers and researchers. Our first finding was that

increasing community coverage led to the most substantial reduction in disease among people receiving

interventions. Second, we found that intervention completeness (i.e., the fraction of disease transmission

along pathways that were intervened on) was an important effect modifier of the impact of community

coverage on intervention effectiveness, with the impact of increased community coverage enhanced when

interventions covered a larger fraction of transmission. Third, our work suggests that interventions are

likely to be more effective when disease burden is low. Finally, we found that multifaceted WASH

interventions (WSH) added value over single component interventions (W, S, or H). Each of these

findings suggest a path forward for policy and program recommendations for WASH investments and

demonstrates how transmission models can be used to design the next generation of WASH interventions

and set location-specific programmatic targets.

The importance of ensuring high community WASH coverage to address health outcomes has been

highlighted in multiple context, including latrines,[29,36] bed nets,[37] and chemotherapy for helminths,[38]

among others. Further work is needed to improve our measures of indirect and direct intervention

effects[29,39] to better determine sanitation targets. Our findings support the call for systems-level WASH

provisioning and improved universal access, underscoring the fundamental push to achieve the 2030

sustainable development targets.[40]

Our finding that the intervenable fraction (completeness) was an important effect modifier, emphasizes

the need to better understand the sources of exposure not impacted by traditional WASH interventions.

For example, contamination of food outside the home or from flies or exposure to feces from animals

living near or inside the home may not be reduced by water quality or latrine interventions.[41–43] Capturing

and reducing transmission through additional targeted interventions would increase the fraction of

transmission intervened on and thereby make increased community coverage even more effective.

Low diarrheal prevalence makes it more difficult to observe a statistically significant reduction in

diarrhea.[14] However, from a mechanistic perspective, we found that intervention effectiveness would have

been lower had the background disease pressure in the community been higher (i.e., higher baseline

disease prevalence) because individual-level interventions can be overwhelmed by higher disease

pressure from the community, including those not covered by the intervention. This finding is supported

by the outcomes of WASH-B Kenya trial, which had higher disease prevalence (27% in the control arm)

and no significant intervention effects on diarrheal prevelance[15] and is consistent with previous literature

that has shown that non-pharmaceutical interventions are more effective for less transmissible pathogens

or when the population has a higher degree of population immunity.[44] This is not to say that individual

improvements would have no effect but that the effects are blunted if disease pressure in the rest of the

community were not also addressed.

Similarly, many have suggested that when preexisting WASH conditions are relatively high, interventions

do not provide a substantial improvement in efficacy and thus health outcomes.[18,19,22,45] However, from

the transmission system perspective reflected by our results, if the preexisting WASH conditions

(particularly among those *not* covered by the intervention) were poorer, the community disease pressure

would be greater, and it would be more difficult to protect study participants from infection, even if the

people covered by the intervention had a greater improvement in protection.

Because enteric pathogens can exploit multiple transmission pathways, many studies have tried to

determine whether combined WASH interventions (WSH) are more effective than single interventions

(W, S, or H).[32,46] Whether or not there is an additional effect of combined interventions depends on

whether the interventions are complementary, that is, whether they each block some of the transmission

that the other interventions would not have blocked.[47] This complementarity is an assumption in our

transmission model framework (as each intervention affects different parts of the disease system), and

because the model can fit the data, we find that complementarity is consistent with the observed trial

results.[33] Other modeling and empirical studies, support that WASH interventions can complement each

other, or even potentially be synergistic.[48,49] In this work, we found that the combined interventions could

have a greater effect than the individual interventions, but that the effects were generally sub-additive,

meaning that the effectiveness of the combined WSH intervention was less than the sum of its parts

(Table 3). Combined interventions offer a substantially better chance of disease elimination, especially at

higher coverage levels (Figure 5).

One challenge that WASH RCTs often face is achieving high compliance through both high fidelity

(providing the interventions as planned) and high adherence of participants to the use of the intervention.

In WASH-B Bangladesh, the intervention compliance, as defined by trial investigators, was high,

generally above 90%.[14,34] Accordingly, our full compliance counterfactual was limited in the impact it

could detect.

The strength of our approach is underscored by the rich and high-quality data collected by the WASH-B

Bangladesh trial (and other RCTs) and in our transmission model framework capturing relative disease

prevalence. RCTs provide the gold standard of evidence about intervention effectiveness in a specific

context, and our approach allows us to generalize RCT results to other contexts, providing a tool for

powerful policy and programmatic guidance. The SISE-RCT model can be customized for local contexts

and interventions and then used to support local decision-making (e.g., to determine whether to invest in

community coverage vs intervention efficacy). Future work may also develop recommendations for

achieving elimination while minimizing costs. One limitation of our study is the high uncertainty in many

of the model parameters, especially the intervenable fraction, which propagates into the counterfactual

scenarios. These uncertainties stem from potential trade-offs in the model, e.g., a low intervenable

fraction and a low intervention efficacy may have similar effects. Fortunately, our framework has the

potential to incorporate additional information about parameters like the intervenable fraction and

efficacy through our Bayesian sampling-importance resampling approach, allowing us to tailor

projections of intervention effectiveness to specific parameter regions based on additional information

(e.g., chlorination efficacy above 75%). One limitation of the data was the inability to distinguish whether

non-target children were members of the same household as the target child or not, which introduced

misspecification into our classification of W and H exposures, likely attenuating the efficacy estimates for

those interventions. Also, we accounted for changes in disease pressure between, but not within, survey

periods; future work may more directly address seasonal changes in disease pressure and even pathway

strength, as a function of precipitation, seasonal flooding, etc. Another limitation of this study is that our

results do not directly address some aspects of the Sustainable Development Goal (SDG) Target 6.2.[40] For

example, the sanitation arm did not move households from no or basic sanitation to improved sanitation

(as defined by the Joint Monitoring Programme). So, the "sanitation" intervention outcomes we estimated

may not directly correspond to the policy-relevant changes required to meet SDG target. Likewise, the

"water" intervention focused on water quality improvements (chlorination) but not water quantity. None

of these issues are limitations of our modeling framework; rather, they are limitations of our specific

application. Applying our methods across other trial datasets could address these limitations by allowing

for modeling of other—and perhaps more policy-relevant—WASH exposure parameters.

Our work contributes to the robust discussion[18–24,50] about the future directions of WASH research and

programming, and our modeling approach is well-suited to reevaluating current evidence during the

"pause for reflection" recommended by a consensus of WASH reserachers.[19] This consensus group said

that "the lesson perhaps lies in not seeking to attribute benefits to individual WASH factors but in that the

public health dividends are paid when comprehensive services are in place." Our work underscores this

conclusion, not only by emphasizing the importance of coverage and completeness of interventions, but

also in its rejection of the hypotheses that greater effectiveness might be found in areas with greater

disease prevalence or lower preexisting WASH infrastructure. Indeed, our findings suggest that the effect

of individual-level WASH improvements will be blunted the further the community is from achieving

herd protection. Accordingly, this analysis provides further evidence supporting community-level

interventions seeking to achieve herd protection through high community coverage.

482 Contributors

JNSE, MCE, MCF, and AFB conceived of the study. JNSE and MCF secured funding for the study. AFB,

MCE, and JNSE developed the model. AFB wrote and implemented the software code, completed formal

analysis and visualization, and curated the data and code. MHZ and ANMK validated the software code.

AFB wrote the original draft with input from JNSE and MCF. BFA, SA, JBC, JMC, AE, SPL, AJP, and

MR contributed equally by aiding in interpretation of the results and providing their expertise in the

WASH Benefits trials. All authors reviewed and edited the manuscript. All authors had full access to all

study data.

490 Acknowledgements

This work was funded by the Bill & Melinda Gates Foundation (grant INV-005081) and the National

Science Foundation (grant DMS-1853032). The original WASH-B Bangladesh trial was also funded by

the Bill & Melinda Gates Foundation (grant OPPGD759). ANMK's contributions were directly funded by

the Bill & Melinda Gates Foundation and not as part of the foundation grant to the authors. ANMK is an

employee of the Bill & Melinda Gates Foundation; however, this study does not necessarily represent the

views of the Bill & Melinda Gates Foundation.

Data sharing

The WASH Benefits Bangladesh data is publicly available at https://osf.io/tprw2/. The data and code

underlying the results of this paper are available at https://doi.org/10.5281/zenodo.10950560. Data

underlying each of the figures is given in the Excel spreadsheet supplement.

References

1.   Vos T, Lim SS, Abbafati C, et al. Global burden of 369 diseases and injuries in 204 countries and
     territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet*.
     2020;396(10258):1204-1222. doi:10.1016/S0140-6736(20)30925-9

2.   Troeger C, Blacker B, Khalil IA, et al. Estimates of the global, regional, and national morbidity,
     mortality, and aetiologies of lower respiratory infections in 195 countries, 1990–2016: a
     systematic analysis for the Global Burden of Disease Study 2016. *Lancet Infect Dis*.
     2018;18(11):1191-1210. doi:10.1016/S1473-3099(18)30310-4

3.   Wolf J, Johnston RB, Ambelu A, et al. Burden of disease attributable to unsafe drinking water,
     sanitation, and hygiene in domestic settings: a global analysis for selected adverse health
     outcomes. *Lancet*. 2023;401(10393):2060-2071. doi:10.1016/S0140-6736(23)00458-0

4.   Humphrey JH. Child undernutrition, tropical enteropathy, toilets, and handwashing. *Lancet*.
     2009;374(9694):1032-1035. doi:10.1016/S0140-6736(09)60950-8

5.   Julian TR. Environmental transmission of diarrheal pathogens in low and middle income
     countries. *Environ Sci Process Impacts*. 2016;18(8):944-955. doi:10.1039/C6EM00222F

6.   Wagner EG, Lanoix JN. Excreta disposal for rural areas and small communities. *Monogr Ser
     World Health Organ*. 1958;39:1-182. http://www.ncbi.nlm.nih.gov/pubmed/13581743.

7.   Kotloff KL, Nataro JP, Blackwelder WC, et al. Burden and aetiology of diarrhoeal disease in
     infants and young children in developing countries (the Global Enteric Multicenter Study, GEMS):
     A prospective, case-control study. *Lancet*. 2013;382(9888):209-222. doi:10.1016/S0140-
     6736(13)60844-2

8.   Liu J, Platts-Mills JA, Juma J, et al. Use of quantitative molecular diagnostic methods to identify
     causes of diarrhoea in children: a reanalysis of the GEMS case-control study. *Lancet*.
     2016;388(10051):1291-1301. doi:10.1016/S0140-6736(16)31529-X

9.   Watson SI, Rego RTT, Hofer T, Lilford RJ. Evaluations of water, sanitation and hygiene
     interventions should not use diarrhoea as (primary) outcome. *BMJ Glob Heal*. 2022;7(5):e008521.
     doi:10.1136/bmjgh-2022-008521

10. Wolf J, Hubbard S, Brauer M, et al. Effectiveness of interventions to improve drinking water,
     sanitation, and handwashing with soap on risk of diarrhoeal disease in children in low-income and
     middle-income settings: a systematic review and meta-analysis. *Lancet*. 2022;400(10345):48-59.

doi:10.1016/S0140-6736(22)00937-0

11.   Clasen T, Boisson S, Routray P, et al. Effectiveness of a rural sanitation programme on diarrhoea,
soil-transmitted helminth infection, and child malnutrition in Odisha, India: A cluster-randomised
trial. *Lancet Glob Heal*. 2014;2(11):e645-e653. doi:10.1016/S2214-109X(14)70307-9

12.   Patil SR, Arnold BF, Salvatore AL, et al. The effect of India's total sanitation campaign on
defecation behaviors and child health in rural Madhya Pradesh: A cluster randomized controlled
trial. *PLoS Med*. 2015;11(8). doi:10.1371/journal.pmed.1001709

13.   Pickering AJ, Djebbari H, Lopez C, Coulibaly M, Alzua ML. Effect of a community-led sanitation
intervention on child diarrhoea and child growth in rural Mali: A cluster-randomised controlled
trial. *Lancet Glob Heal*. 2015;3(11):e701-e711. doi:10.1016/S2214-109X(15)00144-8

14.   Luby SP, Rahman M, Arnold BF, et al. Effects of water quality, sanitation, handwashing, and
nutritional interventions on diarrhoea and child growth in rural Bangladesh: A cluster randomised
controlled trial. *Lancet Glob Heal*. 2018;6(3):e302-e315. doi:10.1016/S2214-109X(17)30490-4

15.   Null C, Stewart CP, Pickering AJ, et al. Effects of water quality, sanitation, handwashing, and
nutritional interventions on diarrhoea and child growth in rural Kenya: a cluster-randomised
controlled trial. *Lancet Glob Heal*. 2018;6(3):e316-e329. doi:10.1016/S2214-109X(18)30005-6

16.   Rogawski McQuade ET, Platts-Mills JA, Gratz J, et al. Impact of Water Quality, Sanitation,
Handwashing, and Nutritional Interventions on Enteric Infections in Rural Zimbabwe: The
Sanitation Hygiene Infant Nutrition Efficacy (SHINE) Trial. *J Infect Dis*. 2020;221(8):1379-1386.
doi:10.1093/infdis/jiz179

17.   Knee J, Sumner T, Adriano Z, et al. Effects of an urban sanitation intervention on childhood
enteric infection and diarrhea in Maputo, Mozambique: A controlled before-and-after trial. *Elife*.
2021;10. doi:10.7554/eLife.62278

18.   Cumming O, Curtis V. Implications of WASH Benefits trials for water and sanitation. *Lancet*
Glob Heal. 2018;6(6):e613-e614. doi:10.1016/S2214-109X(18)30192-X

19.   Cumming O, Arnold BF, Ban R, et al. The implications of three major new trials for the effect of
water, sanitation and hygiene on childhood diarrhea and stunting: A consensus statement. *BMC*
Med. 2019;17(1):1-9. doi:10.1186/s12916-019-1410-x

20.   Levy K, Eisenberg JNS. Moving towards transformational WASH. *Lancet Glob Heal*.
2019;7(11):e1492. doi:10.1016/S2214-109X(19)30396-1

21.   Pickering AJ, Null C, Winch PJ, et al. The WASH Benefits and SHINE trials: interpretation of
562   WASH intervention effects on linear growth and diarrhoea. *Lancet Glob Heal*. 2019;7(8):e1139-
563   e1146. doi:10.1016/S2214-109X(19)30268-2

564   22.   Coffey D, Spears D. Implications of WASH Benefits trials for water and sanitation. *Lancet Glob*
565   *Heal*. 2018;6(6):e615. doi:10.1016/S2214-109X(18)30225-0

566   23.   Haque SS, Freeman MC. The Applications of Implementation Science in Water, Sanitation, and
567   Hygiene (WASH) Research and Practice. *Environ Health Perspect*. 2021;129(6):65002.
568   doi:10.1289/EHP7762

569   24.   Amebelu A, Ban R, Bhagwan J, et al. The Lancet Commission on water, sanitation and hygiene,
570   and health. *Lancet*. 2021;398(10310):1469-1470. doi:10.1016/S0140-6736(21)02005-5

571   25.   Li X, Mukandavire C, Cucunubá ZM, et al. Estimating the health impact of vaccination against ten
572   pathogens in 98 low-income and middle-income countries from 2000 to 2030: a modelling study.
573   *Lancet*. 2021;397(10272):398-408. doi:10.1016/S0140-6736(20)32657-X

574   26.   World Health Organization (WHO). *Impact of the COVID-19 Pandemic on Seven Neglected*
575   *Tropical Diseases: A Model-Based Analysis*. Geneva; 2021.
576   https://apps.who.int/iris/handle/10665/343993.

577   27.   Lofgren E, Halloran ME, Rivers CM, et al. Mathematical models: A key tool for outbreak
578   response. *Proc Natl Acad Sci*. 2015;112(2):E234-E234. doi:10.1073/pnas.1423846112

579   28.   Coffeng LE, Vaz Nery S, Gray DJ, Bakker R, de Vlas SJ, Clements ACAA. Predicted short and
580   long-term impact of deworming and water, hygiene, and sanitation on transmission of soil-
581   transmitted helminths. Walker M, ed. *PLoS Negl Trop Dis*. 2018;12(12):1-17.
582   doi:10.1371/journal.pntd.0006758

583   29.   Fuller JA, Eisenberg JNS. Herd protection from drinking water, sanitation, and hygiene
584   interventions. *Am J Trop Med Hyg*. 2016;95(5):1201-1210. doi:10.4269/ajtmh.15-0677

585   30.   Brouwer AF, Eisenberg MC, Bakker KM, et al. Leveraging infectious disease models to interpret
586   randomized controlled trials: Controlling enteric pathogen transmission through water, sanitation,
587   and hygiene interventions. Lau EH, ed. *PLOS Comput Biol*. 2022;18(12):e1010748.
588   doi:10.1371/journal.pcbi.1010748

589   31.   Brouwer AF, Kraay ANM, Zahid MH, Eisenberg MC, Freeman MC, Eisenberg JNS. A
590   mechanistic modeling approach to assessing the sensitivity of outcomes of water, sanitation, and

hygiene interventions to local contexts and intervention factors. *medRxiv*. 2024.
doi:10.1101/2024.03.09.24304020

32. Arnold BF, Null C, Luby SP, et al. Cluster-randomised controlled trials of individual and
combined water, sanitation, hygiene and nutritional interventions in rural bangladesh and Kenya:
The WASH benefits study design and rationale. *BMJ Open*. 2013;3(8):1-17.
doi:10.1136/bmjopen-2013-003476

33. Brouwer AF, Eisenberg MC, Bakker KM, et al. Leveraging infectious disease models to interpret
randomized controlled trials: controlling enteric pathogen transmission through water, sanitation,
and hygiene interventions. *medRxiv*. January 2022:2022.04.28.22274441.
doi:10.1101/2022.04.28.22274441

34. Ercumen A, Pickering AJ, Kwong LH, et al. Do Sanitation Improvements Reduce Fecal
Contamination of Water, Hands, Food, Soil, and Flies? Evidence from a Cluster-Randomized
Controlled Trial in Rural Bangladesh. *Environ Sci Technol*. 2018;52(21):12089-12097.
doi:10.1021/acs.est.8b02988

35. Garn J V., Sclar GD, Freeman MC, et al. The impact of sanitation interventions on latrine
coverage and latrine use: A systematic review and meta-analysis. *Int J Hyg Environ Health*.
2017;220(2):329-340. doi:10.1016/j.ijheh.2016.10.001

36. Harris M, Alzua ML, Osbert N, Pickering A. Community-Level Sanitation Coverage More
Strongly Associated with Child Growth and Household Drinking Water Quality than Access to a
Private Toilet in Rural Mali. *Environ Sci Technol*. 2017;51(12):7219-7227.
doi:10.1021/acs.est.7b00178

37. Teklehaimanot A, Sachs JD, Curtis C. Malaria control needs mass distribution of insecticidal
bednets. *Lancet*. 2007;369(9580):2143-2146. doi:10.1016/S0140-6736(07)60951-9

38. Anderson R, Truscott J, Hollingsworth TD. The coverage and frequency of mass drug
administration required to eliminate persistent transmission of soil-transmitted helminths. *Philos
Trans R Soc B*. 2014;369(1645):20130435. doi:10.1098/rstb.2013.0435

39. Halloran ME, Auranen K, Baird S, et al. Simulations for designing and interpreting intervention
trials in infectious diseases. *BMC Med*. 2017;15(1):1-8. doi:10.1186/s12916-017-0985-3

40. World Health Organization (WHO) and the United Nations Children's Fund (UNICEF). *Progress
on Household Drinking Water, Sanitation and Hygiene 2000-2020: Five Years into the SDGs*.

Geneva; 2021.

41.   Penakalapati G, Swarthout J, Delahoy MJ, et al. Exposure to Animal Feces and Human Health: A
Systematic Review and Proposed Research Priorities. *Environ Sci Technol*. 2017;51(20):11537-
11552. doi:10.1021/acs.est.7b02811

42.   Khamesipour F, Lankarani KB, Honarvar B, Kwenti TE. A systematic review of human pathogens
carried by the housefly (Musca domestica L.). *BMC Public Health*. 2018;18(1):1049.
doi:10.1186/s12889-018-5934-3

43.   DuPont HL. The Growing Threat of Foodborne Bacterial Enteropathogens of Animal Origin. *Clin*
Infect Dis. 2007;45(10):1353-1361. doi:10.1086/522662

44.   Baker RE, Saad-Roy CM, Park SW, Farrar J, Metcalf CJE, Grenfell BT. Long-term benefits of
nonpharmaceutical interventions for endemic infections are shaped by respiratory pathogen
dynamics. *Proc Natl Acad Sci*. 2022;119(49). doi:10.1073/pnas.2208895119

45.   Arnold BF, Null C, Luby SP, Colford JM. Implications of WASH Benefits trials for water and
sanitation – Authors' reply. *Lancet Glob Heal*. 2018;6(6):e616-e617. doi:10.1016/S2214-
109X(18)30229-8

46.   Fewtrell L, Kaufmann RB, Kay D, Enanoria W, Haller L, Colford JM. Water, sanitation, and
hygiene interventions to reduce diarrhoea in less developed countries: A systematic review and
meta-analysis. *Lancet Infect Dis*. 2005;5(1):42-52. doi:10.1016/S1473-3099(04)01253-8

47.   Eisenberg JNS, Scott JC, Porco T. Integrating disease control strategies: Balancing water
sanitation and hygiene interventions to reduce diarrheal disease burden. *Am J Public Health*.
2007;97(5):846-852. doi:10.2105/AJPH.2006.086207

48.   Eisenberg JNS, Trostle J, Sorensen RJD, Shields KF. Toward a Systems Approach to Enteric
Pathogen Transmission: From Individual Independence to Community Interdependence. *Annu Rev*
Public Health. 2012;33(1):239-257. doi:10.1146/annurev-publhealth-031811-124530

49.   Esrey SA. Water, waste, and well-being: A multicountry study. *Am J Epidemiol*. 1996;143(6):608-
623. doi:10.1093/oxfordjournals.aje.a008791

50.   D'Mello-Guyett L, Yates T, Bastable A, et al. Setting priorities for humanitarian water, sanitation
and hygiene research: A meeting report. *Confl Health*. 2018;12(1):10-14. doi:10.1186/s13031-
018-0159-8

Tables

Table 1: Parameters of the SISE-RCT model. The SISE-RCT model is a compartmental susceptible-infectious-susceptible (SIS) model with transmission through environmental (E) compartments and simulated to steady state to approximate an RCT. The median value column denotes the median values for the multi-intervention model. Parameters ρ_0 and ρ do not have median values because they are determined by the data.

Parameter	Definition	Median value
ρ_0	Baseline WASH conditions (fraction of individuals in the community with intervention-level WASH infrastructure)	—
ρ	Compliance (fraction of individuals in intervention arm using intervention)	—
R_0	Transmission potential (basic reproduction number)	1.10
π_c^*	Baseline disease prevalence (note: this technically a model output, but we treat it as a parameter because it has a 1-to-1 correspondence with R_0 given the other parameters)	6.9%
$R_{0,w}/R_0$	Fraction of transmission along the water pathway	
$R_{0,f}/R_0$	Fraction of transmission along the fomite & hands pathway	
$R_{0,f}/R_0$	Fraction of transmission on pathways not intervened on (i.e., 1 minus the intervenable fraction)	0.48
$1 - \varphi_\alpha$	Intervention efficacy for reducing shedding (S=sanitation intervention)	0.23 (S),
$1 - \varphi_\beta$	Intervention efficacy for reducing transmission (W= water, H=hygiene, N=nutrition interventions)	0.44 (W), 0.33 (H), 0.16 (N)
ω	Community coverage fraction (fraction of community included in the intervention trial)	5.4%

662 **Table 2: WASH-B Bangladesh counterfactual scenarios and implementations.** WASH = water, sanitation, & hygiene. *indicates parameters in the

663 counterfactual simulation.

Category	Definition	What would have happened if…	Implementation
WASH conditions	Quality of WASH infrastructure at baseline	… no households had preexisting WASH conditions substantively equivalent to the intervention.	Pre-existing conditions were removed from individuals not in the corresponding intervention arms. (Setting ρ_0^* to the vector $(1, 0, … , 0)$).
Compliance	The extent to which individuals assigned to an intervention received it (fidelity) and used it (adherence)	… all households assigned an intervention received and used it.	All individuals in each intervention arm were modelled as using the intervention. (Adjusting ρ^* appropriately, e.g., moving the fraction of the population from S to HS in the hygiene arm).
Disease conditions	Disease prevalence at baseline in the absence of preexisting WASH conditions	… the disease pressure was greater	The basic reproduction number is increased such that the baseline prevalence in the absence of preexisting WASH conditions is doubled. (Optimization was used to determine the appropriate value of R_0^* for each parameter set.)
Intervenable fraction	Whether there are transmission pathways that are not affected by the intervention	… more of transmission was along pathways that could be intervened on.	The strength of the other pathway is reduced by 50% and replaced proportionally by the water and fomite pathways. $(R_{0,w}^* = R_{0,w} + \frac{1}{2}R_{0,o} \cdot R_{0,w}/(R_{0,w} + R_{0,f}),\ R_{0,f}^* = R_{0,f} + \frac{1}{2}R_{0,o} \cdot R_{0,f}/(R_{0,w} + R_{0,f}),\ R_{0,o}^* = \frac{1}{2}R_{0,o})$
Efficacy	The extent to which using the intervention reduced transmission along relevant pathways	… the interventions provided a greater reduction in transmission.	The strength of the reduction in transmission from each intervention (and corresponding preexisting condition) is doubled. $(\varphi^* = \min(2\varphi, 1))$
Community coverage	The fraction of the at-risk population in a cluster that was provided the intervention	… a different fraction of the population was provided the intervention.	Study coverage is 20%, …, 90%, 100%. $(\omega^* = 0.2, … , 1.0)$

Table 3. Median intervention effectiveness and median percent point change in intervention effectiveness in each intervention arm for each counterfactual scenario compared to the original scenario. Intervention effectiveness (ε) in intervention effectiveness is 1 minus the relative risk of diarrhea in the intervention arm vs the control arm in each scenario, expressed as a percentage. The column Δε gives the median change in intervention effectiveness in percentage points (*not* the change in median intervention effectiveness); a negative number reflects a decrease in intervention effectiveness.

	W		S		H		WSH		N		WSHN	
	ε	Δε	ε	Δε	ε	Δε	ε	Δε	ε	Δε	ε	Δε
Original scenario	8.1%	—	36.3%	—	33.0%	—	30.2%	—	33.9%	—	34.5%	—
No WASH baseline conditions	8.0%	-0.2%	21.7%	-14.6%	24.6%	-8.6%	20.6%	-9.5%	26.5%	-7.3%	25.7%	-8.4%
Double baseline disease prevalence	6.8%	-1.3%	14.7%	-21.6%	15.7%	-17.5%	18.6%	-11.9%	17.7%	-16.0%	24.1%	-10.3%
Full compliance	11.6%	+3.6%	36.4%	+0.1%	34.2%	+1.1%	34.7%	+4.4%	34.7%	+0.9%	38.6%	+4.1%
Half of other pathway transmission can be intervened on	14.2%	+6.0%	48.4%	+11.8%	43.7%	+10.6%	50.8%	+20.1%	37.8%	+3.8%	53.0%	+18.2%
Double efficacy of chlorination	15.4%	+7.1%		—		—	36.4%	+6.1%		—	40.6%	+5.9%
Double efficacy of latrine water seal		—	55.4%	+18.9%		—	46.0%	+15.7%		—	48.9%	+14.2%
Double efficacy of handwashing		—		—	54.6%	+21.3%	50.2%	+19.8%		—	52.8%	+18.3%
Double efficacy of nutrition		—		—		—		—	45.3%	+11.3%	44.3%	+10.1%
Increase community coverage to 20%	22.9%	+14.6%	43.3%	+6.8%	41.4%	+8.3%	63.8%	+34.0%	49.2%	+15.3%	79.6%	+45.5%

 Figure captions

Figure 1: Single-intervention SISE-RCT model diagram with one attenuated exposure population and a regular exposure population interacting
through shared environments. The SISE-RCT model is a compartmental susceptible-infectious-susceptible (SIS) model with transmission through
environmental (E) compartments and simulated to steady state to approximate an RCT. The black lines denote infection and recovery, the blue lines denote
shedding from infectious individuals into environmental compartments, the grey lines denote pick-up of pathogens from the environment by susceptible
individuals, and the orange lines denote environmental pathogen decay. S_i and I_i denote susceptible and infectious fraction of the ith attenuated exposure
population, and S_0 and I_0 denote susceptible and infectious fraction of the regular exposure population. E_w, E_f, and E_o denote environmental pathways for water,
fomites & hands, and all other pathways. This figure was adapted from Figure 2 of Brouwer et al (2022).[30]

Figure 2: Intervention effectiveness as a function of WASH intervention factors. The SISE-RCT model is a compartmental susceptible-infectious-
susceptible (SIS) model with transmission through environmental (E) compartments and simulated to steady state to approximate an RCT. The model was
simulated at the median values of the model parameters when fit to the WASH Benefits Bangladesh trial and across ranges of counterfactual values of six
contextual and intervention WASH factors. The six WASH factors are A) preexisting WASH conditions (fraction of individuals not enrolled in the intervention
arm that are using preexisting WASH infrastructure), B) baseline disease prevalence (a function of the basic reproduction number R_0), C) compliance (fraction of
individuals enrolled in the intervention arm that are using the intervention), D) intervenable fraction of transmission (how much of the transmission could be
prevented in a perfect intervention), E) the community coverage fraction (fraction of the population enrolled in the trial), and F-I) the intervention efficacy
(fraction reduction in transmission or shedding when using the intervention) of each intervention. The underlying data are provided in Excel Table S1. WASH =
water, sanitation, & hygiene.

Figure 3: Percentage point change in intervention effectiveness compared to the original scenario in each counterfactual scenario. The SISE-RCT model
is a compartmental susceptible-infectious-susceptible (SIS) model with transmission through environmental (E) compartments and simulated to steady state to
approximate an RCT. Here, we applied it to data from the WASH-B Bangladesh trial, selecting 50,000 parameter sets consistent with the trial outcomes. We
simulated each parameter set under each counterfactual scenario (Table 2). The violin plots give the distribution of values across the 50,000 simulations, with
median points. The underlying data are provided in Excel Table S2A-H. W = water, S = sanitation, H = hygiene, N = nutrition.

Figure 4: Percentage point change in intervention effectiveness compared to the original scenario in the 20% coverage counterfactual scenario (a)
overall and (b) considering other parameters as potential effect modifiers (b-h). The SISE-RCT model is a compartmental susceptible-infectious-susceptible
(SIS) model with transmission through environmental (E) compartments and simulated to steady state to approximate an RCT. Here, we applied it to data from

the WASH-B Bangladesh trial, selecting 25,000 parameter sets consistent with the trial outcomes. We simulated each parameter set under the 20% coverage

counterfactual scenario (Table 2). The violin plots give the distribution of values across the 25,000 simulations, with median points. In plots b-h, the five violin

plots give the distributions of the intervention effectiveness across quintiles of the listed potential effect modifier. The underlying data are provided in Excel

Table S3A-H. W = water, S = sanitation, H = hygiene, N = nutrition.

Figure 5: Percentage point change in intervention effectiveness compared to the original scenario in each arm for the lowest and highest quintiles of

intervenable fraction. The SISE-RCT model is a compartmental susceptible-infectious-susceptible (SIS) model with transmission through environmental (E)

compartments and simulated to steady state to approximate an RCT. Here, we applied it to data from the WASH-B Bangladesh trial, selecting 25,000 parameter

sets consistent with the trial outcomes. We simulated each parameter set for coverage counterfactual scenarios ranging from 20% to 100% (Table 2). The violin

plots give the distribution of values across the simulations, with median points and a line connecting the medians, for highest (dark) and lowest (light) quintiles of

the intervenable fraction, i.e., the fraction of transmission that the interventions could directly act on. The underlying data are provided in Excel Table S4A-F. W

= water, S = sanitation, H = hygiene, N = nutrition.