Tcte1 **knockout influence on energy chain transportation, apoptosis and spermatogenesis – implications for male infertility**

Running title: *Tcte1* gene in male infertility – mouse 'knockout' model

Marta Olszewska[1], Agnieszka Malcher[1], Tomasz Stokowy[2], Nijole Pollock[3], Andrea J. Berman[4], Sylwia Budkiewicz[1], Marzena Kamieniczna[1], Hanna Jackowiak[5], Piotr Jedrzejczak[6], Alexander N. Yatsenko[3], Maciej Kurpisz[1*].

[1] Institute of Human Genetics, Polish Academy of Sciences, Poznan, Poland
[2] Scientific Computing Group, IT Division, University of Bergen, Norway
[3] Department of OB/GYN and Reproductive Sciences, School of Medicine, University of Pittsburgh, PA, USA
[4] Department of Biological Sciences, University of Pittsburgh, PA, USA
[5] Department of Histology and Embryology, Poznan University of Life Sciences, 60-625 Poznan, Poland
[6] Division of Infertility and Reproductive Endocrinology, Department of Gynecology, Obstetrics and Gynecological Oncology, Poznan University of Medical Sciences, 60-535 Poznan, Poland

[*] Corresponding author:

        Maciej Kurpisz, PhD, MD
        Institute of Human Genetics, Polish Academy of Sciences
        Strzeszynska 32
        60-479 Poznan, POLAND
        phone: +48616579202
        e-mail: maciej.kurpisz@igcz.poznan.pl

other authors' emails:

Marta Olszewska, PhD	marta.olszewska@igcz.poznan.pl
Agnieszka Malcher, PhD	agnieszka.malcher@igcz.poznan.pl
Tomasz Stokowy, PhD	tomasz.stokowy@uib.no
Nijole Pollock, BSc	Nijole.pollock@gmail.com
Andrea J. Berman, PhD	AJB190@pitt.edu
Sylwia Budkiewicz, MSc	sylwia.budkiewicz@wp.pl
Marzena Kamieniczna, PhD	marzena.kamieniczna@igcz.poznan.pl
Hanna Jackowiak, PhD	hanna.jackowiak@up.poznan.pl
Piotr Jedrzejczak, PhD, MD	piotrjedrzejczak@gmail.com
Alexander N. Yatsenko, PhD, MD	yatsenkoan@mwri.magee.edu

ORCiD:

Marta Olszewska, PhD	0000- 0002- 2337- 4545
Agnieszka Malcher, PhD	0000- 0002- 3433- 6990
Tomasz Stokowy, PhD	0000-0003-0017-8338
Nijole Pollock, BSc	-
Andrea J. Berman, PhD	0000-0003-1217-7412
Sylwia Budkiewicz, MSc	-
Marzena Kamieniczna, PhD	0000-0002-9619-8220
Hanna Jackowiak, PhD	0000-0001-5167-8215
Piotr Jedrzejczak, PhD, MD	0000-0003-2437-7737
Alexander N. Yatsenko, PhD, MD	0000-0002-1690-0172
Maciej Kurpisz, PhD, MD	0000-0003-3275-3245

Abstract

STUDY QUESTION: Is *Tcte1* mutation causative for male infertility?

SUMMARY ANSWER: Collected data underline the complex and devastating effect of the single-gene mutation

on testicular molecular network, leading to male reproductive failure.

WHAT IS KNOWN ALREADY: Latest data revealed mutations in genes related to axonemal dynein arms as

causative for morphology and motility abnormalities in spermatozoa of infertile males, including dysplasia of

fibrous sheath (DFS) and multiple morphological abnormalities in the sperm flagella (MMAF). The nexin-dynein

regulatory complex (N-DRC) coordinates the dynein arm activity, and is built from DRC1-DRC7 proteins. DRC5

(TCTE1) – one of N-DRC element, has been already reported as a candidate for abnormal sperm flagella beating,

however, only in restricted manner with no clear explanation of respective observations.

STUDY DESIGN, SIZE, DURATION: Using CRISPR/Cas9 genome editing technique, mouse knockout line of *Tcte1*

gene was created on the basis of C57Bl/6J strain. Then, the mouse reproductive potential, semen characteristics,

testicular gene expression level, sperm ATP and testis apoptosis level measurements have been performed,

followed by visualization of N-DRC proteins in sperm, and protein modeling *in silico*. Also, a pilot genomic

sequencing study of samples from human infertile males (n=248) was applied for screening of *TCTE1* variants.

PARTICIPANTS/MATERIALS, SETTING, METHODS: To check the reproductive potential of KO mice, adult animals

were crossed for delivery of three litters per caged pair, but no longer than for 6 months, in various combinations

of zygosity. All experiments were performed for wild type (WT – control group), heterozygous *Tcte1*$^{+/-}$, and

homozygous *Tcte1*$^{-/-}$ male mice. Gross anatomy was performed on testis and epididymis, followed by semen

analysis. Sequencing of RNA (RNAseq; Illumina) has been done for mice testis tissues. STRING interactions have

been checked for protein-protein interactions, based on changed expression level of corresponding genes

identified in the mouse testis RNAseq experiments. Immunofluorescence *in situ* staining was performed to detect

the N-DRC complex proteins: Tcte1 (Drc5), Drc7, Fbxl13 (Drc6), and Eps8l1 (Drc3) in mouse spermatozoa. To

determine the ATP amount in spermatozoa, the luminescence level was measured. Also, immunofluorescent *in*

situ staining was performed to check the level of apoptosis via caspase 3 visualization on mouse testis samples.

DNA from whole blood samples of infertile males (n=137 non-obstructive azoospermia or cryptozoospermia,

n=111 samples with spectrum of oligoasthenoteratozoospermia, including n=47 with asthenozoospermia) has

been extracted to perform genomic sequencing (WGS, WES or Sanger). Protein prediction modeling of human

identified variants and the exon 3 structure deleted in mouse knockout has been also performed.

MAIN RESULTS AND THE ROLE OF CHANCE: No progeny at all was found for homozygous males with revealed

oligoasthenoteratozoospermia, while heterozygous animals (fertile) manifested oligozoospermia, suggesting

haploinsufficiency. RNA-sequencing of the testicular tissue showed the influence of *Tcte1* mutations on the

expression pattern of 21 genes responsible for mitochondrial ATP processing, linked with apoptosis, or

spermatogenesis. In *Tcte1*[-/-] males the protein revealed only residual amounts in sperm head nucleus, and was

not transported to sperm flagella, as other N-DRC components. Decreased ATP level (2.4-fold lower) was found

in spermatozoa of homozygous mice, together with disturbed tail:midpiece ratio, leading to abnormal sperm tail

beating. Casp3-positive signals (indicating apoptosis) were observed in spermatogonia only, at similar level in all

three mouse genotypes. Mutation screening of human infertile males revealed 1 novel and 5 ultrarare

heterogeneous variants (predicted as disease causing) in 6.05% of patients studied. Protein prediction modeling

of identified variants revealed changes in the protein surface charge potential, leading to disruption in helix

flexibility or its dynamics, thus, suggesting the disrupted TCTE1 interaction with its binding partners located

within the axoneme.

KEYWORDS

TCTE1, *DRC5*, N-DRC, mouse knockout model, male infertility, asthenozoospermia,
oligoasthenoteratozoospermia, oligozoospermia, spermatogenesis, sperm motility, sperm mitochondria,
mitochondrial respiratory complex, *mt-Co2*, *Fetub*, haploinsufficiency, MMAF, apoptosis

INTRODUCTION

Infertility has been defined by World Health Organization as a social disease concerning approximately

10-18% of couples at reproductive age, which are unable to conceive within 1 year of regular unprotected sexual

intercourse [WHO, 2020]. Approximately 7% of males and ~12% of females worldwide reveal reproductive

problems, while the male factor is responsible for approximately 20-30% of all infertility cases [Barratt *et al.,*

2017; Vander Borght and Wyns, 2018; Agarwal et al., 2021]. Male infertility is a disease with multifactorial

etiology, including: genetic factors, chromosomal aberrations, epigenetic mutations, hormonal abnormalities,

infections or reproductive tract abnormalities [Agarwal et al., 2021; Matzuk and Lamb 2008; Zorrilla and Yatsenko

2013; Krausz and Riera-Escamilla 2018]. Genetic causes of male infertility can be detected in ~10-15% of cases,

while in 25-50% of infertile males the etiology remains unexplained [Agarwal *et al.,* 2021; Matzuk and Lamb

2008; Zorrilla and Yatsenko 2013]. The male infertility phenotype mostly results in decreased semen parameters,

including: low sperm concentration, abnormal morphology or diminished motility. The actual standards for

genetic testing of male infertility rely on chromosomal screening for structural and numerical abnormalities,

followed by analyses of deletions in the AZF (azoospermia factor) regions of the Y chromosome [Hwang et al.,

2018]. However, it is still not sufficient due to the fact, that approximately 2,000 genes may be related to

spermatogenesis, while only ~10% of them seems to have documented fertility role, so far [Krausz and Riera-

Escamilla 2018; Houston *et al.,* 2021] Recent years NGS technology development, including genomic sequencing

of DNA (exome or genome; large cohorts of patients), and RNA (RNAseq, scRNAseq on reproductive tissues),

followed by animal models of infertility (mostly mouse models; more than 860 genes evaluated, so far [Mouse

Genome Informatics, 24 June 2021]) have started the new era of exploration on the massive scale [Houston *et*

al., 2021; Oud *et al.,* 2021; Wyrwoll *et al.,* 2020; Hardy *et al.,* 2021; Xavier *et al.,* 2021].

Beside the genes related to spermatogenesis and low sperm count, also studies of candidates

responsible for diminished sperm motility are pending. Thus, the main structural element responsible for motility

– the sperm flagella, has to be of high-quality to guarantee sperm motility and then fertilization success. It is

known that mutations in genes encoding factors involved in ciliogenesis process, lead to defects in structure

and/or functioning of cilia [Girardet *et al.,* 2019; Sironen *et al.,* 2020; Smith *et al.,* 2021; Kumar and Reiter 2021]

There is a plenty of data concerning respiratory cilia, however, still little is known about the axonemal dynein

arms functioning in sperm tail. Latest data revealed mutations in genes related to axonemal dynein arms as

causative for morphology and motility abnormalities in spermatozoa of infertile males [Girardet *et al.,* 2019;

Sironen *et al.,* 2020; Aprea *et al.,* 2021; Lorès *et al.,* 2021; Touré *et al.,* 2021; Martinez *et al.,* 2022]. Two

definitions were established: (i) for sperm cells with disorganization in axonemal structures, so called dysplasia

of fibrous sheath (DFS) [Chemes *et al.,* 1998] and (ii) for multiple morphological abnormalities in the sperm

flagella (MMAF), concerning all possible defects of sperm tail, mainly short and thick flagella, and a lack of the

central pair (CP) of microtubules or dynein arms [Khelifa *et al.,* 2014]. Spectrum of abnormal sperm tail

phenotypes observed allows qualification of a part of disorders into both listed groups (DFS and MMAF) [Sironen

et al., 2020; Touré *et al.,* 2021]

The basic structural element of motile cilia and flagella is axoneme built from nine microtubules

arranged in doublets positioned radially and one central pair (CP) of singlet microtubules. This central pair is

connected to radial spokes (RS) – protein complexes of mechanic regulatory network responsible for regular

waveform beating of the cilia [Gui *et al.,* 2019]. RS are linked to inner dynein arms (IDA) localized within each

outer dublet. Each outer doublet carries outer dynein arms (ODA) – multi-subunit motor protein complexes,

responsible for generation and regulation of sliding between dynein arms leading to ATP-dependent beating

[Satir *et al.,* 2014; Wilson et al., 2015]. The 9+2 basic structure is evolutionary conservative [Girardet *et al.,* 2019;

Touré *et al.* 2021; Ishikawa 2017]. Doublets are linked with the nexin-dynein regulatory complex (N-DRC), which

coordinates the dynein arm activity and stabilizes the attachment of doublet microtubules. The N-DRC is built

from DRC1-DRC7 proteins, among which DRC1, DRC2 and DRC4 are responsible for physical linkage between A

and B microtubules of doublets [Gui *et al.,* 2019; Morohoshi *et al.,* 2020; Bowar *et al.,* 2018; Wirschell *et al.,*

2013]. In terms of male infertility, the best studied gene among N-DRC is *DRC7* (MIM:618769), which is amenable

for proper assembly of the N-DRC structure (in *Drc7*$^{-/-}$ male mice the length of the sperm tail was reduced,

followed by abnormalities of sperm head morphology) [Morohoshi *et al.,* 2020]. Mouse model of *Drc6* (*Fbxl13*)

gene (MIM:609080) revealed no essential role in sperm formation and motility [Morohoshi *et al.*, 2020],

however, its disruption together with CEP192 (MIM:616426) leads to defects in HEK293T cell motility [Fung et

al., 2018]. Mutations in *Drc3* gene (MIM:618758) indirectly revealed male infertility as a lack of pregnancy success

with wild-type females [Ha et al., 2016].

One of the N-DRC structural genes is *TCTE1* (T-Complex-associated-Testis-Expressed 1), known also as

DRC5, FAP155, *D6S46* in humans (*locus* 6p21.1; MIM:186975) or *Tcte1, D17S*, *Tcte-1*, *D17Sil1* in mouse (*locus* 17

B3; 17 22.54 cM; MGI:98640). *TCTE1* is evolutionary conserved in most eukaryotic organisms, that contain

motility cilia or flagella in their life cycle (Figure 1), and is required for proper flagella functioning. Expression of

TCTE1 is being observed in testis (highest; with expression starting at the stage of spermatid), choroid plexus and

Fallopian tube. TCTE1 contains five leucine rich repeats (LRRs). To date, only one report described a role of *Tcte1*

gene in infertility, where the sperm cells of *Tcte1-null* mice exhibited asthenozoospermia [Castaneda et al., 2017].

However, understanding of the nature of this specific motility type defect in *Tcte1-null* animals has been limited.

In our study we aimed to determine the role of *Tcte1* in male infertility using mouse knockout model.

We created homo- and heterozygous animals, which were next examined for their reproductive potential,

differences in anatomy and histology, and semen parameters. We also performed RNA sequencing (RNAseq) in

testis tissue to obtain an answer about the influence of *Tcte1* null mutation on the expression pattern of other

gonadal genes. Also, the ATP measurement of spermatozoa has been performed. Additionally, screening of

variants in human samples from males with severely decreased sperm count was made, followed by protein

prediction modeling of revealed variants. Thus, this study contains novel and the most complete data concerning

TCTE1 role in male infertility, so far.

MATERIALS AND METHODS

Animals

Using CRISPR/Cas9 genome editing technique, mouse heterozygous knockout line of *Tcte1* gene

(*Tcte1$^{+/}$*) was created on the basis of C57Bl/6J strain in the Cyagen Biosciences company (USA). The mouse *Tcte1*

gene is located on chromosome 17 B3; 17 22.54 cM (GenBank: NM_013688.2; Ensembl:

ENSMUSG00000023949). Exon 3 of the *Tcte1* gene was selected as target site (deletion of 1292 bp) (Figure 2A).

gRNAs targeting vectors (constructed, sequenced and generated by *in vitro* transcription) were injected into

fertilized oocytes (Supplementary file 1). The founders were genotyped and those with positive results were bred

to the next generation. Then, in the animal facility of the Institute of Human Genetics PAS (Poznan, Poland), adult

Tcte1$^{+/-}$ animals were mated to generate homozygous *Tcte1$^{-/-}$* mice. Wild type *Tcte1$^{+/+}$* (WT) animals C57Bl/6J

were purchased from Charles River company (Germany). Relative expression of *Tcte1* gene was verified using

Real-time PCR method (Figure 2B; Supplementary file 2). All mice were housed in pathogen-free animal facility

with food and water *ad libitum*, stable temperature of 22^{o}C, and a light:dark light cycle of 12:12. All procedures

were in accordance with the guidelines and regulations for animal experimentations and restricted usage of

genetically modified organisms (GMO), and were approved by the Local Ethical Committee for Animal

Experiments at the Poznan University of Life Sciences (no. 1/2014, 18/2017), and Ministry of Environment (no.

70/2017). Mouse models with infertility phenotypes were reviewed in the Mouse Genome Informatics database

(Jackson Laboratory; http://www.informatics.jax.org/) and through literature searches (PubMed;

https://pubmed.ncbi.nlm.nih.gov/).

Reproductive potential

To check the reproductive potential of KO mice, adult animals (7-8 weeks old) were crossed for delivery

of three litters per caged pair, but no longer than for 6 months. Crossing combinations included: (i) 4 pairs:

homozygous *Tcte1*$^{-/-}$ male × wild type (WT) female, (ii) 5 pairs: heterozygous *Tcte1*$^{+/-}$ male × heterozygous *Tcte1*$^{+/-}$

female, and (iii) 3 pairs: heterozygous *Tcte1*$^{+/-}$ male × wild type (WT) female. Vaginal plug presence was checked

every morning to confirm mating incidence. Time period from mating start point to first pedigree appearance

was checked. Number of pups was counted on the day after birth. For each crossed pair, the number of litters,

pups, and sex ratio were estimated (also according to pups' genotypes) (Supplementary file 3).

Genotyping

Genotyping of the pups delivered was performed via PCR assay, with two sets of primers: (i) to identify

wildtype and positive KO (het- and homozygous) animals: Tcte1-F: CGTTTTAAGAATGTATTGAGGGTTGGG, Tcte1-

R: CTCCGTAGGCTCCTGCCAATATG; estimated product size: for WT 1915 bp, for KO ~630 bp (deleted ~1300 bp);

PCR results analysis: one band with 1915 bp or 0 bp determined WT, while one band with 630 bp evidenced

hetero- or homozygotes; (ii) to distinguish between hetero- and homozygotes: Tcte1-WT/He-F:

AGCTTGCCACACCCTCAAGGTACTA, Tcte1-R: CTCCGTAGGCTCCTGCCAATATG; estimated product size: 658 bp for

heterozygotes and 0 bp for homozygotes; PCR results analysis: one band with 658 bp determined heterozygotes,

while no one band (0 bp) indicated homozygotes (Figure 2C). The PCR mixture and reaction conditions are

described in Supplementary file 4.

Tissue gonadal preparation/collection

Gross anatomy was performed on testis and epididymis of WT (n=21), *Tcte1*$^{+/-}$ (n=22), and *Tcte1*$^{-/-}$ (n=18)

male animals (10 weeks old). Mice were weighted, then euthanized with carbon dioxide (CO_2) overdose

(euthanasia Minerve Easy-box; Minerve; France) with stable flow rate per minute. Testes and epididymises were

collected, weighted, and measured. For histological evaluation tissues were immersed into 10% formalin

solution, neutral buffered (Sigma Aldrich, USA) for 3 males per zygosity. Then, Paraplast-embedded sections were

prepared (5 μm thickness), using Microtome Leica 2055. To obtain the histological picture, slides were stained

with Masson-Goldner protocol [Romeis et al., 2010]. Thereafter, slides were documented with Leica DM5500

light microscope (×1000 magnification, oil immersion).

Mouse semen samples

Mouse spermatozoa were collected from epididymides *post mortem*. Collected epididymides were

placed immediately in a 2.0 mL test tube with warm 500 μL of a non-capacitating Whitten's HEPES medium (pH

7.2-7.4, 37°C), then cutted into 5-7 fragments, and incubated in 37°C for 15 min. to allow the sperm cells to swim

out. Whittens-HEPES medium contained: 100.0 mM NaCl, 4.4 mM KCl, 1.2 mM KH2PO4, 1.2 mM MgSO4, 5.4 mM

glucose, 0.8 mM pyruvic acid, 4.8 mM lactic acid (hemi-Ca), HEPES 20 mM [Wertheimer et al., 2013]. Then,

spermatozoa were collected and proceeded with seminological analysis (concentration, motility, morphology).

Next, semen samples were fixed in a fresh fixative solution (methanol:acetic acid, 3:1 v/v, -20°C; 3×) and stored

until further use in -20°C.

Semen analysis

10 μl of sperm suspension was applied on the Makler counting chamber (Sefi medical instrument Ltd.,

Israel) to test the sperm concentration and total sperm count. Also, the ratio of progressive (fast and slow), non-

progressive (total and circular) and immotile spermatozoa were calculated [WHO, 2020]. For assessment of

sperm morphology, the Papanicolaou staining was performed. Briefly, the air-dried smears were fixed in

ethanol:diethyl ether (1:1 ratio, vol/vol) solution for 10 min. and left for air-drying. Then, slides were stained by

Papanicolau procedure according to WHO guidelines [WHO, 2020], and mounted in DPX Mountant (Sigma-

Aldrich; USA; cat. no. 06522). A total of 200 spermatozoa were analyzed using Leica DM5500 light microscope

(×1000 magnification, oil immersion). Various morphological forms of spermatozoa were determined including

abnormalities of head, midpiece and tail. Also, the length of sperm tail, midpiece and the tail:midpiece ratio were

measured to check possible differences between males with various zygosity (6 males/per zygosity); for each

male at least 100 spermatozoa were measured with the 'length measurement' option by CellSense Dimension

software (Olympus).

Sequencing of RNA (RNAseq)

For RNA extraction testis tissue was collected into in RNAlater buffer (Invitrogen; Waltham, MA, USA)

and freezed immediately (-80°C). Samples were processed using the AllPrep DNA/RNA/Protein Mini Kit (Qiagen;

Hilden, Germany). Quality and quantity of extracted RNA samples were checked using NanoDrop (Thermo

Scientific) and Quantus (Promega), and 1 μg of RNA per sample has been used for sequencing (Macrogen; South

Korea). RNA sequencing was performed with Illumina NovaSeq 6000 platform using TruSeq Stranded mRNA LT

Sample Prep Kit, with specified conditions of paired-end reads at 100 bp, and throughput 100 M reads per

sample. RNA-Seq raw reads were aligned to mouse reference genome mm10 using hisat 2.0.5 with Gencode

vM13 transcriptome reference. Reads aligned in the coding regions of the genome were counted using Feature

Counts. Then, read counts were normalized using DESeq2 and subjected to differential analysis. The differential

analysis included computing of median based fold change, statistical significance using the Students' t-test and

false discovery rates (correction for multiple testing) in the R/Bioconductor programming environment. Testis

tissue expression was retrieved from GTEx (https://gtexportal.org/home/), BioGPS (http://biogps.org/), and

AceView (https://www.ncbi.nlm.nih.gov/IEB/Research/Acembly/). All RNA-seq expression data have been

deposited in Gene Expression Omnibus (GEO), accession number GSE207805.

STRING interactions

For analysis of protein-protein interactions, based on changed expression level of corresponding genes

identified in the mouse testis RNAseq experiments, the STRING analysis was performed [version 11.0]. STRING

database allows to delineate probability of the interactions between evaluated proteins using different evidence

channels [Mering *et al.,* 2005; Szklarczyk *et al.,* 2019]. This tool enables the integration of all data available in

public resources concerning protein-protein interactions.

Immunofluorescence on sperm – N-DRC visualization

Immunofluorescence *in situ* staining was performed to detect the N-DRC complex proteins: Tcte1 (Drc5),

Drc7, Fbxl13 (Drc6), and Eps8l1 (Drc3) in mouse spermatozoa. Mouse spermatozoa were collected and fixed as

described above. Specific antibodies were applied: primary antibodies – rabbit anti-human-TCTE1, rabbit anti-

mouse-Drc7, anti-mouse-Fbxl13, and anti-mouse-Eps8l1 (all from Biorbyt; Cambridge, UK); secondary antibody

– goat anti-rabbit-AF594 (Abcam; cat. no. ab150160; Cambridge, UK). The UniProt IDs and homology of the

immunization regions of primary antibodies between human and mouse were described in Supplementary file

5. Antibodies were diluted in 0.5% PBST (primary: 1:100, secondary: 1:500). First, slides with fixed sperm smears

were washed in a series of washes in 1× PBST, and then incubated in 25 mM DTT/1 M Tris-HCl, pH 9.5, at room

temperature for 20 min. Then, washing in 1× PBST, and incubation in 6 N HCl for 30 min. was applied, followed

by blocking step with 1% BSA/1× PBST for 30 min. Next, overnight incubation with antibodies was performed at

4°C in a humidified container. After washing, the samples with 1× PBST, secondary AF594-conjugated antibody

was applied for 1 hr. Next, unconjugated antibodies were washed out (4× in 1× PBST, 5 min each, slight rotation).

For the final detection, 20 ul of Fluoroshield with DAPI (Sigma-Aldrich; cat. no. F6057) was added to the samples,

and microscopic analysis was performed. Images were acquired using a fluorescence microscope with a proper

filter set: Leica DM5500, filters: DAPI/TxR/Triple; objectives: 10× and 63× with oil immersion; DFC 7000T camera;

software: LASX.

Sperm ATP measurement

For measurement of ATP in spermatozoa the CellTiter-Glo 2.0 Assay (Promega; USA) was used. This

ready-to-use assay allows to determine the quantitation of the ATP amount, indicating the presence of

metabolically active cells. The assay is based on the thermostability of luciferase (Ultra-GloTM Recombinant

Luciferase), and was performed as described in manufacturers' protocol. Briefly, all laboratory work was done in

the stable temperature of 22°C. After collection of mouse spermatozoa (as described above), the amount of

approximately 1×10^6 of sperm cells was transferred to black opaque 1.5 mL test tube and proper amount of

CellTiter-Glo kit was added (1:1, v:v). Next, 2 min. of orbital mixing of the content was performed to activate cell

lysis. Then, after 10 min. of incubation (to stabilize the luminescence), the measurement of luminescent signal

was completed using Glomax 20/20 luminometer (Promega; USA). For each sample, three measurements

(repeats) were done, each in two time points of 0 and +60 min. Numbers of animals used were: WT (n=7), *Tcte1*$^{+/-}$

(n=7), and *Tcte1*$^{-/-}$ (n=6). Mean luminescence values obtained (represented as RLU – relative luminescence unit)

were compared between tested mice groups.

Immunofluorescence on testis – apoptosis evaluation

Immunofluorescence *in situ* staining was performed to check the level of apoptosis via caspase 3

visualization on mouse testis samples from all the three zygosities. Caspases play important role in the induction,

transduction, and amplification of intracellular apoptotic signals, and thus, caspase-3 can be used as one of the

apoptotic indicators [Lej et al, 2022]. Mouse testes were collected and fixed as described above. Specific

antibodies were applied: primary antibody – rabbit anti-human-Casp3, (Biorbyt; Cambridge, UK), diluted 1:100

in 0.5% PBST; secondary antibody – goat anti-rabbit-AF594 (Abcam; Cambridge, UK), diluted 1:500 in 0.5% PBST.

First, slides with paraffine fixed testis cross sections were deparaffinized in 3 changes of xylene (10 mins. each),

and incubated in a series of ethanol: 100%, 96%, and 70%, twice for 5 mins. in each solution. Next, after rinsing

in distilled water, tissue sections were incubated sequentially in: a freshly prepared 2% $NaBH_4$/1× PBS, and in

0.1M glycine (30 min., room temp. each). Heat-induced epitope retrieval was performed by immersing the slides

in a 0.01M sodium citrate/0.05% Tween 20, pH 6.0 for 30 min., followed by cooling down in a room temp. for

next 20 min. Blocking step was done in an aliquot of freshly prepared 1% BSA/1× PBST, at 37 °C for 1 h. Then,

tissue sections were incubated with primary antibody in a humidified container overnight at 4°C. Secondary

antibody was applied for 1 hr, followed by the washing out of the unconjugated antibody (4× in 1× PBST, 5 min

each, slight rotation). For the final detection, 20 μl of Fluoroshield with DAPI (Sigma-Aldrich; cat. no. F6057) was

added. Images were captured using a fluorescence Leica DM5500 microscope with a proper filter set:

DAPI/SpO/Triple; objectives: 10× and 63× with oil immersion; DFC 7000T camera, LASX software. The analysis

was performed for three independent males per zygosity, 2 cross-sections each time, 15-20 tubules per tissue

section.

Sequencing of the human samples

A pilot genomic sequencing study of samples from human infertile males was also applied. DNA from

whole blood samples (collected into sterile single test tubes with EDTA) was extracted using standard protocols

of MagCore HF16 Automated Nucleic Acid Extractor (RBC Biosciences, Taiwan), or Gentra Puregene kit (Qiagen,

USA). Total number of sequenced patients was n=248, including: 137 with non-obstructive azoospermia (lack of

spermatozoa in ejaculate) or cryptozoospermia (spermatozoa detectable in ejaculate after centrifugation),

followed by 111 samples with spectrum of seminal abnormalities in concentration, and/or motility, and/or

morphology. Among those samples, 47 revealed decreased motility (asthenozoospermia). Semen evaluation was

done according to WHO guidelines [WHO, 2020], followed by AUA and ASRM guidelines for infertile male

diagnosis [Schleger et al., 2021]. All men revealed normal hormonal levels, had no chromosomal abnormalities

and a lack of AZF microdeletions. For whole exome samples (n=76 from Pittsburgh), sequencing libraries were

constructed using the Agilent technologies (Santa Clara, USA), including: SureSelect XT All Exon V4 + UTRs capture

kit and SureSelect XT Target Enrichment System. Libraries were sequenced with average read coverage of 62-

100x (~6Gb per sample), using 100 bp paired-end sequencing with TruSeq PE Cluster Kit v3 cBot HS and 4 × 50

TruSeq SBS V3 HS sequencing kit (Illumina, San Diego, USA) on the HiSeq2000 (Illumina, San Diego, CA). Whole

genome sequencing (n=42, Poznan) was performed using Illumina HiSeqX with the coverage of at least 30x (100-

120 Gb per sample). Sanger sequencing for all exomes of *TCTE1* gene (n=130, Poznan) was performed using

LightRun96 system of Eurofins Genomics (Eurofins Genomics AT GmbH, Vienna, Austria). Samples were prepared

with QIAquick Gel Extraction (cat. no. 28704) or PCR Purification (cat. no. 28104) kits (Qiagen, USA), and analyzed

with CLC Main Workbench 8 software (Qiagen, USA).

Filtering of whole genome sequencing results

The quality of raw NGS data was evaluated using the FastQC and MultiQC packages. Reads were aligned to the

reference human genome GRCh37 using bwa-mem aligner included in the Speedseq package [Chiang et al.,

2015]. For calling of single nucleotide variants (SNVs) and small indels the freeBayes v0.9.21 (speedseq-var) was

implemented, following previously described pipeline [Supernat *et al.,* 2018]. Variant disease association and

missense mutation prediction effects were determined using: dbSNP (https://www.ncbi.nlm.nih.gov/snp/),

OMIM   (https://www.ncbi.nlm.nih.gov/omim/),   ClinVar   (https://www.ncbi.nlm.nih.gov/clinvar/),

MutationTaster   (http://www.mutationtaster.org/),   SIFT   (https://sift.bii.a-star.edu.sg/),   Polyphen

(http://genetics.bwh.harvard.edu/pph2/),   CADD   (https://cadd.gs.washington.edu/snv),   MetaDome

(https://stuart.radboudumc.nl/metadome/), REVEL (rare exome variant ensemble learner) [Ioannidis et al.,

2016], Ensembl Variant Effect Predictor [Mclaren et al., 2016], and Human Gene Mutation Database (HGMD;

Qiagen, Redwood City, CA). SNV pathogenicity was assessed on the basis of American College of Medical Genetics

and Genomics (ACMG) guidelines [Richards et al., 2015]. PhyloP (Cold Spring Harbor Laboratory) and Clustal

Omega were used for determination of conservation (https://www.ebi.ac.uk/Tools/msa/clustalo/). All NGS

variants were confirmed via the Integrative Genomics Viewer (Broad Institute). Variant frequencies have been

obtained from the gnomAD database (frequencies from more than 125k exomes and 15k genomes,

http://gnomad.broadinstitute.org; v2.1.1), and were filtered in search of rare variants (with minor allele

frequency (MAF) <1%).

Protein prediction

The amino acid sequence of TCTE1 was input into Phyre2 [Kelley et al., 2015] for both intensive and fast

mode homology modeling, I-TASSER [Roy *et al.,* 2010; Yang *et al., 2*015, Yang and Zhang 2015], and AlphaFold

[Jumper *et al.,* 2021; Varadi *et al.,* 2022] which utilize sequence alignment-based algorithms for generating

predicted secondary and tertiary structures; AlphaFold additionally uses machine learning to integrate biological

principles of protein folding with the alignment-based algorithms. Confidence scores were obtained from these

programs; AlphaFold additionally generates predicted aligned error plots for assessing the quality of the model's

long-range interactions. Models were inspected manually and aligned using the secondary structure matching

(SSM) [Krissinel and Henrick, 2004] tool in COOT [Emsley *et al.,* 2010] to minimize user input bias. Docking of the

structures into electron tomography data was performed manually in COOT [Emsley *et al.,* 2010]. No refinement

was performed to minimize bias. All structure model figures and mutations without post refinement, were

generated using Pymol (Schrödinger). I-TASSER was used also for prediction of the exon 3 structure deleted in

our mouse KO model (Supplementary file 6).

375   *Statistical analysis*

376   Statistical analyses included the following: normality test (D'Agostino and Pearson or Kolmogorov-

377   Smirnoff), unpaired two-tailed t-test with Welch's correction, one-way ANOVA test and Fisher's exact, and two-

378   tailed Pearson correlation. All tests were performed with a significance level of α=0.05 using GraphPad Prism

379   (v.7.0e) software.

381   **RESULTS**

382   *Reproductive potential of KO mice*

383   Reproductive potential results were shown in Figure 3 and Supplementary file 3. Only pairs with

homozygous *Tcte1*[-/-] males revealed no progeny (*Tcte1*[-/-] male x WT female). Other combinations (*Tcte1*[+/-] male

× *Tcte1*[+/-] female; *Tcte1*[+/-] male × WT female) showed similar results, when considering: number of litters, pups,

and sex ratio. Mean time of caging to delivery of the 1st pedigree was longer in *Tcte1*[+/-] male × *Tcte1*[+/-] female

combination by 25.45% (statistically not significant). When considering genotypes of the pups, statistically

significant differences (p<0.05) were observed in the number of *Tcte1*[-/-] male and female pups delivered – no

homozygous animals were obtained from combination of *Tcte1*[+/-] male × WT female, while in mating of *Tcte1*[+/-]

male × *Tcte1*[+/-] female, homozygous males (33.25% of all pups) and females (24.67%) were received.

391   *Histopathology and seminal parameters*

392   Histopathological evaluation of testicular and epididymal sections revealed no differences between WT,

393   *Tcte1*[+/-], and *Tcte1*[-/-] males (Figure 4A). No odds were found for body mass of the animals and the length of the

394   epididymis (Figure 4B). In the *Tcte1*[-/-] males testis weight was decreased by approximately 41% (p<0.0001)

395   followed by the ~22% decline of its size (p<0.0001), while in *Tcte1*[+/-] males only testis weight was slightly

396   decreased (approx. 7%; p<0.05) when compared to control animals.

397   When considering seminal parameters, both: in *Tcte1*[-/-] as well as in *Tcte1*[+/-] males the sperm

concentration was importantly decreased (3.27, and 2.47-fold, respectively), when collating to control results

(p<0.0001; Figure 4C). Additionally, in *Tcte1*[-/-] males the frequency of progressively motile spermatozoa was

residual, with progressive type of motility in only 0.22% of spermatozoa (decreased ~150-fold vs. control 33.52%,

(p<0.0001)), followed by 4.89% of slow progressive spermatozoa (decreased ~3-fold vs. control 15.50%)

(p<0.0001). Interestingly, ~2.4-fold increase in circular type of motility (around own axis and non-progressive)

was observed for $Tcte1^{-/-}$ males when compared to WT and $Tcte1^{+/-}$ animals (p<0.01) (Supplementary file 7;

Supplementary file 8; Supplementary file 9). The frequency of immotile spermatozoa also increased in $Tcte1^{-/-}$

males (2-fold to control; p<0.0001). The frequency of morphologically normal spermatozoa in $Tcte1^{-/-}$ animals

was significantly decreased (2-fold to control; p<0.0001; Figure 4C, D, E; Supplementary file 10). The majority of

morphological defects was observed for sperm head, mostly of decapitated spermatozoa in $Tcte1^{-/-}$ animals –

52.55% vs. 19.16% for $Tcte1^{+/-}$ and 23.33% for WT, and amorphous head shape with decreased frequency in

$Tcte1^{-/-}$ (26.11%) and $Tcte1^{+/-}$ (33.72%) males vs. WT animals (48.88%) (Supplementary file 10). Midpiece defects

were found in $Tcte1^{+/-}$ (9.71%) and WT (8.20%) animals, while $Tcte1^{-/-}$ mice revealed only low frequency (0.38%)

of such defects. On the other hand, an increased frequency (~5-fold) of coiled tail was observed for $Tcte1^{+/-}$

animals when compared to other mice zygosity (p<0.01) (Supplementary file 10). Summarizing, seminological

changes in mice with $Tcte1$ mutation resulted in oligoasthenoteratozoospermia in homozygous $Tcte1^{-/-}$ or

oligozoospermia in heterozygous $Tcte1^{+/-}$ animals.

Increased mean length of tail and tail:midpiece ratio in $Tcte1^{-/-}$ animals (139.30 ± 7.71 um; 5.16,

respectively) was measured when comparing to WT (122.00 ± 6.375 um; 4.51; p<0.002) and $Tcte1^{+/-}$ (123.50 ±

1.89; 4.83; p<0.008) (Figure 4F). Also, in the $Tcte1^{+/-}$ the midpiece length differed from $Tcte1^{-/-}$ animals

(p=0.0334), followed by tail:midpiece ratio differences when compared to WT (p=0.0435).

Sequencing of RNA (RNAseq) and STRING pathway analysis

RNA sequencing analysis of gene expression in testes of KO mice revealed statistically significant

changes (p<0.05) in 21 genes, including 5 genes both for $Tcte1^{-/-}$ vs. $Tcte1^{+/-}$ and WT, 8 genes only for $Tcte1^{-/-}$ vs.

WT, 7 genes only for $Tcte1^{-/-}$ vs. $Tcte1^{+/-}$, and 1 gene both for $Tcte1^{+/-}$ vs. $Tcte1^{-/-}$ and WT (Figure 5). In $Tcte1^{-/-}$

animals vs. both: $Tcte1^{+/-}$ and WT, the expression level of 4 genes from kallikrein subfamily was decreased 2-8-

fold, including genes involved in semen liquefaction and degradation of extracellular matrix proteins in the

interstitial area surrounding Leydig cells of the adult mouse testes (*Klk1b27*, *Klk1b21*, *Klk1b24*), or neuronal

activity (*Klk1b22*) (Figure 5; Supplementary file 11) [Michaelis *et al.,* 2017; Marques *et al.,* 2016; Du Plessis *et al.,*

2013; Fink *et al.,* 2007]. 2-fold decrease of expression of genes related to mitochondrial cytochrome oxidase

subunits and asthenozoospermia was documented (*mt-Co2*, *mt-Co3*) (Figure 5) [Heidari *et al.,* 2016; Mostafa *et*

al., 2016; Baklouti-Gargouri *et al.,* 2013]. In a scheme: $Tcte1^{-/-}$ vs. WT, 8 genes revealed changes in expression

level, including 4.7-fold rise for *Fetub* gene, which is known as required for egg fertilization (via specific inhibitor

activity of the *zona pellucida* (ZP) ovastacin, the cysteine protease responsible for ZP2 cleveage) [Binsila *et al.,*

2021; Karmilin *et al.,* 2019], and plays also a role of a secreted factor during reprogramming [Bansho *et al.,* 2017].

Two-fold rise for *Igf2* gene (strongly linked to spermatogenesis, sex determination, testis development,

infertility and prostate cancer; [Matsuzaki et al., 2020; Cannarella et al., 2020; Neirijnck et al., 2019; Damaschke

et al., 2017]) has been also noticed. Igf2 and Klk proteins are enclosed in a common network consisted of proteins

directly linked to fertility (Figure 6; Supplementary file 11; Supplementary file 12). One of them is Ambp, a fusion

protein, which proteolysis generates bikunin – a protein which deficiency results in infertility, due to the

diminished formation of the stable cumulus-oocyte complex required for maturation and ovulation [Sato and

Kan,2001; Naz and Rajesh, 2005]. Bikunin is one of the elements of ITIH domains (Inter-α-Trypsin Inhibitor Heavy-

chain), responsible for hyaluronian metabolism [Hull *et al.,* 2012; Salustri *et al.,* 2019]. ITIH is also linked to Plg,

which has the protease activity and activates metalloproteins, such as: Mmp8 and Mmp9, also included in the

network. Mmp8 is engaged in degradation/stabilization of occluding and blood-testis barrier [Chen *et al.,* 2018],

while Mmp9 (with Igf1 and Igf2), is important for normal placentation [Ghosh *et al.,* 2011]. Igf1 and Igf2, are

related to Igfbp3, which interacts with humanin expressed in Leydig cells. Humanin together with IGFBP3 and

Bax prevents the activation of germ cell apoptosis [Jia et al., 2015; Lue et al., 2021]. What is important, expression

of *IGFBP3* is regulated by *HOXD10* gene [Xue *et al.,* 2013; Pan *et al.*, 2021], of which expression has been found

with 2.9-fold rise in our study. *Hoxd10* is involved in prostate cancer processing, cell differentiantion and

morphogenesis [Jonkers *et al.,* 2020; Mo *et al.,* 2017; Zhang *et al.,* 2019].

Another gene with 2.2-fold expression rise is *Slc45a3,* a well-known marker of prostate cancer

progression, when a loss of the protein determined poorer prognosis for cancer patients [Hernández-Llodrà *et*

al. 2017; Pin *et al.,* 2017; Perner *et al.,* 2013]. An overexpression of another gene identified in our study (12-fold

rise for *Usp39* gene) is linked with promotion of prostate or ovarian tumorigenesis. *Usp39* is also involved in pre-

mRNA splicing, cytokinesis and spindle checkpoint working, and is crucial for apoptosis [Huang *et al.,* 2016; Yuan

et al., 2021; Zhao *et al.,* 2016].

Also, 2.2-2.5-fold rise for ncRNA genes (*Gm26848*, *Gm45648*; no data concerning their role) and 2-fold

reduction for one pseudogene (*Gm28661*; also no data) were found (Figure 5). Following the latest data

(November 2021) from Gene NCBI database, the Gm45648 has been annotated as one with the testis-restricted

expression in mouse but no function has been suggested. The second ncRNA *Gm26848* found is suggested as

protein coding organic cation transporter-like protein across membranes in *Drosophila sechellia*. There were also

two other genes with expression change (3.6-fold decrease – pseudogene *Gm44126*, and 2.5-fold increase –

4933430L12Rik probably involved in antisense transcription). 'Gene' database data showed that the *Gm44126*

in mice is rather a protein coding gene with the function of UDP-glucuronic acid decarboxylase, and contains two

domains involved in NAD(P)H processing. Thus, it can be suggested that this gene is as a novel candidate involved

in mitochondrial respiratory chain. Our data with 3.6-fold expression decrease of this gene seem to fit into

observed diminished functioning of mitochondrial machinery in *Tcte1$^{-/-}$* mice.

Another three genes manifested 2-fold decrease of their testis expression when comparing *Tcte1$^{-/-}$* vs.

Tcte1$^{+/-}$ (but not to WT): *Aqp2* (aquaporin-2 involved in osmotic transportation of the water), with well-known

role in spermatogenesis, female endometrium cycle and urinary incontinenece; [Klein *et al.,* 2013; Ramli *et al.,*

2019], *Mt3* (metallothionein-3, committed in homeostasis and detoxification of toxic metals, also in testis)

important in apoptosis regulation of Bax and Caspase pathways [Gao *et al.,* 2021; Honda *et al.,* 2010]), and

Rps27rt (ribosomal protein) (Figure 5; Supplementary file 11). It was also found that Rps27rt may interact

indirectly with Usp39 and mitochondrial respiratory complex linking the mitochondrial energy processing with

apoptosis (Supplementary file 11). Last gene – *Ube2l6*, was found to have 2-fold increased expression level in

testis in *Tcte1$^{+/-}$* animals vs. *Tcte1$^{-/-}$* and WT. Its role has been established as ubiquitin enzyme engaged in: stress

damage response, cell cycle progression, embryo development, immune response, and factors linked to ATP

machinery (Figure 5) [Huang *et al.,* 2016; Yuan *et al.,* 2021; Zhao *et al.,* 2016; Sandy *et al.,* 2020; Cai *et al.,* 2021].

Increased expression of *Ube2l6* was linked to degradation/suppression of cancer cells and affected downstream

apoptotic factors (i.e., Caspase 3, Caspase 9, Bcl-2, Bax) [Li *et al.,* 2018].

Immunofluorescence on sperm

An example of immunofluorescence *in situ* staining on mouse spermatozoa used to detect the N-DRC

proteins has been presented in Figure 7. WT mouse spermatozoa showed Tcte1 protein presence in the sperm

head nucleus, midpiece, and a part of the tail (Figure 7A). Immunofluorescent signal of Tcte1 in *Tcte1$^{+/-}$* mice was

similar to WT animals, while in *Tcte1$^{-/-}$* was localized only in the nucleus (Figure 7A). It seems to confirm that in

Tcte1$^{-/-}$ males the protein has not been produced properly, revealing only residual amounts in sperm head

nucleus, and not transported to sperm flagella. The other N-DRC components (Drc7, Fbxl13 and Eps8l1) were

localized in the sperm nucleus and among the whole sperm tail (Fbxl13, Eps8l1) or only within its fragment (Drc7),

with exception of the midpiece (Figure 7B). Results obtained seem to suggest co-localization of evaluated N-DRC

proteins in various combinations, not common among the midpiece and the terminal part of the sperm tail.

Sperm ATP level

Measurements performed in two time points (0 and +60 min.) revealed that in *Tcte1*[-/-] samples the

luminescence level was approximately 2.4-fold lower when compared to WT and *Tcte1*[+/-] mice (p<0.0001) (Figure

8; Supplementary file 13). There were no differences between WT and *Tcte1*[+/-] males. Additionally, the decrease

of the luminescent signal after 60 min. was similar in all three groups studied – approximately by 35% (WT:

35.27%, *Tcte1*[+/-]: 33.88%, *Tcte1*[-/-] 37.76%).

Apoptosis in testicular tissue

Evaluation of the immunofluorescent staining results revealed Casp3-positive signals (indicating

apoptosis) in spermatogonia only, in all three mouse genotypes (Figure 9). In each zygosity the numbers of

tubules and positively stained spermatogonia (intensity of the signal) were similar suggesting equal levels of

apoptosis.

Sequencing – human male samples

Potentially associated/important 6 heterozygous SNVs for *TCTE1* gene has been found in 15 out of 248

(6.05%) infertile males with drastically decreased semen parameters, ranging from azoospermia to severe

oligoasthenozoospermia (Table 1). 3 variants were identified with ultrarare allele frequency (MAF ≤0.0005 in the

general population; gnomAD) in three males with NOA: c.862C>T (p.Arg288Ter) predicted as stop-gained

mutation leading to non-mediated decay, and thus to protein loss of function with high probability; c.185G>A

(p.Arg62His), or c.1048G>A, (p.Gly350Ser), defined as missense ones and probably damaging. Next two SNVs

were found as the rare missense ones with MAF <0.005: c.397C>T (p.Arg133Cys) (one NOA case) and c.470A>C

(p.Glu157Ala) (four NOA cases), and were defined as probably damaging. The last 1 SNVs (1 NOA, 1 crypto-, and

5 with severe oligoasthenozoospermia) was not found in databases, and thus specified as the novel one

(c.374T>G; p.Ile125Arg) with missense prediction (Table 1; Supplementary file 14).

Protein prediction

To understand the molecular implications of identified mutations on the structure and function of

TCTE1, we performed homology modeling experiments using the Phyre2, I-TASSER, and AlphaFold tools [Kelley

et al., 2015; Roy *et al.,* 2010; Yang *et al.,* 2015; Yang and Zhang 2015; Tunyasuvunakool *et al.*, 2021]. TCTE1 is

predicted to contain at least five tandem leucine-rich repeats (LRR) in its C-terminus [UniProt 2019], while no

known three-dimensional protein folds have been assigned to its N-terminus. All homology searches yielded a

polypeptide with tandem leucine-rich repeats with nearly 100% confidence (Figure 10A; Supplementary file 15).

Modeling the N-terminal region of TCTE1 was more difficult; although both Phyre2 and I-TASSER confidently

predicted a largely α-helical secondary structure for the N-terminal half of TCTE1 (Figure 10A; Supplementary

file 15), neither tertiary structure was predicted with high confidence; indeed, comparison of the two

demonstrates vastly different folds for the N-terminus from the programs. However, the confidence of the

AlphaFold-generated model was high for the region just N-terminal to the LRR. Further, based on the predicted

aligned error plot for calculated for its models, the confidence of the predicted structure of residues 100-200

was high relative to the position of residues 220-480, providing support for the modeled through-space three-

dimensional interactions. That said, we interpreted these models very conservatively.

We attempted to hypothesize how mutations of interest could affect the tandem LRRs and putative

secondary structure of the N-terminus of the protein. Residue R64 resides in a predicted unstructured region

and as such most likely is solvent exposed in the tertiary structure (Figure 10A, B). Mutation of this residue from

arginine to histidine could therefore affect the surface potential of this disordered region, making the it less

positively charged. Indeed, this mutation would be expected to impact the interactions that the N-terminus of

TCTE1 can have with other molecules, including other members of the N-DRC. The interaction of microtubule-

associated proteins, including the N-DRC, with binding partners is known to be largely charge-based [Kubo *et al.,*

2017; Drechsler *et al.,* 2019].

Three mutations of interest occur in the globular region just N-terminal to the LRR. The AlphaFold model

clearly shows how mutation I125R would affect the structure of this region Figure 10a, c). Residue isoleucine 125

is predicted to be nestled within the hydrophobic core including L97 L109, L112, F91, W131, L197, Y130, and the

methyl group of the T121 sidechain. A substitution of positively charged arginine for isoleucine in this position

would not be accommodated in this hydrophobic region, likely resulting in local unfolding that could potentially

propagate beyond the pocket. Likewise, residue R133 supports the structure of this putative globular domain by

forming hydrogen bonds with the backbone of residues 109, 110, and/or 112, thereby possibly contributing to

the overall organization of this domain and its positioning relative to the LRR (Figure 10d). Mutation of this

residue to a cysteine would remove the hydrogen bonding groups supporting these interactions, likely leading

to a local unfolding that could possibly propagate farther. Interestingly, residue E157 is predicted to interact with

residues in the LRR, likely stabilizing positively charged residues around it, including several from the linker

leading up to the first β-strand of the LRR (Figure 10B, E). Mutation of this residue to an alanine would lead to a

tertiary destabilization, most likely affecting the overall structure of the N-terminal region of the LRR and the

globular region just N-terminal to it.

Two mutations of interest affect the LRR helices. Mutation R288* results in premature truncation,

eliminating most of the LRR region from the final protein product, although it is predicted that this premature

termination codon would lead to nonsense mediated decay. Mutation G350S would be expected to result in the

destabilization of inter-repeat interactions (Figure 10F), possibly even unfolding the protein from that position

on; the lack of a side chain in glycine allows for the close approach of these tandem structural elements, whereas

the addition of the serine side chain would abrogate the tight packing of residues, regardless of which rotamer

it adopts. This could result in a restructuring its C-terminus and/or deforming the surface of the module,

therefore affecting its interactions with its binding partners.

Since the N-DRC is conserved from single-celled eukaryotes to mammals, we took advantage of

published cryo-electron tomography studies examining the N-DRC from model organism *Chlamydomonas* (1).

This study was of great interest since it explicitly located the position of the C-terminus of DRC5 with a nanogold

particle; importantly, it was consistent with earlier cryo-ET experiments positioning the N-DRC (2). While the

resolution of this structure is only 40Å, it was still useful for placing the modeled TCTE1 structure in the context

of the entire N-DRC (Supplementary file 15B). Manual docking of the AlphaFold predicted TCTE1 structure is

consistent with the cryo-EM tomography data [Gui *et al.,* 2019; Heuser *et al.,* 2009].

DISCUSSION

In this study we have investigated the role of *Tcte1* in male infertility using mouse knockout model.

Using homo- and heterozygous animals, their reproductive potential has been examined, noting no progeny by

homozygous males. While there were no differences in histology of the testes, the size of the gonads varied

between the zygosity of animals, followed by two semen phenotypes revealed within the sperm quality

evaluation. RNAseq analysis of the mutant testicular tissue demonstrated the influence of *Tcte1* mutations on

the expression pattern of genes related to mitochondrial cytochrome oxidase activity, apoptosis and

spermatogenesis. The ATP level estimate of spermatozoa showed its decrease in homozygous males.

Additionally, protein prediction modeling of identified variants in human samples revealed changes in the protein

surface charge potential, leading to disruption in helix flexibility or dynamics, and thus, suggesting the disrupted

TCTE1 interaction with its binding partners within the axoneme. The major findings concerning observed

abnormal sperm tail beating in KO *Tcte1* mice were summarized in Figure 11. We have also identified and

discussed novel functions of genes with testicular expression significantly disturbed in the *Tcte1* knockout.

Therefore, this study contains novel and the most complete data concerning *TCTE1* role in male infertility, so far.

Phenotypes of homozygotes and heterozygotes

There was only one published study so far, concerning knockout mouse model of *Tcte1* [Castaneda e*t*

al., 2017], with Authors the infertile homozygous males due to asthenozoospermia derived from limited and

circular flagellar beating, and fertile heterozygotes. The study did not find any differences between homozygous

and WT males in the testicular histology, followed by similar testis weight, sperm concentration and sperm

morphology [Castaneda *et al.,* 2017]. We also have not found any differences in the testicular and epididymal

tissues histology. However, the weight of the testes was decreased significantly (~41% in $Tcte1^{-/-}$, 7% in $Tcte1^{+/-}$

), followed also by testis size reduction in $Tcte1^{-/-}$ animals (by 22%) (Figure 4B). When comparing semen

parameters (Figure 4C-E), we have found enormous decline in sperm concentration in both: $Tcte1^{-/-}$ (3.3-fold) or

$Tcte1^{+/-}$ (2.5-fold) males, and double increase in abnormal sperm morphology in $Tcte1^{-/-}$ animals. Similarly, to

Castaneda *et al.* [2017], we have documented the impaired sperm motility representing circular mode of the

sperm tail beating. Also, the increased ratio of immotile spermatozoa in homozygous males was documented in

our study. We can suggest that these discrepancies may result from various number of animals used for

evaluation: 3 males in study of Castaneda *et al.,* [2017] vs. at least 18 per each zygosity in our study, when

comparing semen parameters and tissue characteristics (except of sperm morphology evaluation with at least 5

animals used). Probably, also the differences in knockout construction used could be causative for observed

variances: reporter-tagged insertion with conditional potential vs. the exon 3 deletion. Thus, we have observed

two various phenotypes dependent on the zygosity of the animals: homozygous males manifested

oligoasthenoteratozoospermia (OAT) and were infertile, while heterozygous males demonstrated

oligozoospermia (O) and remained fertile. Data collected in International Mouse Phenotypic Consortium (IMPC;

a repository of all available mouse phenotypes created, followed with the possible linkage to human disease

defects; https://www.mousephenotype.org), concern only one *Tcte1* mouse study [Castaneda *et al.,* 2017], while

we reveal an unmatched phenotypes of decreased sperm count.

We suggest that similar effect – decreased sperm count linked to *TCTE1* mutations, could be observed

in humans. Our pilot mutation screening of 248 human infertile males with severely decreased sperm count

(from oligo-, via crypto-, to azoospermia) revealed 6 heterogeneous SNVs (1 novel, 3 ultrarare, and 2 rare; all

predicted as disease causing) with the total frequency of 6.05% (Table 1). We propose that molecular

epidemiology studies of infertile patients should reveal the existence of homozygous or a compound

heterozygous *TCTE1* variants with clear direct downgrading effect on spermatogenesis. However, we claim that

also heterozygous variants can be causative, similarly to *SPINK2* which deficiency has been documented as

responsible for oligoasthenozoospermia in heterozygous mice vs. azoospermia in homozygous animals, followed

by the azoospermia in human homozygous brothers vs. their heterozygous father with oligozoospermia [Kherraf

et al., 2017]. It can be explained by the probable haploinsufficiency occurrence, when the single copy of the WT

allele in heterozygous combination of which a variant allele is insufficient to produce the WT phenotype. Our

observations, followed by the *SPINK2* data and data related to infertility (i.e.: P1/P2 protamines, *Pde8b*, *Trim28*,

Tcfl5), and to the other diseases (i.e.: *PIKFYVE* in cataract, *NFIA*, *SCN1A* in neurological or epilepsy disturbances,

KLF1 in hemoglobin persistenece, *NR5A2* in pancreatic cancer), seem to confirm that the severity of the

phenotype depends on the gene expression level [Jodar and Oliva, 2014; Mei *et al.,* 2022; Ogura *et al.,* 2022;

Heshusius *et al.,* 2022; Valassina *et al.,* 2022; Sandhu *et al.,* 2021; Leal *et al.,* 2021; Tan *et al.,* 2020; Xu *et al.,*

2022]. We have documented variable phenotypes for a novel *TCTE1* variant c.374T>G (p.Ile125Arg) that has been

documented in seven males, including: azoospermia, crypto-, and severe oligoasthenozoospermics. Thus, it

seems to suggest the reasonable candidacy of haploinsufficiency as the mechanism responsible for observed

phenotypic spectra also in case of the *TCTE1* gene.

Kallikreins, Aquaporin 2

Our pathway analysis revealed a network including two genes with changed expression level in testis:

Igf2 (2-fold increased) and *Klk1b22* (8-fold decreased), supported by an interesting set of proteins linked to

apoptosis and cellular processing (Figure 5; Figure 6; Supplementary file 11; Supplementary file 12). We can

suggest that observed *Igf2* overexpression in our study can be one of the associated factors leading to decreased

sperm count and testis parameters in evaluated KO mice model because of its known relation to spermatogenesis

and testis development [Matsuzaki *et al.,* 2020; Cannarella *et al.,* 2020; Neirijnck *et al.,* 2019; Damaschke *et al.,*

2017]. Documented reductions can be supported by the observed constrained expression of *Klk1b22* – kallikrein

mostly known from its neuronal activity; however, it cannot be excluded that we have revealed additional role

of this protein, similar to the other members of the kallikrein family – role in semen liquefaction and degradation

of extracellular matrix proteins in the interstitial area surrounding Leydig cells of the adult mouse testes (*Klk1b27*,

Klk1b21, *Klk1b24*) [Michaelis et al. 2017; Marques *et al.,* 2016, Du Plessis *et al.,* 2013; Fink *et al.,* 2007]. Notably,

Klk1b22 expression is significantly enriched in testis. While expression of all four kallikreins was decreased 2.5-

8-fold in our *Tcte1* KO model, indicating their participation in the reduction of sperm parameters measured

(Fig.5).

We have also documented 2-fold decrease of expression of *Aqp2* (Aquaporin 2), involved in fluid

resorption and water movement within the duct system, with known role in spermatogenesis [Klein *et al.,* 2013;

Ramli *et al.,* 2019]. It was documented that in stallions Aqp2 is being expressed in Leydig cells, round and

elongated spermatids [Klein *et al.,* 2013]. Also, this protein works within a network of molecules that are linked

to cell membrane trafficking (i.e., Rab11a, Myo5b), followed by activity of protein kinases (Prkacb, Prkaca) that

are known as regulators of cell proliferation, chromatin condensation/decondensation, and nuclear envelope

disassembly/reassembly, and all is supported by the ATP metabolism (Hspa8) [Supplementary file 11] [Markou

et al., 2022; Wagner *et al.,* 2022]. Considering this evidence, we suggest Aqp2 play an important role in reduction

of the spermatids volume during spermatogenesis and in proper sperm morphology assessement and observed

morphological differences between *Tcte1* mice zygosities in our study.

Disruptions in sperm cell structure

TCTE1 protein contains five leucine rich repeats (LRRs) that are known to be directly involved in

receptor-mediated signaling on the structural basis for interaction between proteins [Ng *et al.,* 2011; Kobe and

Kajava 2001]. It was documented that variants in genes encoding LRRs are commonly implicated in the

pathogenesis of more than 60 human diseases [Ng *et al.,* 2011; Matsushima *et al.,* 2019]. LRR mutations mostly

occur within regions that are protective for the hydrophobic core of the LRR domains. Thus, mutations in LRRs

may affect the charge of the surface potential, leading to conformational changes of the protein or induction of

protein cumulation. Our prediction model of potential changes in TCTE1 protein resulting from mutations found

in human samples seems to point out the prospective impact into the interactions of the TCTE1 with other

molecules (binding partners) known to be largely charge-base, including N-DRC elements or microtubule-

associated proteins [Kubo et al., 2017; Drechsler et al., 2019]. In the case of TCTE1 protein observed amino acid

substitutions promote the abnormal aggregation of TCTE1 protein, and decrease its structural stability, then

probably leading to the observed disturbances. Beating motion of flagella is maintained by electrostatic cross-

bridge between negatively charged tubulins and positively charged N-DRC [Kubo *et al.,* 2017]. Thus, changes in

the charge potential of TCTE1 surface, which is one of the N-DRC components, seem to provide modified

(reduced) flagellar beating, similarly as in case of mutations in other N-DRC element – DRC4 protein, of which

changes in charge were essential for proper flagellar motility [Kubo *et al.,* 2017; Lewis *et al.,* 2016]. Also, the

inflexibility of the midpiece observed in spermatozoa of *Tcte1* knockout mice seems to be a consequence of

changed structure of LRRs in Tcte1 protein.

It is also known that the lengths of the sperm midpiece and tail may be crucial in determination of sperm

swimming velocity, followed then by decreased fertilization rate [Firman and Simmons 2010; Gu *et al.,* 2019].

Available data showed that sperm with shorter midpiece may swim slower and were observed in

asthenozoospermic cases [Firman and Simmons 2010; Gu *et al.,* 2019; Anderson *et al.,* 2002]. The meta-analysis

of sperm shape features according to sperm competition theory has also shown, that in majority of vertebrates

the longer sperm cell (including midpiece or flagellum) has been attributed to faster swimming, followed by the

statement that the sperm size may be increased when sperm number is lowered [Lüpold *et al.,* 2020]. In our

study we have documented longer sperm tails and midpieces in *Tcte1*$^{-/-}$ animals, that in fact were infertile. Thus,

we can suggest that: (i) the observed longer sperm tails and midpieces may be co-responsible for aberrant

motility observed, and (ii) the documented longer sperm tails and midpieces may be an indirect result of

constrained sperm count observed, because of some sort of compensation for aberrant motility. However, this

issue remains only a suggestion because of a restricted literature data relative to similar observations. Bennison

et al. [2016] revealed than in a passerine bird the sperm tail velocity was correlated to total sperm length, but

only to some point of the length value at which the velocity started to be decreased. On the other hand, Gu et

al. [2019] have shown that in mammalian species there were some changes in the axonemal slopes when the

flagellar length increases. The results obtained within our study, and supported by other research concerning

sperm tail beating pathways, should be a good start point for creation of further mathematical computations for

better understanding of the asthenozoospermia.

Constrained energy metabolism

The flagellar motility is determined by coordinated activity of asymmetrically distributed dyneins in the

axoneme. The driving factor for this activity is the hydrolysis of ATP released from OXPHOS system (Mitochondrial

Oxidative Phosphorylation System) [Tourmente *et al.,* 2015; Ford 2006]. ATP is being produced in mitochondria

placed within sperm midpiece. In our study ATP level measurements revealed its decreased level in sperm cells

of *Tcte1*$^{-/-}$ mice, what can result in faulty dynein functioning, leading to asthenozoospermia via abnormal sperm

tail beating. We have documented that *Tcte1* KO resulted in significant changes in expression level of testicular

genes involved in mitochondrial energy transport. That leads to the conclusion that homozygous *Tcte1* mutation

results not only in inflexible midpiece leading to asymmetric circular motility, as presented in this study and by

others [Fung *et al.,* 2018], and then to asthenozoospermia, but also to aberrant functioning of mitochondrial

OXPHOS machinery leading to disruption of ATP production. Thus, the sperm tail has also insufficient power for

beating because of the decreased ATP content. Castaneda et al. suggested that the reasons of the observed ATP

decrease may result from disturbances in a glycolytic pathway [Castaneda *et al.,* 2017]. It is already known, that

glycolytic energy is enough for spermatozoa to survive [Nascimento *et al.,* 2008; Zhang *et al.,* 2022]. However,

for sperm motility and then fertilization, the OXPHOS energy production is required [Tourmente *et al.,* 2015;

Ford 2006; Zhang *et al.,* 2022]. In our study, RNAseq results clearly revealed the influence of $Tcte1^{-/-}$ mutation on

genes involved directly in electron transport chain, such as: *mt-Co2* and *mt-Co3*, crucial components of the

respiratory chain complex IV, one of the OXPHOS elements. Observed reduced expression seems to clearly

evidence the decreased ATP production via disturbed electron chain transportation, and then revealing the

sperm motility reduction observed in $Tcte1^{-/-}$ males, leading to observed asthenozoospermia. Detailed STRING

analysis revealed definite interactions between both listed proteins identified in our study, and a list of other

mitochondrial ones (mt-Co1, mt-Cytb, Cox5a, Cox6a, Cox7c, mt-Atp6; Supplementary file 11; Supplementary file

12) clearly related to proper processing of cytochrome energetic activity [Brischigliaro and Zeviani 2021;

Ganetzky *et al.,* 2019]. Overexpression or mutations in *COX1*, *COX2*, and *COX3* were previously observed in

infertile men with varicocele and OAT, or diabetes [Heidari et al., 2016; Mostafa *et al.,* 2016; Perrotta *et al.,*

2012]. Additionally, it was shown that system of COX2 and prostaglandins plays key role in development of male

gonad, spermatogenesis and steroidogenesis, especially in functioning of Sertoli and Leydig cells, and also was

found to be significant in the background basis of idiopathic infertility status [Frungieri *et al.,* 2015]. It seems that

in our study the reason of observed disturbed motility, represented as circular movement of spermatozoa,

supported by increased ratio of immotile sperm cells in $Tcte^{-/-}$ mice, leading to asthenozoospermia, resulted from

two factors: (i) inflexibility of sperm midpiece resulted from changed protein charge level in the Tcte1 protein,

followed by elevated lengths of sperm tail and midpiece, and (ii) decreased amount of ATP produced, that may

be an outcome of inability to get sufficient energy for proper sperm motility.

Protection and/or compensation?

We have also documented an important role of the *Fetub* gene for the first time in the male testis.

Besides of its known role in female fertility [Karmilin *et al.,* 2019; Bansho *et al.,* 2017], the latest study pay

attention also in the context of prostate cancer is a detailed way [Zhan *et al.,* 2020]. It was found that

overexpression of *FETUB* increases the apoptosis of prostate cancer cells, followed by the inhibition of their

proliferation, migration and invasion, and also slower growth of tumor when compared to controls [Zhan *et al.,*

2020]. Thus, in the overexpression mode, the Fetub plays protective role for the prostate. It seems to explain

results obtained in our study, where the *Fetub* expression was increased almost 5-fold in *Tcte1*$^{-/-}$ mice, but no

tumor signs have been observed in the mouse testis at the tissue level. Zhan et al. [2020] have found the

connection between *Fetub* overexpression, and induction of Bax (Bcl-2-associated X protein) and capases,

resulting in promoting the apoptosis. It is known that the Bax/Bcl-2 induction/inhibition is directly related to the

release of cytochrome c (Cyt c) in the caspase-dependent signaling [Dadsena *et al.,* 2021; Samaiya *et al.,* 2021;

Li *et al.,* 2021]. And cytochrome c is one of the two electron carriers in electron transport chain responsible for

transport directly to the respiratory chain complex IV [Yin and O'Neil 2021; Alvarez-Paggi *et al.,* 2017]. Due to

Cyt c release, complex IV is essential for apoptosis and cell fate [Samaiya *et al.,* 2021; Li *et al.,* 2021]. Following

of all of those facts, our results seem to reveal the connection between *Fetub* and diminished mitochondrial

chain IV activity in mice testis, what have not been reported so far. It can be also supported by the fact of the

known relation between FETUB and the inactivation of the PI3K/AKT signaling pathway (key player in regulation

of cell proliferation and apoptosis in cancerogenesis), in which the Bax/Bcl-2 are involved [Zhan *et al.,* 2020].

The novel testis-protective role in our study can be also credited to three other genes with increased

expression level in testicular tissue: *Hoxd10* (3-fold increase), *Slc45a3* (2.2-fold), and *Ube2l* (2-fold).

Overexpression/activation of *HOXD10* has been already documented as inhibitor of tumorigenesis [Jonkers *et*

al., 2020; Mo *et al.,* 2017; Zhang *et al.,* 2019]. Expression of *Slc45a3*, known also as 'prostein', has been

documented as weaker in cancer tissues [Hernández-Llodrà et al. 2017; Pin et al., 2017; Perner et al., 2013].

Elevated expression level of *Ube2l* is known to be connected with the degradation/suppression of tumor cells [Li

et al., 2018]. On the other hand, the 12-fold increased gene expression level for *Usp39* (ubiquitin specific

peptidase 39) – an oncogenic splicing factor which expression is positively linked with cancerogenesis' promotion

in prostate or ovaries, while its silencing induces apoptosis and cell cycle arrest, has been noticed [Huang *et al.,*

2016; Yuan *et al.,* 2021; Zhao *et al.,* 2016].

Taken together data obtained for genes related to cancerogenesis and/or apoptosis described above,

we can hypothesize that we observe an interesting novel network of interactions between two groups of genes

that are reciprocally neutralizing potential influence onto the testis tissue. Mainly, in the first group we have

genes with suggested novel anti-cancer role in the testis (*Fetub, Hoxd10, Slc45a3, Ube2l6*), which increased

expression level inhibits the cancer cell development, supported by protective role of *Mt3*; while in the second

corner of the ring the *Usp39* gene with elevated expression level (the most from all the observed changes) should

lead to reveal some tumor-like changes within the testis tissue. Following the fact, that both: histological

evaluation, as well as caspase 3 staining (Fig.9), have shown no differences between hetero- and homozygous

mutant mice, our hypothesis seems to be justifiable. On the other hand, there is a prediction that many of the

genes (especially so-called cancer-testis associated antigens, CTA) related to spermatogenesis may function in

tumorigenesis [Aurriere *et al.,* 2021; Yang *et al.,* 2020; Babatunde *et al.,* 2017; Wang *et al.,* 2016; Bode *et al.,*

2014; Cheng *et al.,* 2011; Nagirnaja *et al.,* 2018]. Due to known similarities between oncogenic and

spermatogenic processes, especially in the energy metabolism, we can carefully suggest/hypothesize that in our

KO model the genes with changed expression profile found, known as highly involved in tumorigenesis, may also

be important in spermatogenesis. Obviously, there is a need for further studies in this direction to confirm these

suggestions or to reveal the other (still unknown) mechanisms.

Conclusions

We have documented that *TCTE1* gene is the next one that should be added to the 'male infertility list'

of molecular background because of its crucial role in spermatogenesis and proper sperm functioning. The

mutations in mouse *Tcte1* knockout model revealed two phenotypes, depending from zygosity with infertile

oligoasthenoteratozoospermic homozygotes, and fertile oligozoospermic heterozygotic males, suggesting

haploinsufficiency mechanism involvement. All collected data pointing that the Tcte1 protein is functioning in

spermatogenesis and show how its mutations add the level of complexity to the clinical manifestation of

decreased semen parameters. The disrupted energy machinery of the spermatozoa, followed by newly

suggested roles of genes related to cancerogenesis and/or apoptosis (here in the context of their

neutralizing/compensation effect on the testicular tissue), seem to underline the complex and devastating effect

of the single-gene mutation, on the variety of testicular molecular networks, leading to reproductive failure.

ACKNOWLEDGEMENTS

We are grateful to Oliwia Kordyl, MSc, and Katarzyna Kurek, MSc (IHG PAS, Poznan) for their technical help in

preparation of the samples for sequencing, genotyping, and immunofluorescence staining.

Authors' contributions:

MO design of the study, manuscript drafting and editing, coordination of the tasks, data interpretation, gene selection, preparation of human blood and semen samples, mice: mating, genotyping, gross sections and preparation of biological samples, RNAseq analysis, immunofluorescence, Sanger sequencing, ATP measurements; **AM** gene selection, data interpretation, genotyping, RNAseq analysis, collection and preparation of human testis samples, WGS and Sanger sequencing, manuscript editing; **TS** bioinformatic analysis of RNAseq and WGS results; **NP** WES of azoospermia samples (incl. bioinformatics); **AB** protein prediction modeling and its description; **SB** preparation of: DNA for sequencing, RNA for RNAseq, screening of databases for RNAseq results; **MKa** collection of human blood and semen samples, semen analyses of human and mouse samples, detailed morphology of mouse spermatozoa; **HJ** histopathology of mouse tissue sections; **PJ** recruitment of patients and their medical history; **ANY** gene selection, interpretation of azoospermia WES results, data interpretation; **MKu** data interpretation, gene selection, recruitment of patients and their medical history, correction of the manuscript, funds collection and supervision.

All authors critically reviewed and approved the final version of the manuscript.

LARGE SCALE DATA: All data generated or analyzed during this study are included in this published article and its supplemental information files.

RNAseq data are available in GEO database (https://www.ncbi.nlm.nih.gov/geo/) under the accession number: GSE207805.

The results described in the publication are based on whole-genome or exome sequencing data which includes sensitive information in the form of patient specific germline variants. Information regarding such variants must not be shared publicly following the European Union legislation described in the following document: https://publications.jrc.ec.europa.eu/repository/bitstream/JRC113479/policy_report_-_review_of_eu_national_legislation_on_genomics_-_with_identifiers_1.pdf, therefore the access to raw data that support findings of this study are available from the corresponding author upon reasonable request.

LIMITATIONS, REASONS FOR CAUTION: In the study, the *in vitro* fertilization ability of homozygous male mice was not checked.

WIDER IMPLICATIONS OF THE FINDINGS: This study contains novel and the most complete data concerning

TCTE1 role in male infertility. *TCTE1* gene is the next one that should be added to the 'male infertility list' of

molecular background because of its crucial role in spermatogenesis and proper sperm functioning.

STUDY FUNDING/COMPETING INTEREST(S): This work was supported by National Science Centre in Poland,

grants no.: 2015/17/B/NZ2/01157 and 2020/37/B/NZ5/00549 (to MK), 2017/26/D/NZ5/00789 (to AM), and

HD096723, GM127569-03, NIH SAP #4100085736 PA DoH (to AY). The authors declare that there is no conflict

of interest that could be perceived as prejudicing the impartiality of the research reported.

REFERENCES

Agarwal A, Baskaran S, Parekh N, Cho CL, Henkel R, Vij S, Arafa M, Panner Selvam MK, Shah R. Male infertility. *Lancet* 2021;**397**: 319-333 https://doi.org/10.1016/S0140-6736(20)32667-2

Alvarez-Paggi D, Hannibal L, Castro MA, Oviedo-Rouco S, Demicheli V, Tórtora V, Tomasina F, Radi R, Murgida DH. Multifunctional Cytochrome c: Learning New Tricks from an Old Dog. *Chem Rev* 2017;**117**:13382-13460 https://doi.org/10.1021/acs.chemrev.7b00257

Anderson MJ, Dixson A F. Sperm competition: motility and the midpiece in primates. *Nature* 2002;**416**:496 https://doi.org/10.1038/416496a

Aprea I, Raidt J, Höben M, Loges NT, Nöthe-Menchen T, Pennekamp P, Olbrich H, Kaiser T, Biebach L, Tüttelmann F. et al. Aprea I, et al. Defects in the cytoplasmic assembly of axonemal dynein arms cause morphological abnormalities and dysmotility in sperm cells leading to male infertility. *PloS Genet* 2021;17:e1009306 https://doi.org/10.1371/journal.pgen.1009306

Aurrière J, Goudenège D, Baris OR, Boguenet M, May-Panloup P, Lenaers G, Khiati S. Cancer/Testis Antigens into mitochondria: a hub between spermatogenesis, tumorigenesis and mitochondrial physiology adaptation. *Mitochondrion* 2021;**56**:73-81 https://doi.org/10.1016/j.mito.2020.11.002

Babatunde KA, Najafi A, Salehipour P, Modarressi MH, Mobasheri MB. Cancer/Testis genes in relation to sperm biology and function. *Iran J Basic Med Sci* 2017;**20**:967-974 https://doi.org/10.22038/IJBMS.2017.9259

Baklouti-Gargouri S, Ghorbel M, Mahmoud AB, Mkaouar-Rebai E, Cherif M, Chakroun N, Sellami A, Fakhfakh F, Ammar-Keskes L. Mitochondrial DNA mutations and polymorphisms in asthenospermic infertile men. *Mol Biol Rep* 2013;**40**:4705–4712 https://doi.org/10.1007/s11033-013-2566-7

Bansho Y, Lee J, Nishida E, Nakajima-Koyama M. Identification and characterization of secreted factors that are upregulated during somatic cell reprogramming. *FEBS Letters* 2017;**591**:1584-1600 https://doi.org/10.1002/1873-3468.12665

Barratt CLR, Björndahl L, De Jonge CJ, Lamb DJ, Martini FO, McLachlan R, Oates RD, van der Poel S, St John B, Sigman M, et al. The diagnosis of male infertility: an analysis of the evidence to support the development of global WHO guidance-challenges and future research opportunities. *Hum Reprod Update* 2017;**23**:660-680 https://doi.org/10.1093/humupd/dmx021

Bennison C, Hemmings N, Brookes L, Slate J, Birkhead T. Sperm morphology, adenosine triphosphate (ATP)

concentration and swimming velocity: unexpected relationships in a passerine bird. Proc. Royal Soc. B -Biol Sci

2016;**283**:20161558 https://doi.org/10.1098/rspb.2016.1558

Binsila BK, Archana SS, Ramya L, · Swathi D, · Selvaraju S, Shivakumar Gowda NKS, · Pal DT, · Rafay A,· Bhattaet R.

Elucidating the processes and pathways enriched in buffalo sperm proteome in regulating semen quality. Cell

Tissue Res 2021;**383**:881-903 https://doi.org/10.1007/s00441-020-03303-9

Bode PK, Thielken A, Brandt S, Barghorn A, Lohe B, Knuth A, Moch H. Cancer testis antigen expression in testicular

germ cell tumorigenesis. *Modern Pathology* 2014;**27**: 899-905 https://doi.org/10.1038/modpathol.2013

Bower R, Tritschler D, VanderWaal Mills K, Heuser T, Nicastro D, Porter ME. DRC2/CCDC65 is a central hub for

assembly of the nexin-dynein regulatory complex and other regulators of ciliary and flagellar motility. *Mol Biol*

Cell 2018;**29**:137-153 https://doi.org/10.1091/mbc.E17-08-0510

Brischigliaro M, Zeviani M. Cytochrome c oxidase deficiency. *Biochim Biophys Acta - Bioenerg* 2021;**1862**: 148335

https://doi.org/10.1016/j.bbabio.2020.148335

Cai J, Huang H, Hu X, Lang W, Fu W, Xu L, Qiu Z, Zhong H, Chen F. Homoharringtonine Synergized with Gilteritinib

Results in the Downregulation of Myeloid Cell Leukemia-1 by Upregulating UBE2L6 in FLT3-ITD-Mutant Acute

Myeloid (Leukemia) Cell Lines. *J Oncol* 2021, 3766428 https://doi.org/10.1155/2021/3766428

Cannarella R, Condorelli RA, La Vignera S, Bellucci C, Luca G, Calafiore R, Calogeroet AE. IGF2 and IGF1R mRNAs

Are Detectable in Human Spermatozoa. *World J Mens Health* 2020;**38**:545-551

https://doi.org/10.5534/wjmh.190070

Castaneda JM, Hua R, Miyata H, Oji A, Guo Y, Cheng Y, Zhou T, Guo X, Cui Y, Shen B, et al. TCTE1 is a conserved

component of the dynein regulatory complex and is required for motility and metabolism in mouse spermatozoa.

PNAS 2017;**114**,E5370-E5378 https://doi.org/10.1073/pnas.1621279114

Chemes HE, Olmedo SB, Carrere C, Oses R, Carizza C, Leisner M, Blaquier J. Ultrastructural pathology of the

sperm flagellum: association between flagellar pathology and fertility prognosis in severely asthenozoospermic

men. *Hum Reprod* 1998;**13**:2521-2526 https://doi.org/10.1093/humrep/13.9.2521

Chen Y, Wang J, Pan C, Li D, Han X. Microcystin-leucine-arginine causes blood-testis barrier disruption and

degradation of occludin mediated by matrix metalloproteinase-8. *Cell Molecukar Life Sci* 2018;**75**:1117-1132

https://doi.org/10.1007/s00018-017-2687-6

Cheng YH, Wong EWP, Cheng CY. Cancer/testis (CT) antigens, carcinogenesis and spermatogenesis.

Spermatogenesis 2011;**1**:209-220 https://doi.org/10.4161/spmg.1.3.17990

Chiang C, Layer RM, Faust GG, Lindberg MR, Rose DB, Garrison EP, Marth GT, Quinlan AR, Hall IM. SpeedSeq:

Ultra-fast personal genome analysis and interpretation. *Nat Methods* 2015;**12**:966-968

https://doi.org/10.1038/nmeth.3505.

Dadsena S, King LE, Garcia-Saez AJ. Apoptosis regulation at the mitochondria membrane level. *Biochim Biophys*

Acta – Biomembranes 2021;**1863**:183718 https://doi.org/10.1016/j.bbamem.2021.183716

Damaschke NA, Yang B, Bhusari S, Avilla M, Zhong W, Blute ML, Huang W, Jarrard DF. Loss of Igf2 Gene Imprinting

in Murine Prostate Promotes Widespread Neoplastic Growth. Cancer Res 2017;**77**:5236-5247

https://doi.org/10.1158/0008-5472.CAN-16-3089

Drechsler H, Xu Y, Geyer VF, Zhang Y, Diez S. Multivalent electrostatic microtubule interactions of synthetic

peptides are sufficient to mimic advanced MAP-like behavior. *Mol Biol Cell* 2019;**30**:2953-2968

https://doi.org/10.1091/mbc.E19-05-0247

Du Plessis SS, Gokul S, Agarwal A. Semen hyperviscosity: causes, consequences, and cures. *Front Biosci (Elite Ed)*

2013;**5**:224-231 https://doi.org/10.2741/E610

Emsley P, Lohkamp B, Scott WG, Cowtan K. Features and development of Coot. *Acta Crystallogr D Struct Biol*

2010;**66**:486-501 https://doi.org/10.1107/S0907444910007493

Fink E, Bhoola KD, Snyman C, Neth P, Figueroa CD. Cellular expression of plasma prekallikrein in human tissues.

Biol Chem 2007;**388**:957-963 https://doi.org/10.1515/BC.2007.104

Firman RC, Simmons L. W. Sperm midpiece length predicts sperm swimming velocity in house mice. Biol Letters

2010;**6**:513-516 https://doi.org/10.1098/rsbl.2009.1027

Ford WC. Glycolysis and sperm motility: does a spoonful of sugar help the flagellum go round? Hum Reprod

Update 2006;12:269-274 https://doi.org/10.1093/humupd/dmi053

Frungieri MB, Calandra RS, Mayerhofer A, Matzkin ME. Correlation between testicular mast cell count and

spermatogenic epithelium in non-obstructive azoospermia. *Reproduction* 2015;**149**:R169-R180

https://doi.org/10.1111/iep.12261

Fung E, Richter C, Yang HB, Schäffer I, Fischer R, Kessler BM, Bassermann F, D'Angiolella V. FBXL13 directs the

proteolysis of CEP192 to regulate centrosome homeostasis and cell migration. *EMBO Rep* 2018;**19**:e44799

https://doi.org/10.15252/embr.201744799

Ganetzky RD, Stendel C, McCormick EM, Zolkipli-Cunningham Z, Goldstein AC, Klopstock T, Falk MJ. MT-ATP

mitochondrial disease variants: phenotypic and biochemical features analysis in 218 published cases and cohort

of 14 new cases. *Hum Mutation* 2019;**40**:499-515   https://doi.org/10.1002/humu.23723

Gao Y, Hong J, Guo Y, Chen M, Chang AK, Xie L, Ying X. Assessment spermatogenic cell apoptosis and the

transcript levels of metallothionein and p53 in Meretrix meretrix induced by cadmium. *Ecotoxicol Environ Saf*

2021;**217**:112230 https://doi.org/10.1016/j.ecoenv.2021.112230

Ghosh D, Najwa AR, Khan MA, Sengupta J. IGF2, IGF binding protein 1, and matrix metalloproteinases 2 and 9 in

implantation-stage endometrium following immunoneutralization of vascular endothelial growth factor in the

rhesus monkey. *Reproduction* 2011;**141**:501-509 https://doi.org/10.1530

Girardet L, Augière C, Asselin MP, Belleannée C. Primary cilia: biosensors of the male reproductive tract.

Andrology 2019;7:588-602 https://doi.org/10.1111/andr.12650

Gu NH, Zhao WL, Wang GS, Sun F. Comaprative analysis of mammalian sperm ultrastructure reveals relationships

between sperm morphology, mitochondrial functions and motility. *Reprod Biol Endocrinol* 2019;**17**:66

https://doi.org/10.1186/s12958-019-0510-y

Gui L, Song K, Tritschler D, Bower R, Yan S, Dai A, Augspurger K, Sakizadeh J, Grzemska M, Ni T, et al. Scaffold

subunits support associated subunit assembly in the Chlamydomonas ciliary nexin-dynein regulatory complex.

Proc Natl Acad Sci USA 2019;**116**:23152-23162 https://doi.org/10.1073/pnas.1910960116

Ha S, Lindsay AM, Timms AE, Beier DR. Mutations in Dnaaf1 and Lrrc48 Cause Hydrocephalus, Laterality Defects,

and Sinusitis in Mice. *G3* 2016;**6**:2479-2487 https://doi.org/10.1534/g3.116.030791

Hardy JJ, Wyrwoll MJ, Mcfadden W,· Malcher A,· Rotte N, · Pollock NC, Munyoki S, Veroli MV, Houston BJ, · Miguel

J et al. XavierVariants in GCNA, X-linked germ-cell genome integrity gene, identified in men with primary

spermatogenic failure. *Hum Genet* 2021;140:1169-1182 https://doi.org/10.1007/s00439-021-02287-y

Heidari MM, Khatami M, Danafar A, Dianat T, Farahmand G, Talebi AR. 4Mitochondrial Genetic Variation in

Iranian Infertile Men with Varicocele. *Int J Fertil Steril* 2016;**10**:303-309 doi: 10.22074/ijfs.2016.5047.

Hernández-Llodrà S, Juanpere N, de Muga S, Lorenzo M, Gil J, Font-Tello A, Agell L, Albero-González R, Segalés L,

Merino J, et al. ERG overexpression plus SLC45A3 (prostein) and PTEN expression loss: Strong association of the

triple hit phenotype with an aggressive pathway of prostate cancer progression. *Oncotarget* 2017;**8**:74106-74118

https://doi.org/10.18632/oncotarget.18266

Heshusius, S, Grech L, Gillemans N, Brouwer RWW, den Dekker XT, van IJcken WFJ, Nota B, Felice AE, van Dijk

TB, von Lindern M, et al. Epigenomic analysis of KLF1 haploinsufficiency in primary human erythroblasts. *Sci Rep*

2022;**12**:336 https://doi.org/10.1038/s41598-021-04126-6

Heuser T, Raytchev M, Krell J, Porter ME, Nicastro D. The dynein regulatory complex is the nexin link and a major

regulatory node in cilia and flagella. *J Cell Biol* 2009;**187**:921-933 https://doi.org/10.1083/jcb.200908067

Honda A, Komuro H, Shimada A, Hasegawa T, Seko Y, Nagase H, Hozumi I, Inuzuka T, Hara H, Fujiwara Y, et al.

Attenuation of cadmium-induced testicular injury in metallothionein-III null mice. *Life Sci* 2010;**87**:545-550

https://doi.org/10.1016/j.lfs.2010.09.009

Houston BJ, Riera-Escamilla A, Wyrwoll MJ, Salas-Huetos A, Xavier MJ, Nagirnaja L,Friedrich C, Conrad DF, Aston

AI, Krausz C et al. A systematic review of the validated monogenic causes of human male infertility: 2020 update

and a discussion of emerging gene–disease relationships. *Hum Reprod Update* 2021;28:15-29

https://doi.org/10.1093/humupd/dmab030

Huang Y, Huang Y, Pan XW, Li L, Lu Chen, Liu X, Lu JL, Zhu XM, Huang H, Yang QW, Ye JQ et al. Overexpression of

USP39 predicts poor prognosis and promotes tumorigenesis of prostate cancer via promoting EGFR mRNA

maturation and transcription elongation. *Oncotarget* 2016;**7**:22016-22030

https://doi.org/10.18632/oncotarget.7882

Hull RL, Johnson PY, Braun KR, Day AJ, Wight TN. Hyaluronan and hyaluronan binding proteins are normal

components of mouse pancreatic islets and are differentially expressed by islet endocrine cell types. *J Histochem*

Cytochem 2021;**60**:749-760 https://doi.org/10.1369/002215541245704

Hwang K, Smith JF, Coward RM, Penzias A, Bendikson K, Butts S, Coutifaris C, Falcone T, Fossum G, Gitlin S et al

(Practice Committee of the American Society for Reproductive Medicine in collaboration with the Society for

Male Reproduction and Urology) Evaluation of the azoospermic male: a committee opinion. *Fertil Steril*

2018;109:777-782 https://doi.org/10.1016/j.fertnstert.2018.01.043

Ioannidis NM, Rothstein JH, Pejaver V, Middha S, McDonnell SK, Baheti S, Musolf A, Li Q, Holzinger E, Karyadi D,

et al. REVEL: An Ensemble Method for Predicting the Pathogenicity of Rare Missense Variants. *Am J Hum Genet*

2016;**99**:877-885 https://doi.org/10.1016/j.ajhg.2016.08.016

Ishikawa T. Axoneme structure from motile cilia. *Cold Spring Harbor Perspectives in Biology* 2017;**9**,a028076

https://doi.org/10.1101/cshperspect.a028076

Jia Y, Ohanyan A, Lue YH, Swerdloff RS, Liu PY, Cohen P, Wang C. The effects of humanin and its analogues on

male germ cell apoptosis induced by chemotherapeutic drugs. *Apoptosis* 2015;**20**:551-561

https://doi.org/10.1007/s10495-015-1105-5

Jodar M, Oliva R. Protamine alterations in human spermatozoa. *Adv Exp Med Biol* 2014;**791**:83-102

https://doi.org/10.1007/978-1-4614-7783-9_6

Jonkers J, Pai P, Sukumar S. Multiple roles of HOX proteins in Metastasis: Let me count the ways. *Cancer*

Metastasis Rev 2020;**39**:661-679 https://doi.org/10.1007/s10555-020-09908-4

Jumper J Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, Žídek A, Potapenko

A et al. Highly accurate protein structure prediction with AlphaFold. *Nature* 2021;**596**:583-589

https://doi.org/10.1038/s41586-021-03819-2

Karmilin K, Schmitz C, Kuske M, Körschgen H, Olf M, Meyer K, Hildebrand A, Felten M, Fridrich S, Yiallouros I, et

al. Mammalian plasma fetuin-B is a selective inhibitor of ovastacin and meprin metalloproteinases. *Sci Rep*

2019;**9**:546 https://doi.org/10.1038/s41598-018-37024-5

Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJ. The Phyre2 web portal for protein modeling, prediction

and analysis. *Nat Protoc* 2015;**10**:845-858 https://doi.org/10.1038/nprot.2015.053.

Khelifa MB, Coutton C, Zouari R, Karaouzène T, J, Bidart, M, Yassine SV, Delaroche J, Hennebicq S, et al. Mutations

in DNAH1, which encodes an inner arm heavy chain dynein, lead to male infertility from multiple morphological

abnormalities of the sperm flagella. *Am J Hum Genet* 2014;**94**:95-104

http://dx.doi.org/10.1016/j.ajhg.2013.11.017

Kherraf ZE, Christou-Kent M, Karaouzene T, Amiri-Yekta A, Martinez G, S Vargas AS, Lambert E, Borel C, Dorphin

B, Aknin-Seiferet I, et al. SPINK2 deficiency causes infertility by inducing sperm defects in heterozygotes and

azoospermia in homozygotes. *EMBO Mol Med* 2017;**9**:1132-1149 https://doi.org/10.15252/emmm.201607461

Klein C, Troedsson MHT, Rutllant J. Region-Specific Expression of Aquaporin Subtypes in Equine Testis,

Epididymis, and Ductus Deferens. *Anatomical Records (Hoboken)* 2013;**296**:1115-1126

https://doi.org/10.1002/ar.22709

Kobe B, Kajava AV. The leucine-rich repeat as a protein recognition motif. *Curr Opin Struct Biol* 2001;**11**:725-732

https://doi.org/10.1016/S0959-440X(01)00266-4

Krausz C, Riera-Escamilla A. Genetics of male infertility. *Nat Rev Urol* 2018;15: 369-384

https://doi.org/10.1038/s41585-018-0003-3

Krissinel E, Henrick K. Secondary-structure matching (SSM), a new tool for fast protein structure alignment in

three dimensions. *Acta Crystallogr D Biol Crystallogr* 2004;**60**:2256-2268

https://doi.org/10.1107/S0907444904026460

Kubo T, Oda T. Electrostatic interaction between polyglutamylated tubulin and the nexin-dynein regulatory

complex regulates flagellar motility. *Mol Biol Cell* 2017;**28**:2260-2266 https://doi.org/10.1091/mbc.e17-05-0285

Kumar D. Reiter J. How the centriole builds its cilium: of mothers, daughters, and the acquisition of appendages.

Curr Opin Strucl Biol 2021;66:41-48 https://doi.org/10.1016/j.sbi.2020.09.006

Leal LF, Szarek E, Berthon A, Nesterova M, Faucz FR, London E, Mercier C, Abu-Asab M, Starost MF, Dye L, et al.

Pde8b haploinsufficiency in mice is associated with modest adrenal defects, impaired steroidogenesis, and male

infertility, unaltered by concurrent PKA or Wnt activation. *Mol Cellular Endocrinol* 2021;**522**:111117

https://doi.org/10.1016/j.mce.2020.111117

Lei Q, Huanga X, Zhenga L, Zhenga F, Donga J, Chena F, Zeng W. Biosensors for Caspase-3: From chemical

methodologies to biomedical applications. *Talanta* 2022;240:123198

https://doi.org/10.1016/j.talanta.2021.123198

Lewis WR, Malarkey EB, Tritschler D, Bower R, Pasek RC, Porath JD, Birket SE, Saunier S, Antignac C, Knowles MR,

et al. Mutation of Growth Arrest Specific 8 Reveals a Role in Motile Cilia Function and Human Disease. *PLoS Genet*

2016;**12**:e1006220 https://doi.org/10.1371/journal.pgen.1006220

Li H, Xing L, Zhao N, Wang J, Zheng N. Furosine Induced Apoptosis by the Regulation of STAT1/STAT2 and

UBA7/UBE2L6 Genes in HepG2 Cells. *Int J Mol Sci* 2018;**19**:1629 https://doi.org/10.3390/ijms19061629

Li K, van Delft MF, Dewson G. Too much death can kill you: inhibiting intrinsic apoptosis to treat disease. *EMBO*

J 2021;**40**:e107341 https://doi.org/10.15252/embj.2020107341

Lorès P, Kherraf ZE, Amiri-Yekta A, Whitfeld M, Daneshipour A, Stouvenel L, Cazin C, Cavarocchi E, Coutton C,

Llabador MA et al. A missense mutation in IFT74, encoding for an essential component for intraflagellar transport

of Tubulin, causes asthenozoospermia and male infertility without clinical signs of Bardet-Biedl syndrome. *Hum*

Genet 2021;**140**:1031-1043 https://doi.org/10.1007/s00439-021-02270-7

Lue Y, Swerdloff R, Jia Y, Wang C. The emerging role of mitochondrial derived peptide humanin in the testis.

Biochim Biophys Acta Gen Subj 2021;**1865**:130009 https://doi.org/10.1016/j.bbagen.2021.130009

Lüpold S, de Boer RA, Evans JP, Tomkins JL, Fitzpatrick JL. How sperm competition shapes the evolution of testes

and sperm: a meta-analysis. *Philos Trans R Soc Lond B Biol Sci* 2020;**375**:20200064

https://doi.org/10.1098/rstb.2020.0064

Markou, A, Unger L, Abir-Awan M, Saadallah A, Halsey A, Baklava Z, Conner M, Törnroth-Horsefield S, Greenhill

SD, Alex Conneret A, et al. Molecular mechanisms governing aquaporin relocalisation. *Biochim Biophys Acta –*

Biomembr 2022;**1864**:183853 https://doi.org/10.1016/j.bbamem.2021.183853

Marques PI, Fonseca F, Carvalho AS, Puente DA, Damião I, Almeida V, Barros N, Barros A, Carvalho F, Azkargortaet

M et al. Sequence variation at KLK and WFDC clusters and its association to semen hyperviscosity and other male

infertility phenotypes. *Hum Reprod* 2016;**31**:2881-2891 https://doi.org/10.1093/humrep/dew267

Martinez G, Coutton C, Loeuillet C, Cazin C, Muroňová J, Boguenet M, Lambert E, Dhellemmes M, Chevalier G,

Hograindleur JP et al. Oligogenic heterozygous inheritance of sperm abnormalities in mouse. *Elife*

2022;**11**:e75373 https://doi.org/10.7554/eLife.75373

Matsushima N, Takatsuka S, Miyashita H, Kretsinger RH. Leucine rich repeat proteins: sequences, mutations,

structures and diseases. *Protein Pept Lett* 2019;**26**:108-131

https://doi.org/10.2174/0929866526666181208170027

Matsuzaki, H., Hitomi Matsuzaki, Kuramochi D, Okamura E, Hirakawa K, Ushiki A, Tanimoto K. Recapitulation of

gametic DNA methylation and its post-fertilization maintenance with reassembled DNA elements at the mouse

Igf2/H19 locus. *Epigenetics Chromatin* 2020:**13**:2 https://doi.org/10.1186/s13072-019-0326-1

Matzuk MM, Lamb DJ. The biology of infertility: research advances and clinical challenges. *Nat Med*

2008;14:1197-1213 https://doi.org/10.1038/nm.f.1895

McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GRS, Thormann A, Flicek P, Cunningham F. The Ensembl Variant Effect

Predictor. *Genome Biol* 2016;**17**:122 https://doi.org/10.1186/s13059-016-0974-4

Mei S, Wu Y, Wang Y, Cui Y, Zhang M, Zhang T, Huang X, Yu S, Yu T. Disruption of PIKFYVE causes congenital

cataract in human and zebrafish. *Elife* 2022;**11**:e71256 https://doi.org/10.7554/eLife.71256

Michaelis M, Sobczak A, Koczan D, Langhammer M, Reinsch N, Schoen J, WeitzeL JM. Selection for female traits

of high fertility affects male reproductive performance and alters the testicular transcriptional profile. *BMC*

Genomics 2017;**18**:889 https://doi.org/10.1186/s12864-017-4288-z

Mo RJ, Lu JM, Wan YP, Hua W, Liang YX, Zhuo YJ, Kuang QW, Liu YL, He HC, Zhong WD. Decreased HoxD10

Expression Promotes a Proliferative and Aggressive Phenotype in Prostate Cancer. *Curr Mol Med* 2017;**17**:70-78

https://doi.org/10.2174/1566524017666170220104920

Morohoshii A, Miyataid H, Shimadaid K, Nozawa K, Matsumura T, Yanaseid R, Shibaid K, Inabaid K, Ikawa M.

Nexin-dynein regulatory complex component DRC7 but not FBXL13 is required for sperm flagellum formation

and male fertility in mice. *PloS Genet* 2020;**16**:e1008585   https://doi.org/10.1371/journal.pgen.1008585

Mostafa T, Rashed L, Taymour M. Seminal cyclooxygenase relationship with oxidative stress in infertile

oligoasthenoteratozoospermic   men   with   varicocele.   *Andrologia*   2016;**48**:137142

https://doi.org/10.1111/and.12423

Nagirnaja L, Aston KI, Conrad DF. Genetic intersection of male infertility and cancer. *Fertil Steril* 2018;**109**:20-26

https://doi.org/10.1016/j.fertnstert.2017.10.028

Nascimento JM, Shi LZ, Tam J, Chandsawangbhuwana C, Durrant B, Botvinick EL, Berns MW. Comparison of

glycolysis and oxidative phosphorylation as energy sources for mammalian sperm motility, using the combination

of fluorescence imaging, laser tweezers, and real-time automated tracking and trapping. *J Cell Physiol*

2008;**217**:745-51 https://doi.org/10.1002/jcp.21549

Naz RK, Rajesh C. Gene knockouts that cause female infertility: search for novel contraceptive targets. *Front*

Biosci 2005;**10**:2447-2459 https://doi.org/10.2741/1709

Neirijnck Y, Kühne F, Mayère C, Pavlova E, Sararols P, Foti M, Atanassova N, Nef S. Tumor Suppressor PTEN

Regulates Negatively Sertoli Cell Proliferation, Testis Size, and Sperm Production In Vivo. *Endocrinology*

2019;**160**:387-398 https://doi.org/10.1210/en.2018-00892

Ng ACY, Eisenberg JM, Heath RJ, Huett A, Robinson CM, Nau GJ, Xavier RJ. Human leucine-rich repeat proteins:

a genome-wide bioinformatic categorization and functional analysis in innate immunity. *Prot Nat Acad Sci*

2011;**108**:4631-4638 https://doi.org/10.1073/pnas.100009310

Ogura, Y., Uehara T, Ujibe K, Yoshihashi H, Yamada M, Suzuki H, Takenouchi T, Kosaki K, Hirata H. The

p.Thr395Met missense variant of NFIA found in a patient with intellectual disability is a defective variant.

American Journal of Medical Genetics A 2022;**188**:1184-1192 https://doi.org/10.1002/ajmg.a.62638

Oud MS, Houston BJ, Volozonoka L, Mastrorosa FK, Holt GS, Alobaidi BKS, deVries PF, Astuti G, Ramos L,

Mclachlan RI et al. Exome sequencing reveals variants in known and novel candidate genes for severe sperm

motility disorders. *Hum Reprod* 2021;36:2597-2611 https://doi.org/10.1093/humrep/deab099

Pan W, Wang K, Li J, Li H, Cai Y, Zhang M, Wang A, Wu Y, Gao W, Weng W. 1Restoring HOXD10 Exhibits

Therapeutic Potential for Ameliorating Malignant Progression and 5-Fluorouracil Resistance in Colorectal Cancer.

Front Oncol 2021;**11**:771528 https://doi.org/10.3389/fonc.2021.771528

Perner S, Rupp NJ, Braun M, Rubin MA, Moch H, Dietel M, Wernert N, Jung K, Stephan C, Kristiansen G. Loss of

SLC45A3 protein (prostein) expression in prostate cancer is associated with SLC45A3-ERG gene rearrangement

and an unfavorable clinical course. *Int J Cancer* 2013;132;807-812 https://doi.org/10.1002/ijc.27733

Perrotta I, Santoro M, Guido C, Avena P, Tripepi S, De Amicis F, Gervasi MC, Aquila S. Expression of

cyclooxygenase-1 (COX-1) and COX-2 in human male gametes from normal patients, and those with varicocele

and diabetes: a potential molecular marker for diagnosing male infertility disorders. *J Anat* 2012;**221**:209-220

https://doi.org/10.1111/j.1469-7580.2012.01534.x

Pin E, Frauke Henjes F, Hong MG, Wiklund F, Patrik Magnusson P, Anders Bjartell A, Uhlen M, Nilsson P, Schwenk

JM. Identification of a Novel Autoimmune Peptide Epitope of Prostein in Prostate Cancer. *J Proteome Res*

2017;**16**:04-216 https://doi.org/10.1021/acs.jproteome.6b00620

Ramli NSK, Giribabu N, Karim K, Salleh N. Hormonal control of vas deferens fluid volume and aquaporin

expression in rats. *J Mol Histol* 2019;**50**:21-34 https://doi.org/10.1007/s10735-018-9804-1

Richards S, Aziz N, Sherri Bale S, Bick D, Das S, Gastier-Foster J, Grody WW, Hegde M P, Lyon E, Spector E, et al.

Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the

American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med*

2015;**17**:405-424 https://doi.org/doi:10.1038/gim.2015.30

Romeis B, Mulisch M, Welsch U. Romeis Mikroskopische Technik. Dordrecht: Springer. 2010; xii, 550 s. p.

Roy A, Kucukural A, Zhang Y. I-TASSER: a unified platform for automated protein structure and function

prediction. *Nat Protoc* 2010;**5**:725-738 https://doi.org/10.1038/nprot.2010.5

Salustri A, Campagnolo L, Klinger FG, Camaioni A. Molecular organization and mechanical properties of the

hyaluronan matrix surrounding the mammalian oocyte. *Matrix Biol* 2019;78-79:11-23

https://doi.org/10.1016/j.matbio.2018.02.002

Samaiya PK, Krishnamurthy S, Kumar A. Mitochondrial dysfunction in perinatal asphyxia: role in pathogenesis

and potential therapeutic interventions. *Mol Cell Biochem* 2021;**476**:4421-4434 https://doi.org/10.1007/s11010-

021-04253-8

Sandhu N, Rana S, Meena K. Nuclear receptor subfamily 5 group A member 2 (NR5A2): role in health and

diseases. *Mol Biol Rep* 2021;**48**:8155-8170 https://doi.org/10.1007/s11033-021-06784-1

Sandy Z, da Costa IC, Schmidt CK. More than Meets the ISG15: Emerging Roles in the DNA Damage Response and

Beyond. *Biomolecules* 2020;**10**:1557 https://doi.org/10.3390/biom10111557

Satir P, Heuser T, Sale WS. A structural basis for how motile cilia beat. *Bioscience* 2014;**64**:1073-1083

https://doi.org/10.1093/biosci/biu180

Sato JD, Kan M. Media for culture of mammalian cells. *Curr Protoc Cell Biol* 2001; Chapter 1:Unit 1.2.

Schlegel PN, Sigman M, Collura B, De Jonge CJ, Eisenberg ML, Lamb DL, Sandlow JI, Sokol RZ, Spandorfer SD,

Tanrikut C, et al. Diagnosis and treatment of infertility in men: AUA/ASRM guideline part I. *Fertil Steril* 2021;**115**,

54–61 https://doi.org/10.1016/j.fertnstert.2020.11.016

Sironen A, Shoemark A, Patel M, Loebinger MR, Mitchison HM. Sperm defects in primary ciliary dyskinesia and

related causes of male infertility. *Cell Mol Life Sci* 2020;77:2029-2048 https://doi.org/10.1007/s00018-019-

03389-7

Smith CEL, Lake ARV, Johnson CA. Primary Cilia, Ciliogenesis and the Actin Cytoskeleton: A Little Less Resorption,

A Little More Actin Please. *Front Cell Dev Biol* 2020;8:622822 https://doi.org/10.3389/fcell.2020.622822

Supernat A, Vidarsson OV, Steen VM, Stokowy T. Comparison of three variant callers for human whole genome

sequencing. *Sci Rep* 2018;**8**:17851 https://doi.org/10.1038/s41598-018-36177-7

Szklarczyk D, Gable AL, Lyon D, Junge A, Wyder S, Jaie Huerta-Cepas J, Simonovic M, Doncheva NT, Morris JH,

Bork P et al. STRING v11: protein-protein association networks with increased coverage, supporting functional

discovery   in   genome-wide   experimental   datasets.   *Nucleic   Acid   Res*   2019;**47**:D607-D613

https://doi.org/10.1093/nar/gky1131

Tan JHL, Wollmann H, van Pelt AMM, Kaldis P, Messerschmidt DM. Infertility-Causing Haploinsufficiency Reveals

TRIM28/KAP1   Requirement   in   Spermatogonia.   *Stem   Cell   Rep*   2020;**14**:818-827

https://doi.org/10.1016/j.stemcr.2020.03.013

Touré A, Martinez G, Kherraf ZE, Cazin C, Beurois J, Arnoult C, Ray PF, Coutton C. The genetic architecture of

morphological abnormalities of the sperm tail. *Hum Genet* 2021;**140**:21-42 https://doi.org/10.1007/s00439-020-

02113-x

Tourmente M, Villar-Moya P, Rial E, Roldan ER. 2015 Differences in ATP generation via glycolysis and oxidative

phosphorylation and relationships with sperm motility in mouse species. J Biol Chem 2015;290:20613–20626

https://doi.org/10.1074/jbc.M115.664813

Tunyasuvunakool K, Adler J, Wu Z, Green T, Zielinski M, Žídek A, Bridgland A, Cowie A, Meyer C, Laydon A, et al.

Highly accurate protein structure prediction for the human proteome. *Nature* 2021;**596**:590-596

https://doi.org/10.1038/s41586-021-03828-1

UniProt, C. UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Res* 2019;**47**:D506-D15

https://doi.org/10.1093/nar/gky1049

Valassina N, Brusco S, Salamone A, Serra L, Luoni M, Giannelli S, Bido S, Massimino L, Ungaro F, Mazzara PG, et

al. Scn1a gene reactivation after symptom onset rescues pathological phenotypes in a mouse model of Dravet

syndrome. *Nat Commun* 2022*;***13**:161 https://doi.org/10.1038/s41467-021-27837-w

Vander Borght M,Wyns C. Fertility and infertility: Definition and epidemiology. *Clin Biochem* 2018;**62**: 2-10

Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, Yuan D, Stroe O, Wood G, Laydoet A et al.

AlphaFold Protein Structure Database: massively expanding the structural coverage of protein-sequence space

with high-accuracy models. *Nucleic Acids Res* 2022;**50**:D439-D444 https://doi.org/10.1093/nar/gkab1061

von Mering C, Jensen LJ, Snel B, Hooper SD, Krupp M, Foglierini M, Jouffre N, Huynen MA, Bork P. STRING: known

and predicted protein-protein associations, integrated and transferred across organisms. *Nucleic Acid Res*

2005;**33**:D433-443 https://doi.org/10.1093/nar/gki005

Wagner K, Unger L, Salman MM, Kitchen P, Bill RM, Yool AJ. Signaling Mechanisms and Pharmacological

Modulators Governing Diverse Aquaporin Functions in Human Health and Disease. *Int J Mol Sci* 2022;**23**:1388

https://doi.org/10.3390/ijms23031388

Wang C, Gu Y, Zhang K, Xie K, Zhu M, Dai N, Jiang Y, Guo X, Liu M, Dai J, et al. Systematic identification of genes

with a cancer-testis expression pattern in 19 cancer types. *Nat Commun* 2016;**7**:10499

https://doi.org/10.1038/ncomms10499

Wertheimer E, Krapf D, de la Vega-Beltran, Sánchez-Cárdenas C, Navarrete F, Haddad D, Escoffier J, Salicioni AM,

Levin LR, Buck J, Mager J, et al. Compartmentalization of distinct cAMP signaling pathways in mammalian sperm.

J Biol Chem 2013;**288**:35307-35320 https://doi.org/10.1074/jbc.M113.489476

WHO 2020: https://www.who.int/news-room/fact-sheets/detail/infertility

Wilson KS, Gonzalez O, Dutcher SK, Bayly PV. Dynein-deficient flagella respond to increased viscosity with

contrasting changes in power and recovery strokes. *Cytoskeleton (Hoboken)* 2015;**72**:477-490

https://doi.org/10.1002/cm.21252

Wirschell M, Olbrich H, Werner C, Tritschler D, Bower R, Sale WS, Loges NT, Pennekamp P, Lindberg S, Stenram

U, et al. The nexin-dynein regulatory complex subunit DRC1 is essential for motile cilia function in algae and

humans. *Nat Genet* 2013;45:262-268 https://doi.org/10.1038/ng.2533

Wyrwoll MJ, Temel SG, Nagirnaja L, Oud MS, Lopes AM, van der Heijden GW, Heald JS, Rotte N, Wistuba J,Marius

Wöste M et al. Bi-allelic Mutations in M1AP Are a Frequent Cause of Meiotic Arrest and Severely Impaired

Spermatogenesis Leading to Male Infertility. *Am J Hum Genet* 2020;107:342–351

https://doi.org/10.1016/j.ajhg.2020.06.010

Xavier MJ, Salas-Huetos A, Oud MS, Aston KI, Veltman JA. Disease gene discovery in male infertility: past, present

and future. *Hum Genet* 2021;140:7-19 https://doi.org/10.1007/s00439-020-02202-x

Xu W, Zhang Y, Qin D, Gui Y, Wang S, Du G, Yang F, Li L, Yuan S, Wang M, et al. Transcription factor-like 5 is a

potential DNA- and RNA-binding protein essential for maintaining male fertility in mice. *J Cell Sci*

2022;**135**:jcs259036 https://doi.org/10.1242/jcs.259036

Xue M, Xue M, Fang Y, Sun G, Zhuo W, Zhong J, Qian C, Wang L, Wang L, Si J, Shujie Chen. IGFBP3, a transcriptional

target of homeobox D10, is correlated with the prognosis of gastric cancer. *PLoS One* 2013;**8**:e81423

https://doi.org/10.1371/journal.pone.0081423

Yang J, Yan R, Roy A, Xu D, Poisson J, Zhang Y. The I-TASSER Suite: protein structure and function prediction. *Nat*

*Methods 2015;***12;**7-8 https://doi.org/10.1038/nmeth.3213

Yang J, Zhang Y. I-TASSER server: new development for protein structure and function predictions. Nucleic *Acids*

Res 2015;**43**:W174-181 https://doi.org/10.1093/nar/gkv342

Yang LF, Yang F, Zhang FL, Xie YF, Hu ZX, Huang SL, Shao ZM, Li DQ. Discrete functional and mechanistic roles of

chromodomain Y-like 2 (CDYL2) transcript variants in breast cancer growth and metastasis. *Theranostics*

2020;**10**:5242-5258 https://doi.org/10.7150/thno.43744

Yin, M., O'Neil LAJ. The role of the electron transport chain in immunity. *FASEB J* 2021;**35**:e21974

https://doi.org/10.1096/fj.202101161R

Yuan J, Li X, Zhang G, Cheng W, Wang W, Lei Y, Ma Q, Song G. USP39 mediates p21-dependent proliferation and

neoplasia of colon cancer cells by regulating the p53/p21/CDC2/cyclin B1 axis. *Mol Carcinog*enesis 2021;**60**:265-

278 https://doi.org/10.1002/mc.23290

Zhan K, Liu R, Tong H, Gao S, Yang G, Hossain A, Li T, He W. Fetuin B overexpression suppresses proliferation,

migration, and invasion in prostate cancer by inhibiting the PI3K/AKT signaling pathway. *Biomed Pharmacother*

2020;**131**:110689 https://doi.org/10.1016/j.biopha.2020.110689

Zhang J, Liu S, Zhang D, Ma Z, Sun L. Homeobox D10, a tumor suppressor, inhibits the proliferation and migration

of esophageal squamous cell carcinoma. *J Cell Biochem* 2019;**120**:13717-13725

https://doi.org/10.1002/jcb.28644

Zhang Z, Miao J, Wang Y. Mitochondrial regulation on spermatogenesis. Reproduction 2022;**163**:R55-R69

https://doi.org/10.1530/REP-21-0431

Zhao Y, Zhang B, Lei Y, Sun J, Zhang Y, Sen Yang S, Zhang X. Knockdown of USP39 induces cell cycle arrest and

apoptosis in melanoma. *Tumour Biol* 2016;**37**:13167-13176 https://doi.org/10.1007/s13277-016-5212-x

Zorrilla M, Yatsenko AN. The Genetics of Infertility: Current Status of the Field. *Curr Genet Med Rep* 2013;1:247–

260 DOI 10.1007/s40142-013-0027-1

1191

1192

1193

1194

1195

1196   **Figures**

1197

1198   **Figure 1: Characteristics of *Tcte1* gene. A** Putative homologs of *Tcte1* gene conserved in Bilateria, according to

1199   HomoloGene (8434), including collation of proteins and their conserved domains; **B** Sequence alignement

1200   between human, mouse and rat; **C** NCBI Gene schematic representation of *Tcte1* gene in *Mus musculus*.

1201

Figure 2: Characteristics of *Tcte1* mice knockout. A Schematic representation of gRNA localization in

CRISPR/Cas9 KO creation; **B** Relative expression of *Tcte1* gene in wild type (WT), hetero- [+/-] and homozygous

mice [-/-] (Real-time PCR); **** $p<0.0001$, ** $p<0.01$; **C** Example of genotyping result. Lane 1 – 1kb marker

(GeneRuler), lanes 2, 3 – wild type (WT), lanes 4, 5 – hetero- $^{(+/-)}$, and lanes 6, 7 – homozygous mice $^{(-/-)}$; lanes: 2,

4, 6 identified the lack (0 bp) or presence of mutation (630 bp), lanes: 3, 5, 7 differentiated hetero- (658 bp) and

homozygotes (0 bp).

1208

Figure 3: Reproductive potential of KO *Tcte1* mice. Mating combinations and their results including: time of

breeding, number of litters and pups, number of male and female pups, and genotypes observed in pups of each

combination. **A** Mean time: from mating to 1st pedigree, and between particular litters; **B** Ratio of males and

females per litter in each mating combination; **C** Frequency of males and females and their genotype observed

each mating combination; left panel: according to sex of pups; right panel: according to mating combinations; **D**

Ratio of males and females per litter and their genotype observed each mating combination.

*** $p < 0.0001$, ** $0.0001 < p < 0.01$, * $0.01 < p < 0.05$

1216

Figure 4: Characteristics of *Tcte1* wild type (WT), hetero- $^{(+/-)}$ and homozygous mice $^{(-/-)}$. A Histopathology of

testes and epididymises (caput); staining: Masson-Goldner protocol; microscope: Leica DM5500, objective 63×,

magnification 630×; **B** Comparison of body mass, testis weight and size, and epididymis length, followed by gross

morphology of gonads; **C** Sperm parameters: concentration, motility and morphology comparison; statistical

marks indicated differences according to WT animals; **D** Frequency of morphological defects according to three

part of the spermatozoa; **E** Examples of main sperm morphological abnormalities found (all abnormalities

observed available in Supplementary file 7); **F** Comparison of the mean length of: sperm tail, midpiece and ratio

tail:midpiece; *** $p < 0.0001$, ** $0.0001 < p < 0.01$, * $0.01 < p < 0.05$

1225

1226 **Figure 5: RNAseq results of changes in gene expression level found in testes of KO *Tcte1*$^{+/-}$ and *Tcte1*$^{-/-}$ mice vs.**

1227 **control (WT).** Volcano plots: black dots indicated all analyzed genes within the genome, red dots represented

1228 genes with significantly changed expression level (followed by the heatmap and the table). Filtering criteria: p

1229 val <0.05, log ratio >1 or <-1 (adequate to fold change value of 2 (expression level increased) or 0.5 (expression

1230 level decreased)), expression level >8. nd – no data concerning function or disease.

1231

1232 **Figure 6: STRING analyzis of potential interactions between proteins with observed differences in gene**

1233 **expression level in RNAseq for mouse testis of the KO model for *Tcte1* gene.** Genes revealed in RNAseq were

1234 bracketed. (STRING database, 24th Feb 2021).

1235

Figure 7: Immunolocalization of the nexin-dynein regulatory complex (N-DRC) proteins: Tcte1 (Drc5), Drc7,

Fbxl13 (Drc6), and Eps8l1 (Drc3) within mouse sperm cells. A. Localization of Tcte1 in spermatozoa from wild

type (WT), hetero- ($^{+/-}$) and homozygous ($^{-/-}$) animals; **B.** Localization of other proteins (Drc7, Fbxl13, Eps8l1)

building the nexin-dynein regulatory complex (N-DRC), responsible for coordination of the dynein arm activity

and stabilization of the doublet microtubules attachment; wild type (WT) spermatozoa. Antibodies: primary (all

anti-rabbit 1:100; Biorbyt, UK): Tcte1 (orb357083); Drc7 (orb58695); Fbxl13 (orb678278); Eps8l1 (orb382538);

secondary: goat anti-rabbit-AF594 conjugated, 1:500, ab150160, Abcam, UK. Fluorescent microscope: Leica

DM5500, filters – DAPI, TxR, FITC, BGR, magnification 630× (with immersion); software: LASX or CytoVision. Bar

represents 10 μm.

1245

1246 **Figure 8: ATP level in wild type (WT), heterozygous ($^{+/-}$) and homozygous ($^{-/-}$) *Tcte1* mice.** Measurement done

1247 in two time points (0 and +60 min.) showed decrease of approximately 35% of luminescence in all three groups.

1248 Mean measured values in homozygous group were approximately 2.4-fold lower. Dots – particular animals; long

1249 dash – mean value for the group; short dashes – standard deviation; RLU – relative luminescence unit. ***

1250 p<0.0001, ** 0.0001 < p < 0.01.

1251

Figure 9: Immunolocalization of caspase 3 within mouse testicular tissue of KO *Tcte1* model. Caspase 3 is one

of the apoptotic indicators becuase of its engagement in the induction, transduction, and amplification of

intracellular apoptotic signals. Caspase 3 signals were found only on spermatogonia in all three genotypes

evaluated (wild type (WT), hetero- ($^{+/-}$) and homozygous ($^{-/-}$)). No differences between the level of fluorescent

signals were found between evaluated mouse genotypes, indicating similar level of apoptosis. Antibodies:

primary (1:100) rabbit anti-Casp3 (Biorbyt, UK); secondary (1:500): goat anti-rabbit-AF594 conjugated

(ab150160, Abcam, UK). Fluorescent microscope: Leica DM5500, filters – DAPI, TxR, magnification 630× (with

immersion); software: LASX. Bar represents 5 μm. Analysed in 3 independent males per zygosity, on 15-20

tubules each.

1261

Figure 10: Homology modeling of TCTE1. A Secondary structure diagrams and associated confidence schema

generated by I-TASSER [46,47,48], Phyre2[45], and AlphaFold [49,50] intensive mode. Positions of amino acids

of interest are indicated. Green, predicted unstructured regions; yellow rectangles, predicted α-helices; orange

arrows and arrowheads, predicted β−strands. Confidence scale ranges from white to cyan, with the probability

of the secondary structure increasing with increasing intensity of cyan; **B** Cartoon diagram of homology model

of human TCTE1 generated by AlphaFold with residues of interest shown as spheres and labeled. N, amino

terminus; C, carboxy terminus; **C-F** Zoomed views of indicated residues and mutations: **C** Ile-125 resides deep

within a hydrophobic pocket, which would be disturbed upon substitution with arginine; **D** Mutation R133C

would impact the hydrogen bond interactions that organize the architecture of the globular domain N-terminal

to the tandem leucine-rich repeats; **E** Mutation of Glu-157 to alanine would destabilize the charge of the surface

of this globular region, likely leading to unfolding; **F** Gly-350 allows for the compact interactions of tandem α-

helices of the LRR. Depending on the rotamer it adopted, a serine side chain in this position would sterically clash

with residues D321, A324, or V346 in the neighboring helix.

1275

Figure 11: Summary of findings revealed for abnormal sperm tail beating in mouse knockout model of *Tcte1*

gene, a component of the N-DRC complex in sperm tail. Three major groups of factors required for proper sperm

tail beating were marked with bold colour arrows (green, yellow and blue). Documented connections (solid grey

arrows) revealed on the basis of: RNAseq data for testis tissues, protein *in silico* predictioning for mutated Tcte1

protein, and measurements of ATP level and sperm cell components in mouse spermatozoa. Unresolved possible

linkages (cause/effect) marked with question (?) mark.

1282

1283

1284

1285

1286

1287

1288

1289

1290

1291

1292

1293

1294

1295

1296

1297

1298

1299

1300

1301

1302

1303

1304

1305

1306

1307

1308   Figure 1

A

| species | gene symbol | Gene ID/ NCBI | UniProt | Indentity (%) | | conserved domains |
				protein	DNA	
						🟦 Leucine-rich repeats (LRRs), cd00116
						🟪 Leucine-rich repeats (LRRs), cl15309
						🟩 CemA family (cl03855)
H. sapiens	TCTE1	202500	Q5JU00			NP_872345.2, 501 aa
vs.						
P. troglodytes	TCTE1	462732	H2QT36	99.2	99.2	XP_518509.2, 501 aa
M. mulatta	TCTE1	702738	F7CLN3	97.0	96.9	XP_001099848.1, 501 aa
C. lupus	TCTE1	609664	-	85.4	87.0	XP_852058.1, 501 aa
B. taurus	TCTE1	523600	-	77.3	80.3	XP_002697348.1, 246 aa
M. musculus	Tcte1	21645	A6H639	77.3	77.5	NP_038716.2, 498 aa
R. norvegicus	Tcte1	316242	D3ZQW5	78.0	78.0	NP_001101676.1, 498 aa
G. gallus	TCTE1	421437	-	63.8	66.8	XP_004935446.1, 457 aa
X. tropicalis	tcte1	100487694	-	57.6	63.5	XP_002942200.1, 460 aa
D. rerio	tcte1	100331698	I3IS38	55.5	61.0	XP_002667399.3, 448 aa
D. melanogaster	CG14325	42123	Q9VEH4	31.6	43.6	NP_650641.2, 460 aa
A. gambiae	AgaP_AP002751	1273219	Q7QCP9	28.3	45.5	XP_312174.4, 430 aa

1309

1310

1311

1312

1313

1314

1315

1316

1317

1318   Figure 2

1319

1320

1321

1322

1323

1324

1325

1326

1327

1328

1329

1330

1331

1332

1333

1334

1335

1336

1337

1338

1339

1340

1341   Figure 3

1342

1343

1344

1345

1346

1347

1348

1349

1350

1351

1352

1353

1354

1355   Figure 4

1356   Figure 5

gene symbol	gene ID /NCBI	gene type	fold change			known function/characteristics/disease
			Tcte1⁻/⁻ vs. WT	Tcte1⁻/⁻ vs. Tcte1⁺/⁻	Tcte1⁺/⁻ vs. WT	
Klk1b27	16619	protein coding	0.2305	0.2937	-	degradation of extracellular matrix proteins in the interstitial area
Klk1b21	16616	protein coding	0.2333	0.2626	-	surrounding the Leydig cells of the adult mouse testis,
Klk1b24	16617	protein coding	0.3101	0.3102	-	seminal liquefaction
Klk1b16	16615	protein coding	-	0.3277	-	seminal liquefaction
Klk1b22	13646	protein coding	0.1266	0.1494	-	neurons' differentiation
mt-Co2	17709	protein coding	0.4966	0.4734	-	mitochondrial cytochrome oxidase subunit, asthenozoospermia, cancer
mt-Co3	17710	protein coding	-	0.4859	-	diabetes, deafness, apoptosis
Fetub	59083	protein coding	4.6789	-	-	fertilization marker, sperm binding to zona pellucida, diabetes, coronary disease, inhibition and apoptosis of prostate cancer cells
Hoxd10	15430	protein coding	2.8503	-	-	prostate cancer, cell differentiation and morphogenesis
Igf2	16002	protein coding	2.0400	-	-	spermatogenesis, sex determination, testis development, infertility, sperm DNA damage, prostate cancer, apoptosis
Slc45a3	212980	protein coding	2.2164	-	-	prostate cancer, myelin maintenance
Usp39	28035	protein coding	12.0339	-	-	spliceosomal complex, prostate cancer, apoptosis, cytokinesis, mitotic spindle checkpoint, vascular remodeling
Gm28661	-	unprocessed pseudogene	0.4576	-	-	nd
Gm26848	-	protein coding	2.2287	-	-	nd (membrane trasporter-like protein in D. sechellia)
Gm45648	102637934	ncRNA	2.5642	-	-	nd (testis-restricted expression in mouse)
Aqp2	11827	protein coding	-	0.4813	-	spermatogenesis, female endometrium cycle, stress urinary incontinence
Mt3	17751	protein coding	-	0.4868	-	testis protection from cadmium, apoptosis, VEGF induction, retinal damage, Alzheimer disease
Rps27rt	100043813	protein coding	-	0.4842	-	ribosomal protein
Gm44126	-	protein coding (?)	-	0.2745	-	nd (suggested: UDP-glucuronic acid decarboxylase in mouse)
4933430L12Rik	71251	protein coding	-	2.5102	-	nd
Ube2l6	56791	protein coding	-	0.4969	2.1110	cancer, apoptosis, autophagy, obesity

1357   Figure 6

1358

1359

1360

1361

1362

1363

1364

1365

1366

1367

1368

1369

1370

1371

1372

1373   Figure 7

1374

1375

1376

1377

1378

1379

1380

1381

1382 Figure 8

1383

1384

1385

1386

1387

1388

1389 Figure 9

1390

1391

1392

1393

1394

1395   Figure 10

1396

1397

1398

1399

1400

1401

1402

1403

1404

1405

1406   Figure 11

1407

1408 **Table 1** Potentially disease-causing novel and ultrarare heterozygous *TCTE1* variants found in patients with disturbed spermatogenesis. Results obtained from screening of

1409 n=248 participants. Minor allele frequency data obtained from GnomAD v2.1.1.

1410 SOAs – severe oligoasthenozoospermia; NOA – non obstructive azoospermia; SCO – Sertoli Cell-Only syndrome; Crypto – cryptozoospermia; NMD – non mediated decay; NT

1411 – nucleotide; AA – amino acid; SNV – single nucleotide variant; PLoF – Loss of Function probability; rs – Reference SNP accession number; CADD – Combined Annotation

1412 Dependent Depletion tool

Sample ID	Phetotype	origin	NT change	cDNA position	prediction	Mutation T@ster	GnomAD	rs	length of protein	CADD	position(s) of altered AA [HGVSp]	Polyphen/ Sift
novel variants												
P52, P88, P105, P111, P161	SOAs	Poland	c.374T>G	cDNA.497T>G	disease causing	amino acid sequence changed; protein features (might be) affected; splice site changes	not found	-	normal	missense	p.Ile125Arg	unknown
P94	Crypto											
P149	NOA											
ultrarare variants (MAF <0.005)												

U13	NOA	USA	c.185G>A	cDNA.308G>A	disease causing	amino acid sequence changed	0.00007778	rs201144307	normal	missense	p.Arg62His	probably damaging/ deletorious
U17	NOA: meiotic arrest	USA	c.397C>T	cDNA.520C>T	disease causing	amino acid sequence changed; heterozygous in TGP or ExAC; protein features (might be) affected; splice site changes	0.001722	rs145304614	normal	missense	p.Arg133Cys	probably damaging/ deletorious
U50, U57, U117, U235	NOA	USA	c.470A>C	cDNA.593A>C	disease causing	amino acid sequence changed; heterozygous in TGP or ExAC; protein features (might be) affected; splice site changes	0.002681	rs45591736	normal	missense	p.Glu157Ala	probably damaging/ deletorious

U11	NOA: meiotic arrest	USA	c.862C>T	cDNA.985C>T	disease causing	NMD amino acid sequence changed; protein features (might be) affected; splice site changes	0.0002840	rs138414421	NMD	stop_gained	p.Arg288Ter	stopgain SNV; PLoF: high confidence
U76	NOA	USA	c.1048G>A	cDNA.1171G>A	disease causing	amino acid sequence changed; protein features (might be) affected; splice site changes	0.00006382	rs570334918	normal	missense	p.Gly350Ser	probably damaging/ deletorious

1413

1414 **Supplementary files legend**

1415

1416 **Supplementary File 1: Vectors used for knockout creation of Tcte1 gene.**

1417

1418 **Supplementary File 2: Real time PCR conditions used for determination of relative expression level of *Tcte1***

1419 **gene in mouse knockout animals.**

1420

1421 **Supplementary File 3: Reproductive potential of KO *Tcte1* mice.** Mating combinations and their results

1422 including: time of breeding, number of litters and pups, number of male and female pups, and genotypes

1423 observed in pups of each combination.

1424

1425 **Supplementary File 4: Genotyping protocol for *Tcte1* knockout model.**

1426

1427 **Supplementary File 5: Characteristics of antibodies used for N-DRC complex staining in mouse sperm.** Human

1428 antigen sequence identity between human and mouse orthologs was also shown (blast comparison; information

1429 from Biorbyt producer).

1430

1431 **Supplementary File 6: Schematic models of exon 3 deleted in KO mice. A** Tcte1 protein exon 3 (deleted in KO)

1432 prediction of 5 top models (software: I-TASSER). C-score is the confidence score of the prediction. C-score ranges

1433 [0-1], where a higher score indicates a more reliable prediction; TM-score is a measure of global structural

1434 similarity between query and template protein., RMSD - root-mean-square deviation of the average distance

1435 between the atoms. **B** The deletion sequence used for modeling, including exon 3 (yellow).

1436

1437 **Supplementary File 7: An example of circular type movement pathway of spermatozoa from *Tcte1*⁻/⁻ male mice.**

1438

1439 **Supplementary File 8: An example of sperm cells motility in *Tcte1*⁺/⁻ male mice.**

1440

1441 **Supplementary File 9: An example of sperm cells motility in *WT* male mice.**

1442

Supplementary File 10: Detailed parameters of mouse sperm morphology. Statistically significant values: p <

0.05; ns – statistically not significant ($p \geq 0.05$). For samples with normal distribution (positive Shapiro-Wilk

normality test), t-test with Welch's correction was applied; for samples without normal distribution (midpiece

and tail defects), Kolmogorov-Smirnoff test was applied (GraphPad Prism v. 7.0e). n=200 spermatozoa were

analyzed per each mouse; for WT and *Tcte1*[-/-] n=5 animals were evaluated, while for *Tcte1*[-/+] the number of

animals was n=7. Spermatozoa stained with the classic Papanicolau staining procedure. Bright field microscope:

Leica DM5500, magnification 630x (with immersion), software: CytoVision.

1450

1451   **Supplementary File 11: STRING analyzis data for each gene with changed expression pattern in *Tcte1* KO mouse**

1452   **model.**

1453

1454   **Supplementary File 12: STRING analyzis of potential interactions between proteins with observed major**

1455   **differences in gene expression level in mouse testis of the homozygous *Tcte1* mice (Usp39, Fetub, Klk1b22).**

1456   According to genes revealed in RNAseq analysis (marked with arrows), two groups of interactions were formed

1457   for genes with highest expression changes (red arrows); (STRING database, 11th March 2021).

1458

1459   **Supplementary File 13: ATP measurement data obtained for KO Tcte1 male mice.** wt = wild type, het =

1460   heterozygous, ho - homozygous. RLU = relative luminescence unit.

1461

1462   **Supplementary File 14: Examples of potentially disease-causing rare heterozygous *TCTE1* variants found in**

1463   **patients with disturbed spermatogenesis.** Results obtained from screening of n=248 participants. **A** novel

1464   variant (Sanger sequencing); **B** ultra rare variants (WES).

1465

Supplementary File 15: The C-terminal region of TCTE1 was consistently modeled and fits well in published

maps. A Superposition of the cartoon representation of three models of human TCTE1 generated by Phyre2 [45]

intensive (green) or fast mode (grey), I-TASSER [46,47,48] (yellow), and AlphaFold [49,50] (cyan). N- and C-termini

are labeled. **B.** Manual docking of the model of full-length TCTE1 generated by AlphaFold in maps calculated from

a cryo-electron tomography experiment performed on *C. reinhardtii* using nanogold to locate the C-terminus of

DRC5 (EMD-20821) [23].