

**Title:** Research Letter: Association between long COVID symptoms and employment status

**Running Title:** Long COVID and employment

**Authors:** Roy H. Perlis, MD MSc (1,2)\*, Kristin Lunz Trujillo, PhD (3,6), Alauna Safarpour, PhD (3, 6), Mauricio Santillana, PhD (3), Katherine Ognyanova, PhD (4), James Druckman, PhD (5), David Lazer, PhD (3)

**Affiliations:**

- (1) Massachusetts General Hospital, Boston, MA
- (2) Harvard Medical School, Boston, MA
- (3) Northeastern University, Boston, MA
- (4) Rutgers University, New Brunswick, NJ
- (5) Northwestern University, Evanston, IL
- (6) Harvard University, Cambridge, MA

**\* Correspondence:**

Roy H. Perlis

[rperlis@mgh.harvard.edu](mailto:rperlis@mgh.harvard.edu)

Massachusetts General Hospital  
185 Cambridge Street, 6th Floor  
Boston, MA 02114  
617-726-7426

**Keywords:** Post-acute sequelae of COVID-19, chronic, rehabilitation, survey, cross-sectional study

Main Text Word Count: 592

Tables and Figures: 1 Figure

## ABSTRACT

Background: Symptoms of Coronavirus-19 (COVID-19) infection persist beyond 2 months in a subset of individuals, a phenomenon referred to as long COVID, but little is known about its functional correlates and in particular the relevance of neurocognitive symptoms.

Method: We analyzed a previously-reported cohort derived from 8 waves of a nonprobability-sample internet survey called the COVID States Project, conducted every 4-8 weeks between February 2021 and July 2022. Primary analyses examined associations between long COVID and lack of full employment or unemployment, adjusted for age, sex, race and ethnicity, education, urbanicity, and region, using multiple logistic regression with interlocking survey weights.

Results: The cohort included 15,307 survey respondents ages 18-69 with test-confirmed COVID-19 at least 2 months prior, of whom 2,236 (14.6%) reported long COVID symptoms, including 1,027/2,236 (45.9%) reporting either 'brain fog' or impaired memory. Overall, 1,418/15,307 (9.3%) reported being unemployed, including 276/2,236 (12.3%) of those with long COVID and 1,142/13,071 (8.7%) of those without; 8,228 (53.8%) worked full-time, including 1,017 (45.5%) of those with long COVID and 7,211 (55.2%) without. In survey-weighted regression models, presence of long COVID was associated with being unemployed (crude OR 1.44, 95% CI 1.20-1.72; adjusted OR 1.23, 95% CI 1.02-1.48), and with lower likelihood of working full-time (crude OR 0.73, 95% CI 0.64-0.82; adjusted OR 0.79, 95% CI 0.70 -0.90). Among individuals with long COVID, the presence of cognitive

symptoms – either brain fog or impaired memory – was associated with lower likelihood of working full time (crude OR 0.71, 95% CI 0.57-0.89, adjusted OR 0.77, 95% CI 0.61-0.97).

Conclusion: Long COVID was associated with a greater likelihood of unemployment and lesser likelihood of working full time in adjusted models. Presence of cognitive symptoms was associated with diminished likelihood of working full time. These results underscore the importance of developing strategies to respond to long COVID, and particularly the associated neurocognitive symptoms.

## INTRODUCTION

Symptoms of Coronavirus-19 (COVID-19) infection persist beyond 2 months in a subset of individuals, a phenomenon referred to as long COVID.<sup>1</sup> This syndrome has become prevalent<sup>2-4</sup>, but little is known about its functional correlates and in particular the relevance of neurocognitive symptoms<sup>5</sup>. We therefore examined whether long COVID and neurocognitive symptoms in particular are associated with differential rates of employment as a proxy for functional impairment, to better guide development of potential interventions.

## METHOD

We analyzed a previously-reported cohort derived from 8 waves of a nonprobability-sample internet survey called the COVID States Project, conducted every 4-8 weeks between February 2021 and July 2022<sup>2</sup>. Respondents were consenting U.S. residents ages 18 and older, according to a protocol approved by the Institutional Review Board of Harvard University. We report results in accordance with AAPOR guidelines.

The cohort included all individuals who reported a positive COVID-19 test result, either PCR or antigen-based, at least 2 months prior to the survey month. Respondents were asked whether their acute symptoms had resolved; those who reported that they had not completed a checklist of commonly-reported symptoms. Respondents also self-reported sociodemographic variables including race and ethnicity based on U.S. Census categories, to allow for survey weighting.

Primary analyses examined associations between long COVID and lack of full employment or unemployment, adjusted for age, sex, race and ethnicity, education, urbanicity (urban, suburban, or rural), and region, using multiple logistic regression with interlocking survey weights in R 4.0.

## RESULTS

The cohort included 15,307 survey respondents ages 18-69 with test-confirmed COVID-19 at least 2 months prior, of whom 2,236 (14.6%) reported long COVID symptoms, including 1,027/2,236 (45.9%) reporting either 'brain fog' or impaired memory. Mean age was 38.8 (13.5) years; 9,678 (63.2%) identified as women and 5,629 (36.8%) men; 810 (5.3%) were Asian, 1,814 (11.9%) were Black, 1,501 (9.8%) were Hispanic, and 10,719 (70.0%) were White.

Overall, 1,418/15,307 (9.3%) reported being unemployed, including 276/2,236 (12.3%) of those with long COVID and 1,142/13,071 (8.7%) of those without; 8,228 (53.8%) worked full-time, including 1,017 (45.5%) of those with long COVID and 7,211 (55.2%) without. Of the remainder, 207 (14.8%) reported part-time or gig work, 13 (0.9%) being a homemaker, 53 (3.8%) being a student, and 11 (0.8%) being retired. In survey-weighted regression models, presence of long COVID was associated with being unemployed (crude OR 1.44, 95% CI 1.20-1.72; adjusted OR 1.23, 95% CI 1.02-1.48), and with lower likelihood of working full-time (crude OR 0.73, 95% CI 0.64-0.82; adjusted OR 0.79, 95% CI 0.70 -0.90).

Among individuals with long COVID, the presence of cognitive symptoms – either brain fog or impaired memory – was associated with lower likelihood of working full time (crude OR 0.71, 95% CI 0.57-0.89, adjusted OR 0.77, 95% CI 0.61-0.97). (Figure 1).

## **DISCUSSION**

Among 15,307 US adults surveyed between February 2021 and March 2022, long COVID was associated with a greater likelihood of unemployment and lesser likelihood of working full time in adjusted models. Further, among those with long COVID, presence of cognitive symptoms was associated with diminished likelihood of working full time, extending recent reports associating cognitive symptoms with poor quality of life among employed individuals<sup>5</sup>.

A limitation in our analysis is the reliance on cross-sectional data, precluding determinations of causation. Moreover, the extent to which our results generalize remains to be determined, though prior work strongly suggests consistency with other approaches<sup>6</sup>.

These results underscore the importance of developing strategies to respond to long COVID, and particularly the associated neurocognitive symptoms. Whether rehabilitation strategies drawn from neurology and psychiatry can help to ameliorate the impact of such symptoms merits investigation. More generally, while the economic impact of the pandemic

is difficult to estimate, these results suggest the importance of considering persistent effects of lost productivity.

## **Acknowledgements:**

The survey was supported in part by the National Science Foundation (Drs. Ognyanova, Lazer, Druckman) and by the National Institute of Mental Health (Drs. Perlis and Lazer - RF1MH132335). The sponsors did not have any role in design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication. The authors had the final responsibility for the decision to submit for publication. Dr. Perlis had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

## **Author Contributions:**

Roy H. Perlis, MD MSc -- analyzed data, drafted and revised manuscript

Kristin Lunz Trujillo, PhD – revised manuscript

Alauna Safarpour, PhD – revised manuscript

Mauricio Santillana, PhD – revised manuscript

Katherine Ognyanova, PhD – contributed to data collection and analysis, revised manuscript

James Druckman, PhD – revised manuscript

David Lazer, PhD – revised manuscript



### **Competing Interests:**

Dr. Perlis has received consulting fees from Burrage Capital, Genomind, Circular Genomics, and Takeda. He holds equity in Psy Therapeutics and Circular Genomics. The other authors report no disclosures.

## References

1. A clinical case definition of post COVID-19 condition by a Delphi consensus, 6 October 2021 [Internet]. [cited 2022 Jun 21]. Available from: [https://www.who.int/publications-detail-redirect/WHO-2019-nCoV-Post\\_COVID-19\\_condition-Clinical\\_case\\_definition-2021.1](https://www.who.int/publications-detail-redirect/WHO-2019-nCoV-Post_COVID-19_condition-Clinical_case_definition-2021.1)
2. Perlis RH, Santillana M, Ognyanova K, Safarpour A, Lunz Trujillo K, Simonson MD, et al. Prevalence and Correlates of Long COVID Symptoms Among US Adults. *JAMA Netw Open*. 2022 Oct 3;5(10):e2238804.
3. Bull-Otterson L. Post-COVID Conditions Among Adult COVID-19 Survivors Aged 18–64 and ≥65 Years — United States, March 2020–November 2021. *MMWR Morb Mortal Wkly Rep* [Internet]. 2022 [cited 2022 Jun 21];71. Available from: <https://www.cdc.gov/mmwr/volumes/71/wr/mm7121e1.htm>
4. Sudre CH, Murray B, Varsavsky T, Graham MS, Penfold RS, Bowyer RC, et al. Attributes and predictors of long COVID. *Nature Medicine*. 2021 Apr;27(4):626–31.
5. Chatys-Bogacka Z, Mazurkiewicz I, Slowik J, Bociaga-Jasik M, Dzieza-Grudnik A, Slowik A, et al. Brain Fog and Quality of Life at Work in Non-Hospitalized Patients after COVID-19. *Int J Environ Res Public Health*. 2022 Oct 6;19(19):12816.
6. Radford J, Green J, Quintana A, Safarpour A, Simonson MD, Baum M, et al. Evaluating the generalizability of the COVID States survey — a large-scale, non-probability survey [Internet]. *OSF Preprints*; 2022 [cited 2022 Mar 8]. Available from: <https://osf.io/cwkg7/>

## Tables and Figures

**Figure 1.** Among individuals with long COVID, symptoms associated with likelihood of not working full time at time of survey

