Aberrant visual salience in participants with schizophrenia during free-viewing of natural images

Masatoshi Yoshida[1,7,*], Kenichiro Miura[2,*], Michiko Fujimoto[2,3], Hidenaga Yamamori[2,3,4], Yuka Yasuda[2,5], Masao Iwase[3,6], Ryota Hashimoto[2]

[1] Center for Human Nature, Artificial Intelligence, and Neuroscience (CHAIN), Hokkaido University, Sapporo, Japan

[2] Department of Pathology of Mental Diseases, National Institute of Mental Health, National Center of Neurology and Psychiatry, Kodaira, Japan

[3] Department of Psychiatry, Osaka University Graduate School of Medicine, Suita, Japan

[4] Japan Community Health Care Organization, Osaka Hospital, Osaka, Japan

[5] Medical Corporation Foster, Life Grow Brilliant Mental Clinic, Osaka, Japan

[6] Osaka Psychiatric Research Center, Osaka Psychiatric Medical Center, Osaka Prefectural Hospital Organization, Hiarakata, Japan

[7] Lead contact

[*] Correspondence: myoshi@chain.hokudai.ac.jp (M.Y.), miura.kenichiro.2x@ncnp.go.jp (K.M.)

SUMMARY

Abnormalities in visual exploration affect the daily lives of patients with schizophrenia; however, its origin is unknown. In this study, we examined whether

such abnormalities reflect aberrant processing of visual salience. Eye movements

of 82 patients and 252 healthy individuals viewing natural and/or complex images

were examined using saliency maps for static images to determine the

contributions of low-level visual features to salience-guided eye movements. The

results showed that the gazes of the participants with schizophrenia were attracted

to position in the images with high orientation salience but not luminance or color

salience. Further analyses revealed that orientation salience defined by the L+M

channel of the DKL color space is specifically affected in schizophrenia, suggesting

abnormalities in the magnocellular visual pathway. These results suggest aberrant

processing of visual salience in schizophrenia, thereby connecting the dots between

abnormalities in early visual processing and the aberrant salience hypothesis of

psychosis.

Keywords: saccades; salience; aberrant salience hypothesis

INTRODUCTION

Schizophrenia is a mental disorder with psychosis, that is, delusion and

hallucination. Previous studies have successfully identified various behavioral markers

of schizophrenia (Onitsuka et al., 2022). However, to understand the pathological

mechanisms of schizophrenia, it is necessary to find links between these behavioral

markers and theories about how the symptoms develop. Here, we examined the

relevance of exploratory eye movements as a behavioral marker and the aberrant

salience hypothesis of psychosis (Kapur, 2003), which states that aberrant assignment of

salience to elements of one's experience leads to delusion and hallucination.

Various abnormalities in eye movements have been reported in patients with

schizophrenia (Morita et al., 2020; Wolf et al., 2021). Among them, significant

differences in eye movements are observed in a simple free-viewing task. Gaze covers a

relatively large area of the images in healthy participants, whereas participants with

schizophrenia tend to limit their gaze to a narrower area of the images, resulting in

shorter scanpath lengths than healthy participants (Beedie et al., 2012; Benson et al.,

2012; Morita et al., 2017; Sprenger et al., 2013). Visual and visuo-cognitive processing

are affected (Butler et al., 2008; Silverstein, 2016; Silverstein and Keane, 2011), while

the motor aspects of saccadic eye movements are less affected (Miura et al., 2014;

Morita et al., 2017). In schizophrenia, it would be reasonable to assume that

abnormalities in visual exploration result from abnormalities in visual or visuo-

cognitive processing.

We sought to determine whether these abnormalities in eye movements were due

to aberrant assignment of visual salience to objects in the visual scene. To this end, we

examined eye movements during free-viewing of natural images in participants with

schizophrenia (SZ) and healthy controls (HC). The visual salience of the presented

images was quantified by Itti and Koch's computational model of salience, which has

already proven useful for predicting and analyzing the visual exploration behavior of

humans and non-human primates (Berg et al., 2009; Chen et al., 2020; Itti et al., 1998;

Yoshida et al., 2012). In the present study, we first tested whether the gaze sequences of

participants with schizophrenia differed from those of healthy controls in terms of

salience values. Since the results showed case-control differences, we examined in

detail which low-level visual features contributed to the differences in eye movements

and found that the difference stemmed specifically from orientation salience. We then

explored the origin of the effects of orientation salience by examining the stages of the

salience computation in the model.

RESULTS

Visual-oculomotor properties are affected in schizophrenia but oculomotor

properties are not. Eye-tracking data was obtained from 82 participants with

schizophrenia and 252 healthy controls while freely viewing 56 natural and/or complex

images. Table S1 summarizes the demographics of the participants, scores of cognitive

tests, and saccade characteristics during free-viewing. Analysis of saccades revealed

that visual exploration is affected in schizophrenia, as reported in previous studies

(Miura et al., 2014; Morita et al., 2017). The average number of saccades during 8

seconds of the viewing period, the average amplitude of saccades, and the scanpath

length were smaller in participants with schizophrenia than in healthy controls (Table

S1). In contrast, oculomotor properties, as assessed by the fitted parameters of the main

sequence relationship, were relatively unaffected (Table S1).

Salience-guided eye movements are affected in examples of single images. To

investigate the relationship between the gaze positions (defined as endpoints of regular

saccades) and the visual salience of the test images, saliency maps were computed using

the Itti-Koch saliency computational model (Figure 1A) (Harel et al., 2007; Itti et al.,

1998).

In representative examples (for the test image #8 and #38), saccades of healthy

controls were distributed not only in highly salient positions (shown in yellow) but also

in other locations in the image (Figure 1B, left). In contrast, saccades of participants

with schizophrenia remained at salient locations of the image during free-viewing

(Figure 1B, right). To quantify the time course of salience values during the 8-second

viewing period, the salience values for each gaze were averaged across participants for

the test images and were plotted across time (Figure 1C, left). The mean salience values

were higher throughout the viewing period in participants with schizophrenia (SZ; blue

in Figure 1C, left) than in healthy controls (HC; magenta in Figure 1C, left). The

difference between the participant groups becomes more evident when the salience

values were averaged across all saccades during the viewing period (Figure 1C, right).

Visual salience attracts the gaze of participants with schizophrenia. To assess

population averages, we plotted the salience values for each saccade averaged across all

test images and all participants (Figure 2A). Consistent with the single image results

(Figure 1C), the mean saliency values were consistently higher in SZs than in HCs.

For statistical analyses, a linear mixed model with random intercepts and slopes

was used to test the main effect of the participant group (SZ vs. HC) and the interaction

between the participant group and the fixation number. Using the log of saccade

numbers as a factor, it is more reasonable to fit the data with a linear mixed model

(Figure 2B, left). Since the interaction term was not significant ($F(1, 330.11) = 0.17$ and

$P = 0.67$), a model without interaction was selected. The main effect for the participant

group was highly significant ($F(1, 332.00) = 44.29$ and $P = 1.17*10^{-10}$). These results

suggest that the gaze of SZs is more likely to be directed toward visually salient

locations than that of HCs.

When the data from age-matched resampled data for HC were compared with the

data from SZ (Figure 2B, right), the results of the statistical analysis were not affected.

The interaction term was not significant ($F(1, 159.69) = 0.02$ and $P = 0.87$), and a model

without interaction was selected. The main effect for the participant group was highly

significant ($F(1, 162.00) = 25.49$ and $P = 1.18*10^{-6}$). The resampling procedure was

repeated 100 times, one of which was used throughout the following analyses. See also

the supplementary text, "Consideration of resampling scheme" for more details.

The gaze of participants with schizophrenia is attracted by orientation

salience. To assess the contribution of low-level visual features, single-feature saliency

maps for color ("Col"), luminance ("Lum"), and orientation ("Ori") were used for

analysis as in the full salience model (Figure 1A). In all three models, interaction terms

were not statistically significant ($F(1, 160.79) = 0.93$ and $P = 0.34$ for color, $F(1,$

$161.60) = 0.52$ and $P = 0.47$ for luminance, and $F(1, 160.14) = 1.79$ and $P = 0.18$ for

orientation, respectively). In the color (Figure 2C left) and the luminance (Figure 2C

middle) models, the main effects for the participant group were not significant ($F(1,$

$162.01) = 0.16$ and $P = 0.69$ for color and $F(1, 161.97) = 0.12$ and $P = 0.72$ for

luminance, respectively). In contrast, the main effect of the participant group was highly

significant ($F(1, 161.99) = 44.80$ and $P = 3.39 * 10^{-10}$) in the orientation model (Figure

2C right). These results suggest that the main effect in the participant group in the full

model (Figure 2B) is explained sorely by orientation salience.

The specific effect of orientation salience is also evident in time-averaged

data. Since the liner mixed model showed no significant interaction effect between the

participant group and saccade numbers in all four salience models (Figures 2B and C), it

is reasonable to summarize the data by averaging all saccades during the viewing period

(Figure 3A). The mean saliency value from the full model was higher in SZ than in HC

($Z = 4.82$, $P = 5.75 * 10^{-6}$; in Wilcoxon rank-sum test with Bonferroni correction). The

effect size evaluated by Cliff's delta was 0.44. The mean salience value from the

orientation model was higher in SZ than in HC ($Z = 6.41$, $P = 5.64 * 10^{-10}$; in Wilcoxon

rank-sum test with Bonferroni correction). The effect size evaluated by Cliff's delta was

0.58, which is classified as a large effect. On the other hand, the mean salience value

from the color and the luminance models were not significantly different between SZ

and HC ($Z = 0.08$, $P > 1.0$ for color; $Z = 0.26$, $P > 1.0$ for luminance; in Wilcoxon rank-

sum test with Bonferroni correction). Based on these findings, the following analyses

were performed on the time-averaged data.

All of these mean salience values were significantly higher than chance. See also

the supplementary text, "Consideration of random sampling scheme" for more details.

We also tested whether the saliency values for the first saccade were significantly

different between SZ and HC (Figure S1). See Discussion for details.

Image category does not affect the effect of orientation salience. To examine

whether the effects of orientation salience depend on image category, the effect size

(Cliff's delta) that evaluates the difference between the time-averaged saliency values of

HC and SZ for each test image was plotted for all four salience models (Figure 3B).

Positive values in the effect size indicate that saliency values were higher in SZ than in

HC. The results (Figure 3B) show that the effect sizes for the Full and the orientation

models were consistently positive for all image categories except for the Face and Noise

categories. This suggests that the effect of orientation salience is overall robust to image

categories.

The key computational stage that produces differences in orientation salience

is Gabor filtering. Next, we examined which stage of the salience computation

produces differences in orientation salience. In the Itti-Koch salience model, saliency

maps for the orientation feature are generated through five processing stages (Figure

4A): 1) transforming the input image into a grayscale image with various spatial

resolutions (i.e., Gaussian pyramids), 2) filtering the images with Gabor patches with

four different orientations, 3) center-surround inhibition, 4) peak normalization, and

finally, 5) adding all images together to obtain the saliency map of the orientation

feature. These intermediate maps were calculated and used for analysis as was done in

the analysis of Figure 3A. Differences between participant groups were detected not

only in the saliency map for the orientation feature (Figure 4B, rightmost) but also in

peak normalization, center-surround inhibition, and Gabor filter (Figure 4B, middle). In

contrast, no differences between participant groups were detected in intensity (Figure

4B, leftmost). These differences became more evident when each comparison was

evaluated by the effect size (Figure 4C). These results suggest that the differences

between participant groups emerge at the stage of Gabor filtering with four orientations,

a unique computational stage that is not present in the color or luminance salience but

only in the orientation salience.

It is also interesting to note that the effect size of Gabor filtering is larger for low-

pass images (those with higher numbers in Figure 4C). Since it is already known that

the magnocellular pathway conveys relatively low spatial frequency signals (Kauffmann

et al., 2014) and that the magnocellular pathway is specifically impaired in

schizophrenia (Butler et al., 2008), this point is directly examined in the next section.

the L+M channel of the DKL color space is dominant in the effect of

orientation salience. In the Itti-Koch salience model (hereafter referred to as the

"original model"), the orientation salience is calculated from grayscale images. Instead,

it is possible to decompose the test images into images of three channels (L+M, L-M,

and S-(L+M)), based on the DKL color space (Derrington et al., 1984). The

decomposed images were then processed to obtain saliency maps with (orientation map)

or without (intensity map) Gabor filtering (Figure 5A). We refer to this as the

"Extended six-channel model."

Then the salience values for these six maps were compared between SZ and HC

(Figures 5B and C). In the intensity maps (Figure 5B), all differences were not

statistically significant (Z = 0.21, P > 1.0 for the L+M channel; Z = 1.09, P > 1.0 for the

L-M channel; Z = -0.51, P > 1.0 for the S-(L+M) channel; in Wilcoxon rank-sum test

with Bonferroni correction). In the orientation maps (Figure 5C), all of the differences

were statistically significant (Z = 6.25, P = 2.51 $*10^{-9}$ for the L+M channel; Z = 4.20, P

= 1.59 $*10^{-4}$ for the L-M channel; Z = 3.84, P = 7.49 $*10^{-4}$ for the S-(L+M) channel; in

Wilcoxon rank-sum test with Bonferroni correction). These results further support the

finding of a specific effect of orientation salience and suggest that the L+M channel of

the DKL color space is dominant in the effect of orientation salience.

Orientation salience is correlated with the scores of cognitive tests and visual

oculomotor characteristics. Next, we examined whether orientation salience is

correlated with demographic data, cognitive ability, and eye movement characteristics

collected from the same participants (See (Morita et al., 2017, 2019) for details). When

the dependent variables (demographic data, cognitive tests, or eye movement

characteristics) were fitted with orientation salience with the participant group as a

covariate, age was not correlated with orientation salience but WAIS-3, processing

speed (PS), social functioning scale (SFS), and scanpath length (SPL) were correlated

with orientation salience (Figure 6A).

For a more systematic analysis, two regression models were constructed. In the

first model, the dependent variables were fitted with saliency values from the original

model, with the participant group as a covariate. The t-values obtained from the fitting

were then plotted (Figure 6B). Orientation salience is correlated with various scores of

cognitive tests and eye movement characteristics (indicated by the gray bars in Figure

6B). Regression analysis also suggested that these correlations are specific to

orientation.

In the second model in which the dependent variables were fitted with saliency

values from the extended six-channel model, with the participant group as a covariate,

orientation salience in the L+M channel was significantly correlated with a cognitive

test score (SFS) and eye movement characteristics (saccade amplitude and SPL) (Figure

6C). In contrast, orientation saliences in the other channels are not correlated with these

values. These results again suggest that the L+M channel is dominant in the effect of

orientation salience.

Saliency maps for orientation are correlated with fixation maps. In Figure 3A

and Figure 5C, saliency values were evaluated separately for each feature/channel. It is

important to note that saliency maps for the different features/channels are spatially

correlated (Figure 7B). This correlation may affect statistical analyses in Figure 3A and

Figure 5C. To address this issue, we calculated the partial correlation between the

saliency maps and the "Fixation maps" which quantifies gaze distribution as a density

map summed across participants for each test image (Figure 7A). We also created a

fixation map for the difference between SZ and HC ("SZ-HC") (Figure 7A). The partial

correlations of these maps were then calculated for each test image (n=56).

For the original model, we plotted the partial correlations between the fixation

maps and the saliency maps for the three features (color, luminance, and orientation)

(Figure 7C). For the three fixation maps (SZ, HC, and SZ-HC), the partial correlation

with the saliency map for orientation was significantly larger than zero ($P = 4.0*10^{-8}$ for

SZ, $P = 1.7*10^{-7}$ for HC, and $P = 1.6*10^{-7}$ for SZ-HC, respectively; Wilcoxon signed-

rank test with Bonferroni correction). In contrast, the partial correlations with the

saliency map for color and luminance were not significantly larger than zero ($P > 0.05$,

Wilcoxon signed-rank test with Bonferroni correction) except for a negative correlation

in luminance for SZ-HC ($P = 2.7*10^{-4}$).

For the extended six-channel model, three maps for orientation were shown

(Figure 7D). For two fixation maps (SZ, HC), the partial correlations with the saliency

map for the L+M and the L-M channels were significantly larger than zero ($P = 2.0*10^{-5}$

and 0.002 for SZ, $P = 6.0*10^{-5}$ and 0.008 for HC; Wilcoxon signed-rank test with

Bonferroni correction). On the other hand, for the SZ-HC fixation map, the partial

correlation with the saliency map of the L+M channel was significantly larger than zero

(p = $5.2*10^{-4}$, Wilcoxon signed-rank test with Bonferroni correction). These results

suggest that the L+M channel contributes specifically to the effect of orientation

salience, even after accounting for spatial correlations between saliency maps.

DISCUSSION

In this study, we examined whether abnormalities in eye movements result from

aberrant processing of visual salience. For this purpose, we analyzed the eye movement

data during free-viewing with Itti-Koch's salience model (Figure 1). We found that the

saliency values at the gaze of SZs were persistently higher during the viewing period

compared to the HCs (Figures 2A and B). Further analysis using single-feature saliency

maps revealed that this difference was due to orientation salience (Figure 2C). We then

confirmed that these results are robust for various image categories (Figure 3). Next, we

delved into the stages of salience computation and found that differences between SZs

and HCs were found in an early stage of salience computation, where grayscale images

were filtered with Gabor patches with four orientations (Figure 4). We also analyzed the

gaze with an extended salience model that evaluates the channels for the DKL color

space separately and found that the L+M cannel has a dominant role in the effects of

orientation salience (Figure 5). In addition, the saliency values at gazes were not

correlated with symptom-related measures such as PANSS scores but were correlated

with various measures of cognitive functions and saccade-related characteristics (Figure

6). Finally, we evaluated a spatial correlation between fixation maps and saliency maps

and found that orientation salience in the L+M channel correlates specifically with the

difference between the fixation maps of the SZs and those of HCs. These results suggest

that visual processing of orientation salience is affected in participants with

schizophrenia, which is consistent with the aberrant salience hypothesis of psychosis, as

will be discussed shortly.

Relevance to the aberrant salience hypothesis of psychosis. The aberrant

salience hypothesis of psychosis proposes that an aberrant assignment of salience to the

elements of one's experience leads to delusion and hallucination (Kapur, 2003). The

concept of salience in this hypothesis covers not only salience derived from emotion

and motivation (motivational/incentive salience) (Berridge, 2018), but also salience due

to novelty and sensory features (perceptual salience) (Itti and Koch, 2001). The present

study provides direct evidence that visual salience is affected in schizophrenia, thereby

explicitly extending the concept of aberrant salience to the visual domain. In support of

our findings, a recent study revealed that brain responses to images with various forms

of salience such as novelty, negative emotion, targetness, and rarity/deviance are

affected in schizophrenia (Knolle et al., 2018).

Relationship with eye movement abnormalities: visual-oculomotor

properties. As described in the Introduction, visual-oculomotor properties such as

scanpath length, saccade number, and saccade amplitudes are affected in the gaze of

SZs during free-viewing (see also Table S1). We found that these visual-oculomotor

properties are correlated with saliency values (Figure 6). Since the bottom-up salience

of the images is related to target selection, the abnormalities in visual exploration that

have been reported in various studies (Butler et al., 2008) arise, at least in part, from the

affected salience-guided eye movement found in this study.

Relationship with eye movement abnormalities: inhibition-of-return.

Recently, our research group reported the finding that inhibition-of-return is impaired in

SZs (Okada et al., 2021). This might raise the question of whether return saccades

toward salient stimuli are more frequent in schizophrenia (Figure S1A). If this is the

case, salience values at the gaze could be higher in SZ than in HC. However, it is

unlikely because salience values at the first gaze for the viewing period, where it has no

contribution to inhibition-of-return, were significantly higher in SZ than HC in the

orientation model (Figure S1C). Furthermore, if the SZs made more return saccades

toward salient positions than in the HCs, the slope of the time course of saliency values

would be shallower in the SZs than in the HCs (Figure S1B, left). However, this is not

the case because analyses using linear mixed models showed no interaction between the

participant group and saccade numbers. Thus, abnormalities in visual salience found in

this study have a different origin from abnormal inhibition-of-return. For further

analysis and discussion, see also (Okada et al., 2021).

Contribution of low-level visual features, early visual cortex, and the

magnocellular pathway. We found that salience computation for the orientation

feature is specifically affected in schizophrenia. Consistent with this finding, previous

findings have also shown that orientation processing is affected in schizophrenia

(Rokem et al., 2011; Yoon et al., 2009). Other papers have shown that the early visual

cortex is involved in changes in contextual modulation in schizophrenia (Seymour et al.,

2013; Tibber et al., 2013; Yoon et al., 2010). The effect of orientation salience can

explain impairment in contour integration in schizophrenia (Yoon et al., 2009) because

contour integration can be distracted by aberrant processing of orientation salience.

Based on the analysis of Figure 4 and the accompanying text, we argue that the

crucial difference between SZs and HCs stems from their response to Gabor-filtered

images, which resembles responses in the early visual cortex. On the other hand, there

was no difference in gazes toward luminance salience. Computation of luminance

salience involves center-surround inhibition without Gabor filtering. Such processing

resembles the response of the lateral geniculate nucleus (LGN). We propose that

aberrant orientation salience reflects abnormalities at the level of the primary visual

cortex but not at the level of the LGN. Post-mortem morphological studies of the

schizophrenia brains support this possibility. The volume and number of neurons in the

primary visual cortex are reduced in schizophrenia (Dorph-Petersen et al., 2007;

Selemon, 2001) but it is not the case for the LGN (Dorph-Petersen et al., 2009; Selemon

and Begovic´, 2007). Thus, the idea that visual abnormalities in schizophrenia occur

between the LGN and primary visual cortex is consistent with both previous studies and

the present current study.

The analyses in Figures 5, 6, and 7 suggest that orientation salience in the L+M

channel of the DKL color space is specifically affected in schizophrenia. Since the L+M

channel carries information in the magnocellular pathway (Derrington et al., 1984), the

present results are consistent with previous findings that visual information processing

in the magnocellular pathway is specifically affected in schizophrenia (Butler et al.,

2001, 2005; Schechter et al., 2003).

Relationship with clinical/cognitive tests. Figure 6 shows that symptom-related

measures such as the PANSS total score, CPZ equivalent, and duration of illness were

not significantly correlated with orientation salience. These results suggest that

orientation salience is a trait marker rather than a state marker. This is consistent with

the view that eye movement-related measures are trait markers of schizophrenia (Beedie

et al., 2011).

Figure 6 also shows that scores on some of the cognitive tests are correlated with

orientation salience. This is consistent with previous findings that scores of cognitive

tests are correlated with visual oculomotor characteristics such as scanpath length

(Morita 2018, 2019). It is unlikely that the abnormality in scanpath length causes

abnormalities in orientation salience, as oculomotor properties of eye movements are

less affected in schizophrenia. Rather, we propose that aberrant processing of

orientation salience causes abnormalities in visual exploration, which can be assessed

by visual oculomotor properties such as scanpath length, which in turn causes difficulty

in cognitive abilities such as social functioning.

Visual salience in schizophrenia and autism spectrum disorder (ASD).

Analysis of visual salience during free-viewing can be a useful method for detecting

psychiatric or neurological disorders (Itti, 2015; Tseng et al., 2012). The current

findings added another example of the successful application of the computational

salience model to the analysis of eye movements in participants with psychiatric

disorders.

Gazes of participants with autism spectrum disorder (ASD) showed a general

increase in visual salience, defined by the low-level visual features (Wang et al., 2015).

This appears similar to the findings in the current study about schizophrenia but differs

in that the present study reported specific effects of the orientation feature. Also

interesting is the finding that the effect size of orientation salience was lower in the Face

category than in the other image categories in the present study (Figure 3B). This

suggests that faces are relatively ineffective in discriminating between SZs and the HCs,

in contrast to the abnormalities in gaze and processing of faces in ASD (Dalton et al.,

2005).

In terms of the predictive coding/Bayesian brain hypothesis, both schizophrenia

and ASD are explained by difficulties in evaluating prediction errors in sensory signals

(Adams et al., 2013; de Cruys et al., 2014). How to distinguish the two in terms of

computational perspective is an ongoing debate. By extending the concept of visual

salience to Bayesian surprise (Itti and Baldi, 2009) and treating visual salience as spatial

outliers from the current belief on what we see, saliency analysis could help resolve this

issue.

Future projects. In this study, we obtained clues about what is affected in the

brain of schizophrenia, for example, Gabor filtering that is presumably performed in the

magnocellular pathway of the early visual cortex. To understand exactly how the brain

performs such computation, it is necessary to understand the processes at the neuronal

level. To this end, neurophysiological studies using animal models of schizophrenia are

needed. Since eye-tracking during free-viewing and analysis of visual salience is an

experimental paradigm that has been successfully used in non-human primates such as

macaque monkeys (Yoshida et al., 2012) and marmosets (Chen et al., 2020), replicating

the present results in a non-human primate model of schizophrenia would open the door

to understanding the precise brain mechanism of schizophrenia.

ACKNOWLEDGMENTS

This research was supported in part by AMED under Grant Numbers

JP19dm0207069 (MY and KM), JP21dk0307103 (RH), JP21wm0425012 (KM and

RH), JP18dm0307002 (RH), and JP21uk1024002 (KM and RH); JSPS KAKENHI

grant numbers 22H02936 (MY), JP20H03611 (RH), JP19H05467 (RH), JP20K06920

(KM); and Intramural Research Grant (3-1 and 4-6) for Neurological and Psychiatric

Disorders of NCNP (KM and RH).

12　**AUTHOR CONTRIBUTIONS**

21　Conceptualization, M.Y.; Methodology, M.Y. and K.M.; Software, M.Y.;

22　Investigation, M.F., H.Y., Y.Y. and M.I.; Resources, K.M. and R.H.; Data Curation:

23　M.Y., K.M. M.F., H.Y., Y.Y. and M.I; Writing – Original Draft: M.Y.; Writing –

24　Review & Editing: M.Y. and K.M.; Funding Acquisition, M.Y., K.M., and R.H.;

25　Supervision, K.M. and R.H.

DECLARATION OF INTERESTS

The authors declare no competing financial interests.

FIGURE TITLES AND LEGENDS

Figure 1. Salience-guided eye movements are affected in examples of single images.

(A) The saliency map was calculated from the Itti-Koch model. Visual salience for low-

level visual features (color, "Col"; luminance, "Lum"; orientation, "Ori") was also

computed in this model.

(B) Gaze positions of two control subjects (left) and two SZs (right) represented by

numbers and superimposed on the saliency map of test images. Numbers indicate

saccade order.

(C) Left: The saliency values averaged across participants are plotted over time for test

images #8 (top) and #38 (bottom). Error bars denote standard errors across participants.

Magenta, HC; blue, SZ. Right: the saliency values averaged across participants and

saccades plotted as a violin plot. Symbols denote the median. Error bars denote the first

and the third quartile.

Figure 2. The gaze of participants with schizophrenia is attracted by orientation salience.

(A) The saliency values averaged across test images and participants are plotted over time after image onset. Magenta, the healthy controls (HC; n=252); blue, the participants with schizophrenia (SZ; n = 82). Each point denotes the mean salience value for each saccade (1st through 16th). Error bars denote the standard errors across participants. Inset: number of images obtained for each number of saccades (up to 56) during the 8-sec viewing period. The dotted line denotes the cutoff point (16th saccades). See Methods for details.

(B) (Left) Same as (A) but the data are plotted across saccade numbers on a log scale. (Right) Age-matched resampled data are plotted for healthy controls. Numbers on the plots denote P values for the main effect of the participant group.

(C) As in (B), but for single feature salience models.

Figure 3. Image category does not affect the effect of orientation salience.

(A) Mean saliency values for all images and saccades are plotted for the healthy controls (HC) and the schizophrenia group (SZ). Data from four salience models are plotted. Symbols denote median values. Error bars denote the first and the third quartile. Numbers on the plots denote P values ("p") and the effect sizes ("Δ", Cliff's delta) of the Wilcoxon rank-sum test.

(B) Effect sizes (Cliff's delta) for the test of difference in time-averaged saliency values. Top: histograms of effect sizes.

Figure 4. The key computational stage that produces differences in orientation salience is Gabor filtering.

(A) The Itti-Koch saliency model computes the saliency map of the orientation feature through five stages. Intermediate images for a given image are shown in grayscale.

(B) Averaged map values for intermediate images are plotted as in Figure 3A.

(C) Effect sizes (Cliff's delta) for the tests in (B). Numbers indicate the spatial scales of the Gaussian pyramids. In the three plots in the center, the mean effect size for four orientations (bar), as well as the effect size for four orientations (symbols) were plotted. "o," 0 degrees; "x," 45 degrees; triangle, 90 degrees, and square, 135 degrees. The horizontal lines in the rightmost plot ("Saliency map (Orientation)") indicate the common guidelines for the effect size (Cliff's delta). See Methods for details.

Figure 5. the L+M channel of the DKL color space is dominant in the effect of orientation salience.

(A) To construct the extended six-channel model, the original image was decomposed into three channels in the DKL color space: the magnocellular L+M channel, the parvocellular L-M channel, and the koniocellular S-(L+M) channel. Then saliency maps for intensity and orientation were obtained for each of the three channels (six maps in total).

(B and C) As in Figure 3A, the salience values averaged across test images and saccades were plotted, but for six saliency maps. Symbols are the same as in Figure 3A.

2 **Figure 6. Orientation salience is correlated with the scores of cognitive tests and**

3 **visual oculomotor characteristics.**

(A) Scattered plots for the mean orientation salience (averaged across images and

saccades) and age, WAIS-3 processing speed (PS), social functioning scale (SFS), and

scanpath length (SPL). Each dot represents the value for one participant. Magenta:

healthy controls (HC); blue, participants with schizophrenia (SZ). Lines indicate

regression lines for each participant group.

(B) Absolute value of t-values from regression analysis 1, where the dependent

variables were fitted individually with the saliency values of the original model. Gray

bars indicate statistical significance ($P < 0.05$) after correction of multiple comparisons

by FDR.

(C) As in (B), but those from regression analysis 2, where the dependent variables were

fitted individually with the saliency values of the extended six-channel model.

Figure 7. Saliency maps for orientation are correlated with fixation maps.

(A) Examples of fixation maps for test images #8 and #38. SZ, the participants with schizophrenia; HC, the healthy controls (age-matched resamples); SZ-HC, the difference between SZ and HC.

(B) (Top) Colors indicate the median of partial correlation coefficients between saliency maps of the original model (top) and the extended six-channel model (bottom).

(C) Violin plots of the partial correlation between the fixation maps and saliency maps

(original model). Symbols are the same as in Figure 3A.

(D) Violin plots of the partial correlation between the fixation maps and saliency maps

(the extended six-channel). Only the correlations with orientation maps are shown.

Numbers in (C) and (D) indicate P values for a significant difference from zero

(Wilcoxon signed-rank test after Bonferroni correction). Only the P values less than

0.05 are shown.

1　**STAR METHODS**

2　**KEY RESOURCES TABLE**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Deposited data		
Human eye-tracking data	This paper	N/A
The International Affective Pictures System (IAPS)	(Lang et al., 2008)	https://csea.phhp.ufl.edu/Media.html
Japanese and Caucasian Facial Expressions of Emotion (JACFEE) and Neutral Faces (JACNeuF)	(Matsumoto and Ekman, 1988)	N/A
Software and algorithms		
MATLAB 2022a	MathWorks	https://www.mathworks.com/
Psychophysics Toolbox for MATLAB	(Brainard, 1997)	http://psychtoolbox.org/
GBVS toolbox for MATLAB	(Harel et al., 2007)	http://www.klab.caltech.edu/~harel/share/gbvs.php
R ver 4.2.1	The R Foundation	https://www.r-project.org/
RStudio 2022.07.1	RStudio PBC	https://www.rstudio.com/
R package lme4	(Bates et al., 2015)	https://github.com/lme4/lme4
R package LmerTest	(Kuznetsova et al., 2017)	https://github.com/runehaubo/lmerTestR
Custom MATLAB scripts	This paper	https://github.com/pooneil68/saliency_analysis DOI: 10.5281/zenodo.7312329
Other		
EyeLink 1000 Eye tracker	SR Research	https://www.sr-research.com/

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources should be addressed to and will be

fulfilled by the lead contact of the study, Masatoshi Yoshida, Ph.D.,

myoshi@chain.hokudai.ac.jp.

Materials Availability

This study did not generate new unique reagents.

Data and code availability

Codes are available from GitHub (https://github.com/pooneil68/saliency_analysis DOI:

10.5281/zenodo.7312329).

All data reported in this study cannot be deposited in a public repository because of

ethical prohibitions. To request access, contact the lead contact.

Any additional information required to reanalyze the data reported in this paper is

available from the lead contact upon request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Participants

Eye movement data were sampled from 82 SZs (male, 42; female, 40) and 252 HCs

(male, 144; female, 108) as part of a large-scale cohort recruited at Osaka University

(Table S1) (Miura et al., 2014; Morita et al., 2017, 2019). There was an overlap in data

with a previous study on eye movement abnormalities in schizophrenia (Morita et al.,

2017). All participants were biologically unrelated, were of Japanese descent, and had

no history of the ophthalmologic disease, or neurological/medical conditions that could

influence the central nervous system. Specific exclusion criteria included atypical

headaches, head trauma with loss of consciousness, chronic lung disease, kidney

disease, chronic hepatic disease, thyroid disease, active cancer, cerebrovascular disease,

epilepsy, seizures, substance-related disorders, or mental retardation (Miura et al., 2014;

Morita et al., 2017).

SZs were recruited from Osaka University Hospital and had been diagnosed by two or

more trained psychiatrists according to criteria from the DSM-IV based on the

Structured Clinical Interview for DSM-IV (SCID). Estimated cognitive decline was

calculated by the methods described by Fujino (Fujino et al., 2017). The current

symptoms of the SZs were assessed using the Positive and Negative Syndrome Scale

(PANSS) (Kay et al., 1987), and daily antipsychotic use was calculated using

chlorpromazine (CPZ) equivalents (mg/day) (Inada and Inagaki, 2015).

HCs were assessed for psychiatric, medical, and neurological concerns using a non-

patient version of the SCID to exclude individuals with current or past contact with

psychiatric services or who had received psychiatric medication.

All participants provided written consent to the study after a full explanation of the

study procedures. Anonymity was preserved for all participants. The study was

performed in accordance with the World Medical Association's Declaration of Helsinki

and was approved by the Research Ethical Committee of Osaka University, the National

Center of Neurology and Psychiatry, and Center for Experimental Research in Social

Sciences, Hokkaido University.

METHOD DETAILS

Task and stimuli

In the free-viewing task, the participants faced a 19-inch liquid crystal display monitor

(1280 × 1024 pixels) placed 70 cm from the observer's eyes. Presentation of the visual

stimuli was done using the Psychophysics Toolbox extension (Brainard, 1997) in

MATLAB (The Mathworks, Natick, MA, USA). Each trial began with the presentation

of a fixation point on the center of the display. Once the participant fixated on the

fixation spot for a random time, a test image was presented for 8 seconds. The

participant was instructed to view the image as they like. One task consisted of 56

images, the order of which was randomly shuffled for each participant. The images

were chosen from eight categories: natural environments, buildings, everyday items,

foods, faces, animals, fractal patterns, and noise (seven images for each). The images of

natural environments and animals were selected from the International Affective

Pictures System (IAPS) (Lang et al., 2008), and the face images from Matsumoto and

Ekman (Matsumoto and Ekman, 1988). Since IAPS images are not allowed to be

published in scientific journals, only the saliency maps for these images (#8 and #38)

are shown in the figures. In cases when images needed to be shown as examples

(Figures 1A, 5A, and 6A), a photograph taken by one of the authors was used.

Recording and preprocessing of eye movement data

Recording and preprocessing of eye movement data were performed as follows (Morita

et al., 2017). Eye position and pupil area of the left eye were measured at 1 kHz using

EyeLink1000 (SR Research, Ontario, Canada). Eye position data (in degrees) were

smoothed with a digital FIR filter (-3 dB at 30 Hz), and eye velocity and acceleration

traces were derived from a two-point difference algorithm. Eye movement recordings

were segmented into blink, saccade, and fixation periods. Detected saccades included

both regular saccades and microsaccades. Here, following previous papers on

microsaccades during free-viewing such as (Otero-Millan et al., 2008), saccades with

amplitudes greater than one degree were selected as regular saccades.

Computational models and saliency analysis

To assess salience-guided eye movements, we used a validated computational model of

visual attention and compared it with individual eye movements. The saliency maps for

the test images were computed with the Itti-Koch saliency model for static images (Itti

et al., 1998) implemented in the GBVS toolbox for Matlab (Harel et al., 2007). The Itti

and Koch model is a neurobiologically inspired model which computes salient locations

for low-level visual features (Figure 1A). Since we are interested in neurobiological

mechanisms of saliency-guided eye movements, the Itti-Koch model was chosen.

In the full model, saliency maps of the three features were summed with equal weights.

When evaluating the contribution of low-level visual features, single-feature saliency

maps were used (Figure 1A). For a test image of 640*512 pixels, we obtained a saliency

map of 80*64 pixels. To treat the saliency maps as density maps, all maps were

normalized so that the sum of saliency values of all pixels was one.

Intermediate files for salience computation were used for the analysis in Figures 4 and

the accompanying text. These files were also generated by the GBVS toolbox.

Extended six-channel model for visual salience

An "extended six-channel model" for visual salience was constructed for the analysis in

Figures 5, 6, and 7 and the accompanying text. For this purpose, the test images were

decomposed into three-channel images (L+M, L-M, and S-(L+M)) based on the DKL

color space (Derrington et al., 1984). The decomposed images were then subjected to

salience computation separately with (orientation map) or without (intensity map)

Gabor filtering (Figure 5A). The luminance salience in the original model corresponds

approximately to the intensity map of the L+M channel. The color salience in the

original model corresponds approximately to the intensity map of the L-M channel plus

the S-(L+M) channel. The orientation salience in the original model corresponds

approximately to the orientation map of the L+M channel. The other two maps (the

orientation map of the L-M channel and the orientation map of the S-(L+M) channel)

are new components not found in the original model.

Age-matched resampling

To account for the effect of the age difference between the healthy control group (HC,

28.8 ± 11.5 years, mean $\pm$SD) and the schizophrenia group (SZ, 35.1 ± 12.4 years, mean

$\pm$SD), resampling of the data for the HC group was performed to match the mean age of

the HC participants with that of the SZ group. For this purpose, the ages of both groups

were grouped into 5-year bins. The HC participants in each bin were then randomly

selected to match the number of SZ participants in each bin. This procedure was

repeated 100 times to obtain 100 sets of resampled HC participants. The mean age of

the resampled HC participants ranges from 34.9 – 35.2 years, in close agreement with

that of the SZ group (35.1 years). The standard deviation of the age of the resampled

HC participants ranges from 12.0 – 12.5 years, in close agreement with that of the SZ

group (12.4 years old). We confirmed that the results of statistical analysis were not

affected by the choice of the resampled dataset (see supplemental text, "Consideration

of resampling scheme"). Therefore, we selected one of the resampled datasets and used

it throughout our analysis.

Linear mixed models

The procedure for the statistical analysis of the time course of the mean salience values

at saccade endpoints (Figures 1 and 2) is as follows. First, a cut point for the number of

saccades was set because the number of saccades during the 8-second viewing time

varied between trials and between the participant groups (see Table S1 for the mean

numbers of saccades; 16.1 in SZ and 20.9 in HC). The cut point was set to 16 because

the mean number of images obtained for each saccade number (up to 56) is less than

half (Figure 2A, inset).

We then fit linear mixed models using the R package lme4 (Bates et al., 2015) to obtain

estimates and statistics. For this purpose, the salience values for each saccade number

and each participant were averaged over 56 test images. Saccades that landed off-screen

were treated as not a number. The mean salience values ("Sal") were then treated as the

dependent variable and as a linear function of participant groups (HC and SZ as

"SubjectGroup") and saccade numbers (1st, 2nd…, 16th as "SaccadeNum"). The

random effects were modeled using random intercepts and random slopes for individual

differences (ranging from 1 to 334 as "SubjectID") nested under the participant group.

The model formulae in R format are as follows:

1) Sal ~ SubjectGroup + SaccadeNum + (SaccadeNum |SubjectID)

2) Sal ~ SubjectGroup * SaccadeNum + (SaccadeNum |SubjectID)

Model #1 is a model with the main effects of participant group and saccade number.

Model #2 is a model with an interaction between the participant group and saccade

number. The two-sided probability values and degrees of freedom associated with each

statistic were then determined using the Satterthwaite approximation implemented in the

R package LmerTest (Kuznetsova et al., 2017).

General linear models

To examine whether the salience values are corelated to other measures such as

demographic data, cognitive tests, and eye movement properties, general linear models

were constructed (Figure 6). The model formulae in R format are as follows:

1) Dependent variables ~ SubjectGroup + Sal(Col) + Sal(Lum) + Sal(Ori)

2) Dependent variables ~ SubjectGroup + Sal(Int, L+M) + Sal(Int, L-M) +

Sal(Int, S-(L+M)) + Sal(Ori, L+M) + Sal(Ori, L-M) + Sal(Ori, S-(L+M))

In Model #1, the dependent variables (demographic data, cognitive tests, and saccade-

related properties) were fitted with the participant group ("SubjectGroup"), and the

salience values were calculated by the original saliency model consisting of three

features. In Model #2, the dependent variables were fitted with the participant group

("SubjectGroup") and the salience values were calculated by the extended six-channel

model. P values were adjusted for multiple comparisons using FDR.

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical analyses were performed using MATLAB (Mathworks, NY), except for the

linear mixed models, which were performed by the R packages. The significance level

was set at p < 0.05. Non-parametric Wilcoxon rank-sum tests after Bonferroni

correction were performed in a post-hoc analysis to compare SZ and HC for the median

salience values or other indices (Figure 3A, Figure 5B and C, Figure S1C). Cliff's delta

(Δ) was used to quantify the effect size of the estimated difference. For interpretation,

we followed general guidelines: negligible for $|\Delta| < 0.147$, small for $0.147 <= |\Delta| < 0.33$,

medium for $0.33 <= |\Delta| < 0.474$, and large for $|\Delta| >= 0.474$ (Romano et al., 2006).

SUPPLEMENTAL INFORMATION TITLES AND LEGENDS

• Supplemental text

• Figure S1

• Table S1

REFERENCES

Adams, R.A., Stephan, K.E., Brown, H.R., Frith, C.D., and Friston, K.J. (2013). The
computational anatomy of psychosis. Front. Psychiatry *4*, 47.
Bates, D., Mächler, M., Bolker, B.M., and Walker, S.C. (2015). Fitting linear mixed-
effects models using lme4. J. Stat. Softw. *67*.
Beedie, S.A., Clair, D.M.S., and Benson, P.J. (2011). Atypical scanpaths in
schizophrenia : Evidence of a trait- or state-dependent phenomenon ? *36*, 150–164.
Beedie, S.A., Benson, P.J., Giegling, I., Rujescu, D., and St. Clair, D.M. (2012).
Smooth pursuit and visual scanpaths: Independence of two candidate oculomotor risk
markers for schizophrenia. World J. Biol. Psychiatry *13*, 200–210.
Benson, P.J., Beedie, S.A., Shephard, E., Giegling, I., Rujescu, D., and St. Clair, D.
(2012). Simple viewing tests can detect eye movement abnormalities that distinguish
schizophrenia cases from controls with exceptional accuracy. Biol. Psychiatry *72*, 716–
724.

Berg, D.J., Boehnke, S.E., Marino, R.A., Munoz, D.P., and Itti, L. (2009). Free viewing of dynamic stimuli by humans and monkeys. J. Vis. *9*, 1–15.

Berridge, K.C. (2018). Evolving concepts of emotion and motivation. Front. Psychol. *9*, 1–20.

Brainard, D.H. (1997). The Psychophysics Toolbox. Spat. Vis. *10*, 433–436.

Butler, P.D., Schechter, I., Zemon, V., Schwartz, S.G., Greenstein, V.C., Gordon, J., Schroeder, C.E., and Javitt, D.C. (2001). Dysfunction of early-stage visual processing in schizophrenia. Am. J. Psychiatry *158*, 1126–1133.

Butler, P.D., Zemon, V., Schechter, I., Saperstein, A.M., Hoptman, M.J., Lim, K.O., Revheim, N., Silipo, G., and Javitt, D.C. (2005). Early-stage visual processing and cortical amplification deficits in schizophrenia. Arch. Gen. Psychiatry *62*, 495–504.

Butler, P.D., Silverstein, S.M., and Dakin, S.C. (2008). Visual Perception and Its Impairment in Schizophrenia. Biol. Psychiatry *64*, 40–47.

Chen, C.-Y., Matrov, D., Veale, R.E., Onoe, H., Yoshida, M., Miura, K., and Isa, T. (2020). Properties of visually-guided saccadic behavior and bottom-up attention in marmoset, macaque, and human. J. Neurophysiol. jn.00312.2020.

de Cruys, S., Evers, K., der Hallen, R., Van Eylen, L., Boets, B., de-Wit, L., and Wagemans, J. (2014). Precise minds in uncertain worlds: Predictive coding in autism. Psychol. Rev. *121*, 649–675.

Dalton, K.M., Nacewicz, B.M., Johnstone, T., Schaefer, H.S., Gernsbacher, M.A., Goldsmith, H.H., Alexander, A.L., and Davidson, R.J. (2005). Gaze fixation and the neural circuitry of face processing in autism. Nat. Neurosci. *8*, 519–526.

Derrington, A.M., Krauskopf, J., and Lennie, P. (1984). Chromatic mechanisms in lateral geniculate nucleus of macaque. J. Physiol. *357*, 241–265.

Dorph-Petersen, K.-A., Pierri, J.N., Wu, Q., Sampson, A.R., and Lewis, D.A. (2007). Primary visual cortex volume and total neuron number are reduced in schizophrenia. J. Comp. Neurol. *501*, 290–301.

Dorph-Petersen, K.-A., Caric, D., Saghafi, R., Zhang, W., Sampson, A.R., and Lewis, D.A. (2009). Volume and neuron number of the lateral geniculate nucleus in schizophrenia and mood disorders. Acta Neuropathol. *117*, 369–384.

Fujino, H., Sumiyoshi, C., Yasuda, Y., Yamamori, H., Fujimoto, M., Fukunaga, M., Miura, K., Takebayashi, Y., Okada, N., Isomura, S., et al. (2017). Estimated cognitive decline in patients with schizophrenia: A multicenter study. Psychiatry Clin. Neurosci. *71*, 294–300.

Harel, J., Koch, C., and Perona, P. (2007). Graph-Based Visual Saliency. In Advances in Neural Information Processing Systems 19 (NIPS 2006), (Cambridge, MA: MIT Press), pp. 545–552.

Inada, T., and Inagaki, A. (2015). Psychotropic dose equivalence in Japan. Psychiatry Clin. Neurosci. *69*, 440–447.

Itti, L. (2015). New Eye-Tracking Techniques May Revolutionize Mental Health Screening. Neuron *88*, 442–444.

Itti, L., and Baldi, P. (2009). Bayesian surprise attracts human attention. Vision Res. *49*, 1295–1306.

Itti, L., and Koch, C. (2001). Computational modelling of visual attention. Nat. Rev. Neurosci. *2*, 194–203.

Itti, L., Koch, C., and Niebur, E. (1998). A model of saliency-based visual attention for rapid scene analysis. IEEE Trans. Pattern Anal. Mach. Intell. *20*, 1254–1259.

Kapur, S. (2003). Psychosis as a state of aberrant salience: A framework linking biology, phenomenology, and pharmacology in schizophrenia. Am. J. Psychiatry *160*, 13–23.

Kauffmann, L., Ramanoël, S., and Peyrin, C. (2014). The neural bases of spatial frequency processing during scene perception. Front. Integr. Neurosci. *8*, 1–14.

Kay, S.R., Fiszbein, A., and Opler, L.A. (1987). The Positive and Negative Syndrome Scale (PANSS) for Schizophrenia. Schizophr. Bull. *13*, 261–276.

Knolle, F., Ermakova, A.O., Justicia, A., Fletcher, P.C., Bunzeck, N., Düzel, E., and Murray, G.K. (2018). Brain responses to different types of salience in antipsychotic naïve first episode psychosis: An fMRI study. Transl. Psychiatry *8*, 1–13.

Kuznetsova, A., Brockhoff, P.B., and Christensen, R.H.B. (2017). lmerTest Package: Tests in Linear Mixed Effects Models . J. Stat. Softw. *82*.

Lang, P.J., Bradley, M.M., and Cuthbert, B.N. (2008). International Affective Picture System (IAPS): Digitized Photographs, Instruction Manual and Affective Ratings, Technical Report A-8. Gainesville: The Center for Research in Psychophysiology, University of Florida.

Matsumoto, D., and Ekman, P. (1988). Japanese and Caucasian facial expressions of emotion (JACFEE) [Slides]. San Francisco, CA: Intercultural and Emotion Research Laboratory, Department of Psychology, San Francisco State University.

Miura, K., Hashimoto, R., Fujimoto, M., Yamamori, H., Yasuda, Y., Ohi, K., Umeda-Yano, S., Fukunaga, M., Iwase, M., and Takeda, M. (2014). An integrated eye movement score as a neurophysiological marker of schizophrenia. Schizophr. Res. *160*, 228–229.

Morita, K., Miura, K., Fujimoto, M., Yamamori, H., Yasuda, Y., Iwase, M., Kasai, K., and Hashimoto, R. (2017). Eye movement as a biomarker of schizophrenia: Using an integrated eye movement score. Psychiatry Clin. Neurosci. *71*, 104–114.

Morita, K., Miura, K., Fujimoto, M., Yamamori, H., Yasuda, Y., Kudo, N., Azechi, H., Okada, N., Koshiyama, D., Ikeda, M., et al. (2019). Eye movement abnormalities and their association with cognitive impairments in schizophrenia. Schizophr. Res. *209*, 255–262.

Morita, K., Miura, K., Kasai, K., and Hashimoto, R. (2020). Eye movement characteristics in schizophrenia: A recent update with clinical implications. Neuropsychopharmacol. Reports *40*, 2–9.

Okada, K., Miura, K., Fujimoto, M., Morita, K., Yoshida, M., Yamamori, H., Yasuda, Y., Iwase, M., Inagaki, M., Shinozaki, T., et al. (2021). Impaired inhibition of return during free-viewing behaviour in patients with schizophrenia. Sci. Rep. *11*, 3237.

Onitsuka, T., Hirano, Y., Nemoto, K., Hashimoto, N., Kushima, I., Koshiyama, D., Koeda, M., Takahashi, T., Noda, Y., Matsumoto, J., et al. (2022). Trends in big data analyses by multicenter collaborative translational research in psychiatry. Psychiatry Clin. Neurosci. *76*, 1–14.

Otero-Millan, J., Troncoso, X.G., Macknik, S.L., Serrano-Pedraza, I., and Martinez-Conde, S. (2008). Saccades and microsaccades during visual fixation, exploration, and search: foundations for a common saccadic generator. J. Vis. *8*, 21.1--18.

Rokem, A., Yoon, J.H., Ooms, R.E., Maddock, R.J., Minzenberg, M., and Silver, M.A. (2011). Broader Visual Orientation Tuning in Patients with Schizophrenia. Front. Hum. Neurosci. *5*.

Romano, J., Kromrey, J.D., Coraggio, J., Skowronek, J., and Devine, L. (2006). Exploring methods for evaluating group differences on the NSSE and other surveys:

Are the t-test and Cohen's d indices the most appropriate choices? Annu. Meet. South. Assoc. Institutional Res. 14–17.

Schechter, I., Butler, P.D., Silipo, G., Zemon, V., and Javitt, D.C. (2003). Magnocellular and parvocellular contributions to backward masking dysfunction in schizophrenia. Schizophr. Res. *64*, 91–101.

Selemon, L.D. (2001). Regionally diverse cortical pathology in schizophrenia: Clues to the etiology of the disease. Schizophr. Bull. *27*, 349–377.

Selemon, L.D., and Begovic′, A. (2007). Stereologic analysis of the lateral geniculate nucleus of the thalamus in normal and schizophrenic subjects. Psychiatry Res. *151*, 1–10.

Seymour, K., Stein, T., Sanders, L.L.O., Guggenmos, M., Theophil, I., and Sterzer, P. (2013). Altered Contextual Modulation of Primary Visual Cortex Responses in Schizophrenia. Neuropsychopharmacology *38*, 2607 EP----2612.

Silverstein, S.M. (2016). Visual Perception Disturbances in Schizophrenia: A Unified Model. In Nebraska Symposium on Motivation, pp. 77–132.

Silverstein, S.M., and Keane, B.P. (2011). Perceptual organization impairment in schizophrenia and associated brain mechanisms: Review of research from 2005 to 2010. Schizophr. Bull. *37*, 690–699.

Sprenger, A., Friedrich, M., Nagel, M., Schmidt, C.S., Moritz, S., and Lencer, R. (2013). Advanced analysis of free visual exploration patterns in schizophrenia. Front. Psychol. *4*, 1–10.

Tibber, M.S., Anderson, E.J., Bobin, T., Antonova, E., Seabright, A., Wright, B., Carlin, P., Shergill, S.S., and Dakin, S.C. (2013). Visual Surround Suppression in Schizophrenia. Front. Psychol. *4*.

Tseng, P.-H., Cameron, I.G.M., Pari, G., Reynolds, J.N., Munoz, D.P., and Itti, L. (2012). High-throughput classification of clinical populations from natural viewing eye movements. J. Neurol. *260*, 1–10.

Wang, S., Jiang, M., Duchesne, X.M., Laugeson, E.A., Kennedy, D.P., ADOLPHS, R., and Zhao, Q. (2015). Atypical visual saliency in autism spectrum disorder quantified through model-based eye tracking. Neuron *88*, 1–13.

Wolf, A., Ueda, K., and Hirano, Y. (2021). Recent updates of eye movement abnormalities in patients with schizophrenia: A scoping review. Psychiatry Clin. Neurosci. *75*, 82–100.

Yoon, J.H., Rokem, A.S., Silver, M.A., Minzenberg, M.J., Ursu, S., Ragland, J.D., and Carter, C.S. (2009). Diminished orientation-specific surround suppression of visual processing in schizophrenia. Schizophr. Bull. *35*, 1078–1084.

Yoon, J.H., Maddock, R.J., Rokem, A., Silver, M.A., Minzenberg, M.J., Ragland, J.D., and Carter, C.S. (2010). GABA concentration is reduced in visual cortex in schizophrenia and correlates with orientation-specific surround suppression. J. Neurosci. *30*, 3777–3781.

Yoshida, M., Itti, L., Berg, D.J., Ikeda, T., Kato, R., Takaura, K., White, B.J., Munoz, D.P., and Isa, T. (2012). Residual attention guidance in blindsight monkeys watching complex natural scenes. Curr. Biol. *22*, 1429–1434.

SUPPLEMENTAL INFORMATION

SUPPLEMENTAL TEXT

Consideration of resampling scheme. Considering the possibility that resampled

data could affect the results of the statistical analysis, the resampling procedure was

repeated 100 times in the statistical analysis with the full model using age-matched

control data (Figure 2B, right). For the interaction term, the range of F values was

$5.8*10^{-5}$ - 1.07 and that of P values was 0.30 - 0.99. For the main effect for the

participant group in the model without interaction, the range of F values was 19.0 - 36.9

and that of P values was $8.4*10^{-9}$ - $2.2*10^{-5}$. Thus, differences in the choice of

resampled data do not affect the result of the statistical analysis. Thus, one of the

resampled data was selected and used throughout the analysis in the main text.

Consideration of random sampling scheme. In Figure 3A, we also examined

whether the mean salience values were higher than the chance level, here defined as the

expected value from random sampling, i.e., $1.95 * 10^{-4}$ (= $1 / (80*64)$). In all cases of

the four models and two participant groups, they were significantly higher than chance

($p < 10^{-12}$, Wilcoxon signed-rank test with Bonferroni's correction for multiple

comparisons). This confirms that in all models and all participant groups, the

participants' gazes were directed to the salient locations in the test images.

Figure S1 (Related to Figure 3). Inhibition-of-return (IOR) cannot explain the effect of orientation salience.

(A) Possible scheme for a relationship between the return saccades (dotted line) and the visual salience of the test images (shown in brightness).

(B) Possible time courses of salience values when IOR contributes to salience-guided eye movements (left) and when IOR does not contribute to salience-guided eye movements (right). Our results in Figure 2 support the latter.

(C) Violin plots of the mean orientation salience at the first gaze for each 8-second viewing time. The same notation as Figure 3A.

	Subjects with schizophrenia (n = 82)		Age-matched healthy controls (n = 82)		Statistics	
	Mean ± SD		Mean ± SD		P value	Effect size
Sex (male/female)	42 / 40		35 / 47		0.28	0.09
Age (years)	35.1 ± 12.4		35.2 ± 12.3		0.94	-0.01
Education (years)	13.8 ± 2.6		15.1 ± 2.1		5.0×10^{-4}	0.30
Symptom-related						
Onset age (years)	24.4 ± 11.4		NA		NA	NA
Duration of illness (years)	10.7 ± 8.9		NA		NA	NA
CPZ equivalent (mg/day)	600.6 ± 545.2		NA		NA	NA
Cognitive decline	13.4 ± 12.8		NA		NA	NA
PANSS positive symptoms	19.1 ± 6.1		NA		NA	NA
PANSS negative symptoms	20.8 ± 6.1		NA		NA	NA
PANSS general psychopathology	43.8 ± 12.0		NA		NA	NA
PANSS total	83.7 ± 23.2		NA		NA	NA
Cognitive tests						
WAIS-3 FIQ	88.0 ± 17.9		112.6 ± 10.2		8.1×10^{-18}	0.81
WAIS-3 VC	94.3 ± 17.2		112.5 ± 12.9		8.5×10^{-12}	0.71
WAIS-3 PO	88.1 ± 18.0		106.1 ± 11.6		1.2×10^{-10}	0.67
WAIS-3 WM	89.6 ± 17.2		111.1 ± 14.1		1.4×10^{-13}	0.76
WAIS-3 PS	82.3 ± 18.3		110.1 ± 12.4		3.7×10^{-18}	0.87
WMS-R VerM	89.0 ± 21.5		114.8 ± 13.6		1.7×10^{-13}	0.73
WMS-R ViM	86.3 ± 18.9		105.5 ± 8.2		1.7×10^{-12}	0.70
WMS-R GM	86.4 ± 21.6		114.0 ± 12.4		2.0×10^{-15}	0.78
WMS-R AC	96.3 ± 15.0		112.0 ± 10.9		1.5×10^{-10}	0.64
WMS-R DR	83.7 ± 21.7		109.7 ± 12.2		8.0×10^{-14}	0.74
UPSA-B social	68.7 ± 16.9		83.0 ± 10.0		1.2×10^{-08}	0.55
SFS	101.2 ± 31.0		148.5 ± 19.6		3.5×10^{-19}	0.85
Work hours	13.0 ± 17.2		36.2 ± 16.9		1.4×10^{-13}	0.71
Saccade related properties						
FV MainSeq a	481.1 ± 116.5		452.6 ± 101.1		0.09	-0.15
FV MainSeq b	9.8 ± 4.9		9.7 ± 3.1		0.55	0.05
FV MainSeq c	28.7 ± 9.8		28.4 ± 7.6		0.84	0.02
FV Sac number	16.1 ± 5.0		21.3 ± 4.6		9.5×10^{-11}	0.59
FV Sac amplitude	2.9 ± 1.2		3.3 ± 0.8		**0.02**	0.21
FV Scanpath Length	72.6 ± 29.1		109.5 ± 22.9		5.9×10^{-14}	0.68

2   **Table S1. Demographic data (Related to STAR methods)**

The "Statistics" column denotes statistical tests for the difference between SZs and

healthy controls. All P values are the results of the Wilcoxon rank-sum test for two

independent samples, except for the "sex" row, where a χ^2 test for a 2 x 2 contingency

table is performed. P values in bold denote significant differences after correction of

multiple comparisons by FDR (P < 0.05). Cliff's delta is shown as the effect size.

Cramer's V is shown as the effect size for the χ^2 test. PANSS: Positive and negative

syndrome scale. WAIS-3: Wechsler Adult Intelligence Scale 3rd edition; WMS-R:

Wechsler Memory Scale-Revised; UPSA-B: UCSD Performance-based Skills

Assessment - Brief; SFS: Social Functioning Scale; FV: free-viewing; Sac: saccade.

MainSeq a, b, and c: parameters obtained from fitting to the main sequence relationship.

See Methods for details.

12