Aberrant visual salience in participants with schizophrenia
during free-viewing of natural images

Masatoshi Yoshida[1,*], Kenichiro Miura[2,*], Michiko Fujimoto[2,3], Hidenaga

Yamamori[2,3,4], Yuka Yasuda[2,5], Masao Iwase[3,6], Ryota Hashimoto[2]

[1] Center for Human Nature, Artificial Intelligence, and Neuroscience (CHAIN),

Hokkaido University, Sapporo, Japan

[2] Department of Pathology of Mental Diseases, National Institute of Mental Health,

National Center of Neurology and Psychiatry, Kodaira, Japan

[3] Department of Psychiatry, Osaka University Graduate School of Medicine, Suita,

Japan

[4] Japan Community Health Care Organization, Osaka Hospital, Osaka, Japan

[5] Medical Corporation Foster, Life Grow Brilliant Mental Clinic, Osaka, Japan

[6] Osaka Psychiatric Research Center, Osaka Psychiatric Medical Center, Osaka

Prefectural Hospital Organization, Hiarakata, Japan

[*] Correspondence: myoshi@chain.hokudai.ac.jp (M.Y.),

miura.kenichiro.2x@ncnp.go.jp (K.M.)

Abstract

Abnormalities in visual exploration affect the daily lives of patients with

schizophrenia; however, its origin is unknown. In this study, we examined whether

such abnormalities reflect aberrant processing of visual salience. Eye movements

of 82 patients and 252 healthy individuals viewing natural and/or complex images

were examined using saliency maps for static images to determine the contributions of low-level visual features to salience-guided eye movements. The results showed that the gazes of the participants with schizophrenia were attracted to position in the images with high orientation salience but not luminance or color salience. Further analyses revealed that orientation salience defined by the L+M channel of the DKL color space is specifically affected in schizophrenia, suggesting abnormalities in the magnocellular visual pathway. These results suggest aberrant processing of visual salience in schizophrenia, thereby connecting the dots between abnormalities in early visual processing and the aberrant salience hypothesis of psychosis.

Schizophrenia is a mental disorder with psychosis, that is, delusion and hallucination. Previous studies have successfully identified various behavioral markers of schizophrenia [1]. However, to understand the pathological mechanisms of schizophrenia, it is necessary to find links between these behavioral markers and theories about how the symptoms develop. Here, we examined the relevance of exploratory eye movements as a behavioral marker and the aberrant salience hypothesis of psychosis [2–4], which states that aberrant assignment of salience to elements of one's experience leads to delusion and hallucination.

Various abnormalities in eye movements have been reported in patients with schizophrenia [5,6]. Among them, significant differences in eye movements are observed in a simple free-viewing task. Gaze covers a relatively large area of the images in healthy participants, whereas participants with schizophrenia tend to limit their gaze to a narrower area of the images, resulting in shorter scanpath lengths than healthy participants [7–10]. Visual and visuo-cognitive processing are affected [11–13], while the

motor aspects of saccadic eye movements are less affected [10,14]. In schizophrenia, it

would be reasonable to assume that abnormalities in visual exploration result from

abnormalities in visual or visuo-cognitive processing.

We sought to determine whether these abnormalities in eye movements were due

to aberrant assignment of visual salience to objects in the visual scene. To this end, we

examined eye movements during free-viewing of natural images in participants with

schizophrenia (SZ) and healthy controls (HC). The visual salience of the presented

images was quantified by Itti and Koch's computational model of salience, which has

already proven useful for predicting and analyzing the visual exploration behavior of

humans and non-human primates [15–20]. In the present study, we first tested whether the

gaze sequences of participants with schizophrenia differed from those of healthy

controls in terms of salience values. Since the results showed case-control differences,

we examined in detail which low-level visual features contributed to the differences in

eye movements and found that the difference stemmed specifically from orientation

salience. We then explored the origin of the effects of orientation salience by examining

the stages of the salience computation in the model.

Results

Visual-oculomotor properties are affected in schizophrenia but oculomotor

properties are not

Eye-tracking data was obtained from 82 participants with schizophrenia and 252

healthy controls while freely viewing 56 natural and/or complex images. Supplementary

Table 1 summarizes the demographics of the participants, scores of cognitive tests, and

saccade characteristics during free-viewing. Analysis of saccades revealed that visual

exploration is affected in schizophrenia, as reported in previous studies [10,14]. The

average number of saccades during 8 seconds of the viewing period, the average

amplitude of saccades, and the scanpath length were smaller in participants with

schizophrenia than in healthy controls (Supplementary Table 1). In contrast, oculomotor

properties, as assessed by the fitted parameters of the main sequence relationship, were

relatively unaffected (Supplementary Table 1).

Salience-guided eye movements are affected in examples of single images

To investigate the relationship between the gaze positions (defined as endpoints of

regular saccades) and the visual salience of the test images, saliency maps were

computed using the Itti-Koch saliency computational model (Fig. 1A) [15,21].

In representative examples (for the test image #8 and #38), saccades of healthy

controls were distributed not only in highly salient positions (shown in yellow) but also

in other locations in the image (Fig. 1B, left). In contrast, saccades of participants with

schizophrenia remained at salient locations of the image during free-viewing (Fig. 1B,

right). To quantify the time course of salience values during the 8-second viewing

period, the salience values for each gaze were averaged across participants for the test

images and were plotted across time (Fig. 1C, left). The mean salience values were

higher throughout the viewing period in participants with schizophrenia (SZ; blue in

Fig. 1C, left) than in healthy controls (HC; magenta in Fig. 1C, left). The difference

between the participant groups becomes more evident when the salience values were

averaged across all saccades during the viewing period (Fig. 1C, right).

1   Visual salience attracts the gaze of participants with schizophrenia

To assess population averages, we plotted the salience values for each saccade

averaged across all test images and all participants (Fig. 2A). Consistent with the single

image results (Fig. 1C), the mean saliency values were consistently higher in SZs than

in HCs.

For statistical analyses, a linear mixed model with random intercepts and slopes

was used to test the main effect of the participant group (SZ vs. HC) and the interaction

between the participant group and the fixation number. Using the log of saccade

numbers as a factor, it is more reasonable to fit the data with a linear mixed model (Fig.

2B, left). Since the interaction term was not significant ($F(1, 330.11) = 0.17$ and $P =$

0.67), a model without interaction was selected. The main effect for the participant

group was highly significant ($F(1, 332.00) = 44.29$ and $P = 1.17*10^{-10}$). These results

suggest that the gaze of SZs is more likely to be directed toward visually salient

locations than that of HCs.

When the data from age-matched resampled data for HC were compared with the

data from SZ (Fig. 2B, right), the results of the statistical analysis were not affected.

The interaction term was not significant ($F(1, 159.69) = 0.02$ and $P = 0.87$), and a model

without interaction was selected. The main effect for the participant group was highly

significant ($F(1, 162.00) = 25.49$ and $P = 1.18*10^{-6}$). The resampling procedure was

repeated 100 times, one of which was used throughout the following analyses. See also

the supplementary text, "Consideration of resampling scheme" for more details.

The gaze of participants with schizophrenia is attracted by orientation

salience

To assess the contribution of low-level visual features, single-feature saliency

maps for color ("Col"), luminance ("Lum"), and orientation ("Ori") were used for

analysis as in the full salience model (Fig. 1A). In all three models, interaction terms

were not statistically significant ($F(1, 160.79) = 0.93$ and $P = 0.34$ for color, $F(1,$

$161.60) = 0.52$ and $P = 0.47$ for luminance, and $F(1, 160.14) = 1.79$ and $P = 0.18$ for

orientation, respectively). In the color (Fig. 2C left) and the luminance (Fig. 2C middle)

models, the main effects for the participant group were not significant ($F(1, 162.01) =$

0.16 and $P = 0.69$ for color and $F(1, 161.97) = 0.12$ and $P = 0.72$ for luminance,

respectively). In contrast, the main effect of the participant group was highly significant

($F(1, 161.99) = 44.80$ and $P = 3.39 *10^{-10}$) in the orientation model (Fig. 2C right).

These results suggest that the main effect in the participant group in the full model (Fig.

2B) is explained sorely by orientation salience.

The specific effect of orientation salience is also evident in time-averaged

data

Since the liner mixed model showed no significant interaction effect between the

participant group and saccade numbers in all four salience models (Fig. 2B and C), it is

reasonable to summarize the data by averaging all saccades during the viewing period

(Fig. 3A). The mean saliency value from the full model was higher in SZ than in HC (Z

$= 4.82$, $P = 5.75 *10^{-6}$; in Wilcoxon rank-sum test with Bonferroni correction). The

effect size evaluated by Cliff's delta was 0.44. The mean salience value from the

orientation model was higher in SZ than in HC ($Z = 6.41$, $P = 5.64 *10^{-10}$; in Wilcoxon

rank-sum test with Bonferroni correction). The effect size evaluated by Cliff's delta was

0.58, which is classified as a large effect. On the other hand, the mean salience value

from the color and the luminance models were not significantly different between SZ

and HC (Z = 0.08, P > 1.0 for color; Z = 0.26, P > 1.0 for luminance; in Wilcoxon rank-

sum test with Bonferroni correction). Based on these findings, the following analyses

were performed on the time-averaged data.

All of these mean salience values were significantly higher than chance. See also

the supplementary text, "Consideration of random sampling scheme" for more details.

We also tested whether the saliency values for the first saccade were significantly

different between SZ and HC (Supplementary Fig. 1). See Discussion for details.

Image category does not affect the effect of orientation salience

To examine whether the effects of orientation salience depend on image category,

the effect size (Cliff's delta) that evaluates the difference between the time-averaged

saliency values of HC and SZ for each test image was plotted for all four salience

models (Fig. 3B). Positive values in the effect size indicate that saliency values were

higher in SZ than in HC. The results (Fig. 3B) show that the effect sizes for the Full and

the orientation models were consistently positive for all image categories except for the

Face and Noise categories. This suggests that the effect of orientation salience is overall

robust to image categories.

The key computational stage that produces differences in orientation salience

is Gabor filtering

Next, we examined which stage of the salience computation produces differences

in orientation salience. In the Itti-Koch salience model, saliency maps for the orientation

feature are generated through five processing stages (Fig. 4A): 1) transforming the input

image into a grayscale image with various spatial resolutions (i.e., Gaussian pyramids),

2) filtering the images with Gabor patches with four different orientations, 3) center-

surround inhibition, 4) peak normalization, and finally, 5) adding all images together to

obtain the saliency map of the orientation feature. These intermediate maps were

calculated and used for analysis as was done in the analysis of Fig. 3A. Differences

between participant groups were detected not only in the saliency map for the

orientation feature (Fig. 4B, rightmost) but also in peak normalization, center-surround

inhibition, and Gabor filter (Fig. 4B, middle). In contrast, no differences between

participant groups were detected in intensity (Fig. 4B, leftmost). These differences

became more evident when each comparison was evaluated by the effect size (Fig. 4C).

These results suggest that the differences between participant groups emerge at the stage

of Gabor filtering with four orientations, a unique computational stage that is not

present in the color or luminance salience but only in the orientation salience.

It is also interesting to note that the effect size of Gabor filtering is larger for low-

pass images (those with higher numbers in Fig. 4C). Since it is already known that the

magnocellular pathway conveys relatively low spatial frequency signals [22] and that the

magnocellular pathway is specifically impaired in schizophrenia [11], this point is directly

examined in the next section.

the L+M channel of the DKL color space is dominant in the effect of

orientation salience

In the Itti-Koch salience model (hereafter referred to as the "original model"), the

orientation salience is calculated from grayscale images. Instead, it is possible to

decompose the test images into images of three channels (L+M, L-M, and S-(L+M)), based on the DKL color space [23]. The decomposed images were then processed to obtain saliency maps with (orientation map) or without (intensity map) Gabor filtering (Fig. 5A). We refer to this as the "Extended six-channel model."

Then the salience values for these six maps were compared between SZ and HC (Fig. 5B and C). In the intensity maps (Fig. 5B), all differences were not statistically significant ($Z = 0.21$, $P > 1.0$ for the L+M channel; $Z = 1.09$, $P > 1.0$ for the L-M channel; $Z = -0.51$, $P > 1.0$ for the S-(L+M) channel; in Wilcoxon rank-sum test with Bonferroni correction). In the orientation maps (Fig. 5C), all of the differences were statistically significant ($Z = 6.25$, $P = 2.51 *10^{-9}$ for the L+M channel; $Z = 4.20$, $P = 1.59 *10^{-4}$ for the L-M channel; $Z = 3.84$, $P = 7.49 *10^{-4}$ for the S-(L+M) channel; in Wilcoxon rank-sum test with Bonferroni correction). These results further support the finding of a specific effect of orientation salience and suggest that the L+M channel of the DKL color space is dominant in the effect of orientation salience.

Orientation salience is correlated with the scores of cognitive tests and visual oculomotor characteristics

Next, we examined whether orientation salience is correlated with demographic data, cognitive ability, and eye movement characteristics collected from the same participants (See [10,24] for details). When the dependent variables (demographic data, cognitive tests, or eye movement characteristics) were fitted with orientation salience with the participant group as a covariate, age was not correlated with orientation salience but WAIS-3, processing speed (PS), social functioning scale (SFS), and scanpath length (SPL) were correlated with orientation salience (Fig. 6A).

For a more systematic analysis, two regression models were constructed. In the

first model, the dependent variables were fitted with saliency values from the original

model, with the participant group as a covariate. The t-values obtained from the fitting

were then plotted (Fig. 6B). Orientation salience is correlated with various scores of

cognitive tests and eye movement characteristics (indicated by the gray bars in Fig. 6B).

Regression analysis also suggested that these correlations are specific to orientation.

In the second model in which the dependent variables were fitted with saliency

values from the extended six-channel model, with the participant group as a covariate,

orientation salience in the L+M channel was significantly correlated with a cognitive

test score (SFS) and eye movement characteristics (saccade amplitude and SPL) (Fig.

6C). In contrast, orientation saliences in the other channels are not correlated with these

values. These results again suggest that the L+M channel is dominant in the effect of

orientation salience.

Saliency maps for orientation are correlated with fixation maps

In Fig. 3A and Fig. 5C, saliency values were evaluated separately for each

feature/channel. It is important to note that saliency maps for the different

features/channels are spatially correlated (Fig. 7B). This correlation may affect

statistical analyses in Fig. 3A and Fig. 5C. To address this issue, we calculated the

partial correlation between the saliency maps and the "Fixation maps" which quantifies

gaze distribution as a density map summed across participants for each test image (Fig.

7A). We also created a fixation map for the difference between SZ and HC ("SZ-HC")

(Fig. 7A). The partial correlations of these maps were then calculated for each test

image (n=56).

For the original model, we plotted the partial correlations between the fixation

maps and the saliency maps for the three features (color, luminance, and orientation)

(Fig. 7C). For the three fixation maps (SZ, HC, and SZ-HC), the partial correlation with

the saliency map for orientation was significantly larger than zero (P = $4.0*10^{-8}$ for SZ,

P = $1.7*10^{-7}$ for HC, and P = $1.6*10^{-7}$ for SZ-HC, respectively; Wilcoxon signed-rank

test with Bonferroni correction). In contrast, the partial correlations with the saliency

map for color and luminance were not significantly larger than zero (P > 0.05,

Wilcoxon signed-rank test with Bonferroni correction) except for a negative correlation

in luminance for SZ-HC (P = $2.7*10^{-4}$).

For the extended six-channel model, three maps for orientation were shown (Fig.

7D). For two fixation maps (SZ, HC), the partial correlations with the saliency map for

the L+M and the L-M channels were significantly larger than zero (P = $2.0*10^{-5}$ and

0.002 for SZ, P = $6.0*10^{-5}$ and 0.008 for HC; Wilcoxon signed-rank test with

Bonferroni correction). On the other hand, for the SZ-HC fixation map, the partial

correlation with the saliency map of the L+M channel was significantly larger than zero

(p = $5.2*10^{-4}$, Wilcoxon signed-rank test with Bonferroni correction). These results

suggest that the L+M channel contributes specifically to the effect of orientation

salience, even after accounting for spatial correlations between saliency maps.

Discussion

In this study, we examined whether abnormalities in eye movements result from

aberrant processing of visual salience. For this purpose, we analyzed the eye movement

data during free-viewing with Itti-Koch's salience model (Fig. 1). We found that the

saliency values at the gaze of SZs were persistently higher during the viewing period

compared to the HCs (Fig. 2A and B). Further analysis using single-feature saliency

maps revealed that this difference was due to orientation salience (Fig. 2C). We then

confirmed that these results are robust for various image categories (Fig. 3). Next, we

delved into the stages of salience computation and found that differences between SZs

and HCs were found in an early stage of salience computation, where grayscale images

were filtered with Gabor patches with four orientations (Fig. 4). We also analyzed the

gaze with an extended salience model that evaluates the channels for the DKL color

space separately and found that the L+M cannel has a dominant role in the effects of

orientation salience (Fig. 5). In addition, the saliency values at gazes were not correlated

with symptom-related measures such as PANSS scores but were correlated with various

measures of cognitive functions and saccade-related characteristics (Fig. 6). Finally, we

evaluated a spatial correlation between fixation maps and saliency maps and found that

orientation salience in the L+M channel correlates specifically with the difference

between the fixation maps of the SZs and those of HCs. These results suggest that

visual processing of orientation salience is affected in participants with schizophrenia,

which is consistent with the aberrant salience hypothesis of psychosis, as will be

discussed shortly.

Relevance to the aberrant salience hypothesis of psychosis. The aberrant

salience hypothesis of psychosis proposes that an aberrant assignment of salience to the

elements of one's experience leads to delusion and hallucination [2–4]. The concept of

salience in this hypothesis covers not only salience derived from emotion and

motivation (motivational/incentive salience) [25], but also salience due to novelty and

sensory features (perceptual salience) [26]. The present study provides direct evidence that

visual salience is affected in schizophrenia, thereby explicitly extending the concept of

aberrant salience to the visual domain. In support of our findings, a recent study

revealed that brain responses to images with various forms of salience such as novelty,

negative emotion, targetness, and rarity/deviance are affected in schizophrenia [27].

Relationship with eye movement abnormalities: visual-oculomotor properties. As described in the Introduction, visual-oculomotor properties such as scanpath length, saccade number, and saccade amplitudes are affected in the gaze of SZs during free-viewing (see also Supplementary Table 1). We found that these visual-oculomotor properties are correlated with saliency values (Fig. 6). Since the bottom-up salience of the images is related to target selection, the abnormalities in visual exploration that have been reported in various studies [11] arise, at least in part, from the affected salience-guided eye movement found in this study.

Relationship with eye movement abnormalities: inhibition-of-return. Recently, our research group reported the finding that inhibition-of-return is impaired in SZs [28]. This might raise the question of whether return saccades toward salient stimuli are more frequent in schizophrenia (Supplementary Fig. 1A). If this is the case, salience values at the gaze could be higher in SZ than in HC. However, it is unlikely because salience values at the first gaze for the viewing period, where it has no contribution to inhibition-of-return, were significantly higher in SZ than HC in the orientation model (Supplementary Fig. 1C). Furthermore, if the SZs made more return saccades toward salient positions than in the HCs, the slope of the time course of saliency values would be shallower in the SZs than in the HCs (Supplementary Fig. 1B, left). However, this is not the case because analyses using linear mixed models showed no interaction between the participant group and saccade numbers. Thus, abnormalities in visual salience found in this study have a different origin from abnormal inhibition-of-return. For further analysis and discussion, see also Okada et al., 2021[28].

Contribution of low-level visual features, early visual cortex, and the magnocellular pathway. We found that salience computation for the orientation feature is specifically affected in schizophrenia. Consistent with this finding, previous findings have also shown that orientation processing is affected in schizophrenia [29,30].

Other papers have shown that the early visual cortex is involved in changes in

contextual modulation in schizophrenia [31–33]. The effect of orientation salience can

explain impairment in contour integration in schizophrenia [30] because contour

integration can be distracted by aberrant processing of orientation salience.

Based on the analysis of Fig. 4 and the accompanying text, we argue that the

crucial difference between SZs and HCs stems from their response to Gabor-filtered

images, which resembles responses in the early visual cortex. On the other hand, there

was no difference in gazes toward luminance salience. Computation of luminance

salience involves center-surround inhibition without Gabor filtering. Such processing

resembles the response of the lateral geniculate nucleus (LGN). We propose that

aberrant orientation salience reflects abnormalities at the level of the primary visual

cortex but not at the level of the LGN. Post-mortem morphological studies of the

schizophrenia brains support this possibility. The volume and number of neurons in the

primary visual cortex are reduced in schizophrenia [34,35] but it is not the case for the LGN

[36,37]. Thus, the idea that visual abnormalities in schizophrenia occur between the LGN

and primary visual cortex is consistent with both previous studies and the present

current study.

The analyses in Figs. 5, 6, and 7 suggest that orientation salience in the L+M

channel of the DKL color space is specifically affected in schizophrenia. Since the L+M

channel carries information in the magnocellular pathway [23], the present results are

consistent with previous findings that visual information processing in the

magnocellular pathway is specifically affected in schizophrenia [38–40].

Relationship with clinical/cognitive tests. Fig. 6 shows that symptom-related

measures such as the PANSS total score, CPZ equivalent, and duration of illness were

not significantly correlated with orientation salience. These results suggest that

orientation salience is a trait marker rather than a state marker. This is consistent with

the view that eye movement-related measures are trait markers of schizophrenia [41].

Fig. 6 also shows that scores on some of the cognitive tests are correlated with

orientation salience. This is consistent with previous findings that scores of cognitive

tests are correlated with visual oculomotor characteristics such as scanpath length[24]. It is

unlikely that the abnormality in scanpath length causes abnormalities in orientation

salience, as oculomotor properties of eye movements are less affected in schizophrenia.

Rather, we propose that aberrant processing of orientation salience causes abnormalities

in visual exploration, which can be assessed by visual oculomotor properties such as

scanpath length, which in turn causes difficulty in cognitive abilities such as social

functioning.

Visual salience in schizophrenia and autism spectrum disorder (ASD).

Analysis of visual salience during free-viewing can be a useful method for detecting

psychiatric or neurological disorders [42,43]. The current findings added another example

of the successful application of the computational salience model to the analysis of eye

movements in participants with psychiatric disorders.

Gazes of participants with autism spectrum disorder (ASD) showed a general

increase in visual salience, defined by the low-level visual features [44]. This appears

similar to the findings in the current study about schizophrenia but differs in that the

present study reported specific effects of the orientation feature. Also interesting is the

finding that the effect size of orientation salience was lower in the Face category than in

the other image categories in the present study (Fig. 3B). This suggests that faces are

relatively ineffective in discriminating between SZs and the HCs, in contrast to the

abnormalities in gaze and processing of faces in ASD [45].

In terms of the predictive coding/Bayesian brain hypothesis, both schizophrenia

and ASD are explained by difficulties in evaluating prediction errors in sensory signals

[46,47]. How to distinguish the two in terms of computational perspective is an ongoing

debate. By extending the concept of visual salience to Bayesian surprise [48] and treating

visual salience as spatial outliers from the current belief on what we see, saliency

analysis could help resolve this issue.

Future projects. In this study, we obtained clues about what is affected in the

brain of schizophrenia, for example, Gabor filtering that is presumably performed in the

magnocellular pathway of the early visual cortex. To understand exactly how the brain

performs such computation, it is necessary to understand the processes at the neuronal

level. To this end, neurophysiological studies using animal models of schizophrenia are

needed. Since eye-tracking during free-viewing and analysis of visual salience is an

experimental paradigm that has been successfully used in non-human primates such as

macaque monkeys [16] and marmosets [18,19], replicating the present results in a non-human

primate model of schizophrenia[19,49] would open the door to understanding the precise

brain mechanism of schizophrenia.

17
18

METHODS

Participants

Eye movement data were sampled from 82 SZs (male, 42; female, 40) and 252 HCs

(male, 144; female, 108) as part of a large-scale cohort recruited at Osaka University

(Supplementary Table 1) [10,14,24]. There was an overlap in data with a previous study on

eye movement abnormalities in schizophrenia [10]. All participants were biologically

unrelated, were of Japanese descent, and had no history of the ophthalmologic disease,

or neurological/medical conditions that could influence the central nervous system.

Specific exclusion criteria included atypical headaches, head trauma with loss of

consciousness, chronic lung disease, kidney disease, chronic hepatic disease, thyroid

disease, active cancer, cerebrovascular disease, epilepsy, seizures, substance-related

disorders, or mental retardation [10,14].

SZs were recruited from Osaka University Hospital and had been diagnosed by two or

more trained psychiatrists according to criteria from the DSM-IV based on the

Structured Clinical Interview for DSM-IV (SCID). Estimated cognitive decline was

calculated by the methods described by Fujino [55]. The current symptoms of the SZs

were assessed using the Positive and Negative Syndrome Scale (PANSS) [56], and daily

antipsychotic use was calculated using chlorpromazine (CPZ) equivalents (mg/day) [57].

HCs were assessed for psychiatric, medical, and neurological concerns using a non-

patient version of the SCID to exclude individuals with current or past contact with

psychiatric services or who had received psychiatric medication.

All participants provided written consent to the study after a full explanation of the

study procedures. Anonymity was preserved for all participants. The study was

performed in accordance with the World Medical Association's Declaration of Helsinki

and was approved by the Research Ethical Committee of Osaka University, the National

Center of Neurology and Psychiatry, and Center for Experimental Research in Social

Sciences, Hokkaido University.

Task and stimuli

In the free-viewing task, the participants faced a 19-inch liquid crystal display monitor

(1280 × 1024 pixels) placed 70 cm from the observer's eyes. Presentation of the visual

stimuli was done using the Psychophysics Toolbox extension [52] in MATLAB (The

Mathworks, Natick, MA, USA). Each trial began with the presentation of a fixation

point on the center of the display. Once the participant fixated on the fixation spot for a

random time, a test image was presented for 8 seconds. The participant was instructed to

view the image as they like. One task consisted of 56 images, the order of which was

randomly shuffled for each participant. The images were chosen from eight categories:

natural environments, buildings, everyday items, foods, faces, animals, fractal patterns,

and noise (seven images for each). The images of natural environments and animals

were selected from the International Affective Pictures System (IAPS) [50], and the face

images from Matsumoto and Ekman [51]. Since IAPS images are not allowed to be

published in scientific journals, only the saliency maps for these images (#8 and #38)

are shown in the figures. In cases when images needed to be shown as examples (Figs.

1A, 5A, and 6A), a photograph taken by one of the authors was used.

Recording and preprocessing of eye movement data

Recording and preprocessing of eye movement data were performed as follows [10]. Eye

position and pupil area of the left eye were measured at 1 kHz using EyeLink1000 (SR

Research, Ontario, Canada). Eye position data (in degrees) were smoothed with a digital

FIR filter (-3 dB at 30 Hz), and eye velocity and acceleration traces were derived from a

two-point difference algorithm. Eye movement recordings were segmented into blink,

saccade, and fixation periods. Detected saccades included both regular saccades and

microsaccades. Here, following previous papers on microsaccades during free-viewing

such as [58], saccades with amplitudes greater than one degree were selected as regular

saccades.

Computational models and saliency analysis

To assess salience-guided eye movements, we used a validated computational model of

visual attention and compared it with individual eye movements. The saliency maps for

the test images were computed with the Itti-Koch saliency model for static images [15]

implemented in the GBVS toolbox for Matlab [21]. The Itti and Koch model is a

neurobiologically inspired model which computes salient locations for low-level visual

features (Fig. 1A). Since we are interested in neurobiological mechanisms of saliency-

guided eye movements, the Itti-Koch model was chosen.

In the full model, saliency maps of the three features were summed with equal weights.

When evaluating the contribution of low-level visual features, single-feature saliency

maps were used (Fig. 1A). For a test image of 640*512 pixels, we obtained a saliency

map of 80*64 pixels. To treat the saliency maps as density maps, all maps were

normalized so that the sum of saliency values of all pixels was one.

Intermediate files for salience computation were used for the analysis in Figs. 4 and the

accompanying text. These files were also generated by the GBVS toolbox.

Extended six-channel model for visual salience

An "extended six-channel model" for visual salience was constructed for the analysis in

Figs. 5, 6, and 7 and the accompanying text. For this purpose, the test images were

decomposed into three-channel images (L+M, L-M, and S-(L+M)) based on the DKL

color space [23]. The decomposed images were then subjected to salience computation

separately with (orientation map) or without (intensity map) Gabor filtering (Fig. 5A).

The luminance salience in the original model corresponds approximately to the intensity

map of the L+M channel. The color salience in the original model corresponds

approximately to the intensity map of the L-M channel plus the S-(L+M) channel. The

orientation salience in the original model corresponds approximately to the orientation

map of the L+M channel. The other two maps (the orientation map of the L-M channel

and the orientation map of the S-(L+M) channel) are new components not found in the

original model.

Age-matched resampling

To account for the effect of the age difference between the healthy control group (HC,

28.8 ± 11.5 years, mean $\pm$SD) and the schizophrenia group (SZ, 35.1 ± 12.4 years, mean

$\pm$SD), resampling of the data for the HC group was performed to match the mean age of

the HC participants with that of the SZ group. For this purpose, the ages of both groups

were grouped into 5-year bins. The HC participants in each bin were then randomly

selected to match the number of SZ participants in each bin. This procedure was

repeated 100 times to obtain 100 sets of resampled HC participants. The mean age of

the resampled HC participants ranges from $34.9 - 35.2$ years, in close agreement with

that of the SZ group (35.1 years). The standard deviation of the age of the resampled

HC participants ranges from $12.0 - 12.5$ years, in close agreement with that of the SZ

group (12.4 years old). We confirmed that the results of statistical analysis were not

affected by the choice of the resampled dataset (see supplemental text, "Consideration

of resampling scheme"). Therefore, we selected one of the resampled datasets and used

it throughout our analysis.

Statistical analysis

Statistical analyses were performed using MATLAB (Mathworks, NY), except for the

linear mixed models, which were performed by the R packages. The significance level

was set at $p < 0.05$. Non-parametric Wilcoxon rank-sum tests after Bonferroni

correction were performed in a post-hoc analysis to compare SZ and HC for the median

salience values or other indices (Fig. 3A, Fig. 5B and C, Supplementary Fig. 1C).

Cliff's delta (Δ) was used to quantify the effect size of the estimated difference. For

interpretation, we followed general guidelines: negligible for $|\Delta| < 0.147$, small for

$0.147 <= |\Delta| < 0.33$, medium for $0.33 <= |\Delta| < 0.474$, and large for $|\Delta| >= 0.474$[59].

Linear mixed models

The procedure for the statistical analysis of the time course of the mean salience values

at saccade endpoints (Figs. 1 and 2) is as follows. First, a cut point for the number of

saccades was set because the number of saccades during the 8-second viewing time

varied between trials and between the participant groups (see Supplementary Table 1 for

the mean numbers of saccades; 16.1 in SZ and 20.9 in HC). The cut point was set to 16

because the mean number of images obtained for each saccade number (up to 56) is less

than half (Fig. 2A, inset).

We then fit linear mixed models using the R package lme4 [53] to obtain estimates and

statistics. For this purpose, the salience values for each saccade number and each

participant were averaged over 56 test images. Saccades that landed off-screen were

treated as not a number. The mean salience values ("Sal") were then treated as the

dependent variable and as a linear function of participant groups (HC and SZ as

"SubjectGroup") and saccade numbers (1st, 2nd…, 16th as "SaccadeNum"). The

random effects were modeled using random intercepts and random slopes for individual

differences (ranging from 1 to 334 as "SubjectID") nested under the participant group.

The model formulae in R format are as follows:

1) Sal ~ SubjectGroup + SaccadeNum + (SaccadeNum |SubjectID)

2) Sal ~ SubjectGroup * SaccadeNum + (SaccadeNum |SubjectID)

Model #1 is a model with the main effects of participant group and saccade number.

Model #2 is a model with an interaction between the participant group and saccade

number. The two-sided probability values and degrees of freedom associated with each

statistic were then determined using the Satterthwaite approximation implemented in the

R package LmerTest [54].

General linear models

To examine whether the salience values are corelated to other measures such as

demographic data, cognitive tests, and eye movement properties, general linear models

were constructed (Fig. 6). The model formulae in R format are as follows:

1) Dependent variables ~ SubjectGroup + Sal(Col) + Sal(Lum) + Sal(Ori)

2) Dependent variables ~ SubjectGroup + Sal(Int, L+M) + Sal(Int, L-M) +

12   Sal(Int, S-(L+M)) + Sal(Ori, L+M) + Sal(Ori, L-M) + Sal(Ori, S-(L+M))

In Model #1, the dependent variables (demographic data, cognitive tests, and saccade-

related properties) were fitted with the participant group ("SubjectGroup"), and the

salience values were calculated by the original saliency model consisting of three

features. In Model #2, the dependent variables were fitted with the participant group

("SubjectGroup") and the salience values were calculated by the extended six-channel

model. P values were adjusted for multiple comparisons using FDR.

Data availability

All data reported in this study cannot be deposited in a public repository because of

ethical prohibitions. To request access, contact the lead contact.

Code availability

Custom-written codes used for the analysis of this study are available from GitHub

(https://github.com/pooneil68/saliency_analysis DOI: 10.5281/zenodo.7312329).

REFERENCES

1. Onitsuka, T. *et al.* Trends in big data analyses by multicenter collaborative translational research in psychiatry. *Psychiatry Clin. Neurosci.* **76**, 1–14 (2022).

2. Kapur, S. Psychosis as a state of aberrant salience: A framework linking biology, phenomenology, and pharmacology in schizophrenia. *Am. J. Psychiatry* **160**, 13–23 (2003).

3. Howes, O. D., Hird, E. J., Adams, R. A., Corlett, P. R. & McGuire, P. Aberrant Salience, Information Processing, and Dopaminergic Signaling in People at Clinical High Risk for Psychosis. *Biol. Psychiatry* **88**, 304–314 (2020).

4. Howes, O. D. & Murray, R. M. Schizophrenia: an integrated sociodevelopmental-cognitive model. *Lancet* **383**, 1677–1687 (2014).

5. Morita, K., Miura, K., Kasai, K. & Hashimoto, R. Eye movement characteristics in schizophrenia: A recent update with clinical implications. *Neuropsychopharmacol. Reports* **40**, 2–9 (2020).

6. Wolf, A., Ueda, K. & Hirano, Y. Recent updates of eye movement abnormalities in patients with schizophrenia: A scoping review. *Psychiatry Clin. Neurosci.* **75**, 82–100 (2021).

7. Sprenger, A. *et al.* Advanced analysis of free visual exploration patterns in schizophrenia. *Front. Psychol.* **4**, 1–10 (2013).

8. Benson, P. J. *et al.* Simple viewing tests can detect eye movement abnormalities that distinguish schizophrenia cases from controls with exceptional accuracy. *Biol. Psychiatry* **72**, 716–724 (2012).

9. Beedie, S. A., Benson, P. J., Giegling, I., Rujescu, D. & St. Clair, D. M. Smooth pursuit and visual scanpaths: Independence of two candidate oculomotor risk markers for schizophrenia. *World J. Biol. Psychiatry* **13**, 200–210 (2012).

10. Morita, K. *et al.* Eye movement as a biomarker of schizophrenia: Using an integrated eye movement score. *Psychiatry Clin. Neurosci.* **71**, 104–114 (2017).

11. Butler, P. D., Silverstein, S. M. & Dakin, S. C. Visual Perception and Its Impairment in Schizophrenia. *Biol. Psychiatry* **64**, 40–47 (2008).

12. Silverstein, S. M. Visual Perception Disturbances in Schizophrenia: A Unified Model. in *Nebraska Symposium on Motivation* vol. 63 77–132 (2016).

13. Silverstein, S. M. & Keane, B. P. Perceptual organization impairment in schizophrenia and associated brain mechanisms: Review of research from 2005 to 2010. *Schizophr. Bull.* **37**, 690–699 (2011).

14. Miura, K. *et al.* An integrated eye movement score as a neurophysiological marker of schizophrenia. *Schizophr. Res.* **160**, 228–229 (2014).

15. Itti, L., Koch, C. & Niebur, E. A model of saliency-based visual attention for rapid scene analysis. *IEEE Trans. Pattern Anal. Mach. Intell.* **20**, 1254–1259 (1998).

16. Yoshida, M. *et al.* Residual attention guidance in blindsight monkeys watching complex natural scenes. *Curr. Biol.* **22**, 1429–1434 (2012).

17. Berg, D. J., Boehnke, S. E., Marino, R. A., Munoz, D. P. & Itti, L. Free viewing of dynamic stimuli by humans and monkeys. *J. Vis.* **9**, 1–15 (2009).

18. Chen, C.-Y. *et al.* Properties of visually-guided saccadic behavior and bottom-up attention in marmoset, macaque, and human. *J. Neurophysiol.* jn.00312.2020 (2020) doi:10.1152/jn.00312.2020.

19. Polyakova, Z., Iwase, M., Hashimoto, R. & Yoshida, M. The effect of ketamine on eye movement characteristics during free-viewing of natural images in common marmosets. *Front. Neurosci.* **16**, 1–12 (2022).

20. Borji, A., Cheng, M.-M., Jiang, H. & Li, J. Salient Object Detection: A Benchmark. *IEEE Trans. Image Process.* **24**, 5706–5722 (2015).

21. Harel, J., Koch, C. & Perona, P. Graph-Based Visual Saliency. in *Advances in Neural Information Processing Systems 19 (NIPS 2006)* 545–552 (MIT Press, 2007).

22. Kauffmann, L., Ramanoël, S. & Peyrin, C. The neural bases of spatial frequency processing during scene perception. *Front. Integr. Neurosci.* **8**, 1–14 (2014).

23. Derrington, A. M., Krauskopf, J. & Lennie, P. Chromatic mechanisms in lateral geniculate nucleus of macaque. *J. Physiol.* **357**, 241–265 (1984).

24. Morita, K. *et al.* Eye movement abnormalities and their association with cognitive impairments in schizophrenia. *Schizophr. Res.* **209**, 255–262 (2019).

25. Berridge, K. C. Evolving concepts of emotion and motivation. *Front. Psychol.* **9**, 1–20 (2018).

26. Itti, L. & Koch, C. Computational modelling of visual attention. *Nat. Rev. Neurosci.* **2**, 194–203 (2001).

27. Knolle, F. *et al.* Brain responses to different types of salience in antipsychotic naïve first episode psychosis: An fMRI study. *Transl. Psychiatry* **8**, 1–13 (2018).

28. Okada, K. *et al.* Impaired inhibition of return during free-viewing behaviour in patients with schizophrenia. *Sci. Rep.* **11**, 3237 (2021).

29. Rokem, A. *et al.* Broader Visual Orientation Tuning in Patients with Schizophrenia. *Front. Hum. Neurosci.* **5**, (2011).

30. Yoon, J. H. *et al.* Diminished orientation-specific surround suppression of visual processing in schizophrenia. *Schizophr. Bull.* **35**, 1078–1084 (2009).

31. Seymour, K. *et al.* Altered Contextual Modulation of Primary Visual Cortex Responses in Schizophrenia. *Neuropsychopharmacology* **38**, 2607 EP----2612 (2013).

32. Yoon, J. H. *et al.* GABA concentration is reduced in visual cortex in schizophrenia and correlates with orientation-specific surround suppression. *J. Neurosci.* **30**, 3777–3781 (2010).

33. Tibber, M. S. *et al.* Visual Surround Suppression in Schizophrenia. *Front. Psychol.* **4**, (2013).

34. Selemon, L. D. Regionally diverse cortical pathology in schizophrenia: Clues to the etiology of the disease. *Schizophr. Bull.* **27**, 349–377 (2001).

35. Dorph-Petersen, K.-A., Pierri, J. N., Wu, Q., Sampson, A. R. & Lewis, D. A. Primary visual cortex volume and total neuron number are reduced in schizophrenia. *J. Comp. Neurol.* **501**, 290–301 (2007).

36. Dorph-Petersen, K.-A. *et al.* Volume and neuron number of the lateral geniculate nucleus in schizophrenia and mood disorders. *Acta Neuropathol.* **117**, 369–384 (2009).

37. Selemon, L. D. & Begovic´, A. Stereologic analysis of the lateral geniculate nucleus of the thalamus in normal and schizophrenic subjects. *Psychiatry Res.* **151**, 1–10 (2007).

38. Schechter, I., Butler, P. D., Silipo, G., Zemon, V. & Javitt, D. C. Magnocellular and parvocellular contributions to backward masking dysfunction in schizophrenia. *Schizophr. Res.* **64**, 91–101 (2003).

39. Butler, P. D. *et al.* Dysfunction of early-stage visual processing in schizophrenia. *Am. J. Psychiatry* **158**, 1126–1133 (2001).

40. Butler, P. D. *et al.* Early-stage visual processing and cortical amplification deficits in schizophrenia. *Arch. Gen. Psychiatry* **62**, 495–504 (2005).

41. Beedie, S. A., Clair, D. M. S. & Benson, P. J. Atypical scanpaths in schizophrenia : Evidence of a trait- or state-dependent phenomenon ? **36**, 150–164 (2011).

42. Tseng, P.-H. *et al.* High-throughput classification of clinical populations from natural viewing eye movements. *J. Neurol.* **260**, 1–10 (2012).

43. Itti, L. New Eye-Tracking Techniques May Revolutionize Mental Health Screening. *Neuron* **88**, 442–444 (2015).

44. Wang, S. *et al.* Atypical visual saliency in autism spectrum disorder quantified through model-based eye tracking. *Neuron* **88**, 1–13 (2015).

45. Dalton, K. M. *et al.* Gaze fixation and the neural circuitry of face processing in autism. *Nat. Neurosci.* **8**, 519–526 (2005).

46. Adams, R. A., Stephan, K. E., Brown, H. R., Frith, C. D. & Friston, K. J. The computational anatomy of psychosis. *Front. Psychiatry* **4**, 47 (2013).

47. de Cruys, S. *et al.* Precise minds in uncertain worlds: Predictive coding in autism. *Psychol. Rev.* **121**, 649–675 (2014).

48. Itti, L. & Baldi, P. Bayesian surprise attracts human attention. *Vision Res.* **49**, 1295–1306 (2009).

49. Selvanayagam, J., Johnston, K. D., Wong, R. K., Schaeffer, D. & Everling, S. Ketamine disrupts gaze patterns during face viewing in the common marmoset. *J. Neurophysiol.* **126**, 330–339 (2021).

50. Lang, P. J., Bradley, M. M. & Cuthbert, B. N. *International Affective Picture System (IAPS): Digitized Photographs, Instruction Manual and Affective Ratings, Technical Report A-8. Gainesville: The Center for Research in Psychophysiology, University of Florida.* (2008).

51. Matsumoto, D. & Ekman, P. *Japanese and Caucasian facial expressions of emotion (JACFEE) [Slides]. San Francisco, CA: Intercultural and Emotion Research Laboratory, Department of Psychology, San Francisco State University.* (1988).

52. Brainard, D. H. The Psychophysics Toolbox. *Spat. Vis.* **10**, 433–436 (1997).

53. Bates, D., Mächler, M., Bolker, B. M. & Walker, S. C. Fitting linear mixed-effects models using lme4. *J. Stat. Softw.* **67**, (2015).

54. Kuznetsova, A., Brockhoff, P. B. & Christensen, R. H. B. lmerTest Package: Tests in Linear Mixed Effects Models . *J. Stat. Softw.* **82**, (2017).

55. Fujino, H. *et al.* Estimated cognitive decline in patients with schizophrenia: A multicenter study. *Psychiatry Clin. Neurosci.* **71**, 294–300 (2017).

56. Kay, S. R., Fiszbein, A. & Opler, L. A. The Positive and Negative Syndrome Scale (PANSS) for Schizophrenia. *Schizophr. Bull.* **13**, 261–276 (1987).

57. Inada, T. & Inagaki, A. Psychotropic dose equivalence in Japan. *Psychiatry Clin. Neurosci.* **69**, 440–447 (2015).

58. Otero-Millan, J., Troncoso, X. G., Macknik, S. L., Serrano-Pedraza, I. & Martinez-Conde, S. Saccades and microsaccades during visual fixation, exploration, and search: foundations for a common saccadic generator. *J. Vis.* **8**, 21.1--18 (2008).

59. Romano, J., Kromrey, J. D., Coraggio, J., Skowronek, J. & Devine, L. Exploring methods for evaluating group differences on the NSSE and other surveys: Are the t-test and Cohen's d indices the most appropriate choices? *Annu. Meet. South. Assoc. Institutional Res.* 14–17 (2006).

Acknowledgments

This research was supported in part by AMED under Grant Numbers JP19dm0207069 (MY and KM), JP21dk0307103 (RH), JP21wm0425012 (KM and RH), JP18dm0307002 (RH), and JP21uk1024002 (KM and RH); JSPS KAKENHI grant numbers 22H02936 (MY), JP20H03611 (RH), JP19H05467 (RH), JP20K06920 (KM); and Intramural Research Grant (3-1 and 4-6) for Neurological and Psychiatric Disorders of NCNP (KM and RH).

Author contributions

Conceptualization, M.Y.; Methodology, M.Y. and K.M.; Software, M.Y.; Investigation, M.F., H.Y., Y.Y. and M.I.; Resources, K.M. and R.H.; Data Curation: M.Y., K.M. M.F., H.Y., Y.Y. and M.I; Writing – Original Draft: M.Y.; Writing – Review & Editing: M.Y. and K.M.; Funding Acquisition, M.Y., K.M., and R.H.; Supervision, K.M. and R.H.

Competing interests

The authors declare no competing financial interests.

Figures and legends

Fig. 1. Salience-guided eye movements are affected in examples of single images.

(A) The saliency map was calculated from the Itti-Koch model. Visual salience for low-level visual features (color, "Col"; luminance, "Lum"; orientation, "Ori") was also computed in this model.

(B) Gaze positions of two control subjects (left) and two SZs (right) represented by numbers and superimposed on the saliency map of test images. Numbers indicate saccade order.

(C) Left: The saliency values averaged across participants are plotted over time for test images #8 (top) and #38 (bottom). Error bars denote standard errors across participants. Magenta, HC; blue, SZ. Right: the saliency values averaged across participants and saccades plotted as a violin plot. Symbols denote the median. Error bars denote the first and the third quartile.

Fig. 2. The gaze of participants with schizophrenia is attracted by orientation salience.

(A) The saliency values averaged across test images and participants are plotted over time after image onset. Magenta, the healthy controls (HC; n=252); blue, the participants with schizophrenia (SZ; n = 82). Each point denotes the mean salience value for each saccade (1st through 16th). Error bars denote the standard errors across participants. Inset: number of images obtained for each number of saccades (up to 56) during the 8-sec viewing period. The dotted line denotes the cutoff point (16th saccades). See Methods for details.

(B) (Left) Same as (A) but the data are plotted across saccade numbers on a log scale. (Right) Age-matched resampled data are plotted for healthy controls. Numbers on the plots denote P values for the main effect of the participant group.

(C) As in (B), but for single feature salience models.

Fig. 3. Image category does not affect the effect of orientation salience.

(A) Mean saliency values for all images and saccades are plotted for the healthy controls (HC) and the schizophrenia group (SZ). Data from four salience models are plotted. Symbols denote median values. Error bars denote the first and the third quartile. Numbers on the plots denote P values ("p") and the effect sizes ("Δ", Cliff's delta) of the Wilcoxon rank-sum test.

(B) Effect sizes (Cliff's delta) for the test of difference in time-averaged saliency values. Top: histograms of effect sizes.

Fig. 4. The key computational stage that produces differences in orientation salience is Gabor filtering.

(A) The Itti-Koch saliency model computes the saliency map of the orientation feature through five stages. Intermediate images for a given image are shown in grayscale.

(B) Averaged map values for intermediate images are plotted as in Fig. 3A.

(C) Effect sizes (Cliff's delta) for the tests in (B). Numbers indicate the spatial scales of the Gaussian pyramids. In the three plots in the center, the mean effect size for four orientations (bar), as well as the effect size for four orientations (symbols) were plotted. "o," 0 degrees; "x," 45 degrees; triangle, 90 degrees, and square, 135 degrees. The horizontal lines in the rightmost plot ("Saliency map (Orientation)") indicate the common guidelines for the effect size (Cliff's delta). See Methods for details.

Fig. 5. the L+M channel of the DKL color space is dominant in the effect of

orientation salience.

(A) To construct the extended six-channel model, the original image was decomposed

into three channels in the DKL color space: the magnocellular L+M channel, the

parvocellular L-M channel, and the koniocellular S-(L+M) channel. Then saliency maps

for intensity and orientation were obtained for each of the three channels (six maps in

total).

(B and C) As in Fig. 3A, the salience values averaged across test images and saccades

were plotted, but for six saliency maps. Symbols are the same as in Fig. 3A.

Fig. 6. Orientation salience is correlated with the scores of cognitive tests and visual oculomotor characteristics.

(A) Scattered plots for the mean orientation salience (averaged across images and

saccades) and age, WAIS-3 processing speed (PS), social functioning scale (SFS), and

scanpath length (SPL). Each dot represents the value for one participant. Magenta:

healthy controls (HC); blue, participants with schizophrenia (SZ). Lines indicate

regression lines for each participant group.

(B) Absolute value of t-values from regression analysis 1, where the dependent

variables were fitted individually with the saliency values of the original model. Gray

bars indicate statistical significance ($P < 0.05$) after correction of multiple comparisons

by FDR.

(C) As in (B), but those from regression analysis 2, where the dependent variables were

fitted individually with the saliency values of the extended six-channel model.

Fig. 7. Saliency maps for orientation are correlated with fixation maps.

(A) Examples of fixation maps for test images #8 and #38. SZ, the participants with schizophrenia; HC, the healthy controls (age-matched resamples); SZ-HC, the difference between SZ and HC.

(B) (Top) Colors indicate the median of partial correlation coefficients between saliency maps of the original model (top) and the extended six-channel model (bottom).

(C) Violin plots of the partial correlation between the fixation maps and saliency maps

(original model). Symbols are the same as in Fig. 3A.

(D) Violin plots of the partial correlation between the fixation maps and saliency maps

(the extended six-channel). Only the correlations with orientation maps are shown.

Numbers in (C) and (D) indicate P values for a significant difference from zero

(Wilcoxon signed-rank test after Bonferroni correction). Only the P values less than

0.05 are shown.

8