Epidemiology of Extrapulmonary Tuberculosis in Brunei Darussalam

Liling Chaw [1], Lena Mat Salleh[1,2], Rafizah Abdul Hamid[2], Kyaw Thu[2]

[1]PAPRSB Institute of Health Sciences, Universiti Brunei Darussalam
[2]Disease Control Division, Department of Environmental Health Services, Ministry of Health, Brunei Darussalam

Corresponding author
Liling Chaw, PhD
PAPRSB Institute of Health Sciences, Universiti Brunei Darussalam,
Jalan Tungku Link BE1410, Brunei Darussalam
Email: liling.chaw@ubd.edu.bn

Abstract word count: 200 words

Main text word count: 3235 words

Keywords:

Epidemiology, tuberculosis, extrapulmonary, risk factors, Brunei

Highlights:

- About 1/5 of all TB cases (19.0%) are EPTB
- Common EPTB types are lymphatic (44.8%), pleural (19.4%), and skeletal (11.6%).
- Females and younger age are associated with lymphatic EPTB
- Males and older age are associated with pleural EPTB
- No significant trend changes for EPTB cases during 18-year study period

Funding

This study is funded by Universiti Brunei Darussalam's University Research Grant (Ref: UBD/RSCH/URC/RG(b)/2019/011).

Conflict of interest

All authors have no conflict of interest to declare.

Ethics statement

Ethical approval was given by the Medical and Health Research and Ethics Committee, Ministry of Health, Brunei Darussalam (Ref: MHREC/UBD/2019/2).

CRediT authorship contribution statement

Liling Chaw: Conceptualization, Methodology, Data curation, Supervision, Formal analysis, Funding acquisition, Writing - original draft, Writing - review & editing.

Lena Mat Salleh: Conceptualization, Methodology, Data curation, Formal analysis, Writing - original draft, Writing - review & editing.

Rafizah Abdul Hamid: Supervision, Writing - review & editing.

Kyaw Thu: Data curation, Supervision, Writing - review & editing.

Abstract

Objectives: We reported the epidemiology of extrapulmonary tuberculosis (EPTB) in Brunei Darussalam from January 2001 to December 2018.

Design or methods: A retrospective cohort study was conducted where sociodemographic and clinical data for all active TB cases were collected. Factors associated with developing EPTB were investigated using logistic regression. Chi-square trend test was used to determine any trends during the 18-year study period.

Results: We identified 3,916 TB cases, among which 743 (19.0%) were EPTB cases. Lymphatic (44.8%) and pleural (19.4%) EPTB were most common. The main modes of diagnosis were tissue biopsy (73.6%) and radiologic assessment (18.3%). Treatment success and mortality rate were 79.7% and 7.0%, respectively. Associations with specific EPTB types varies with age-group and gender. Younger age-group (aOR ≥1.94) and females (aOR: 2.45 [95%CI: 1.94,3.11]) had higher adjusted odds of developing lymphatic EPTB, but had lower adjusted odds of developing pleural EPTB [younger age-group (aOR ≤0.54) and female (aOR: 0.41 [95%CI: 0.17,0.90])]. No significant trend differences were observed for overall and specific EPTB types.

Conclusions: Understanding EPTB epidemiology is important as it also contributes to the overall TB burden in a country. Future studies could be done to analyse the association between EPTB and presence of co-morbidities.

Introduction

Tuberculosis (TB) remains as a major public health concern globally, with an estimated 9.9 million cases and 1.5 million deaths in 2020 (World Health Organization 2018). Caused by a bacilli, *Mycobacterium tuberculosis*, TB can affect the lungs (defined as pulmonary TB [PTB]) and also affect other organs in the body (defined as extrapulmonary TB [EPTB]) (World Health Organization 2018). Although EPTB is generally less prevalent and also less infectious than PTB, it also contributes to the overall TB burden (Houda Ben et al. 2018) particularly in terms of TB-related morbidity, complications and lifelong sequelae as a consequence of the disease (Sandgren et al. 2013).

EPTB accounts for an estimate of 15% of the 7.0 million of global TB cases in 2018 (World Health Organization 2018), which is likely an underestimate due to diagnostic challenges (Houda Ben et al. 2018). EPTB may present with a myriad of symptoms, which can be specific to the anatomic site affected and may mimic symptoms of other more common diseases. Also, diagnosing EPTB often requires site-specific microbiological evaluations which could be difficult to access depending on affected site(s). As such, EPTB rates at country level tend to vary depending on the socio-economic situation and if adequate resources was provided to the national TB programmes (Houda Ben et al. 2018).

Recent epidemiological studies from other countries have reported increasing trends of EPTB prevalence accompanied by decreasing trends of PTB prevalence (Sandgren et al. 2013; Gomes et al. 2014; Wang et al. 2014; Lee 2015). The common anatomic sites of EPTB reported were lymphatic and pleural (Lee 2015), while other studies also reported high incidence of urogenital, skeletal, gastrointestinal and meningeal EPTB (Lee 2015; Pang et al. 2019). Conflicting evidence exist on the risk factors associated with EPTB; While risk factors such as being female, younger age, Asian and African ethnicity, and comorbidities such diabetes and HIV were found to be associated with EPTB (Yang et al. 2004; Lee 2015; Pang et al. 2019; Arega et al. 2020; Omar et al. 2021), there were also other reports leaning towards being male (Gomes et al. 2014; Wang et al. 2014) and increased age (Wang et al. 2014).

Brunei Darussalam [pop. 440,715 (Department of Economic Planning and Development 2021)] is a Southeast Asian country with intermediate TB burden, where the TB incidence was 57 per 100,000 population in 2017 (Ministry of Health Brunei Darussalam 2019). Annual TB incidence for the country has remained stagnant since 2004 (Ministry of Health Brunei Darussalam 2013). Non-communicable diseases is a major issue for Brunei, particularly diabetes mellitus which is a known risk factor for TB (Omar et al. 2021).

Despite being generally non-infectious and less recognised than PTB, EPTB cases also do contribute to the overall TB burden in a country. This is thus important to understand the epidemiology of EPTB, particularly in settings whose TB incidence rates remain stagnant. Determining the factors associated with EPTB could assist in improving its early detection and/or diagnosis, thereby allowing early treatment, prevent progression to infectious state and improving patient prognosis. Hence, we undertook this study to determine the epidemiological characteristics of EPTB cases in Brunei and the factors associated with developing EPTB when compared to PTB cases. Our study findings could provide more understanding on the burden of EPTB in the country, and could also contribute towards Brunei's goal of achieving TB elimination by 2050.

Methods

This is a retrospective cohort study involving all registered TB cases (including PTB and EPTB) identified in Brunei Darussalam between January 2001 and December 2018. These cases were registered by the National Tuberculosis Coordination Centre (NTCC), a centre established in 2000 to implement, monitor, coordinate and evaluate TB prevention and control activities in the country (Ministry of Health Brunei Darussalam 2013). All confirmed TB cases (initially presented at both public and private sector) are reported to NTCC or to its 3 directly observed treatment (DOTS) centres located in each of the 3 remaining districts for diagnosis, treatment and follow-up. Information obtained include year of diagnosis, sociodemographic details (age, gender, nationality, and district of residence), and details specific to the disease (type of TB, anatomic sites affected, mode of diagnosis and treatment outcome).

A PTB case is defined as one with TB involving lung parenchyma, while a EPTB case is one with TB involving other organs except lungs (Ministry of Health Brunei Darussalam 2013). The type of EPTB is defined based on the anatomical site where TB bacilli was found. In cases where TB bacilli was found in more than 1 anatomic site, the EPTB case is classified based on the main anatomical site affected. Following the national guidelines, miliary TB (defined as cases with TB involving >1 organ including the lung) was classified as a PTB case. As such, it was not possible to classify cases as with concurrent PTB and EPTB.

We classified type of EPTB into 7 categories based on the main anatomical site affected: lymphatic, pleural, skeletal, gastrointestinal, genitourinary, meningeal, and other (consists of uncommon sites such as breast, skin, brain and eye). The mode of diagnosis was divided into 6

categories: tissue biopsy, cytology, radiology, clinical diagnosis alone, mycobacterium TB DNA (MTB DNA) and unknown. Treatment outcomes were classified into 6 categories (World Health Organization 2013; Omar et al. 2021): (1) cured, defined as smear- or culture- positive TB patient at the start of treatment who was smear- or culture-negative in the last month of treatment; (2) completed, defined as TB patient who has completed treatment without failure but without record that shows smear- or culture- negative result in the last month of treatment; (3) failure, defined as TB patient whose smear or culture is positive on or after the fifth month of treatment; (4) died, defined as TB patient who died for any reason before or during treatment; (5) lost to follow-up, defined as TB patient who did not start treatment or whose treatment was interrupted for ≥2 consecutive months; or (6) not evaluated, defined as TB patient whose treatment outcome is unknown (including those who are still undergoing treatment) or who decided to seek treatment elsewhere, other than in Brunei. Not evaluated cases were mostly foreign workers who were diagnosed with active TB (either PTB or EPTB). In accordance with Brunei's Immigration Act (Laws of Brunei, 2002), such workers would have their employment terminated and asked to return to their home country to continue their TB treatment.

Statistical analysis

Socio-demographic characteristics of all cases (overall, PTB and EPTB) were described. Socio-demographic characteristics, modes of diagnosis and treatment outcome were described for all 7 types of EPTB. Annual proportions of PTB and EPTB cases were calculated, using the total number of cases reported in each respective year as the denominator. Similarly, annual proportions of each EPTB type were calculated using the total number of EPTB cases reported in each respective year as the denominator. Group comparison tests were conducted using Kruskal-Wallis, Chi-square or Fisher's Exact tests, whichever is appropriate. Logistic regression analysis was done to determine any factors associated with EPTB diagnosis. Variables with p-value of $p<0.1$ at the univariate analysis stage were included into the multiple logistic regression model; the exception was for age-group and gender which were included *a priori*. Chi-square trend test was used to analyse trends between the proportion of cases by EPTB types. In all analyses, a p-value of <0.05 was considered statistically significant. All analyses were performed using Microsoft Excel and R (ver. 3.6.3) (R Core Team 2021).

Results

A total of 3,916 TB cases were included in the study, out of which 19.0% (n=743) were diagnosed as EPTB. Among all TB cases (Table 1), their median age (IQR) was 44.0 years (IQR: 30.0 – 60.0), and mainly consisted of males (59.7%, n= 2338), locals (73.7%, n= 2886), and residing at Brunei-Muara district (63.7%, n= 2494). Among the EPTB cases, their median age (IQR) was 40.0 years (IQR: 27.5 – 56.0), about half were females (50.6%, n= 376), and mainly consisted of locals (78.6%, n= 582), and residing at Brunei-Muara district (70.6%, n= 525). When compared to PTB cases (Table 1), EPTB cases tend to be locals (adjusted Odds Ratio, aOR = 1.65 [95% Confidence Interval (95% CI): 1.35, 2.03]), and residing at Brunei-Muara district (Table 1).

Among the 743 EPTB cases, the common type diagnosed was lymphatic (44.8%, n=333), followed by pleural (19.4%, n=144) and skeletal (11.6%, n=86; Figure 1). The male to female ratio varied slightly between EPTB types, with greater disproportion towards females (for lymphatic EPTB) and males (for pleural and meningeal EPTB; Table 2). The main modes of diagnosis were tissue biopsy (73.6% for overall EPTB) and radiologic assessment (18.3% for overall EPTB). High proportions of treatment completed (72.0%) or cured (7.7%) cases were also observed among all EPTB types. The overall mortality rate was 7.0% (n= 52), with disproportionately higher mortality for meningeal EPTB cases (6 out of 22 cases, 27.3%).

Multiple logistic regression analysis (Table 3) showed that age is associated with specific EPTB types: adjusted odds of lymphatic EPTB affecting younger people were at least 1.94 times higher when compared to those >65 years, and those aged between 25-64 years had at least 3.15 times more odds to develop EPTB in other sites. On the other hand, younger people had lower adjusted odds of developing pleural (aOR for 25-44: 0.54 [95% CI: 0.32, 0.89] and aOR for 45-64: 0.42 [95% CI: 0.24, 0.73]) and gastrointestinal EPTB (aOR for 0-24: 0.26 [95% CI: 0.06, 0.78]), when compared to those >65 years. Females had higher adjusted odds of developing lymphatic (aOR: 2.45 [95% CI: 1.94, 3.11]) and skeletal (aOR: 1.91 [95% CI: 1.24, 2.95]) EPTB but had lower adjusted odds of developing pleural EPTB (aOR: 0.41 [95% CI: 0.17, 0.90]) when compared to males. Also, locals had higher adjusted odds of developing skeletal (aOR: 4.44 [95% CI: 2.04, 11.69]), gastrointestinal (aOR: 3.91 [95% CI: 1.84, 9.66]) and other types of EPTB (aOR: 3.42 [95% CI: 1.53, 9.14]) when compared to foreign residents. Lastly, residents of Brunei-Muara district have higher odds of developing lymphatic and pleural EPTB when compared to other smaller districts.

We analysed trends by types of EPTB during the 18-year study period. The proportions of both PTB and EPTB cases remained steady with no significant trend differences (Figure 2), where

EPTB cases accounted between 12.1% and 24.4% of the annual TB cases. Chi-square trend tests revealed no significant trend differences for all EPTB types, when compared with PTB cases (Figure 3).

Discussion

In this 18-year retrospective analysis, we found that about one-fifth of all the TB cases in Brunei (19.0%) were diagnosed as EPTB. This finding is line with that previously reported in population-based studies from other regions (Peto et al. 2009; Sandgren et al. 2013; Gomes et al. 2014; Wang et al. 2014; Lee 2015; Khan et al. 2019; Ohene et al. 2019), which ranges between 10.4% and 21.8% but lower than the 41% reported from England and Wales (Kruijshaar and Abubakar 2009).

We also found that lymphatic EPTB is the most dominant type in Brunei, which is similar in countries with low (Kruijshaar and Abubakar 2009; Peto et al. 2009), intermediate (Khan et al. 2019), and high (Arega et al. 2020) TB-burden. Differences in the dominant EPTB types can be seen in other similar studies particularly from high TB-burden countries, such as disseminated EPTB in Ghana (Ohene et al. 2019), skeletal (Pang et al. 2019) and pleural (Wang et al. 2014) EPTB in China. This point further adds on to the known observation that dominant EPTB types can vary across different geographical areas, possibly due (but not limited) to differences in socio-demographics, diagnostic capability and access to health care.

In addition, we observed interesting age and gender differences in the association between specific EPTB types. On one hand, younger age groups were associated with lymphatic and other EPTB while female gender was associated with lymphatic, skeletal and GI. Both factors (younger age and females) have previously been reported as factors of developing EPTB in high, intermediate and low TB-burden countries (Yang et al. 2004; Peto et al. 2009; Gomes et al. 2014; Khan et al. 2019; Pang et al. 2019; Arega et al. 2020). On the other hand, we found that pleural EPTB was associated with males and older age; a finding consistent with another study in a Chinese city (Wang et al. 2014), but contrasts with another study of Chinese inpatients where pleural EPTB was associated with males and younger age (Pang et al. 2019). Taken together, it is likely that factors associated with EPTB is much dependent on the specific type of EPTB as the main anatomic site. It would thus be more informative to analyse the factors associated with specific types EPTB, rather than for overall EPTB cases.

Another obstacle to clearly determine the epidemiological factors associated with EPTB is the presence of multi-site TB infections, with or without lung involvement. Recent studies have

reported that concurrent EPTB with other EPTB types or PTB was common among EPTB inpatients (Kang et al. 2020), and that the degree of association between socio-demographic factors (age and gender) and the different types of concurrent EPTB tends to vary (Kang et al. 2022).

We also observed that local Bruneian residents had at least 3 times higher odds of developing skeletal, gastrointestinal, and other EPTB. This finding contrasts with previous studies from low TB-burden countries (United States (Peto et al. 2009) and European Union (Sandgren et al. 2013)) that reported foreign residents as a factor associated with EPTB. This finding could reflect the true situation, albeit less precise due to the quite wide 95% confidence intervals. BCG vaccination is also routinely given to local Bruneians, thus ruling out BCG vaccination history as a factor. Another explanation is that it is possible foreign residents may opt to refuse to undergo further diagnostic tests to confirm an EPTB diagnosis due to the monetary costs involved. Similar to the latter countries, Brunei has a significant proportion of foreign residents working in the country, particularly employed as low-skilled workers in construction, wholesale and retail trade (Department of Economic Planning and Statistics Ministry of Finance and Economy Brunei Darussalam 2020; Chaw et al. 2022). Foreign workers are required to undergo medical check-up upon start and renewal of their employment contract, and those diagnosed with active TB (PTB or EPTB) would be asked to return to their home country for treatment (Laws of Brunei 2002). This is apparent from our study population, where 90.5% (86/95) of EPTB cases whose treatment outcome was classified as Not evaluated were foreign residents. From the employer's point of view, it could be more cost-effective for them to just send their foreign workers home upon initial suspicion of EPTB diagnosis during medical check-up. This could occur particularly in EPTB types that are difficult to obtain diagnosis confirmation and that require follow-up testing, such as skeletal EPTB. While TB-related testing and treatment is provided free of charge for everyone, foreign nationals are charged fees for diagnostic tests conducted prior to EPTB diagnosis confirmation; this is free for locals. This could be one reason why locals instead made up the majority of all EPTB cases in this study.

In addition, we found that those living in Brunei-Muara district were positively associated with EPTB (for overall, lymphatic and pleural). One explanation relates to population density; Brunei-Muara district is the smallest but most populated district in Brunei, and where the capital city is located and where 72.3% of the country's population resides (Department of Economic Planning and Development 2021). This district is also generally more urbanised than other districts; residing in urban areas was reported as one factor associated with EPTB in other countries (Gomes et al. 2014; Wang et al. 2014; Khan et al. 2019).

It should be noted that access to healthcare is not a major issue in Brunei, as all 4 districts have their own DOTS centres and district hospitals to enable adequate patient follow-up and diagnostic testing. This is quite evident with the high treatment success rate observed in our study population, where 79.7% of all EPTB cases were either cured or completed their treatment. Our treatment success rate is slightly better than that reported in Malaysia (67.6%) (Khan et al. 2019) and Ghana (70.1%) (Ohene et al. 2019). Efforts were also made to confirm EPTB diagnosis; 73.6% of all EPTB cases in our study population were diagnosed through tissue biopsy and only 0.3% was diagnosed through clinical judgement. These observations suggest that the national TB programme is doing well in EPTB case diagnosis, follow-up and treatment. Despite this, however, no trend changes could be observed for EPTB (and also for PTB) cases during the study period. This finding suggests that the known stagnant situation for the overall TB incidence in Brunei since 2004 (Ministry of Health Brunei Darussalam 2013) is also extended to both PTB and EPTB. One plausible explanation for this could be that people are simply unaware one can get TB in other parts of the body and could manifest in form of non-painful bumps; early symptoms that could be ignored and only seek medical attention when it becomes bigger or painful. One way is to promote public awareness of EPTB together and with the same rigour as for PTB, using existing mechanisms already in place for health information dissemination.

There are several limitations of this study that needs to be addressed. Firstly, data entry errors are possible due to the study's retrospective nature and in some cases, it is not possible to confirm misdiagnosis or to clarify uncertainties in the data. However, efforts were made to review any suspected data entry error by re-checking from either hardcopy or electronic health records (EHR). Secondly, we were unable to collect data on known factors associated with EPTB (such as low immune status, diabetic and HIV positive), as there were not available for the whole study period. This is because data were mainly collected through hardcopy records (with viable degree of completion for presence of co-morbidities) until 2013 when a national EHR system was established. However, this limitation may not greatly affect our results due to potentially small case numbers with co-morbidities. Using data from 2013 to 2018 (shorter but overlapping time period with this study), we have previously reported that 20.1% (53/264) of all EPTB cases were also diagnosed with diabetes (Omar et al. 2021). Also, annual HIV incidence in Brunei is low; the annual number of HIV case notifications from 2001 to 2018 ranged between 0 to 41 cases (unpublished data). Leveraging from the EHR system already in place, co-morbidity information can be added the data collection checklist for TB cases and digitally recorded for future epidemiological studies. Despite these limitations, the strength of the study is

the generalisability of the results to the Bruneian population, as all EPTB notifications in the country were captured in our dataset.

In conclusion, we observed 19.0% of all active TB cases during the 18-year study period had EPTB. Lymphatic EPTB was the most common type, followed by pleural and skeletal EPTB. We also observed that associations with specific EPTB types varies with age-group and gender. More attention should be given to EPTB as it also contributes to the overall TB burden in a country. Future studies could be done to analyse the association between EPTB and presence of co-morbidities.

Acknowledgements

We are grateful for all staff involved in data collection and reviewing process at NTCC and DOTS clinics.

References

Arega B, Mersha A, Minda A, Getachew Y, Sitotaw A, Gebeyehu T, et al. Epidemiology and the
     diagnostic challenge of extra-pulmonary tuberculosis in a teaching hospital in Ethiopia.
     PLoS One [Internet]. 2020;15(12 December):1–15. Available from:
     http://dx.doi.org/10.1371/journal.pone.0243945

Chaw L, Abdul Hamid R, Koh KS, Thu K. Contact investigation of tuberculosis in Brunei
     Darussalam: Evaluation and risk factor analysis. BMJ Open Respir Res.
     2022;9(1):e001224.

Department of Economic Planning and Development. Population estimates for Brunei
     Darussalam, 2020 [Internet]. Ministry of Finance and Economy Brunei Darussalam. 2021.
     Available from: https://deps.mofe.gov.bn/SitePages/Population.aspx

Department of Economic Planning and Statistics Ministry of Finance and Economy Brunei
     Darussalam. Labour Force Survey 2020 [Internet]. Bandar Seri Begawan; 2020. Available
     from: http://www.deps.gov.bn/SitePages/Labour Force.aspx

Gomes T, Reis-santos B, Bertolde A, Johnson JL, Riley LW, Maciel EL. Epidemiology of
     extrapulmonary tuberculosis in Brazil: a hierarchical model. BMC Infect Dis [Internet].
     2014;14(9). Available from: https://doi.org/10.1186/1471-2334-14-9

Houda Ben A, Makram K, Chakib M, Khaoula R, Fatma H, Fatma S, et al. Extrapulmonary
     Tuberculosis: Update on the Epidemiology, Risk Factors and Prevention Strategies. Int J
     Trop Dis. 2018;1(1):1–6.

Kang W, Liu S, Du J, Tang P, Chen H, Liu J, et al. Epidemiology of concurrent extrapulmonary
     tuberculosis in inpatients with extrapulmonary tuberculosis lesions in China: a large-scale
     observational multi-centre investigation. Int J Infect Dis [Internet]. 2022;115:79–85.
     Available from: https://doi.org/10.1016/j.ijid.2021.11.019

Kang W, Yu J, Du J, Yang S, Chen H, Liu J, et al. The epidemiology of extrapulmonary
     tuberculosis in China: A large-scale multicenter observational study. PLoS One [Internet].
     2020;15(8 August 2020):1–15. Available from:
     http://dx.doi.org/10.1371/journal.pone.0237753

Khan AH, Sulaiman SAS, Laghari M, Hassali MA, Muttalif AR, Bhatti Z, et al. Treatment
     outcomes and risk factors of extra-pulmonary tuberculosis in patients with co-morbidities.
     BMC Infect Dis. 2019 Aug;19(1).

Kruijshaar ME, Abubakar I. Increase in extrapulmonary tuberculosis in England and Wales 1999 – 2006. Thorax. 2009;1090–5.

Laws of Brunei. Chapter 17 Immigration [Internet]. 2002 [cited 2021 Jan 12]. Available from: https://www.ilo.org/wcmsp5/groups/public/@ed_protect/@protrav/@ilo_aids/documents/legaldocument/wcms_117279.pdf

Lee JY. Diagnosis and Treatment of Extrapulmonary Tuberculosis. Tuberc Respir Dis (Seoul). 2015;78(2):47–55.

Ministry of Health Brunei Darussalam. Guidelines for Tuberculosis Control in Brunei Darussalam [Internet]. 2013. Available from: http://www.moh.gov.bn/SitePages/Downloads.aspx

Ministry of Health Brunei Darussalam. Health Information Booklet 2017 (Second Revision) [Internet]. 2019 [cited 2020 Feb 11]. Available from: http://moh.gov.bn/SitePages/Health Information Booklet.aspx

Ohene S-A, Bakker MI, Ojo J, Toonstra A, Awudi D, Klatser P. Extra-pulmonary tuberculosis: A retrospective study of patients in Accra, Ghana. Cardona P-J, editor. PLoS One [Internet]. 2019 Jan 9;14(1):e0209650. Available from: https://dx.plos.org/10.1371/journal.pone.0209650

Omar N, Wong J, Thu K, Alikhan F, Chaw L. Prevalence and associated factors of diabetes mellitus among tuberculosis patients in Brunei Darussalam: A 6-year retrospective cohort study. Int J Infect Dis [Internet]. 2021;105(April 2021):267–73. Available from: https://doi.org/10.1016/j.ijid.2021.02.064

Pang Y, An J, Shu W, Huo F, Chu N, Gao M, et al. Epidemiology of extrapulmonary tuberculosis among inpatients, China, 2008-2017. Emerg Infect Dis. 2019 Mar;25(3):457–64.

Peto HM, Pratt RH, Harrington TA, Lobue PA, Armstrong LR. Epidemiology of Extrapulmonary Tuberculosis in the United States , 1993 – 2006. 2009;30333:1350–7.

R Core Team. R: A language and environment for statistical computing [Internet]. Vienna, Austria; 2021. Available from: http://www.r-project.org/

Sandgren A, Hollo V, van der Werf MJ. Extrapulmonary tuberculosis in the European union and European economic area, 2002 to 2011. Eurosurveillance. 2013;18(12):1–9.

Wang X, Yang Z, Fu Y, Zhang G, Wang X, Zhang Y, et al. Insight to the epidemiology and risk factors of extrapulmonary tuberculosis in Tianjin, China during 2006-2011. PLoS One.

2014 Dec;9(12).

World Health Organization. Definitions and reporting framework for tuberculosis 2013 revision
(updated December 2014) [Internet]. Vol. 2019. 2013. Available from:
https://www.who.int/tb/publications/definitions/en/

World Health Organization. Global tuberculosis report 2019 [Internet]. Vol. 2018. 2018.
Available from: http://www.who.int/tb/publications/global_report/en/

Yang Z, Kong Y, Wilson F, Foxman B, Fowler AH, Marrs CF, et al. Identification of Risk Factors
for Extrapulmonary Tuberculosis. Extrapulmonary Tuberculosis Risk Factors • CID. 2004.

Table 1. Characteristics of pulmonary TB (PTB) and extrapulmonary TB (EPTB) cases in Brunei, 2001-2018

Characteristics	Total (n = 3916)	Pulmonary TB (n = 3173)	Extrapulmonary TB (n = 743)	Crude OR	95% CI	Adjusted OR*	95% CI
Age Group in years							
0-24	551 (14.1)	423 (13.3)	128 (17.2)	**1.61**	**1.23, 2.12**	1.4	0.96, 2.04
25-44	1448 (37.0)	1152 (36.3)	296 (39.8)	**1.37**	**1.09, 1.73**	1.18	0.86, 1.62
45-64	1139 (29.1)	943 (29.7)	196 (26.4)	1.11	0.87, 1.42	0.97	0.71, 1.34
>65	778 (19.8)	655 (20.7)	123 (16.6)	1		1	
Gender							
Male	2338 (59.7)	1971 (62.2)	367 (49.4)	1		1	
Female	1578 (40.3)	1202 (37.8)	376 (50.6)	**1.68**	**1.43,1.98**	1.06	0.70,1.58
Nationality							
Local	2886 (73.7)	2304 (72.6)	582 (78.3)	**1.36**	**1.13, 1.65**	**1.65**	**1.35,2.03**
Foreign	1030 (26.3)	869 (27.4)	161 (21.7)	1		1	
District of residence							
Brunei Muara	2494 (63.7)	1969 (62.0)	525 (70.6)	1		1	
Belait	736 (18.8)	624 (19.7)	112 (15.1)	**0.67**	**0.54,0.84**	**0.69**	**0.55,0.87**
Tutong	540 (13.8)	450 (14.2)	90 (12.1)	**0.75**	**0.58.0.95**	**0.74**	**0.57,0.95**
Temburong	146 (3.7)	130 (4.1)	16 (2.2)	**0.46**	**0.26,0.76**	**0.46**	**0.26,0.76**

*Model was also adjusted with interaction term (age group & gender). Significant positive interaction observed between gender & age group 25-44 years

Table 2. Distribution of extrapulmonary TB (EPTB) by demographic and clinical characteristics, Brunei 2001-2018

Characteristic	Total (n = 743) n(%)	Lymphatic (n = 333) n(%)	Pleural (n = 144) n(%)	Skeletal (n = 86) n(%)	Gastrointestinal (n = 68) n(%)	Genitourinary (n = 39) n(%)	Meningeal (n = 22) n(%)	Others (n = 51) n(%)	p value
Median age in years (IQR)	40.0 (27.5 - 56.0)	32.0 (24.0 - 45.0)	46.0 (31.8 - 68.0)	52.5 (34.8 - 64.0)	45.0 (31.0 - 66.0)	53.0 (41.0 - 60.5)	41.0 (29.3 - 56.8)	45.0 (33.5 - 53.5)	<0.001*
Age range (years)	1 to 95	1 to 84	1 to 97	15 to 85	18 to 83	22 to 81	17 to 85	8 to 80	
Age group (years)									
0-24	128 (17.2)	85 (25.5)	18 (12.6)	12 (14.0)	3 (4.4)	3 (7.7)	3 (13.6)	5 (9.8)	
25-44	296 (39.8)	159 (47.8)	49 (34.2)	19 (22.1)	30 (44.1)	10 (25.6)	10 (45.5)	19 (37.3)	<0.001
45-64	196 (26.4)	66 (19.8)	35 (24.5)	34 (39.5)	15 (22.1)	19 (48.7)	4 (18.2)	22 (43.1)	
>65	123 (16.6)	23 (6.9)	41 (28.7)	21 (24.4)	20 (29.4)	7 (18.0)	5 (22.7)	5 (9.8)	
Gender									
Male	367 (49.4)	126 (37.8)	100 (69.4)	40 (46.5)	35 (51.5)	22 (56.4)	14 (63.6)	30 (58.8)	<0.001
Female	376 (50.6)	207 (62.2)	44 (30.6)	46 (53.5)	33 (48.5)	17 (43.6)	8 (36.4)	21 (41.2)	
Nationality									
Local	582 (78.3)	237 (71.2)	109 (75.7)	80 (93.0)	61 (89.7)	33 (84.6)	17 (77.3)	45 (88.2)	<0.001
Foreign	161 (21.7)	96 (28.8)	35 (24.3)	6 (7.0)	7 (10.3)	6 (15.4)	5 (22.7)	6 (11.8)	
District									
Brunei-Muara	525 (70.6)	246 (73.9)	102 (70.8)	56 (65.1)	45 (66.2)	28 (71.8)	13 (59.1)	35 (68.6)	
Belait	112 (15.1)	47 (14.1)	29 (20.1)	10 (11.6)	10 (14.7)	5 (12.8)	3 (13.6)	8 (15.7)	0.02
Tutong	90 (12.1)	38 (11.4)	10 (7.0)	15 (17.5)	12 (17.6)	4 (10.3)	6 (27.3)	5 (9.8)	
Temburong	16 (2.2)	2 (0.6)	3 (2.1)	5 (5.8)	1 (1.5)	2 (5.1)	0 (0.0)	3 (5.9)	
Mode of diagnosis									
Tissue biopsy	547 (73.6)	261 (78.4)	91 (63.2)	63 (73.2)	51 (75.0)	31 (79.5)	17 (77.3)	33 (64.7)	<0.001
Radiology	136 (18.3)	59 (17.7)	34 (23.6)	12 (16.3)	11 (16.2)	5 (12.8)	1 (4.5)	12 (23.5)	
Clinical alone	2 (0.3)	1 (0.3)	1 (0.7)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	
Cytology	15 (2.0)	0 (0.0)	12 (8.3)	0 (0.0)	3 (4.4)	0 (0.0)	0 (0.0)	0 (0.0)	
MTB DNA	30 (4.0)	9 (2.7)	2 (1.4)	5 (5.8)	3 (4.4)	2 (5.1)	4 (18.2)	5 (9.8)	
Unknown	13 (1.8)	3 (0.9)	4 (2.8)	4 (4.7)	0 (0.0)	1 (2.6)	0 (0.0)	1 (2.0)	
Outcome of treatment									

Cured	57 (7.7)	29 (8.7)	11 (7.7)	6 (7.0)	4 (5.8)	2 (5.1)	1 (4.5)	4 (7.8)	
Completed	535 (72.0)	235 (70.6)	93 (65.0)	67 (77.9)	56 (82.4)	35 (89.7)	10 (45.5)	39 (76.5)	
Died	52 (7.0)	14 (4.2)	16 (11.2)	8 (9.3)	5 (7.4)	1 (2.6)	6 (27.3)	2 (3.9)	
Defaulted	3 (0.4)	1 (0.3)	0 (0.0)	1 (1.2)	1 (1.5)	0 (0.0)	0 (0.0)	0 (0.0)	<0.001
Treatment failure	1 (0.1)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (2.0)	
Not evaluated	95 (12.8)	54 (16.2)	23 (16.0)	4 (4.6)	2 (2.9)	1 (2.6)	5 (22.7)	5 (9.8)	

*Post-hoc Dunn test revealed that age-group of lymphatic EPTB patients were significantly different than all other EPTB types (except for meningeal)

Table 3. Multiple logistic regression results for 7 types of extrapulmonary TB (EPTB) compared with pulmonary TB (PTB), Brunei 2001-2018

Characteristic	Lymphatic (n = 333) OR (95% CI)	*Pleural (n = 144) OR (95% CI)	Skeletal (n = 86) OR (95% CI)	Gastrointestinal (n = 68) OR (95% CI)	Genitourinary (n = 39) OR (95% CI)	Meningeal (n = 22) OR (95% CI)	Others (n = 51) OR (95% CI)
Age group in years							
0-24	**4.88 (3.06, 8.07)**	0.72 (0.37, 1.35)	0.99 (0.46, 2.02)	**0.26 (0.06, 0.78)**	0.64 (0.14, 2.32)	0.94 (0.19, 3.86)	1.81 (0.50, 6.57)
25-44	**3.48 (2.26, 5.60)**	**0.54 (0.32, 0.89)**	0.76 (0.39, 1.45)	1.24 (0.69, 2.28)	0.79 (0.30, 2.20)	1.15 (0.40, 3.70)	**3.15 (1.23, 9.65)**
45-64	**1.94 (1.21, 3.22)**	**0.42 (0.24, 0.73)**	1.26 (0.73, 2.23)	0.59 (0.29, 1.15)	1.87 (0.82, 4.82)	0.56 (0.14, 2.11)	**3.45 (1.40, 10.37)**
>65	1	1	1	1	1	1	1
Gender							
Male	1	1	1	1	1	1	1
Female	**2.45 (1.94, 3.11)**	**0.41 (0.17, 0.90)**	**1.91 (1.24, 2.95)**	1.60 (0.98, 2.59)	1.34 (0.70, 2.53)	0.92 (0.36, 2.16)	1.15 (0.65, 2.01)
Nationality							
Local	-	-	**4.44 (2.04, 11.69)**	**3.91 (1.84, 9.66)**	-	-	**3.42 (1.53, 9.14)**
Foreign	1	1	1	1	1	1	1
District of residence							
Brunei Muara	1	1	1	1	1	1	1
Belait	**0.67 (0.47, 0.92)**	0.89 (0.57, 1.34)	0.56 (0.27, 1.06)	-	-	-	-
Tutong	0.80 (0.55, 1.13)	**0.39 (0.19, 0.73)**	1.00 (0.54, 1.75)	-	-	-	-
Temburong	**0.15 (0.02, 0.46)**	0.43 (0.11, 1.17)	1.07 (0.37, 2.48)	-	-	-	-

Figure 1. Types of extrapulmonary TB (EPTB) disease among 743 EPTB cases in Brunei, 2001 - 2018

The following data accompanies the top chart:

	2001-2	2003-4	2005-6	2007-8	2009-10	2011-12	2013-14	2015-16	2017-18
PTB	83.6	81.1	82.6	78.2	79.6	82.3	79.5	81.1	81.3
EPTB	16.4	18.9	17.4	21.8	20.4	17.7	20.5	18.9	18.7

Figure 2. Relative proportion of (top) pulmonary TB (PTB) and extrapulmonary TB (EPTB) cases, and (bottom) of all 7 types extrapulmonary TB (EPTB) by year of diagnosis (reported in 2-year intervals).