Wastewater genomic surveillance captures early detection of Omicron in Utah

Pooja Gupta[1,2]*, Stefan Liao[1], Maleea Ezekiel[1], Nicolle Novak[1], Alessandro Rossi[1], Nathan

LaCross[2], Kelly Oakeson[1], and Andreas Rohrwasser[1]

[1] Utah Public Health Laboratory, Utah Department of Health and Human Services, Salt Lake

City, UT 84129, USA

[2] Utah Department of Health and Human Services, Salt Lake City, UT 84129, USA

*Corresponding author:

Pooja Gupta

Utah Public Health Laboratory, 4431 South 2700 West, Salt Lake City, Utah 84129

Phone: 706-308-3189

Email: pgupta@utah.gov

Abstract

Wastewater-based epidemiology has emerged as a powerful public health tool to trace new

outbreaks, detect trends in infection and provide an early warning of COVID-19 community

spread. Here, we investigated the spread of SARS-CoV-2 infections across Utah by

characterizing lineages and mutations detected in wastewater samples. We sequenced over 1,200

samples from 32 sewersheds collected between November 2021 and March 2022. Wastewater

sequencing confirmed the presence of Omicron (B.1.1.529) in Utah in samples collected on

November 19, 2021, up to ten days before its corresponding detection via clinical sequencing.

Analysis of diversity of SARS-CoV-2 lineages revealed Delta as the most frequently detected

lineage during November, 2021 (67.71%), but it started declining in December, 2021 with the

onset of Omicron (B.1.1529) and its sub-lineage BA.1 (6.79%). Proportion of Omicron increased

to ~58% by January 4th 2022 and completely displaced Delta by February 7th, 2022. Wastewater

genomic surveillance revealed the presence of Omicron sub-lineage BA.3, a lineage that is yet to

be identified from Utah's clinical surveillance. Interestingly, several Omicron-defining mutations

began to appear in early November, 2021 and increased in prevalence across sewersheds from

December to January. Our study suggests that tracking epidemiologically relevant mutations is

critical in detecting emerging lineages in the early stages of an outbreak. Wastewater genomic

epidemiology provides an unbiased representation of community-wide infection dynamics and is

an excellent complementary tool to SARS-CoV-2 clinical surveillance, with the potential of

guiding public health action and policy decisions.

Keywords

SARS-CoV-2, Omicron, wastewater surveillance, whole genome sequencing, Utah, public health

1. Introduction

Since the beginning of the COVID-19 pandemic, the world has witnessed the evolution of the

SARS-CoV-2 virus in real time, with the emergence of new variants and the rapid spread of

COVID-19 affecting millions of lives. So far, five variants of concern (VOCs) have been verified

by the World Health Organization (WHO): Alpha, Beta, Gamma, Delta and Omicron. Globally,

many countries have now lifted COVID-related restrictions, limited vaccine mandates and

transitioned to normality. Thus, there remains an increasing concern over the emergence of novel

SARS-CoV-2 variants with increased virulence, transmissibility and immune escape potential

affecting the efficacy of currently available vaccines (1,2). Over the course of the pandemic,

COVID-19 surveillance strategies have relied on testing of symptomatic individuals and their

contacts to slow the spread of the pandemic. However, symptoms seem to be less severe in the

ongoing Omicron wave, with many affected individuals remaining asymptomatic and often not

seeking clinical testing, thereby rendering traditional surveillance systems less reliable.

Furthermore, increased adoption of at-home testing results in systematic underreporting and less

efficient surveillance. Thus, as the pandemic shifts, relying solely on clinical testing may be

insufficient for timely detection of outbreaks and emerging variants. To circumvent these

challenges, there is an urgent need to incorporate alternative testing strategies to detect low,

circulating levels of SARS-CoV-2 variants.

Wastewater surveillance of SARS-CoV-2 is one such approach that can be used to track

transmission and spread of new and existing variants across communities (3–7). Although,

wastewater-based epidemiology (WBE) has traditionally been used for surveillance of a variety

of targets, including poliovirus (8), antimicrobial resistance genes (9) and monitoring illicit drug-

use (10), WBE has recently gained traction as a promising epidemiological tool to track SARS-

CoV-2 infections. The Centers for Disease Control and Prevention (CDC) developed a national

database - the National Wastewater Surveillance System (NWSS) - in late 2020 to monitor the

spread of COVID-19 in the United States (CDC, 2020). WBE of SARS-CoV-2 complements

existing clinical surveillance as it can capture both symptomatic and asymptomatic cases. As

SARS-CoV-2 is known to be shed in human feces post-infection, usually well before symptoms

appear, the viral loads in sewersheds provide a good indicator of SARS-CoV-2 case burden and

transmission. Previous studies have found strong correlation between SARS-CoV-2

concentrations in wastewater and reported COVID-19 cases (11–14).

WBE of SARS-CoV-2 is useful in monitoring infections when clinical testing is limited,

expensive or overwhelmed due to case surges, especially during the onset of a new variant. It is

also not subject to biases related to health care seeking behavior and willingness to be tested.

Previous work has shown that WBE could be essential to conducting long-term, population-wide

SARS-CoV-2 surveillance in a rapid and cost-effective manner (15). Importantly, WBE can

provide an early warning of rising COVID-19 infections in communities (16,17) and could be an

important tool for guiding public health interventions in combating the ongoing COVID-19

pandemic.

Despite the advantages of WBE for SARS-CoV-2 surveillance, fewer studies implement

genomic analysis of wastewater samples. A wastewater sample represents pooled genomic

content from multiple individuals and presents a mix of different viral lineages. Additionally,

wastewater samples are often degraded and may contain fragmented viral genomes (18). Thus, it

is often challenging to confidently assign lineages in a wastewater sample and assess viral

genetic diversity. Consequently, early research efforts have focused mainly on measuring virus

titers, usually via qPCR, to track trends in COVID-19 infection (3,12,13) or digital droplet PCR

(ddPCR) to characterize variants of interest (VOI) and VOCs (19,20). However, PCR-based

approaches are limited in their application due to mutations being shared across VOI/VOCs and

ongoing evolution of SARS-CoV-2 virus leading to the occurrence of novel mutations. It also

requires continual updating and validation of existing methods to avoid primer dropouts. As

ddPCR targets signature mutations, it may miss amplification of other possible variants which

may be epidemiologically relevant. To circumvent these issues, recent studies have adapted the

use of next generation sequencing approaches to comprehensively scan the entire SARS-CoV-2

genome for potential mutations (21–24). This enables detection of unique and functionally

important mutations and VOC-defining clusters of mutations. Novel computational approaches

are also being developed to parse the different viral lineages in wastewater samples (21,25,26).

Genomic sequencing of SARS-CoV-2 in wastewater offers a universal and powerful tool to

assess the prevalence of circulating SARS-CoV-2 variants and help detect new emerging variants

in a population.

In this study, we sequenced wastewater samples collected from 32 sewersheds across

Utah. Our goal was to track SARS-CoV-2 spread by characterizing lineages and mutations

detected in wastewater communities. We also examined whether lineages detected in wastewater

were comparable to lineage diversity observed in clinical samples. We collaborated with the

Centre for Genomic Pathogen Surveillance to develop an interactive Micoreact dashboard (27) to

visualize our wastewater sequencing data and inform local public health investigations. This

internal dashboard provides an intuitive summary of wastewater sequencing results to our state

epidemiologists and helps in keeping track of various SARS-CoV-2 mutations and associated

lineages over time and across various sampling sites.

2. Materials and methods

2.1. Sample collection

Sewage samples were collected twice per week at 32 wastewater facilities between November

2021 and March 2022 (n = 1,235; Fig. 1; Supplementary Table S1). All samples were raw

influent wastewater collected from municipal treatment systems. Estimated resident populations

varied widely, from a low of 1,322 to a high of 515,494. While the specific type of sample

collected was dependent on sampling equipment available at each partner facility, all but three

facilities provided nominally 24-hour time- or flow-weighted composite samples (mean duration

of 23.9 hours, range of 16 - 28.3 hours). The remaining three facilities provided either grab

samples (Roosevelt City SD) or 6-hour manual composite samples (Snyderville Basin East

Canyon and Snyderville Basin Silver Creek; four grab samples collected every two hours over a

total of six hours then mixed). Samples were stored in a refrigerator or on ice and transported to

the Utah Public Health Laboratory (UPHL) within 24 hours.

2.2. Sample processing and qPCR testing

20mL of raw influent wastewater sample was processed following the steps outlined in

the Promega Maxwell® RSC Enviro Total Nucleic Acid Kit (Promega Corp.). Briefly, the

samples were lysed, the solids were separated, and the samples were buffered. Total nucleic

acids were captured on the Promega PureYield™ Midicolumn, purified and eluted to 600 µL.

The 600 µL concentrate was then extracted by Promega Maxwell RSC 48 automated extraction

and stored at -80°C until RT-qPCR analysis. We conducted RT-qPCR using TaqPath mastermix

and the primers and probes contained within the 2019-nCoV CDC EUA kit (28) to detect the

presence of the targeted gene sequences (N1 and N2) on a ThermoScientific ABI 7500 Fast

instrument. We quantified SARS-CoV-2 RNA against a 5-point standard curve made using serial

dilutions of the CDC EUA 2019-nCoV positive control with N1 and N2 analyzed in

combination.

2.3. Whole-genome Sequencing

We sequenced the SARS-CoV-2 genome using amplicon-based sequencing following the

Illumina COVIDSeq™ RUO protocol (Illumina, Inc. San Diego, CA, USA). Library preparation

was automated via Tecan liquid handlers and pooled, amplified libraries were loaded onto the

S4-flow cell following NovaSeq workflow as per manufacturer's instructions (Illumina Inc).

Dual indexed single-end sequencing with a 75bp read length was performed on the NovaSeq

6000 platform.

2.4. Bioinformatics Analysis

After sequencing, we generated FASTQ files from the raw data using the DRAGEN

COVIDSeq Test Pipeline (Illumina Inc.) on the Illumina DRAGEN Bio-IT platform.

We used viralrecon (v.2.4.1) to analyze the sequencing data and detect SARS-CoV-2 variants

(29). Briefly, we performed data QC, trim and quality filtering on raw FASTQ files using fastp.

We discarded reads with an average Phred quality score below 30 and length shorter than 50 bp.

We mapped the trimmed, human-filtered reads to the SARS-CoV-2 (Wuhan-Hu-1 reference,

RefSeq ID: NC_045512.2) genome using bowtie (v.2.4.4) (30). We soft-clipped the ARTIC

primers using iVar (v.1.3.1) (31) using default settings. Subsequently, we performed variant

calling using iVar and generated consensus sequences using BCFtools (v.1.14) (32). We retained

variants with a minimum base quality of 20, allele frequency of 0.25 and coverage of 10 reads.

We annotated variants with SnpEff (v.5.0e) (33) and SnpSift (v.4.3) (34). All filtered variants

were then compared against a manually curated list of known and epidemiologically important

SARS-CoV-2 variants (including VOCs and VOIs) built using the 'outbreakinfo' R package

(35).

We used Pangolin COVID-19 Lineage Assignment tool (v.4.0) to detect potential SARS-

CoV-2 lineages from consensus sequences (36). We obtained a summary of raw reads, sample

QC, alignment and variant calling metrics via MultiQC (v.1.11) (37). Unlike a clinical sample,

each wastewater sample can contain multiple different SARS-CoV-2 virus lineages shed by

many infected individuals. We used Freyja (v1.3.8) (25) to delineate and ascertain the relative

proportion of different SARS-CoV-2 lineages in each wastewater sample.

3. Results

3.1. Viral RNA concentrations

Viral RNA concentrations varied widely over the study period from non-detect to a

maximum of over 22,000 copies/mL. The maximum concentrations collected from partner

facilities also varied by over an order of magnitude. The highest concentrations were observed at

the height of the Omicron wave (mid to late January 2022), while the lowest concentrations were

observed shortly thereafter in mid to late March 2022 (Supplementary Fig. S2).

3.2. SARS-CoV-2 lineage diversity in wastewater *vs* clinical samples

Of the 1,235 samples collected from 32 sewersheds between November 2021 and March 2022, 1,067 had sufficient RNA for sequencing. The DRAGEN COVIDSeq pipeline successfully generated FASTQ files for 1,034 of those samples. We obtained a total of 9.86 ± 6.08 million raw reads and retained an average of 8.75 ± 5.37 million reads after quality control and read trimming. We found that on average $62.57 \pm 25.45\%$ of total reads mapped to the SARS-CoV-2 reference genome, and covered an average $87.37 \pm 18.94\%$ of the SARS-CoV-2 genome at 1X across all wastewater samples (Supplementary Fig. S3 and Fig. S4).

We detected a total of 70 different lineages (including sub-lineages and with frequency > 5) from 1,034 consensus sequences following PANGO nomenclature. Analysis of diversity of SARS-CoV-2 lineages revealed Delta as the most frequently detected lineage in our dataset (frequency, n= 1,621, 67.71%) during November, prior to the emergence of Omicron (Fig. 3a). However, we first detected Omicron (B.1.1.529) in a wastewater sample collected on November 19th, 2021 from Hyrum City WWTP (HCWWTP21) at an abundance of 0.1% (Fig. 3b). The first clinical case of Omicron in Utah was reported on November 24th, 2021 with a specimen collection date of November 18th, 2021, though it was travel-related. Two additional clinical cases of Omicron were reported on November 29th, 2021 but community transmission of Omicron in Utah was observed around mid-December (Fig. 3c-d).

We observed a steady decrease in the proportion of the Delta (B.1.617.2) lineages and the onset of Omicron (B.1.1529) and its sub-lineage BA.1 (n=112, 6.79%) in wastewater by December. A simple timeline of the detection of Omicron lineages is shown in Fig. 2. We found BA.1.9 (lineage abundance: 8.35%) from Snyderville Basin Silver Creek (SBWRDSC26) on December 7th, 2021, followed by detection of BA.1.17 (lineage abundance: 4.60%) and

BA.1.1.11 (lineage abundance: 27.93%) from Central Valley WRF (CVWRF13) on December

10th (Fig. 4). By December 14th, Omicron spread to two other communities, Snyderville Basin

East Canyon (SBWRDEC27; lineages BA.1.1: 15.8%, BA.1.9: 9.11%) and Brigham City

(BCSD20; lineage BA.1.9: 9.82%), along with increased prevalence in Snyderville Basin Silver

Creek (SBWRDSC26; lineages B.1.1.529: 9.41%, BA.1.9: 52.75% and BA.1.1.16: 14.58%).

Over the next two weeks, several BA.1 sub-lineages of Omicron had spread to nearly all

communities (28/32), including the then common BA.1.15 (Fig. 4). By January 4th, prevalence

of Omicron in wastewater samples had reached 57.58%, and it completely displaced Delta by

February 7th, 2022 (Fig. 3a). Between January and March, Omicron accounted for 95% of total

lineages in circulation (Fig. 3b), with BA.1 being the most common (n=3,037, 82.93%),

followed by BA.2 (n=156, 4.26%) and BA.3 (n=7, 0.19%). Overall, we detected a comparatively

rapid shift from the earlier predominant Delta lineage to Omicron in our wastewater data.

Our wastewater genomic surveillance also identified changes in abundance of other

Omicron sub-lineages over time. We first detected the Omicron sub-lineage BA.2 in our

wastewater sequencing data from Ash Creek SSD (ACSSD32) on January 21st, 2022 (lineage

abundances: BA.2.3, 11.28% and BA.2, 1.79%; Fig. 4). By March 29th, the proportion of BA.2

reached about 45% including its sub-lineages BA.2.1, BA.2.2, BA.2.3, BA.2.4, BA.2.5, BA.2.6,

BA.2.7 and BA.2.8. On February 22nd, 2022, we first detected Omicron sub-lineage BA.3

(lineage abundance: 6.69%) in Orem WRF (OWRF7). Interestingly, Omicron BA.3 persisted at

low abundance in wastewater samples through March, but the BA.3 sub-lineage has not yet been

identified in clinical surveillance. The patterns observed in wastewater sequencing data were

reflected in clinical sequencing (Fig. 3c) and resulted in a huge surge in Omicron cases in

December-February (Supplementary Fig. S5b). The increased transmission and spike in clinical

cases due to Omicron was also evident in wastewater RNA viral loads (Supplementary Fig. S5a)

and later also observed in our analysis of the proportion of mutations associated with the

Omicron lineage.

3.3. Mutational spectra of SARS-CoV-2 in wastewater

We found S:D614G, ORF1ab:T3255I, S:T478K, S:N969K and E:T9I as the most

common mutations among all wastewater samples (Supplementary Fig. S6). All five mutations

were associated with known VOCs. Notably, all wastewater samples prior to the onset of

Omicron (during November 2021) had signature mutations linked to the predominant Delta

variant (e.g., S:P681R, S:D950N, T19R), followed by an increase in frequency of mutations

shared between Delta and Omicron lineages (e.g., S:L452R, S:T478K; Supplementary Fig. S7).

An in-depth analysis of mutations from wastewater samples revealed an increase in

frequency of several Omicron specific mutations (e.g., ORF1ab:T3255I, S:T95I, S:A67V,

S:N856K, S:G142D; allele frequency >0.25) in late December 2021 to January 2022 (Fig. 5).

Genomic sequencing for wastewater samples collected from Roosevelt City (ROOSCD30) on

November 2nd, 2021 confirmed the presence of five signature Omicron mutations (E:T9I,

S:G339D, S:371L, S:373P, S:375P). We also identified two Spike protein mutations, A67V and

T95I (allele frequency 1.00), specific to the Omicron lineage BA.1 from Provo City (PCWRF8)

on November 30th, 2021. Later, we identified six signature mutations of Omicron

(ORF1ab:T3255I, S:T95I, S:G142D, S:A67V, S:N856K, S:D796Y) in Snyderville Basin Silver

Creek (SBWRDSC26) sampled on December 7th, 2021, supporting the assignment of Omicron

BA.1 in our lineage analysis (Fig. 4). We also identified 25 signature mutations of Omicron from

Central Valley WRF (CVWRF13) sampled on December 10th, 2021, aligning with its lineage

assignment as Omicron BA.1 (Fig. 5).

4. Discussion

As the SARS-CoV-2 virus continues to spread and evolve rapidly to form new variants with increased virulence or immune escape potential, it is important to monitor and characterize VOI/VOCs. This is especially important now when pandemic fatigue has become widespread, clinical testing is reduced and more at-home tests are utilized. These factors all contribute towards delays in detecting new emerging lineages when VOCs may be already circulating in communities. In such scenarios, wastewater genomic surveillance could be essential to capture community transmission of COVID in a rapid, cost-effective and unbiased manner. It can provide a more comprehensive picture of SARS-CoV-2 genomic diversity at the community level than individual level data obtained by clinical sequencing. This will allow timely detection of newly emerging lineages and support public health efforts in mitigating COVID-19 spread (38).

Our wastewater sequencing data provides a first detailed view of the diversity of SARS-CoV-2 lineages circulating between November 2021 and March 2022 in Utah, encompassing the displacement of Delta by Omicron lineages. The trends in SARS-CoV-2 RNA levels across sewersheds accurately captured the surge in clinical cases among Utah's population (Supplementary Fig. S5). Notably, we found that the prevalence and diversity of SARS-CoV-2 lineages derived from wastewater samples closely aligned with the lineage proportions derived from clinical genomic surveillance. Sequencing of both wastewater samples and clinical samples between December and January revealed a notable increase in the prevalence of Omicron BA.1 lineages and corresponding decrease in the prevalence of Delta lineages (Fig. 3). Omicron BA.1 was the predominant lineage from January through early March, which aligned with its global surge in cases.

One of the main advantages of wastewater surveillance is that it can be used for early

detection of emerging lineages and help implement public health interventions to limit variant

spread (16,17). In our study, comparing the collection date of Omicron-positive samples from

wastewater with the collection dates of samples from clinical genomic surveillance, we detected

Omicron lineages in wastewater genomic surveillance on November 19th, 2021, up to 10 days

before corresponding detections through genomic clinical surveillance. Notably, the first clinical

case was detected in Utah on November 18th, 2021 but was a result of someone with

international travel history to a region with high community transmission of Omicron. Two

additional clinical cases were identified on November 29th, 2021. Subsequent increases in

abundance of Omicron lineages in wastewater on December 7th, 2021 were followed by an

increase in the number of clinical cases by December 14th, supporting the week ahead early

warning signal detected in wastewater data. The lead time between the detection of a new lineage

in wastewater compared to clinical surveillance is still difficult to estimate in advance and has

been shown to vary from 5-14 days (12,16,17,39). The lead time can potentially be influenced by

environmental factors– temperature, seasons (40), biological factors such as the shedding rates

associated with a lineage (41), severity of infection and whether an individual is symptomatic or

asymptomatic (39) when contributing to a particular catchment.

In our study, analysis of wastewater samples by Freyja, which provides relative

abundance of multiple lineages in a sample, we first detected Omicron (B.1.1.529) from a Hyrum

City (HCWWTP) sample collected on November 19th, 2021. However, based on Pangolin

lineage assignment on consensus sequence, we first detected the Omicron lineage (BA.1.1) on

December 13th, 2021 in Snyderville Basin Silver Creek. This highlights the uncertainty in

lineage assignment based on consensus sequences derived from wastewater samples. Pangolin

lineage assignment tools are perhaps better at identifying the predominant viral lineage in a

community but unlikely to represent the true lineage diversity as is expected in a wastewater

sample, an important consideration for future studies.

Throughout the duration of our study encompassing the peak transmission of Omicron,

wastewater sequencing identified three sub-lineages of Omicron: BA.1, BA.2 and BA.3.

Omicron sub-lineage BA.1 was detected more frequently during our study compared to BA.2

across all sampled communities. By the end of March, as Omicron spread progressed, we

observed an increased prevalence of BA.2 (45% prevalence on March 29th), similar to its

prevalence in other states in the US. While BA.2 was still dominant across communities, we first

detected BA.3 from Orem WRF on February 22nd, 2022. Surprisingly, BA.3 has not been

detected in clinical surveillance in Utah. This could possibly be due to its low prevalence as was

observed in our wastewater data (Fig. 3b) and has been observed throughout the US (<0.5%

prevalence, last checked Sep 19, 2022 on outbreak.info; (42)), suggesting its cryptic circulation

in Utah's wastewater communities. These findings are in agreement with previous studies that

highlight the utility of wastewater surveillance in detection of cryptic lineages – novel SARS-

CoV-2 lineages that are rarely observed in humans (4,25).

Interestingly, the shift from Delta to Omicron was also quite clear from SARS-CoV-2

mutations in wastewater. We found predominance of Delta mutations in November-early

December, followed by an increase in the number of mutations common between Delta and

Omicron lineages and then an increase in Omicron defining mutations (Fig. 4 and Supplementary

Fig. S7). Thus, our ongoing wastewater genomic surveillance allowed us to detect Omicron

lineage even when based on its signature mutations (e.g., S:A67V, S:N856K, S:N969K. E:T9I).

Our study suggests that early detection of a lineage is feasible based on signature mutations,

however, we caution against the interpretation of lineages based on individual mutations,

especially during the early stages of arrival of a new lineage. This is because frequencies of

signature mutations associated with a new lineage are generally low, making it difficult to filter

out the noise from the actual signal in the sequencing data.

Additionally, confident assignment of lineages for a wastewater sample requires

availability of a comprehensive and well-curated lineage database. Novel lineages that have

recently diverged from its ancestral lineage and not yet been curated in the SARS-CoV-2 lineage

databases (e.g. Pangolin) will be challenging to identify via wastewater sequencing. However,

analysis of key mutations from a clinically important region of the SARS-CoV-2 genome (e.g.

Spike gene) that can influence transmissibility, immune or vaccine escape could help us identify

a potential new lineage in wastewater. Recent studies have reported the presence of several rare,

cryptic lineages in wastewater – SARS-CoV-2 lineages that did not match to any known lineages

and had not been reported in humans (4,43). These novel lineages contained several rare

mutations that had not been identified in routine clinical surveillance and shared many mutations

with the circulating VOCs. Routine monitoring of such mutations of concern from wastewater

communities could reveal new SARS-CoV-2 lineages before they become epidemiologically

relevant and spread to the wider human population, thus guiding public health decision-making.

5. Conclusion

Our study illustrates that genomic sequencing of wastewater samples can be used to

detect, monitor and provide insights into the diversity of SARS-CoV-2 variants. Of note is the

cost-effectiveness of wastewater sequencing as a pooled community sample can be analyzed to

monitor disease spread. Wastewater-based genomic epidemiology can be used to track the

prevalence of SARS-CoV-2 variants in near real-time, often earlier than information obtained by

clinical sample sequencing. Early detection of emerging lineages can provide the time needed to

allocate more testing resources to communities showing early detection of an emerging variant

and implement public health interventions to limit variant spread. As we move beyond Omicron

and new variants emerge, wastewater-based surveillance will be a valuable tool in the timely

identification of novel lineages, complementing clinical sequencing and aiding in public health

decision-making. While wastewater surveillance has proven to be an essential tool for SARS-

CoV-2 genomic surveillance, it would be worthwhile expanding its application to detect other

emerging infections and aid surveillance of future emerging infectious diseases.

Acknowledgments

This work was supported by the Centers for Disease Control and Prevention's (CDC)

Epidemiology and Laboratory Capacity for Prevention and Control of Emerging Infectious

Diseases (ELC) Enhanced Detection Cooperative Agreement. We thank our partner wastewater

treatment facilities across the State of Utah who collected and sent us the wastewater samples.

We also thank the UPHL NGS Sequencing Team for conducting the sequencing of wastewater

samples.

Conflicts of Interest

All authors declare that they have no conflicts of interest.

References

1. Cameroni E, Bowen JE, Rosen LE, Saliba C, Zepeda SK, Culap K, et al. Broadly neutralizing

antibodies overcome SARS-CoV-2 Omicron antigenic shift. Nature. 2022 Feb

24;602(7898):664–70.

2. Plante JA, Mitchell BM, Plante KS, Debbink K, Weaver SC, Menachery VD. The variant

gambit: COVID-19's next move. Cell Host Microbe. 2021 Apr 14;29(4):508–15.

3. Kumar M, Patel AK, Shah AV, Raval J, Rajpara N, Joshi M, et al. First proof of the capability

of wastewater surveillance for COVID-19 in India through detection of genetic material of

SARS-CoV-2. Sci Total Environ. 2020 Dec 1;746:141326.

4. Smyth DS, Trujillo M, Gregory DA, Cheung K, Gao A, Graham M, et al. Tracking cryptic

SARS-CoV-2 lineages detected in NYC wastewater. Nat Commun. 2022 Dec;13(1):635.

5. Ahmed W, Angel N, Edson J, Bibby K, Bivins A, O'Brien JW, et al. First confirmed detection

of SARS-CoV-2 in untreated wastewater in Australia: A proof of concept for the wastewater

surveillance of COVID-19 in the community. Sci Total Environ. 2020 Aug 1;728:138764.

6. Haramoto E, Malla B, Thakali O, Kitajima M. First environmental surveillance for the

presence of SARS-CoV-2 RNA in wastewater and river water in Japan. Sci Total Environ.

2020 Oct 1;737:140405.

7. Rothman JA, Loveless TB, Kapcia J, Adams ED, Steele JA, Zimmer-Faust AG, et al. RNA

Viromics of Southern California Wastewater and Detection of SARS-CoV-2 Single-

Nucleotide Variants. Nojiri H, editor. Appl Environ Microbiol. 2021 Nov 10;87(23):e01448-

21.

8. Asghar H, Diop OM, Weldegebriel G, Malik F, Shetty S, El Bassioni L, et al. Environmental

Surveillance for Polioviruses in the Global Polio Eradication Initiative. J Infect Dis. 2014 Nov

1;210(suppl_1):S294–303.

9. The Global Sewage Surveillance project consortium, Hendriksen RS, Munk P, Njage P, van

Bunnik B, McNally L, et al. Global monitoring of antimicrobial resistance based on

metagenomics analyses of urban sewage. Nat Commun. 2019 Dec;10(1):1124.

10. González□Mariño I, Baz□Lomba JA, Alygizakis NA, Andrés□Costa MJ, Bade R,

Bannwarth A, et al. Spatio□temporal assessment of illicit drug use at large scale: evidence

from 7 years of international wastewater monitoring. Addiction. 2020 Jan;115(1):109–20.

11. Peccia J, Zulli A, Brackney DE, Grubaugh ND, Kaplan EH, Casanovas-Massana A, et al.

Measurement of SARS-CoV-2 RNA in wastewater tracks community infection dynamics. Nat

Biotechnol. 2020 Oct;38(10):1164–7.

12. Medema G, Heijnen L, Elsinga G, Italiaander R, Brouwer A. Presence of SARS-

Coronavirus-2 RNA in Sewage and Correlation with Reported COVID-19 Prevalence in the

Early Stage of the Epidemic in The Netherlands. Environ Sci Technol Lett. 2020 Jul

14;7(7):511–6.

13. Acosta N, Bautista MA, Waddell BJ, McCalder J, Beaudet AB, Man L, et al. Longitudinal

SARS-CoV-2 RNA wastewater monitoring across a range of scales correlates with total and

regional COVID-19 burden in a well-defined urban population. Water Res. 2022

Jul;220:118611.

14. Weidhaas J, Aanderud ZT, Roper DK, VanDerslice J, Gaddis EB, Ostermiller J, et al.

Correlation of SARS-CoV-2 RNA in wastewater with COVID-19 disease burden in

sewersheds. Sci Total Environ. 2021 Jun;775:145790.

15. Shrestha S, Yoshinaga E, Chapagain SK, Mohan G, Gasparatos A, Fukushi K. Wastewater-

Based Epidemiology for Cost-Effective Mass Surveillance of COVID-19 in Low- and

Middle-Income Countries: Challenges and Opportunities. Water. 2021 Oct 15;13(20):2897.

16. Ahmed W, Tscharke B, Bertsch PM, Bibby K, Bivins A, Choi P, et al. SARS-CoV-2 RNA

monitoring in wastewater as a potential early warning system for COVID-19 transmission in

the community: A temporal case study. Sci Total Environ. 2021 Mar;761:144216.

17. Karthikeyan S, Ronquillo N, Belda-Ferre P, Alvarado D, Javidi T, Longhurst CA, et al.

High-Throughput Wastewater SARS-CoV-2 Detection Enables Forecasting of Community

Infection Dynamics in San Diego County. Cristea IM, editor. mSystems [Internet]. 2021 Apr 27 [cited 2021 Aug 5];6(2). Available from: https://journals.asm.org/doi/10.1128/mSystems.00045-21

18. Wurtzer S, Waldman P, Ferrier-Rembert A, Frenois-Veyrat G, Mouchel JM, Boni M, et al. Several forms of SARS-CoV-2 RNA can be detected in wastewaters: Implication for wastewater-based epidemiology and risk assessment. Water Res. 2021 Jun;198:117183.

19. Wolfe M, Hughes B, Duong D, Chan-Herur V, Wigginton KR, White BJ, et al. Detection of SARS-CoV-2 Variants Mu, Beta, Gamma, Lambda, Delta, Alpha, and Omicron in Wastewater Settled Solids Using Mutation-Specific Assays Is Associated with Regional Detection of Variants in Clinical Samples. Buan NR, editor. Appl Environ Microbiol. 2022 Apr 26;88(8):e00045-22.

20. Heijnen L, Elsinga G, Graaf M de, Molenkamp R, Koopmans MPG, Medema G. Droplet digital RT-PCR to detect SARS-CoV-2 signature mutations of variants of concern in wastewater. Sci Total Environ. 2021;799:149456.

21. Amman F, Markt R, Endler L, Hupfauf S, Agerer B, Schedl A, et al. Viral variant-resolved wastewater surveillance of SARS-CoV-2 at national scale. Nat Biotechnol [Internet]. 2022 Jul 18 [cited 2022 Sep 1]; Available from: https://www.nature.com/articles/s41587-022-01387-y

22. Resende PC, Motta FC, Roy S, Appolinario L, Fabri A, Xavier J, et al. SARS-CoV-2 genomes recovered by long amplicon tiling multiplex approach using nanopore sequencing and applicable to other sequencing platforms [Internet]. Molecular Biology; 2020 May [cited 2021 Aug 5]. Available from: http://biorxiv.org/lookup/doi/10.1101/2020.04.30.069039

23. Crits-Christoph A, Kantor RS, Olm MR, Whitney ON, Al-Shayeb B, Lou YC, et al. Genome Sequencing of Sewage Detects Regionally Prevalent SARS-CoV-2 Variants. 2021;12(1):9.

24. Rehn A, Braun P, Knüpfer M, Wölfel R, Antwerpen MH, Walter MC. Catching SARS-CoV-

2 by Sequence Hybridization: a Comparative Analysis. mSystems. 0(0):e00392-21.

25. Karthikeyan S, Levy JI, De Hoff P, Humphrey G, Birmingham A, Jepsen K, et al.

Wastewater sequencing reveals early cryptic SARS-CoV-2 variant transmission. Nature. 2022

Sep 1;609(7925):101–8.

26. Baaijens JA, Zulli A, Ott IM, Nika I, van der Lugt MJ, Petrone ME, et al. Lineage abundance

estimation for SARS-CoV-2 in wastewater using transcriptome quantification techniques.

Genome Biol. 2022 Nov 8;23(1):236.

27. Argimón S, Abudahab K, Goater RJE, Fedosejev A, Bhai J, Glasner C, et al. Microreact:

visualizing and sharing data for genomic epidemiology and phylogeography. Microb

Genomics [Internet]. 2016 Nov 30 [cited 2022 Nov 30];2(11). Available from:

https://www.microbiologyresearch.org/content/journal/mgen/10.1099/mgen.0.000093

28. Centers for Disease Control and Prevention. CDC 2019-novel coronavirus (2019-nCoV) real-

time RT-PCR diagnostic panel. 2021;

29. Patel H, Varona S, Monzon S, Espinosa-Carrasco; J, L Heuer M, Underwood A, et al. nf-

core/viralrecon: nf-core/viralrecon v2.4.1 [Internet]. 2022. (Zenodo). Available from:

https://zenodo.org/record/6320980#.Y3a7sezMLJx

30. Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient alignment of

short DNA sequences to the human genome. Genome Biol. 2009;10(3):R25.

31. Grubaugh ND, Gangavarapu K, Quick J, Matteson NL, De Jesus JG, Main BJ, et al. An

amplicon-based sequencing framework for accurately measuring intrahost virus diversity

using PrimalSeq and iVar. Genome Biol. 2019 Dec;20(1):8.

32. Li H. A statistical framework for SNP calling, mutation discovery, association mapping and

population genetical parameter estimation from sequencing data. Bioinformatics. 2011 Nov

1;27(21):2987–93.

33. Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, et al. A program for

annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the

genome of Drosophila melanogaster strain w $^{1118}\square$; iso-2; iso-3. Fly (Austin). 2012

Apr;6(2):80–92.

34. Cingolani P, Patel VM, Coon M, Nguyen T, Land SJ, Ruden DM, et al. Using Drosophila

melanogaster as a Model for Genotoxic Chemical Mutational Studies with a New Program,

SnpSift. Front Genet [Internet]. 2012 [cited 2022 Oct 7];3. Available from:

http://journal.frontiersin.org/article/10.3389/fgene.2012.00035/abstract

35. Alkuzweny M, Hughes L, Gangavarapu K. outbreakinfo: outbreak.info R Client [Internet].

2022. Available from: https://outbreak-info.github.io/R-outbreak-info/

36. O'Toole Á, Scher E, Underwood A, Jackson B, Hill V, McCrone JT, et al. Assignment of

Epidemiological Lineages in an Emerging Pandemic Using the Pangolin Tool. Virus Evol.

2021 Jul 5;veab064.

37. Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: summarize analysis results for

multiple tools and samples in a single report. Bioinformatics. 2016 Oct 1;32(19):3047–8.

38. Kirby AE, Walters MS, Jennings WC, Fugitt R, LaCross N, Mattioli M, et al. Using

Wastewater Surveillance Data to Support the COVID-19 Response — United States, 2020–

2021. MMWR Morb Mortal Wkly Rep. 2021 Sep 10;70(36):1242–4.

39. D'Aoust PM, Graber TE, Mercier E, Montpetit D, Alexandrov I, Neault N, et al. Catching a

resurgence: Increase in SARS-CoV-2 viral RNA identified in wastewater 48 h before COVID-

19 clinical tests and 96 h before hospitalizations. Sci Total Environ. 2021 May;770:145319.

40. Kumar M, Jiang G, Kumar Thakur A, Chatterjee S, Bhattacharya T, Mohapatra S, et al. Lead

time of early warning by wastewater surveillance for COVID-19: Geographical variations and

impacting factors. Chem Eng J. 2022 Aug;441:135936.

41. Singanayagam A, Hakki S, Dunning J, Madon KJ, Crone MA, Koycheva A, et al.

Community transmission and viral load kinetics of the SARS-CoV-2 delta (B.1.617.2) variant

in vaccinated and unvaccinated individuals in the UK: a prospective, longitudinal, cohort

study. Lancet Infect Dis. 2022 Feb;22(2):183–95.

42. Hughes L, Gangavarapu K, Latif AA, Mullen J, Alkuzweny M, Hufbauer E, et al.

Outbreak.info genomic reports: scalable and dynamic surveillance of SARS-CoV-2 variants

and mutations [Internet]. In Review; 2022 Jun [cited 2022 Oct 4]. Available from:

https://www.researchsquare.com/article/rs-1723829/v1

43. Gregory DA, Trujillo M, Rushford C, Flury A, Kannoly S, San KM, et al. Genetic Diversity

and Evolutionary Convergence of Cryptic SARS-CoV-2 Lineages Detected Via Wastewater

Sequencing [Internet]. 2022 Jun [cited 2022 Oct 3]. Available from:

http://medrxiv.org/lookup/doi/10.1101/2022.06.03.22275961

Figures

Fig. 1. Geographic map showing the 32 sampled wastewater treatment sites in Utah.

Fig. 2. Timeline of the important events in the detection of lineages and key mutations of

Omicron from wastewater sequencing.

Fig. 3. Proportion of all Sars-CoV-2 lineages in a) wastewater samples, c) clinical samples and

proportion of Omicron lineages in b) wastewater samples, d) clinical samples from November

2021 to March 2022.

Fig. 4. Abundance of Omicron lineages and its sub-lineages in each sewershed between

December 2021 and January 2022 to highlight the introduction and spread of Omicron in Utah.

Circles are color coded by sewersheds and the size of each circle corresponds to the relative

abundance of lineages.

491   Fig. 5. Heatmap showing the temporal changes in the frequency of SARS-CoV-2 amino acid

492   changes in the Spike and ORF1ab genes, detected in each sewershed from November 2021 to

493   March 2022, encompassing the onset of Omicron. Amino acid changes that were detected only

494   once during the study period are not shown. Colors in the heatmap range from cool yellows/reds

495   representing low frequency to warm blues/violets representing high frequency.