1   Suppression of classical renin-angiotensin system and increase of soluble ACE2 in pregnancies

2   with adverse outcomes

4   [1]Robin Shoemaker, [2]Marko Poglitsch, [3]Hong Huang, [4]Katherine Vignes, [4]Aarthi Srinivasan,

5   [4]Cynthia Cockerham, [3]Aric Schadler, [3]John A. Bauer, [4]John M. O'Brien

7   [1] Department of Dietetics and Human Nutrition, University of Kentucky, Lexington KY, USA

8   [2] Attoquant Diagnostics GmbH, Vienna, Austria

9   [3]Department of Pediatrics, University of Kentucky, Lexington KY, USA

10   [4]Division of Maternal Fetal Medicine, Department of Obstetrics and Gynecology, University of

11   Kentucky, Lexington KY, USA

13   Running Title: Classical and alternative RAS in adverse outcomes of pregnancy

First and Corresponding Author:
Robin Shoemaker, PhD
Assistant Professor
University of Kentucky
Department of Dietetics and Human Nutrition
119 Funkhouser Building
Lexington, KY 40306
859-257-1031
Robin.shoemaker@uky.edu

Abstract

Adverse pregnancy outcomes are major health risks to pregnant women and can increase maternal risk for cardiovascular disease. However, clinical risk factors are not always well-correlated with development of adverse pregnancy outcomes. The renin-angiotensin system (RAS) is a hormone system composed of two arms with opposing actions that plays a major role in cardiovascular function in pregnant and non-pregnant individuals. In the current study, we investigated whether the normal activity of the RAS during pregnancy was dysregulated in pregnancies complicated by adverse outcomes. In a retrospective study of 74 women followed prospectively for pregnancy-hypertension and related adverse outcomes, we used a novel mass spectrometry-based methodology to quantify serum concentrations of multiple angiotensin peptides and enzymatic activity of angiotensin-converting enzyme 2 (ACE2) and neprilysin in the first and third trimesters of pregnancy. There were n=46 women meeting the primary endpoint of a composite of adverse outcomes, and n=28 women without pregnancy complications. Serum concentrations of components of the "classical" RAS, those involved in the formation and action of the main bioactive peptide, angiotensin II, were increased during healthy pregnancy, but markedly suppressed in adverse outcome pregnancies. In contrast, components of the counter-regulatory, or "alternative", RAS, such as ACE2, neprilysin, and angiotensin-1-5, were moderately increased in healthy pregnancy but more robustly increased in pregnancies with adverse outcomes. Our results indicate that a shift in the activation of the classical arm to the alternative arm of the RAS portends the development of adverse outcomes of pregnancy.

Keywords: angiotensin, maternal, pregnancy complications, ACE2, ACE2 shedding, biomarkers, mass spectrometry

Introduction

Adverse outcomes of pregnancy, such as pre-eclampsia, intrauterine growth restriction (IUGR), preterm birth, and maternal organ damage, pose major maternal and fetal health risks. Nearly one in five women in the United States experience adverse outcomes of pregnancy[1], which are associated with long-term health risks for both the mother and the offspring. These include cardiovascular diseases, metabolic syndrome, cognitive and developmental impacts, and early mortality[2]. The risk for adverse outcomes is greatly affected by race and ethnicity, socioeconomic factors, and the presence of cardiometabolic and other clinical risk factors[3]. Clinical parameters, such as blood pressure or proteinuria, poorly predict adverse outcomes[4], underscoring an unmet and growing need for knowledge leading to the detection of patients at risk for adverse outcomes of pregnancy.

The renin angiotensin system (RAS) is a master regulator of cardiovascular function and blood pressure control in non-pregnant and pregnant states. Activity of the RAS increases during pregnancy to prompt vascular and hemodynamic changes necessary to accommodate the growing fetoplacental unit. The activity of the RAS is controlled by the rate-limiting enzyme renin, which cleaves a ten amino acid fragment, angiotensin I, from the precursor protein, angiotensinogen. Angiotensin I is then metabolized by angiotensin-converting enzyme (ACE) to generate the primary effector peptide, angiotensin II. The components involved in the formation and subsequent effects of angiotensin II from angiotensin I are collectively termed the "classical" RAS. An appreciable body of literature demonstrates that the classical RAS is consistently upregulated in pregnancy and suppressed in preeclampsia[5-7]. Angiotensin II can be metabolized by angiotensin converting enzyme 2 (ACE2) to generate angiotensin-(1-7)[8]. Effects of angiotensin-(1-7) are reported to oppose that of angiotensin II[9], and the ACE2/angiotensin-(1-7) axis is termed the "alternative" RAS.

Limited and conflicting studies suggest the alternative RAS may also play a role in pregnancy, and that alterations in these components are associated with pre-eclampsia and

small-for-gestational age infants[10-12]. Recent studies demonstrate that soluble ACE2, a

catalytically active ectodomain released into the circulation via proteolysis[13], is a biomarker for

cardiovascular disease[14-16] and elevated with severity of COVID-19 disease[17]. Given the role of

the RAS in the maternal adaptation to pregnancy, we hypothesized that dysregulation of the

RAS and ACE2 could be a molecular link portending complications in pregnancy. In the current

study, we quantified serum components of the RAS at two time points during human pregnancy

using a novel mass spectrometry-based methodology. Our objective was to determine

associations of serum components of the classical and/or alternative RAS with adverse

outcomes of pregnancy.

Methods

Study Population

This is a retrospective analysis of n=74 women enrolled in an ongoing study at the University of

Kentucky. The cohort is comprised of pregnant women meeting the United States Preventative

Services Task Force (USPTFS) criteria for the recommendation of aspirin for the prevention of

preeclampsia[18] (n=38) and a control population considered to be at low-risk for developing pre-

eclampsia (n=36). Inclusion criteria for the former were more than one of the following clinical

risk factors: history of preterm preeclampsia, chronic hypertension, type 1 or 2 diabetes, renal or

autoimmune disease; or more than two of the following clinical risk factors: nulliparity, obesity

(BMI > 30), family history of pre-eclampsia, Black race, lower income, age 35 or older, previous

pregnancy with small birth weight or adverse outcome, or in-vitro conception. Pregnant women

not meeting these criteria were recruited as controls. Exclusion criteria included age less than

18 or greater than 45 years, multifetal gestation, history of allergy to aspirin, gastrointestinal

bleeding, severe peptic ulcer or liver dysfunction, patients on anticoagulant medications, and

women with anomalous fetus. All subjects gave informed consent to participate in the study, and

for the use of samples for this research, which was approved by the University of Kentucky

Institutional Review Board.

Study Design and Outcomes

Pregnant women between 18 and 45 years of age in the first trimester of pregnancy were

recruited at the time of their routine prenatal visits or ultrasound appointments in the first

trimester screening and followed prospectively for pregnancy outcomes. Patients were treated

according to standard clinical guidelines and data collected as part of routine clinical care.

Demographic information was collected upon enrollment. Clinical data and maternal blood were

collected by trained study personnel at the first (11-16 weeks) and third (28-32 weeks)

trimesters of pregnancy and outcomes were recorded at delivery. Blood pressure was

measured following the American Heart Association Guidelines[19], and was determined as the

average of two consecutive measurements per arm assessed in a seated, resting position by

one observer. Doppler investigations of the uterine artery were performed according to the

standards of the Fetal Medicine Foundation at 18-22 weeks gestation and reported as mean

pulsatility index of the left and right arteries[20]. Clinical data and outcomes were obtained from

electronic medical records (EMR), and input into a REDcap database by study personnel.

Maternal blood was collected into red-top serum tubes and rested at room temperature for 60

minutes followed by centrifugation, transfer into 200 uL single-use aliquots, and barcoding.

Samples were stored at -80 C until analysis.

The primary endpoint of the current study was a composite outcome of maternal adverse

outcomes (AO). The composite outcome was met if one of the following AOs was recorded:

gestational hypertension (new onset >20 weeks), pre-eclampsia (early onset, late onset, or

postpartum), intrauterine growth restriction (IUGR, defined as fetal weight below the 10[th]

percentile), postpartum hemorrhage, preterm premature rupture of membranes (PPROM), pre-

term labor or pre-term delivery (<34 weeks), presence of polyhydramnios, or fetal death.

Quantification of angiotensin peptides and angiotensin-based biomarkers in serum.

Equilibrium levels of six different RAS angiotensin peptide metabolites and aldosterone in serum

samples were quantified by LC-MS/MS using previously validated and described methods (RAS

Fingerprint[TM], Attoquant Diagnostics GmbH, Vienna, Austria)[21]. Previous studies demonstrate

controlled sample conditioning followed by analysis of angiotensin peptides under equilibrium

conditions to be highly correlated with circulating peptide profiles, and effective for prediction of

outcomes[22] [23]. Equilibration was performed at 37 C for one hour, followed by stabilization

through addition of an enzyme inhibitor cocktail. Samples were spiked with stable isotope-

labeled internal standards at 200 pg/mL for each angiotensin metabolite. The samples were

subjected to C-18-based solid-phase-extraction followed by LC-MS/MS analysis using a

reversed-phase analytical column operating in line with a Xevo TQ-S triple quadruple mass

spectrometer (Waters). Internal standards were used to correct for peptide recovery of the

sample preparation procedure for each analyte in each individual sample. Analyte

concentrations are reported in pmol/L. The lower limits of quantification (LLOQs) for each

peptide were: angiotensin I: 5 pmol/L; angiotensin II: 4 pmol/L; angiotensin III: 4 pmol/L;

angiotensin IV: 2 pmol/L; angiotensin-(1-5): 2 pmol/L; angiotensin-(1-7): 3 pmol/L, and

aldosterone: 10 pmol/L. Equilibrium-based biomarkers were calculated as previously reported[22]:

renin activity (pmol/L): [Ang I + Ang I] and ACE activity (pmol/L/pmol/L): [Ang II / Ang I]. For

values below the LLOQ, means for each analyte and calculation of equilibrium-based

biomarkers were calculated using half the LLOQ[22].

Quantification of active ACE2 and neprilysin concentration

LC-MS/MS based quantification of ACE2 and NEP concentrations, respectively, in serum

samples was performed using natural substrate-based conversion assays, as previously

described[24,25]. Briefly, an excess of substrate (angiotensin I or angiotensin II) was spiked to a

diluted sample in the presence of angiotensin-(1-7)-stabilizing protease inhibitor mix. Following

a one-hour incubation at 37°C, angiotensin I, II, and (1-7) levels were quantified by LC-MS/MS

using internal standardization with stable isotopes. The assay was validated by spiking

recombinant human ACE2 or NEP, respectively, into human serum in the presence and

absence of respective inhibitors (MLN-4760 and thiorphan). The angiotensin-(1-7) formation

rate was calculated in (pmol/L)/h and converted to ng/mL of active enzyme concentration based

on the standard curve obtained from assay calibration.

Statistical Analysis

Bivariate statistical analyses were performed via GraphPad Prism version 9 (San Diego, CA,

USA). Independent continuous variables were analyzed using independent samples t-tests or

independent samples difference of medians as appropriate. Statistical significance of

continuous variables between first and third trimester were analyzed using paired samples *t*-test

or Wilcoxon signed rank test as appropriate. A p-value less than 0.05 was considered

statistically significant.

Results

Subject characteristics. Paired clinical data and biological samples from first and third trimester

visits were obtained for 74 women between April 2019 and May 2020. There were 46 women

(62.2%) that developed AOs, and 28 (37.8%) that did not. Subject demographics and clinical

characteristics are described in Table 1. The cohort was primarily Caucasian, with Black,

Hispanic, and mixed race women distributed fairly equally among those who developed AOs

and those who did not. Women with AO had slightly elevated age, first trimester BMI, systolic

and diastolic blood pressures, and uterine artery pulsatility index (PI) compared to those with no

AO, although none of these parameters were statistically significant. Women who experienced

AOs had significantly shorter gestation (P<0.0001) and smaller infant body weights (P<0.05)

compared to those without (Table 1). There were 31 women with one adverse outcome, 9 with

two adverse outcomes, and 6 with three adverse outcomes. All recorded outcomes for this

group are listed in Table 2.

Serum components of the classical RAS were increased in healthy pregnancies, and

suppressed in pregnancies with AOs. In women with no AOs, components of the classical RAS

(angiotensin I, angiotensin II, and renin activity) robustly increased from first to third trimester of

pregnancy (Table 3). In contrast, only angiotensin I was increased during gestation in

pregnancies with AOs. Further, the change in angiotensin I, angiotensin II, and renin activity

over gestation was lower in pregnancies complicated by AOs (Figure 1A). In addition,

concentrations of angiotensin I and angiotensin II and renin activity were generally lower in

pregnancies complicated by AOs compared to those without, and this was statistically

significant at the third trimester time point (Table 3). Angiotensin IV was marginally, but not

significantly, increased with pregnancy in women with and without AOs, with a trend of lower

concentrations in AO pregnancies. Interestingly, activity of ACE was significantly reduced from

first to third trimester in both groups, with no significant differences between groups at either

time point.

ACE2 and the alternative RAS is increased in pregnancy, and more pronounced in pregnancies

complicated by AOs. ACE2 activity was detected in all patients, and was significantly increased

from the first to the third trimester in patients with and without AOs (Table 3). However, ACE2

activity in the first trimester was significantly lower in AO compared to no AO patients (Table 3;

P<0.05), and the change in ACE2 activity over gestation was significantly greater in patients

with AOs (Figure 1B; P<0.05).

In contrast to the observed rise in ACE2 activity during pregnancy, concentrations of

angiotensin-(1-7) (the product of enzymatic action of ACE2) in serum were not similarly

increased over pregnancy. In fact, median concentrations of angiotensin-(1-7) at both time

points were below the lower limit of quantification (LLOQ) in both AO and non-AO groups.

Angiotensin (1-7) concentrations in non-AO patients were above the LLOQ in only 6 and 13

patients in the first and third trimesters, respectively, and in 11 and 12 patients with AOs in the

first and third trimesters, respectively, and were therefore not reported in Table 3. However,

angiotensin-(1-7) can be cleaved at the c-terminus by ACE to generate angiotensin-(1-5)[20], and

this peptide was detected in all patients. Angiotensin-(1-5) concentrations increased during

gestation, but this was only significant in AO patients (Table 3; P<0.05 in AO, versus p=0.0627

in non-AO patients). In addition to the conversion from angiotensin II by ACE2, angiotensin-(1-7)

can generated from angiotensin I via the enzymatic action of neprilysin[26]. Neprilysin activity was

increased from first to third trimester of pregnancy in both groups (Table 3), with the effect more

pronounced in AO (P<0.001) versus non-AO patients (P<0.05). There was no difference in

neprilysin activity at either time point in AO versus non-AO patients.

Discussion

Detection of adverse outcomes of pregnancy is an emergent and unmet need in the fields of

obstetric and maternal-fetal-medicine. Because these conditions can have long-term effects on

cardiovascular health, detection (and ultimately prevention) of adverse outcomes of pregnancy

is an important strategy for the prevention of cardiovascular disease[2]. In the current study, we

investigated the association of serum markers of the RAS, a hormone system with key roles in

both cardiovascular function and maternal adaptation to pregnancy, with adverse outcomes in a

cohort of pregnant women. This is the first study to perform comprehensive molecular profiling

of the RAS (e.g. classical and alternative arms) at multiple time points during pregnancy via

mass spectrometry. The major findings of this study are: 1) both the classical and alternative

arms of the RAS are robustly increased in healthy pregnancy; 2) pregnancies with adverse

outcomes are characterized by attenuated activation of the classical RAS (both compared to

healthy pregnancy and across gestation), and 3) in contrast, the alternative RAS, and release of

soluble ACE2 into the circulation, is more active in pregnancies with adverse outcomes. These

data demonstrate that alterations in the normal activity of the RAS in pregnancy are associated

with adverse outcomes in pregnancy.

Published literature demonstrate elevated plasma renin activity, angiotensin I and/or

angiotensin II concentrations, either longitudinally throughout healthy pregnancy, or compared

to a non-pregnant population[11,27-32]. These data are derived using plasma renin activity assays

(where radioimmunoassay is used to quantify formation of angiotensin I from angiotensinogen,

and values are reported as concentration per unit time[20]), or other antibody-based capture

techniques for quantification of angiotensin I or II, reported as concentration (ng/mL). Data in

our study are derived via RAS Fingerprint[TM] (developed by Attoquant Diagnostics GmBH) a

novel LC-MS/MS-based method for simultaneous measurement of multiple angiotensin peptides

in blood, which are also used to calculate surrogate biomarkers for renin and ACE activity.

Comparison of this method to classical renin activity and antibody-based methods has been

reported in non-pregnant states [33], and reported a body of literature demonstrates RAS

biomarkers generated using this method are associated with clinical conditions and outcomes

related to cardiovascular and hypertensive diseases[22,33-37]. This is the first study to report direct

quantification of angiotensin peptides via LC-MS/MS at multiple time points of pregnancy. In

agreement with published literature, we report a marked increase in renin activity, angiotensin I,

and II over gestation in healthy pregnancy.

Published literature indicates that activation of the classical RAS observed in healthy

pregnancy is suppressed in women with pre-eclampsia[11,28]. Similarly, we demonstrate that in

pregnancies complicated by adverse outcomes, renin activity and angiotensin II concentrations

do not increase over gestation are reduced compared to healthy pregnancies in the third

trimester. In contrast, limited and conflicting findings exist regarding the activity of the alternative

RAS in pregnancy, and few studies provide a comprehensive assessment of both the classical

and alternative RAS. Merrill et al performed a comprehensive assessment of the RAS in women

in the third trimester of uncomplicated pregnancy compared to a non-pregnant group and a

preeclamptic group[11]. Elevated renin activity and concentrations of angiotensins I, II, and (1-7)

in plasma of pregnant versus non-pregnant women, and a reduction in these parameters with

preeclampsia, were reported. The authors hypothesized that angiotensin-(1-7) may function in

healthy pregnancy to oppose potentially adverse (vasoconstrictive) effects of angiotensin II and

that reductions in angiotensin-(1-7) may contribute to hypertension in preeclampsia, further

supported by the observation that plasma concentrations of angiotensin-(1-7) were negatively

associated with blood pressure. Similarly, in a recent study by Tamanna et al, plasma

angiotensin-(1-7) concentrations were elevated throughout pregnancy compared to a non-

pregnant control group, and reduced in preeclampsia[10]. Compared to normal pregnancy,

plasma ACE2, quantified via ELISA, was elevated in the third trimester in small-for-gestational

age (SGA) pregnancies and reduced in preeclamptic pregnancies.

Findings from the current study agree with those of Merrill et al, where the classical RAS

is increased during pregnancy, and with Tamanna et al. that ACE2 activity is increased during

pregnancy. However, using mass spectrometry, we report limited detection of serum

angiotensin-(1-7) in either the first or third trimester; in samples where angiotensin-(1-7) was

detected, concentrations were significantly lower than those reported via radioimmunoassay. A

possible explanation for discrepancies in findings is methodology for detection of angiotensin

peptides, an analyte posing several analytic challenges. These include extremely short half-lives

(e.g seconds) of peptides, inefficient inhibition of angiotensin-metabolizing enzymes during

sampling, and cross-reactivity in antibody-based assays, (as peptide sequences may only differ

by one or two amino acids) [38]. Notably, reliability of ELISA for quantification of angiotensin-(1-7)

has recently been called into question[39]. However, in the current study, angiotensin-(1-5) was

readily detected. This peptide is generated from angiotensin-(1-7) by the actions of ACE[28], and

is thus considered part of the alternative RAS. Although the absolute concentrations of this

analyte were very low, angiotensin-(1-5) levels were significantly reduced in the third trimester in

pregnancies with adverse outcomes, comparted to healthy pregnancies (median value of 5.6

compared to 7.9 pmol/L).

In a recent case control study, plasma levels of ACE2 in mid-pregnancy were elevated in

296 pregnant women prior to the development of preeclampsia compared to 333 healthy

pregnancies using proximity extension assay technology (Olink PEA CVD-II panel; Uppsala,

Sweden)[14]. Similarly, in the current study, we report that soluble ACE2 is significantly increased

in pregnancy, and that this increase is greater in women who subsequently developed adverse

outcomes. ACE2 is a membrane-bound protein, but conditions of tissue damage or stress (e.g.

heart or renal failure) result in shedding of a soluble portion of the protein, which can be

detected in the urine or circulation[13]. Soluble ACE2 is normally not detected, or found in low

levels in healthy individuals (at concentrations 100-fold lower than that of ACE)[40], but a growing

number epidemiological and clinical studies report patients with cardiovascular disease[15,16,41],

and in some reports of severe COVID-19, exhibit higher circulating levels of soluble ACE2.

However, it is not clear why pregnancy is associated with enhanced ACE2 shedding, or from

what organ(s) ACE2 was shed. A dysfunctional placental RAS has been implicated in

preeclampsia[42,43], but in a small cohort, gene expression of ACE2 in placenta was not different

in preeclamptic versus normotensive patients[44]. Regardless of the tissue origin, elevated

shedding of ACE2 is likely reflective of local tissue damage in pregnancies with adverse

outcomes.

Identification and characterization of pathophysiologies related to adverse outcomes of

pregnancy is necessary for the development of novel, targeted risk assessment and intervention

strategies. Results from the current study indicate temporal activity of biomarkers of the RAS as

a potential candidate for identification of women at high risk for adverse outcomes of pregnancy.

Acknowledgements

We are most grateful for the team of caring clinicians and clinical staff that collect and manage

patient samples and data.

Sources of funding. The work was supported in part by a grant from the Kentucky BIRCWH

Program, NIDA 5K12DA035150 (RS) and by the Kentucky Children's Hospital Children's

Miracle Network Research Fund.

Competing interests. M. Poglitsch is an employee at Attoquant Diagnostics, a company

developing angiotensin-based biomarkers for hypertension. The other authors report no

conflicts.

Author contributions. Study conception and design, clinical (JAB, JOB, CM), analytical (RS);

data collection (KV, AS, HH); analysis and interpretation of results (RS, MP, JAB, JOB);

manuscript preparation (RS).

References

330   1.   Cho L, Davis M, Elgendy I, Epps K, Lindley KJ, Mehta PK, Michos ED, Minissian M, Pepine C,
331   Vaccarino V, et al. Summary of Updated Recommendations for Primary Prevention of
332   Cardiovascular Disease in Women: JACC State-of-the-Art Review. *J Am Coll Cardiol*.
333   2020;75:2602-2618. doi: 10.1016/j.jacc.2020.03.060
334   2.   Yee LM, Miller EC, Greenland P. Mitigating the Long-term Health Risks of Adverse Pregnancy
335   Outcomes. *Jama*. 2022;327:421-422. doi: 10.1001/jama.2021.23870
336   3.   Parikh NI, Gonzalez JM, Anderson CAM, Judd SE, Rexrode KM, Hlatky MA, Gunderson EP, Stuart
337   JJ, Vaidya D. Adverse Pregnancy Outcomes and Cardiovascular Disease Risk: Unique
338   Opportunities for Cardiovascular Disease Prevention in Women: A Scientific Statement From the
339   American Heart Association. *Circulation*. 2021;143:e902-e916. doi:
340   10.1161/cir.0000000000000961
341   4.   Zhang J, Klebanoff MA, Roberts JM. Prediction of adverse outcomes by common definitions of
342   hypertension in pregnancy. *Obstet Gynecol*. 2001;97:261-267. doi: 10.1016/s0029-
343   7844(00)01125-x
344   5.   Irani RA, Xia Y. The functional role of the renin-angiotensin system in pregnancy and
345   preeclampsia. *Placenta*. 2008;29:763-771. doi: 10.1016/j.placenta.2008.06.011
346   6.   Verdonk K, Visser W, Van Den Meiracker AH, Danser AH. The renin-angiotensin-aldosterone
347   system in pre-eclampsia: the delicate balance between good and bad. *Clin Sci (Lond)*.
348   2014;126:537-544. doi: 10.1042/CS20130455
349   7.   Verdonk K, Saleh L, Lankhorst S, Smilde JE, van Ingen MM, Garrelds IM, Friesema EC, Russcher H,
350   van den Meiracker AH, Visser W, et al. Association studies suggest a key role for endothelin-1 in
351   the pathogenesis of preeclampsia and the accompanying renin-angiotensin-aldosterone system
352   suppression. *Hypertension*. 2015;65:1316-1323. doi: 10.1161/HYPERTENSIONAHA.115.05267
353   8.   Vickers C, Hales P, Kaushik V, Dick L, Gavin J, Tang J, Godbout K, Parsons T, Baronas E, Hsieh F, et
354   al. Hydrolysis of biological peptides by human angiotensin-converting enzyme-related
355   carboxypeptidase. *J Biol Chem*. 2002;277:14838-14843. doi: 10.1074/jbc.M200581200
356   9.   Ferrario CM, Chappell MC, Tallant EA, Brosnihan KB, Diz DI. Counterregulatory actions of
357   angiotensin-(1-7). *Hypertension*. 1997;30:535-541.
358   10.   Tamanna S, Clifton VL, Rae K, van Helden DF, Lumbers ER, Pringle KG. Angiotensin Converting
359   Enzyme 2 (ACE2) in Pregnancy: Preeclampsia and Small for Gestational Age. *Front Physiol*.
360   2020;11:590787. doi: 10.3389/fphys.2020.590787
361   11.   Merrill DC, Karoly M, Chen K, Ferrario CM, Brosnihan KB. Angiotensin-(1-7) in normal and
362   preeclamptic pregnancy. *Endocrine*. 2002;18:239-245. doi: 10.1385/ENDO:18:3:239
363   12.   Sykes SD, Pringle KG, Zhou A, Dekker GA, Roberts CT, Lumbers ER, consortium S. Fetal sex and
364   the circulating renin-angiotensin system during early gestation in women who later develop
365   preeclampsia or gestational hypertension. *J Hum Hypertens*. 2014;28:133-139. doi:
366   10.1038/jhh.2013.51
367   13.   Lambert DW, Yarski M, Warner FJ, Thornhill P, Parkin ET, Smith AI, Hooper NM, Turner AJ.
368   Tumor necrosis factor-alpha convertase (ADAM17) mediates regulated ectodomain shedding of
369   the severe-acute respiratory syndrome-coronavirus (SARS-CoV) receptor, angiotensin-
370   converting enzyme-2 (ACE2). *J Biol Chem*. 2005;280:30113-30119. doi:
371   10.1074/jbc.M505111200
372   14.   Junus K, Björk Ragnarsdóttir I, Nordlöf Callbo P, Bergman L, Lager S, Wikström AK. Elevated mid-
373   pregnancy plasma levels of angiotensin-converting enzyme 2 in women prior to the
374   development of preeclampsia. *Sci Rep*. 2022;12:4109. doi: 10.1038/s41598-022-08081-8

15. Narula S, Yusuf S, Chong M, Ramasundarahettige C, Rangarajan S, Bangdiwala SI, van Eikels M,
Leineweber K, Wu A, Pigeyre M, et al. Plasma ACE2 and risk of death or cardiometabolic
diseases: a case-cohort analysis. *Lancet*. 2020;396:968-976. doi: 10.1016/s0140-6736(20)31964-
4
16. Ramchand J, Patel SK, Kearney LG, Matalanis G, Farouque O, Srivastava PM, Burrell LM. Plasma
ACE2 Activity Predicts Mortality in Aortic Stenosis and Is Associated With Severe Myocardial
Fibrosis. *JACC Cardiovasc Imaging*. 2020;13:655-664. doi: 10.1016/j.jcmg.2019.09.005
17. Rieder M, Wirth L, Pollmeier L, Jeserich M, Goller I, Baldus N, Schmid B, Busch HJ, Hofmann M,
Kern W, et al. Serum ACE2, Angiotensin II, and Aldosterone Levels Are Unchanged in Patients
With COVID-19. *Am J Hypertens*. 2021;34:278-281. doi: 10.1093/ajh/hpaa169
18. LeFevre ML, Force USPST. Low-dose aspirin use for the prevention of morbidity and mortality
from preeclampsia: U.S. Preventive Services Task Force recommendation statement. *Ann Intern
Med*. 2014;161:819-826. doi: 10.7326/M14-1884
19. Pickering TG, Hall JE, Appel LJ, Falkner BE, Graves JW, Hill MN, Jones DH, Kurtz T, Sheps SG,
Roccella EJ, et al. Recommendations for blood pressure measurement in humans: an AHA
scientific statement from the Council on High Blood Pressure Research Professional and Public
Education Subcommittee. *J Clin Hypertens (Greenwich)*. 2005;7:102-109. doi: 10.1111/j.1524-
6175.2005.04377.x
20. Bhide A, Acharya G, Bilardo CM, Brezinka C, Cafici D, Hernandez-Andrade E, Kalache K, Kingdom
J, Kiserud T, Lee W, et al. ISUOG practice guidelines: use of Doppler ultrasonography in
obstetrics. *Ultrasound Obstet Gynecol*. 2013;41:233-239. doi: 10.1002/uog.12371
21. Haschke M, Schuster M, Poglitsch M, Loibner H, Salzberg M, Bruggisser M, Penninger J,
Krähenbühl S. Pharmacokinetics and pharmacodynamics of recombinant human angiotensin-
converting enzyme 2 in healthy human subjects. *Clin Pharmacokinet*. 2013;52:783-792. doi:
10.1007/s40262-013-0072-7
22. Basu R, Poglitsch M, Yogasundaram H, Thomas J, Rowe BH, Oudit GY. Roles of Angiotensin
Peptides and Recombinant Human ACE2 in Heart Failure. *J Am Coll Cardiol*. 2017;69:805-819.
doi: 10.1016/j.jacc.2016.11.064
23. Wang K, Basu R, Poglitsch M, Bakal JA, Oudit GY. Elevated Angiotensin 1-7/Angiotensin II Ratio
Predicts Favorable Outcomes in Patients With Heart Failure. *Circ Heart Fail*. 2020;13:e006939.
doi: 10.1161/circheartfailure.120.006939
24. Zoufaly A, Poglitsch M, Aberle JH, Hoepler W, Seitz T, Traugott M, Grieb A, Pawelka E, Laferl H,
Wenisch C, et al. Human recombinant soluble ACE2 in severe COVID-19. *Lancet Respir Med*.
2020;8:1154-1158. doi: 10.1016/s2213-2600(20)30418-5
25. Reindl-Schwaighofer R, Hödlmoser S, Eskandary F, Poglitsch M, Bonderman D, Strassl R, Aberle
JH, Oberbauer R, Zoufaly A, Hecking M. Angiotensin-Converting Enzyme 2 (ACE2) Elevation in
Severe COVID-19. *Am J Respir Crit Care Med*. 2021. doi: 10.1164/rccm.202101-0142LE
26. Domenig O, Manzel A, Grobe N, Konigshausen E, Kaltenecker CC, Kovarik JJ, Stegbauer J, Gurley
SB, van Oyen D, Antlanger M, et al. Neprilysin is a Mediator of Alternative Renin-Angiotensin-
System Activation in the Murine and Human Kidney. *Sci Rep*. 2016;6:33678. doi:
10.1038/srep33678
27. Langer B, Grima M, Coquard C, Bader AM, Schlaeder G, Imbs JL. Plasma active renin, angiotensin
I, and angiotensin II during pregnancy and in preeclampsia. *Obstet Gynecol*. 1998;91:196-202.
28. Malha L, Sison CP, Helseth G, Sealey JE, August P. Renin-Angiotensin-Aldosterone Profiles in
Pregnant Women With Chronic Hypertension. *Hypertension*. 2018;72:417-424. doi:
10.1161/HYPERTENSIONAHA.118.10854
29. Leanos-Miranda A, Inova C-G, Mendez-Aguilar F, Molina-Perez CJ, Ramirez-Valenzuela KL, Sillas-
Pardo LJ, Uraga-Camacho NC, Isordia-Salas I, Berumen-Lechuga MG. Lower circulating

423		angiotensin II levels are related to the severity of preeclampsia and its risk as disclosed by a
424		specific bioassay. *Medicine (Baltimore)*. 2018;97:e12498. doi: 10.1097/MD.0000000000012498
425	30.	Zitouni H, Raguema N, Gannoun MBA, Hebert-Stutter M, Zouari I, Maleh W, Faleh R, Letaifa DB,
426		Almawi WY, Fournier T, et al. Impact of obesity on the association of active renin and plasma
427		aldosterone concentrations, and aldosterone-to-renin ratio with preeclampsia. *Pregnancy*
428		*Hypertens*. 2018;14:139-144. doi: 10.1016/j.preghy.2018.09.007
429	31.	Brosnihan KB, Merrill DC, Yamaleyeva LM, Chen K, Neves L, Joyner J, Givner C, Lanier K,
430		Moorefield C, Westwood B. Longitudinal study of angiotensin peptides in normal and pre-
431		eclamptic pregnancy. *Endocrine*. 2020;69:410-419. doi: 10.1007/s12020-020-02296-3
432	32.	Wiegel RE, Jan Danser AH, Steegers-Theunissen RPM, Laven JSE, Willemsen SP, Baker VL,
433		Steegers EAP, von Versen-Hoynck F. Determinants of Maternal Renin-Angiotensin-Aldosterone-
434		System Activation in Early Pregnancy: Insights From 2 Cohorts. *J Clin Endocrinol Metab*.
435		2020;105. doi: 10.1210/clinem/dgaa582
436	33.	Burrello J, Buffolo F, Domenig O, Tetti M, Pecori A, Monticone S, Poglitsch M, Mulatero P. Renin-
437		Angiotensin-Aldosterone System Triple-A Analysis for the Screening of Primary Aldosteronism.
438		*Hypertension*. 2020;75:163-172. doi: 10.1161/HYPERTENSIONAHA.119.13772
439	34.	Pavo N, Goliasch G, Wurm R, Novak J, Strunk G, Gyongyosi M, Poglitsch M, Saemann MD,
440		Hulsmann M. Low- and High-renin Heart Failure Phenotypes with Clinical Implications. *Clin*
441		*Chem*. 2018;64:597-608. doi: 10.1373/clinchem.2017.278705
442	35.	Binder C, Poglitsch M, Agibetov A, Duca F, Zotter-Tufaro C, Nitsche C, Aschauer S, Kammerlander
443		AA, Oeztuerk B, Hengstenberg C, et al. Angs (Angiotensins) of the Alternative Renin-Angiotensin
444		System Predict Outcome in Patients With Heart Failure and Preserved Ejection Fraction.
445		*Hypertension*. 2019;74:285-294. doi: 10.1161/HYPERTENSIONAHA.119.12786
446	36.	Bernstone L, Adaway JE, Keevil BG. An LC-MS/MS assay for analysis of equilibrium angiotensin II
447		in human serum. *Ann Clin Biochem*. 2021:45632211008923. doi: 10.1177/00045632211008923
448	37.	Guo M, Zhou C, Xu G, Tang L, Ruan Y, Yu Y, Lin X, Wu D, Chen H, Yu P, et al. An alternative
449		splicing variant of mineralocorticoid receptor discovered in preeclampsia tissues and its effect
450		on endothelial dysfunction. *Sci China Life Sci*. 2020;63:388-400. doi: 10.1007/s11427-018-9535-9
451	38.	Chappell MC. Biochemical evaluation of the renin-angiotensin system: the good, bad, and
452		absolute? *Am J Physiol Heart Circ Physiol*. 2016;310:H137-152. doi:
453		10.1152/ajpheart.00618.2015
454	39.	Chappell MC, Pirro NT, South AM, Gwathmey TM. Concerns on the Specificity of Commercial
455		ELISAs for the Measurement of Angiotensin (1-7) and Angiotensin II in Human Plasma.
456		*Hypertension*. 2021;77:e29-e31. doi: 10.1161/hypertensionaha.120.16724
457	40.	Rice GI, Jones AL, Grant PJ, Carter AM, Turner AJ, Hooper NM. Circulating activities of
458		angiotensin-converting enzyme, its homolog, angiotensin-converting enzyme 2, and neprilysin in
459		a family study. *Hypertension*. 2006;48:914-920. doi: 10.1161/01.Hyp.0000244543.91937.79
460	41.	Úri K, Fagyas M, Kertész A, Borbély A, Jenei C, Bene O, Csanádi Z, Paulus WJ, Édes I, Papp Z, et al.
461		Circulating ACE2 activity correlates with cardiovascular disease development. *J Renin*
462		*Angiotensin Aldosterone Syst*. 2016;17. doi: 10.1177/1470320316668435
463	42.	Delforce SJ, Lumbers ER, Ellery SJ, Murthi P, Pringle KG. Dysregulation of the placental renin-
464		angiotensin system in human fetal growth restriction. *Reproduction*. 2019;158:237-245. doi:
465		10.1530/rep-18-0633
466	43.	Nonn O, Fischer C, Geisberger S, El-Heliebi A, Kroneis T, Forstner D, Desoye G, Staff AC, Sugulle
467		M, Dechend R, et al. Maternal Angiotensin Increases Placental Leptin in Early Gestation via an
468		Alternative Renin-Angiotensin System Pathway: Suggesting a Link to Preeclampsia.
469		*Hypertension*. 2021;77:1723-1736. doi: 10.1161/hypertensionaha.120.16425

44. Sun Y, Tan L, Neuman RI, Broekhuizen M, Schoenmakers S, Lu X, Danser AHJ. Megalin, Proton
Pump Inhibitors and the Renin-Angiotensin System in Healthy and Pre-Eclamptic Placentas. *Int J*
Mol Sci. 2021;22. doi: 10.3390/ijms22147407

Figure Legends

Figure 1: Adverse outcomes of pregnancies (AO) are associated with suppression of the

classical components of the RAS, and activation of the counter-regulatory protein, ACE2. A)

Renin activity, angiotensin I (Ang I) and angiotensin II (Ang II) peptides, and B) concentration of

soluble ACE2 in serum of pregnant women. Data are the change in concentration over gestation

(third – first trimester). Data are represented as median with 95% confidence interval. *, $P < 0.05$

analyzed by Wilcoxon signed rank test; n=74 pairs.

Table 1. Patient Demographics and Clinical Characteristics of Pregnant Women With and Without Adverse Outcomes

Parameter	No AO N = 28	AO N = 46
Patient Demographics, n (%)		
Race		
Caucasian	18 (64)	34 (74)
Black or African	5 (18)	8 (17)
American	4 (14)	3 (7)
Hispanic	1 (4)	1 (2)
Mixed race		
Primiparous	11 (39)	20 (42)
Chronic hypertension	8 (29)	9 (20)
Previous pre-eclampsia	5 (18)	9 (20)
Type 1 DM	1 (4)	7 (15)
Type 2 DM	2 (7)	2 (4)
Renal disease	0	0
Autoimmune disease	3 (11)	1 (2)
Medications		
Aspirin	15 (54)	25 (54)
Labetalol	2 (7)	8 (17)
Nifedipine	0	3 (7)
Clinical Characteristics, mean $\pm$ SD		
Age, years	28.9 $\pm$ 4.1	29.5 $\pm$ 5.7
BMI, kg/m^2	30.0 $\pm$ 7.6	32.3 $\pm$ 8.3
SBP, mmHg		
1st Trimester	117 $\pm$ 15	120 $\pm$ 9
3rd Trimester	119 $\pm$ 13	124 $\pm$ 14
DBP, mmHg		
1st Trimester	75 $\pm$ 10	79 $\pm$ 6
3rd Trimester	76 $\pm$ 9	80 $\pm$ 9
PI (uterine), %	1.08 $\pm$ 0.36	1.26 $\pm$ 0.47
Gestational age at delivery, weeks	38.7 $\pm$ 1.1	36.9 $\pm$ 2.2[***]
Infant body weight, grams	3383 $\pm$ 429	3083 $\pm$ 648[*]

[*], P< 0.05; [***], P<0.0001

Table 2. Recorded Pregnancy Outcomes

Pregnancy Outcomes	No AO N = 28	AO N = 46 n (%)
Gestational hypertension	0	19 (41)
Pre-eclampsia	0	7 (15)
Post-partum pre-eclampsia	0	3 (7)
IUGR	0	3 (7)
IAEDF	0	2 (4)
Polyhydramnios	0	5 (11)
Pre-term labor	0	3 (7)
PPROM	0	5 (11)
PROM	0	1 (2)
Pre-term birth	0	1 (2)
Stillbirth	0	1 (2)
Postpartum hemorrhage	0	10 (22)

Table 3. Serum Concentrations of Components of the Classical and Alternative Renin Angiotensin System in Pregnancies With and Without Adverse Outcomes

RAS Component	No AO N=28	AO N=46	P value
Classical RAS, median (95% CI)			
Renin activity			
1st Trimester	266.5 (232.2 – 303.9)	199.5 (147.0 – 275.9)	0.070
3rd Trimester	349.5 (294.4 – 476.9)**	239.3 (196.9 – 303.4)	0.011
ACE activity			
1st Trimester	2.18 (1.84 – 2.65)	2.49 (2.22 – 3.08)	0.237
3rd Trimester	1.61 (1.32 – 2.07)****	2.07 (1.76 – 2.26)****	0.140
Ang I			
1st Trimester	86.3 (69.5 – 100.9)	60.9 (40.9 – 97.6)	0.154
3rd Trimester	135.1 (92.2 – 183.0)***	78.4 (62.7 – 119.0)**	0.020
Ang II			
1st Trimester	179.3 (151.5 – 197.0)	147.5 (102.2 – 185.4)	0.107
3rd Trimester	228.2 (177.2 – 292.5)**	150.7 (123.3 – 198.0)	0.010
Ang IV			
1st Trimester	8.3 (6.5 – 9.7)	6.9 (5.0 – 10.0)	0.174
3rd Trimester	9.6 (6.8 – 12.8)	7.6 (5.2 – 9.8)	0.108
Alternative RAS, median (95% CI)			
ACE2			
1st Trimester	1.43 (1.27 – 1.76)	1.23 (1.14 – 1.37)	<0.05
3rd Trimester	1.68 (1.56 – 2.10)**	1.69 (1.5 – 2.09)****	0.844
Ang-1-5			
1st Trimester	5.7 (5.1 – 7.0)	3.9 (3.0 – 5.1)	0.308
3rd Trimester	7.9 (6.6 – 9.3)	5.6 (3.9 – 7.6)**	0.017
NEP			
1st Trimester	10.77 (6.-6 – 16.2)	9.43 (6.69 – 11.29)	0.373
3rd Trimester	14.52 (10.54 – 26.33)*	13.10 (11.18 – 18.55)***	0.415

*, $P<0.05$, **, $P<0.01$, ***, $P<0.001$, ****, $P<0.0001$ for third trimester vs first trimester within group analyzed by Wilcoxon signed rank test

A

B