Single-cell RNA sequencing reveals the distinctive roles of CD4[+] and CD8[+] T cells in autoimmune uveitis

Hao Kang[1#], Hongjian Sun[2,3#], Yang Yang[2,3], Minglei Shu[3], Yunbo Wei[4], Yu Zhang[4], Di Yu[2,5*] and Yong Tao[1*]

1. Department of Ophthalmology, Beijing Chaoyang Hospital, Capital Medical University
2. Frazer Institute, Faculty of Medicine, The University of Queensland, Brisbane, QLD, Australia
3. Shandong Artificial Intelligence Institute, Qilu University of Technology (Shandong Academy of Sciences), Jinan, China
4. School of Pharmaceutical Sciences, Laboratory of Immunology for Environment and Health, Shandong Analysis and Test Center, Qilu University of Technology (Shandong Academy of Sciences), Jinan, China
5. Ian Frazer Centre for Children's Immunotherapy Research, Children's Health Research Centre, Faculty of Medicine, The University of Queensland, Brisbane QLD, Australia

Hao Kang and Hongjian Sun contribute equally to this study.
* Di Yu and Yong Tao contribute equally to this study.

Correspondence:
Prof. Yong Tao
Department of Ophthalmology, Beijing Chaoyang Hospital, Capital Medical University;
No. 8, South Road of Worker's Stadium, Chaoyang District, Beijing 100020, China.
Phone: +86-10-85231718.
Fax: +86-10-85231405.
E-mail: taoyong@bjcyh.com

Prof. Di Yu
Frazer Institute, Faculty of Medicine, The University of Queensland.
Translational Research Institute (TRI), 37 Kent Street, Woolloongabba, QLD 4102, Australia.
Phone: +61-07-34436954.
E-mail: di.yu@uq.edu.au

Abstract

Uveitis is a common cause of blindness and classified as infectious or non-infectious types. Non-infectious uveitis is often secondary to systemic autoimmune diseases, with Behçet's disease (BD) and Vogt-Koyanagi-Harada disease (VKHD) as the two most common causes. Uveitis in BD and VKHD often show similar clinical manifestations, but the underlying immunopathogenesis remains unclear. We collected immune cell infiltrates in aqueous humour of BD and VKHD uveitis patients and analysed them by single-cell RNA paired with T cell receptor (TCR) sequencing. T cells were found as the dominant immune cell infiltration. Intriguingly, a majority of CD4$^+$ in T cell infiltrates was only observed in VKHD but not in BD. In agreement with this discrepancy between VKHD and BD, T cell clonality analysis and Clonotype Neighbour Graph Analysis (CoNGA) also revealed a selective expansion of pro-inflammatory CD4$^+$ T cell clones in VKHD. In contrast, BD showed a selective expansion of pro-inflammatory CD8$^+$ T cell clones. Our study reveals distinct immunopathogenesis of uveitis driven by CD4$^+$ T cells in VKHD and by CD8$^+$ T cells in BD, and therefore suggests differential approaches for their diagnosis and therapy.

Keywords:

Uveitis

single-cell RNA sequencing

Behcet disease

Vogt-Koyanagi-Harada disease

T cell clonality

1. Introduction

Uveitis is an inflammatory disorder of the uveal tract of the eye that can be caused by multiple reasons. This vision-threatening disease can cause serious complications such as visual impairment and blindness in 35% of patients [1]. Uveitis can be divided into infectious and non-infectious types according to its etiology. As a more common type, non-infectious uveitis is often related to autoimmune diseases, such as Behçet's disease (BD) and Vogt-Koyanagi-Harada disease (VKHD). Although BD and VKHD show the characteristics of familial aggregation, they differ in geographical distribution. BD is more prevalent in Asia, the Middle East and the Mediterranean than other regions and is characterised by three main clinical features of nongranulomatous uveitis, oral ulcers and genital ulcers [2]. VKHD is considered to distribute worldwide, with a higher occurrence in people with darker skin [3]. The clinical features of VKHD are bilateral granulomatous pancreatitis, polio, vitiligo, and central nervous system abnormalities [4].

The immunological pathogenesis of BD and VKHD remains elusive although the dysregulated function of T cells, including cytotoxic CD8$^+$ T cells or CD4$^+$ T cell subset Th1 and Th17 cells, have been implicated in the development of both BD and VKHD [5, 6]. To gain insight into the immunological pathogenesis of human uveitis, we decided to perform single-cell RNA sequencing (scRNA-seq) of cellular infiltrates in aqueous humour from uveitis patients. Due to the significant involvement of T cells in immunopathogenesis, we included T cell receptor (TCR) sequencing. We discovered that uveitis in VKHD was associated with a strong clonal expansion of effector CD4$^+$ T cells expressing pro-inflammatory cytokines *IFNG* and *TNF*. In contrast, uveitis in BD was associated with a selective clonal expansion of effector CD8+ T cells expressing cytotoxic molecules *GZMB* and *PRF1*. The different immunopathogenesis of non-infectious uveitis distinctively driven by CD4$^+$ T cells in VKHD and CD8$^+$ T cells in BD strongly suggests differential approaches for the diagnosis and therapy of uveitis in VKHD and BD.

2. Material and methods

2.1 Resource availability

2.1.1 Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the corresponding authors Di Yu and Yong Tao.

2.1.2 Data and code availability

The datasets during this study are available at the *Science DataBase*: https://www.scidb.cn/s/bAF7fe. Codes are available at https://github.com/HongjianSun/Uveitis-project.

2.2 Experiment model and subject details

This study was performed at Beijing Chaoyang Hospital, Capital Medical University, Beijing, China. The ethics

approval was obtained from the Ethics Committee of Beijing Chaoyang Hospital. Written informed consent

was obtained from all patients. Two VKHD patients and one BD patient (30-40 years old, > 3 years of disease

duration) were enrolled at the Ophthalmology Department of Beijing Chaoyang Hospital affiliated with Capital

Medical University. All patients had bilateral or unilateral vision loss and had cataract surgeries.

2.3 Method details

<u>**2.3.1 Single-cell collection**</u>

Aqueous humour of about 100 µL was taken from individual patients after an anterior chamber paracentesis

(ACP). The ACP was carried out by using a 30-gauge needle, on a 1 mL insulin syringe, via the temporal

limbal approach and a rolling technique. Aqueous humour was aspirated and immediately aliquoted into a

microfuge tube. All the aqueous humour samples were centrifuged at 800 g for 15 minutes. After removing the

supernatant, the cells were suspended in red blood cell lysis buffer (Solarbio) and incubated on ice for 2 min

to lyse red blood cells. The pellets were washed twice with PBS.

<u>**2.3.2 Single-cell RNA sequencing**</u>

10X Genomics Platform was used for single-cell RNA sequencing. Single-cell suspension with gel beads

(containing the pre-made 10X primers) and Master Mix together formed Gel Bead-In-Emulsions (GEMs).

Cleavage and reverse transcription were performed in GEMs. After cDNA amplification was completed, we

performed a quality inspection and finally constructed the library. The Illumina Novaseq 6000 sequencing

platform was used for sequencing, obtaining sequencing data, and performing subsequent data analysis.

Fastq reads were initially filtered using *Trimmomatic*[7]. Next, we used Feature Counts software to quantify the

expression of each gene, and counts were obtained for each sample. In our analysis, a qualified gene

expression profile from a single cell should have one or more counts in a sample. The expression level of

each gene was converted to a transcript per million (TPM) value. Then, the log-normalized expression values

were transformed.

<u>**2.3.3 Data processing**</u>

Seurat[8] was used for scRNA-seq data processing. *Doubletfinder* R package was used to identify doublets[9]

and 7.5% was removed as the recommendation. Subsequently, cells from three patient samples were

integrated using the *IntegrateData* package to eliminate batch effects.

2.3.4 Data analyzing

The data were normalized and scaled by NormalizeData and *ScaleData* in *Seurat.* Unsupervised clustering

was performed using *FindClusters* in *Seurat* by a shared nearest neighbour modularity optimization.

Differential gene expression analysis was performed using *FindMarkers* and *FindAllMarkers* in *Seurat* with the

parameter of adjusted P-value threshold<0.05. Trajectory analysis was performed by *learn_graph, order_cells*

and *plot_cells* in *Monocle3* R package and based on UMAP reduction method[10].

2.3.5 Visualization

scRNA-seq data were visualized using the dimensionality reduction method of UMAP *(*Uniform Manifold

Approximation and Projection*).* Stacked violin plots were made by a modified function of *VInplot* in *Seurat.*

Heat map showing the top 5 highest differential expressed genes in each cluster was made by *DoHeatmap*

function.

2.3.6 Enrichment Analysis

Gene set enrichment analysis (*GSEA*) was performed[11] with gene sets from *Molecular Signatures Database*

v7.4. The enrichment scores (ES) for a gene set reflect the extent to which a transcriptome is overrepresented

at the top or bottom of a sequence list of genes, while Normalized Enrichment Scores (NES) are used for

comparing the enrichment results through the gene sets. The P-value and the False Discovery Rate (FDR)

are both parameters indicating the confidence level of gene set enrichment.

2.3.7 V(D)J sequencing

V(D)J sequencing were analyzed by *CellRanger*. The *CellRanger* VDJ function was used for Contig assembly,

cross-cell Consensus sequence, CDR3 Clonotype typing of Loupe V(D)J Browser analysis results on TCR

gene V(D)J region sequences of a single cell. *Seurat* and *scRepertoire*[12] were used to add VDJ sequencing

data as additional data to merge with single-cell sequencing data.

2.3.8 Clonotype neighbour graph analysis

Clonotype neighbour graph analysis(*CoNGA*) identifies the correlation between T cell gene expression(GEX)

and TCR sequence by constructing a GEX similarity graph based on the gene expression, and a TCR

sequence similarity graph based on *TCRdist* for each clonotypes' CDR3 sequence, and then find a statistically

significant overlap between them. Overlap is assessed on a score that reflects the likelihood of seeing equal

or greater overlap by chance, named the CoNGA score. The more the overlap between GEX and TCR graph,

the smaller the CONGA score will be. Then, Clonotypes with CoNGA scores below a threshold are grouped

based on shared GEX and TCR cluster assignments into CoNGA clusters. Clonotypes within each CoNGA

cluster carry their initial GEX and TCR cluster identities, which are combined together and used as a group ID

for the CoNGA cluster.

2.3.9 Statistical analysis

The *Circos* [13] was created to illustrate the correlations between two groups. Pie charts and histograms were

made by *Prism V9* and *Microsoft Excel.*

3.　Result

3.1 T cells are the dominant immune cell type infiltrating in uveitis lesions in both BD and

VKHD

Three uveitis patients at the age of 30-40 years old, one diagnosed with BD and two diagnosed with chronic

recurrent VKHD volunteered to participate in this study. They all had decreased vision and had cataract surgery

more than two years before the sample collection in 2019 (**Fig. 1a**). We choose 5' whole transcriptome gene

expression (10×Genomics) to gain the information for both transcriptomes and paired full-length TCR for

immune repertoire analysis. The library preparation, sequencing, read mapping, gene count normalization and

doublet removal were conducted using a standard pipeline (**Fig. 1b**) [14].

After the integration of data from three samples, a total of 17,961 cells underwent the downstream analysis.

After doublet removing and data integration, the single cells were visualized by UMAP in two-dimensional space.

Using KNN method for clustering, the immune cells in uveitis were classified into 12 clusters (**Fig. 1c, S1a**).

CD3+ T cells (cluster 0-4, 6, 8-11) were the dominant immune cell population (11,983 cells) accounting for 44.6%

(856/1919), 72.7% (4557/6268) and 67.2% (6570/9774) of total cells from BD, VKHD1 and VKHD2 patients.

CD19+ B cells (cluster 5) were approximately 10% in all samples. Interestingly, *CD14*+ monocytes (cluster 7)

showed a higher percentage in BD (20.6%) than those in VKHD (VKHD1: 4.0%; VKHD2: 0.4%) (**Fig. 1d, S1b**).

3.2 T cells in uveitis lesions are composed of distinct functional subsets

We then focused on T cells as the major immune cell population and identified 11 clusters (**Fig. 2a**), which were

predominantly αβ T cells (**Fig. 2b**). We didn't observe significant enrichment of genes for TCR δ and γ,

Vα24/Vβ11 associated with natural killer T (NKT) cells, or Vα2/Vα7/Vβ2/13 associated with mucosal-associated

invariant T (MAIT) cells in particular subsets [15, 16] (**Fig.S2a**). Clusters 1, 2, 4, 5, 7, 8, 9 and 10 expressed *CD4*

while clusters 3 and 6 expressed *CD8A* and *CD8B*. Cluster 0 are CD4$^-$CD8$^-$ (double negative, DN) T cells, a

population which is rare in a normal condition but expands in inflammation and autoimmune diseases [17] (**Fig.**

2b). Notably, VKHD samples showed CD4$^+$ T cells as the clear majority (VKHD1: 68%; VKHD2: 57%) with

CD4/CD8 ratios as 4.3 and 2.6. In contrast, CD4$^+$ and CD8$^+$ T cell numbers were largely comparable in BD (**Fig.**

2c), with the CD4/CD8 ratio of 1.4. This difference suggests a potential difference in immunopathogenesis of

non-infectious uveitis of BD and VKHD.

Using differential gene expressions (DGE) among different clusters (**Fig. 2d**), we annotated their states and

functions. CD4$^-$CD8$^-$ T cells (Cluster 0, C0) were resting T cells with low expression of effector molecules (**Fig.**

2e). Within the CD8$^+$ T cells, C6 was a central memory population with the higher expression of the chemotaxis

and trafficking receptor genes *CCR7* and *SELL* (encoding CD62L) for recirculation, while C3 was the effector

subset with significant expression of effector function genes for C-C chemokine ligands (*CCL5*), granzymes

(*GZMA* and *GZMB*) and perforin (*PRF1*) (**Fig. 2e**). Gene Set Enrichment Analysis (GSEA) demonstrated that

DGE between C3 and C6 was positively associated with gene expression in effector CD8$^+$ T cells as compared

to memory or naïve CD8$^+$ T cells in the model of acute infection (**Fig. S2b**). The trajectory analysis suggested

that CD8$^+$ T cells underwent a relatively linear differentiation with a likely progression from memory to effector

(**Fig. s2c**).

Within the CD4$^+$ T cells, C8 showed a high expression of the proliferation marker *MKI67* (encoding KI-67). Such

a highly proliferative population also expressed CD38 and proinflammatory effector genes *IFNG* and *TNF* (**Fig.**

2e), which were recently shown to be correlated with disease severity in COVID-19 [18]. Compared to C8, C4

expresses higher amounts of *IFNG* and *TNF*. They reduced proliferation but underwent terminal differentiation

with an upregulation of *PDCD1* (encoding PD-1). Notably, C1 showed the highest levels of *PDCD1* and *CTLA4*,

suggesting an exhaustion state with the downregulation of effector molecules (**Fig. 2e**). DGE between C4 and

C1 was enriched with the signatures of T cell exhaustion by GSEA (**Fig. S2d**).

C9 is tissue-residential memory T (T$_{RM}$) cells with the expression of chemokine receptor and integrin *CXCR6*

and IGTB1. C2 and C5 express a high level of inhibitory receptor *KLRB1* (encoding CD161). C10 upregulated

the initiator of the complement classical pathway *C1QA* and *C1QB*, which were reported to restrain autoimmune

response [19]. C7 was regulatory T (T$_{REG}$) cells with the high expression of markers *CTLA4*, *FOXP3* and *IL2RA*

(encoding CD25) (**Fig. 2e**). The trajectory analysis of conventional CD4$^+$ T cells (C8, CD4, C1, C9, C2, C5 and

C10) suggested that recently activated and proliferating CD4$^+$ T cells (C8) may further differentiate and branch

into effector (C4), exhausted (C1) and KLRB1$^+$ (C2 and C5) clusters (**Fig. S2e**).

With the annotation of T cell functional clusters, we noticed major differences in the frequencies of CD4$^+$ and

CD8$^+$ T cell clusters between VKHD and BD whereas the frequencies of DN T cells were largely comparable

among these patients. In agreement with the selectively high CD4/CD8 ratios in VKHD patients, effector (C4)

and exhausted (C1) CD4$^+$ T cells were in much larger amounts in VKHD but not in BD patients. In contrast,

effector (C3) CD8+ T cells were particularly expanded in the BD patient (**Fig. 2f**). These results suggest a notion

that uveitis in VKHD and BD is preferentially associated with effector CD4$^+$ T cells and effector CD8$^+$ T cells,

respectively.

3.3 Divergent T cell clonal expansion in BD and VKHD uveitis

The antigen is recognised by a specific TCR sequence and its stimulation critically drives T cell proliferation and

functional differentiation. We then analysed TCR sequences of T cells in all three patient samples, with an

emphasis on clonality by quantifying the frequencies of T cells with a common CDR3 sequence. According to

the percentages of the T cell numbers of each T cell clone in the total T cells from a given sample, significantly

expanded clones (>0.2% of total T cells) were classified into three groups: Large (> 5% of total T cells), medium

(1~5%), small (0.2-1%) clonotypes. Notably, large clonal types are enriched in effector CD8$^+$ T cells (C3) and

effector (C4) and exhausted (C1) CD4$^+$ T cells (**Fig. 3a**), indicating that effector T cells-mediated inflammation

in uveitis is driven by antigen stimulation. Intriguingly, there was no large or medium clonal expansion for Treg

cells (C7).

A total of 13 large and medium clonaltypes (>1% of total T cells) were detected in VKHD and BD samples,

including 5 in VKHD1, 6 in VKHD2 and 2 in BD (**Fig. 3b, S3**). There were no identical CDR3 sequences among

clonotypes from different samples. We then generated topographic maps to visualise the distribution of cells

from large and medium clonotypes in each sample. Furthermore, to understand the relationship between clonal

expansion and functional T cell clusters, we calculated the frequencies of large, medium and small clonotypes

in each functional T cell cluster and also the frequencies of each cluster in large, medium or small clonotypes.

In VKHD, T cell clonal expansions were centred by effector (C4) and exhausted (C1) CD4$^+$ T cells (left panels,

Fig. 3c). Indeed, effector (C4) and exhausted (C1) CD4$^+$ T cells accounted for about 70% large and medium

clonotypes (middle panels, **Fig. 3c**). Conversely, large and medium clonotypes only accounted for 10-20% of

total effector (C4) and exhausted (C1) CD4$^+$ T cells (right panels, **Fig. 3c**). The inclusion of small clonotypes

could increase the frequencies to about 50%. These results suggest a key role of (auto)antigen-driven CD4$^+$ T

cell activation in the immunopathogenesis in VKHD.

In BD, T cell clonal expansion was exclusively in effector (C3) CD8$^+$ T cells (left panel, **Fig. 3d**). Strikingly,

effector (C3) CD8$^+$ T cells accounted for about 80% of large clonotypes (middle panel, **Fig. 3d**). Conversely,

large clonotypes accounted for >40% of total effector (C3) CD8$^+$ T cells (right panel, **Fig. 3d**). These results

strongly argue an essential role of (auto)antigen-driven CD8$^+$ T cell activation in the immunopathogenesis in BD,

which is distinct from VKHD.

In both VKHD and BD, T$_{REG}$ cells (C7) only accounted for a small fraction of T cells (3-5%, **Fig. 2f**). A striking

observation was that T$_{REG}$ cells (C7) underwent the least clonal expansion in both VKHD and BD, suggesting

that insufficient expansion of T_{REG} cells might posit in the root of losing immune tolerance in non-infectious
uveitis. Our previous clinical trials of low-dose IL-2 therapy in systems lupus erythematosus and Sjogren's
syndrome have demonstrated that it could enhance antigen-independent expansion of T_{REG} cells [20, 21], thus
providing a strategy to improve the poor clonal expansion of T_{REG} cells in non-infectious uveitis.

3.4 CoNGA identifies similar clonotypes with minor contribution to the inflammation in uveitis

About 50% of effector (C4) and exhausted (C1) CD4$^+$ T cells in VKHD (right panels, **Fig. 3c**) and about 40% of
effector (C3) CD8$^+$ T cells in BD (right panels, **Fig. 3d**) didn't carry significantly expanded TCR sequences, e.g.
not in large, medium or small clonaltypes. This may be explained by TCR-independent activation of T cells, a
phenomenon termed 'bystander activation' that has been reported in infection and inflammation [22]. An
alternative explanation could be that (auto)antigen can activate T cells with similar TCR but such related
clonotypes were not identified by identical CDR3 sequences. To test the latter possibility, we applied Clonotype
neighbour graph analysis (*CoNGA*), which is a latest method for identifying related clonotypes with similarities
in both gene expression and TCR sequences [23] (**Fig. 4a**).

We performed *CoNGA* on merged single-cell and TCR sequence data. Noticeably, *CoNGA* transformed the
scRNA-seq and TCR sequence data and reduced multiple T cells with identical TCR to one cell for *CoNGA* plots
[23]. *CoNGA* identified four *CoNGA* clonotypes/clusters (similar gene expression and similar TCR sequences)
(**Fig. 4b**). Intriguingly, *CoNGA* clonotypes/clusters 0-10 (GEX Cluster0-TCR Cluster10, **Fig. 4b**) and 0-3 were
composed of cells from both VKHD1 and VKHD2. *CoNGA* clonotypes/clusters 4-2 and 5-7 were composed of
cells solely from VKHD1 and BD respectively (**Fig. 4c**). We labelled cells identified by *CoNGA*
clonotypes/clusters in the UMAP plot and found the majority of such cells were effector (C4) and exhausted (C1)
CD4$^+$ T cells, or effector (C3) CD8$^+$ T cells, except for some cells in the *CoNGA* clonotype/cluster 5-7 belowed
to DN (C0) T cells (**Fig. 4d**).

These results suggest that (auto)antigen can activate CD4$^+$ or CD8$^+$ T cells with similar rather than identical
TCR to generate effector cells. We next calculated the contribution of 'similar' clonotypes and they only very
modestly increased the clonotype frequencies in clusters **(Fig. S4)**. Taken together, 'similar' clonotypes can
contribute to the immunopathogenesis of uveitis, but 'identical' clonotypes appeared to be the major drivers.

4. Discussion

Ocular immune privilege has been supported by multiple facets of evidence [24]. The prominent presence of T
cell infiltration in non-infectious uveitis suggests they may play a significant role in the immunopathogenesis,
which is also supported by results from mouse models of experimental autoimmune uveitis and clinical benefits
observed from T cell-targeting therapies [25]. However, the results from human studies were not always
consistent, at least partially due to the heterogeneity nature of non-infectious uveitis. Systems immunology
approaches have been taken to stratify uveitis patients, such as by peripheral blood transcriptomic analyses [26].

The latest advancement of scRNA-seq technology allows us to directly investigate immune responses in the

ocular microenvironment, thus providing unprecedented opportunities to understand both immunopathogenesis

and the heterogeneity among patients [27, 28].

Our scRNA-seq of cells in aqueous humour from non-infectious uveitis patients confirmed a dominant presence

and pro-inflammatory phenotypes of T cells. More importantly, the analyses revealed that T cell phenotypes in

VKHD-associated uveitis were different from those in BD-associated uveitis. Uveitis in VKHD patients showed

the dominant role of clonally expanded effector CD4$^+$ T cells expressing *IFNG* and *TNF*. In contrast, uveitis in

the BD patient had few of these effector CD4$^+$ T cells but was characterised by clonally expanded effector CD8$^+$

T cells expressing *CCL5, GZMA, GZMB* and *PRF1*.

The results strongly suggest the distinctive roles of CD4$^+$ and CD8$^+$ T cells in the pathogenesis of non-infectious

uveitis in VKHD and BD. Such a notion is supported by genetic studies including genome-wide association

studies (GWAS). In multiple populations, BD was found associated with MHC class I-specific allele HLA-B*51,

while VKHD was associated with MHC class II-specific alleles HLA-DR4/HLA-DRB1 [29, 30]. The important

question following our study is what (auto)antigens drive the clonal expansion of effector CD4$^+$ T cells in VKHD

and the clonal expansion of effector CD8$^+$ T cells in BD, with ocular antigens as candidates. If such antigen can

be identified to specifically activate individual patient's T cell clones, the therapy to specifically induce the

tolerance to the antigen will be promising as the precedent clinical trial [31]. However, given that identical

clonotypes and similar clonotypes identified by CoNGA accounted for only half of effector CD4$^+$ and CD8$^+$ T

cells, our study suggests TCR-independent 'bystander' activation of T cells in both VKHD and BD-associated

uveitis. This should be taken into account for designing therapeutic strategies for non-infectious uveitis. Another

potential application derived from our study is to combine immunological signatures with genetic signatures to

stratify patients for therapies that selectively target CD4$^+$ or CD8$^+$ T cells.

Similar to two recent scRNA-seq studies of aqueous immune cells from 4-5 samples [27, 28], a major limitation of

our study was the limitation of sample numbers. Our results should be validated using a larger sample number.

Another limitation was the lack of paired blood samples. If the signatures of expanded T cell clonotypes could

be detected in blood samples, a strategy could be considered to monitor blood samples rather than invasive

ocular sampling.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (Grant No.81900849, 82070948,

82071792), Beijing Talent Project (No.2020027), Shunyi District "Beijing Science and technology achievements

transformation coordination and service platform" construction fund (SYGX202010) and Capital Health

Development Scientific Research Project Grant (to Y.T.). Natural Science Foundation of Shandong Province
(Major Basic Program, ZR2020ZD41 to M.S.). Australian National Health Medical Research Council Investigator
Fellowship (GNT2009554) and Bellberry-Viertel Senior Medical Research Fellowship (to D.Y.) and The
University of Queensland Postgraduate Scholarship (to H.S.).
298

299 **Author Contributions**

300 **Hao Kang:** Methodology, Resources, Writing – Original Draft **Hongjian Sun:** Methodology**,** Software, Formal
301 Analysis, Writing – Original Draft, Visualization **Yang Yang:** Methodology **Minglei Shu:** Supervision **Yunbo Wei:**
302 Methodology **Yu Zhang:** Supervision **Di Yu:** Conceptualization, Writing – Review & Editing, Supervision,
303 Funding Acquisition **Yong Tao:** Conceptualization, Writing – Review & Editing, Supervision, Funding Acquisition
304

305 **Additional Information**

306 Supplementary material can be accessed through the online version of this article.
307

308 **Competing interests**

309 The authors declare no competing interest.
310

Figure legends

Figure 1. T cells are the dominant immune cell type infiltrating in lesions of non-infectious uveitis

(a) Demographic and clinical information of VKHD and BD patients. **(b)** Schematics of the research design. **(c)** UMAP plot showing unsupervised clustering of cells in aqueous humour analysed by scRNA-seq (pooled data from three patients); subgraphs showing the gene expression of *CD3*, *CD14* and *CD19*. **(d)** UMAP plots showing single cells and clusters in samples from each patient sample (left panels) and frequencies of CD3⁺ T cells, CD19⁺ B cells and CD14⁺ monocytes (right panels).

Figure 2. T cell functional subsets in non-infectious uveitis differ between VKHD and BD.

(a) UMAP plot showing unsupervised clustering of CD3⁺ T cells (pooled from three patients); subgraphs showing the gene expression of *CD4* and *CD8* (both *CD8A* and *CD8B*). **(b)** Violin plots showing the expression of indicated TCR and co-receptor genes in each cluster. **(c)** UMAP plots showing single cells and clusters in samples from each patient (left panels) and frequencies of CD4⁺ T cells, CD8⁺ T cells and CD4⁻CD8⁻ (double negative, DN) T cells (right panels). **(d)** Heatmap showing top 5 differentially expressed genes in each cluster. **(e)** Violin plots of key marker genes for annotating clusters, with the annotation of clusters to the left. **(f)** Pie charts showing the frequencies of each T cell cluster in samples from individual patients.

Figure 3. *Divergent T cell clonal expansion in BD and VKHD uveitis*.

(a) UMAP plots visualising T cell clonal expansion (pooled data from three patients). *Large*: clonotypes > 5 % of total T cells from the same sample; *Medium*: clonotypes 1-5 % of total T cells from the same sample. *Small*: 0.2-1 % of total T cells from the same sample. **(b)** Treemap plots showing the frequencies of large and medium clonotypes (>1%) in each patient sample. **(c, d)** Contour plots overlayed with UMAP plots showing the distribution of T cells in large and medium clonotypes (left), frequencies of T cell clusters in each category of significantly expanded clonotypes (middle) and frequencies of significantly expanded clonotypes in each T cell cluster (right) for VKHD (c) and BD (d) patients. N.A., not available.

Figure 4. Similar T cell clonotypes in non-infectious uveitis.

(a) Schematics of clonotype neighbour graph analysis (CoNGA). **(b)** (Top) CoNGA graph-graph relationship plots showing GEX (gene expression) & TCR in pooled data from three patients and unveiling clonotype clusters with similar GEX and TCR. Each dot represents one clonotype. The GEX plot was generated from the gene expression of representative cells from each clonotype. The TCR plot was generated from the TCR distance for each clonotype. (Middle), the expression of feature genes. (Bottom), DEG and TCR sequence logos showing V and J gene usage and CDR3 sequences. DEG and TCR sequence logos are scaled by the adjusted P value of the associations. DEGs with fold-changes less than 2 are shown in grey. **(c)** Table showing the numbers of T cells belonging to the four biggest CONGA clusters. The CoNGA cluster name was based on its GEX (left) and TCR (right) clusters with a dash in the middle. **(d)** UMAP plots showing T cells identified in the top 4 biggest

349   CoNGA clusters in each patient sample. T cells belonging to individual CoNGA clusters were highlighted in

350   different colours.

References

[1]   L. Gamalero, G. Simonini, G. Ferrara, S. Polizzi, T. Giani, R. Cimaz. Evidence-Based Treatment for Uveitis. Isr Med Assoc J, 2019;21:475-9.

[2]   H. Yazici, E. Seyahi, G. Hatemi, Y. Yazici. Behcet syndrome: a contemporary view. Nat Rev Rheumatol, 2018;14:107-19.

[3]   Y. R. Hsu, J. C. Huang, Y. Tao, T. Kaburaki, C. S. Lee, T. C. Lin *et al.* Noninfectious uveitis in the Asia-Pacific region. Eye (Lond), 2019;33:66-77.

[4]   P. Yang, S. Liu, Z. Zhong, L. Du, Z. Ye, W. Zhou *et al.* Comparison of Clinical Features and Visual Outcome between Sympathetic Ophthalmia and Vogt-Koyanagi-Harada Disease in Chinese Patients. Ophthalmology, 2019;126:1297-305.

[5]   L. Du, A. Kijlstra, P. Yang. Vogt-Koyanagi-Harada disease: Novel insights into pathophysiology, diagnosis and treatment. Progress in Retinal and Eye Research, 2016;52:84-111.

[6]   B. Tong, X. Liu, J. Xiao, G. Su. Immunopathogenesis of Behcet's Disease. Frontiers in Immunology, 2019;10.

[7]   A. M. Bolger, M. Lohse, B. Usadel. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics (Oxford, England), 2014;30:2114-20.

[8]   Y. Hao, S. Hao, E. Andersen-Nissen, W. M. Mauck, S. Zheng, A. Butler *et al.* Integrated analysis of multimodal single-cell data. bioRxiv, 2020:2020.10.12.335331.

[9]   C. S. McGinnis, L. M. Murrow, Z. J. Gartner. DoubletFinder: Doublet Detection in Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. Cell Syst, 2019;8:329-+.

[10] C. Trapnell, D. Cacchiarelli, J. Grimsby, P. Pokharel, S. Li, M. Morse *et al.* The dynamics and regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells. Nat Biotechnol, 2014;32:381-6.

[11] A. Subramanian, P. Tamayo, V. K. Mootha, S. Mukherjee, B. L. Ebert, M. A. Gillette *et al.* Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. Proc Natl Acad Sci U S A, 2005;102:15545-50.

[12] N. Borcherding, N. L. Bormann, G. Kraus. scRepertoire: An R-based toolkit for single-cell immune receptor analysis. F1000Res, 2020;9:47.

[13] M. Krzywinski, J. Schein, I. Birol, J. Connors, R. Gascoyne, D. Horsman *et al.* Circos: an information aesthetic for comparative genomics. Genome Res, 2009;19:1639-45.

[14] T. Stuart, A. Butler, P. Hoffman, C. Hafemeister, E. Papalexi, W. M. Mauck *et al.* Comprehensive Integration of Single-Cell Data. Cell, 2019;177:1888-+.

[15] D. G. Pellicci, H.-F. Koay, S. P. Berzins. Thymic development of unconventional T cells: how NKT cells, MAIT cells and γδ T cells emerge. Nature Reviews Immunology, 2020;20:756-70.

[16] H. Y. Greenaway, B. Ng, D. A. Price, D. C. Douek, M. P. Davenport, V. Venturi. NKT and MAIT invariant TCRα sequences can be produced efficiently by VJ gene recombination. Immunobiology, 2013;218:213-24.

[17] D. Brandt, C. M. Hedrich. TCRαβ(+)CD3(+)CD4(-)CD8(-) (double negative) T cells in autoimmunity. Autoimmun Rev, 2018;17:422-30.

[18] Z. Chen, E. John Wherry. T cell responses in patients with COVID-19. Nature Reviews Immunology, 2020;20:529-36.

[19] G. S. Ling, G. Crawford, N. Buang, I. Bartok, K. Tian, N. M. Thielens *et al.* C1q restrains autoimmunity and
viral infection by regulating CD8(+) T cell metabolism. Science, 2018;360**:**558-63.
[20] J. He, X. Zhang, Y. Wei, X. Sun, Y. Chen, J. Deng *et al.* Low-dose interleukin-2 treatment selectively
modulates CD4(+) T cell subsets in patients with systemic lupus erythematosus. Nat Med, 2016;22**:**991-3.
[21] J. He, R. Zhang, M. Shao, X. Zhao, M. Miao, J. Chen *et al.* Efficacy and safety of low-dose IL-2 in the
treatment of systemic lupus erythematosus: a randomised, double-blind, placebo-controlled trial. Ann
Rheum Dis, 2020;79**:**141-9.
[22] C. H. Shim, S. Cho, Y. M. Shin, J. M. Choi. Emerging role of bystander T cell activation in autoimmune
diseases. BMB Rep, 2022;55**:**57-64.
[23] S. A. Schattgen, K. Guion, J. C. Crawford, A. Souquette, A. M. Barrio, M. J. T. Stubbington *et al.* Integrating
T cell receptor sequences and transcriptional profiles by clonotype neighbor graph analysis (CoNGA). Nat
Biotechnol, 2022;40**:**54-63.
[24] A. W. Taylor. Ocular immune privilege. Eye, 2009;23**:**1885-9.
[25] V. L. Perez, R. R. Caspi. Immune mechanisms in inflammatory and degenerative eye disease. Trends
Immunol, 2015;36**:**354-63.
[26] J. T. Rosenbaum, C. A. Harrington, R. P. Searles, S. S. Fei, A. Zaki, S. Arepalli *et al.* Identifying RNA
Biomarkers and Molecular Pathways Involved in Multiple Subtypes of Uveitis. Am J Ophthalmol,
2021;226**:**226-34.
[27] M. Paley, L. M. Hassman, E. Esaulova, G. L. Paley, J. Laurent, L. Springer *et al.* Highly expanded clones
representing different lymphocyte lineages are present in individual patients with granulomatous uveitis.
Investigative Ophthalmology & Visual Science, 2020;61**:**3661-.
[28] M. Kasper, M. Heming, D. Schafflick, X. Li, T. Lautwein, M. Meyer Zu Horste *et al.* Intraocular dendritic cells
characterize HLA-B27-associated acute anterior uveitis. Elife, 2021;10.
[29] M. Takeuchi, N. Mizuki, S. Ohno. Pathogenesis of Non-Infectious Uveitis Elucidated by Recent Genetic
Findings. Frontiers in Immunology, 2021;12.
[30] X. F. Huang, M. A. Brown. Progress in the genetics of uveitis. Genes Immun, 2022;23**:**57-65.
[31] R. B. Nussenblatt, I. Gery, H. L. Weiner, F. L. Ferris, J. Shiloach, N. Remaley *et al.* Treatment of uveitis by
oral administration of retinal antigens: results of a phase I/II randomized masked trial. Am J Ophthalmol,
1997;123**:**583-92.

Patient	VKHD1	VKHD2	BD
Age range at sample collection	31-35	36-40	36-40
Gender	female	male	female
Year range for Onset of disease	2015-2019	2005-2009	2015-2019
Sample collection	2019	2019	2019
Ocular Symptom	Decreased vision in bilateral eyes with accompanying blurred vision	Decreased vision in bilateral eyes	Decreased vision in left eye with accompanying red eye
Visual acuity	20/125 OD and 20/50 OS;	20/320 OD and 20/30 OS	No light perception OD and 20/32 OS
Uveitis status	Bilateral recurrent anterior uveitis (Mutton fat keratic precipitates, aqueoue flare 2+ and anterior chamber cells 3+ OU);	Bilateral recurrent anterior uveitis (Mutton fat keratic precipitates, aqueoue flare + and anterior chamber cells 2+ OU);	Bilateral recurrent anterior uveitis (fine, non-pigmented keratic precipitates OD and no KP was observed in OS, aqueoue flare 2+ and anterior chamber cells 3+ OU);
Ophthalmic Examination	Sunset glow fundus, Dalen-Fuchs nodules and retinal pigment epithelium clumping and migration.	Sunset glow fundus.	Optic atrophy, vascular occlusion and gliotic sheathing in OD; retinal vasculitis and retinal oedema in OS.
Surgery	Cataract surgery in bilateral eyes were done.	Cataract surgery in bilateral eyes were done.	Complicated cataract was observed in left eye, and cataract surgery in right eyes were done.
Systemic symptoms	Headache, tinnitus, neck stiffness, hypoacusis, alopecia and poliosis	Headache	Recurrent oral ulcerations, recurrent genital ulcerations and erythema nodosum

c

CONGA cluster (GEX-TCR)	BD T Cells (number)	VKHD1 T Cells (number)	VKHD2 T Cells (number)
0-10	0	21	5
0-3	0	9	5
4-2	0	18	0
5-7	15	0	0

d

VKHD1   VKHD2   BD

- 0-10
- 0-3
- 4-6
- 5-7