



The importance of vaccinated individuals to population-level evolution of pathogens

Maria A. Gutierrez^{1,*} , Julia R. Gog^{1,2,†} 

¹Department of Applied Mathematics and Theoretical Physics, University of Cambridge

²JUNIPER – Joint UNiversities Pandemic and Epidemiological Research

*mag84@cam.ac.uk (corresponding author), †jrg20@cam.ac.uk

This version: November 29, 2022

Abstract

Virus evolution shapes the epidemiological patterns of infectious disease, particularly via evasion of population immunity. At the individual level, host immunity itself may drive viral evolution towards antigenic escape. Using compartmental SIR-style models with imperfect vaccination, we allow for the probability of immune escape to differ in vaccinated and unvaccinated hosts. As the relative contribution to selection in these different hosts varies, the overall effect of vaccination on the antigenic escape pressure at the population level changes.

We find that this relative contribution to escape is important for understanding the effects of vaccination on the escape pressure and we draw out some fairly general patterns. If vaccinated hosts do not contribute much more than unvaccinated hosts to the escape pressure, then increasing vaccination always reduces the overall escape pressure. In contrast, if vaccinated hosts contribute significantly more than unvaccinated hosts to the population level escape pressure, then the escape pressure is maximised for intermediate vaccination levels.

We also find that these results depend on the vaccine efficacy against transmission, particularly through the partial protection against infection. This work highlights the potential value of understanding better how the contribution to antigenic escape pressure depends on individual host immunity.

Keywords: SARS-CoV-2, COVID-19, antigenic escape, vaccine escape pressure, phylodynamics

1 Introduction

Vaccines against infectious pathogens reduce incidence and deaths, although there is a risk that they could favour new antigenic variants [1]. For example, since population immunity against circulating SARS-CoV-2 variants controls virus spread [2], the long-term dynamics of the COVID-19 pandemic depend on the evolutionary trajectory of the virus [1]. Here we investigate how vaccination changes the appearance probability of strains capable of evading host immunity, relative to the baseline of no vaccination. Antigenic escape is complex, with strains generated and selected on various scales [3]. We focus exclusively on the *generation* of escape strains [4], [5], rather than on their establishment after they first appear [6], [7]. Our work builds on other mathematical studies of COVID-19 immune

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

escape risk [4], [5], [8]. We pay particular attention to selection in vaccinated hosts, relative to the unvaccinated.

Evolution of antigenic traits in SARS-CoV-2 (severe acute respiratory syndrome coronavirus 2) [9] generates new variants [10]. Some of these variants (B.1.1.7, *alpha*; B.617.2, *delta*) appear to have a transmission advantage over their precursors [11]. B.1.1.529 —*omicron*—significantly escapes existing immunity [12], but variants with immune escape, at least to some extent, date back to 2020, before the roll-out of vaccines (B.1351, *beta*; B.617.2, *delta*) [11]. Therefore it is important to understand how vaccination impacts antigenic drift. Vaccines may affect the evolution of other pathogen traits, such as virulence [13], but we focus on immune escape.

To the best of our knowledge, there is not yet evidence to assess how COVID-19 vaccines impact virus genomic diversity in an individual host [14]. Some COVID-19 models assume that escape occurs only or primarily in hosts with previous immunity [4], [5]. On the contrary, others assume that vaccination entirely prevents infection [8], and thus escape can only occur in unvaccinated hosts.

Models for pathogens other than SARS-CoV-2 often assume that viruses mutate at the same rate in all hosts [7], [15], [16], [17]. However, in human Influenza A, host immunity can select for antigenic mutations [18]. This effect may be important [19] for more ambitious vaccination campaigns, such as a universal influenza vaccine [20].

Given that we do not generally know the relative selective pressures exerted by infections in vaccinated hosts compared to the unvaccinated, we explore a range of values for this contribution to escape. We define a parameter (Section 2.1) that measures this relative contribution to escape. We use this parameter to express the escape pressure in terms of the infections in vaccinated and unvaccinated hosts, as in (1). We then combine this with deterministic compartmental models for vaccination (Section 2.2) that provides lifelong imperfect immunity. We explore two scenarios: a single wave and endemic disease. We calculate the resultant escape pressure in each of these contexts and describe how it changes as the vaccination coverage varies (Section 2.3). We find the trade-off between selection and the vaccine efficacies (VEs) in the escape pressure of a fully-vaccinated population and the vaccination coverage that maximises the escape pressure (Section 3).

We find that the relative selection strength from infections in vaccinated and unvaccinated hosts shapes the escape pressure. In particular, we find two different qualitative behaviours for how the escape pressure depends on the vaccination coverage. These two behaviours are separated by a threshold value of this relative contribution to escape, in which vaccinated hosts contribute individually somewhat more to escape than the unvaccinated. If the relative contribution to escape is below this threshold, vaccination always reduces antigenic escape, as in [15]. However, above this threshold, intermediate vaccination levels are the most likely to generate escape strains, as in [4]. We also find that the susceptibility reduction provided by the vaccines has a large effect on the escape pressure. The reduction in infectiousness is less important, but still lowers escape by reducing prevalence.

2 Methods

2.1 Measure of escape pressure

We take a simple approach building on [4] to study the generation of strains at the population-level. We assume that the escape pressure P depends linearly on the number of cases, as

$$P = N_U + \theta_E N_V \quad (1)$$

Underlying this factor is the assumption that each infection contributes very slightly to escape potential. We weight the (breakthrough) infections in vaccinated hosts N_V , relative to the infections in naïve

hosts N_U , by a factor θ_E . Only the relative value of P is important (see Section 2.3), so we only need the relative weighting θ_E amongst infection in the vaccinated and unvaccinated rather than their absolute contribute to escape. θ_E is comparable to the vaccine efficacies θ_* of [4] (see Section 2.2), in that it is a factor that reflects a change in a single host induced by vaccination. For example, if $\theta_E > 1$, vaccinated hosts infected with the resident strain would be more likely to pass on escape strains than the unvaccinated. Lower viral loads and shorter infections in the vaccinated [21] could lead to fewer mutations [22]. But in the vaccinated, these escape strains can avoid the vaccine-induced immunity that the original strain faces [23], and may be selected within a host enough to be transmitted to others [22]. There is no prior reason to exclude any value $\theta_E \geq 0$, so we explore the full range.

2.2 Epidemic models

We use systems of ordinary differential equations (ODEs) to generate N_U and N_V under various scenarios. We assume that the population is well-mixed and a single-strain infectious disease circulates unaffected by any escape strain it may have generated. We consider two epidemiological scenarios. In scenario (a) the outbreak is transient, which can be interpreted as a single epidemic wave. In scenario (b) the system is modified so that it reaches a non-zero equilibrium, corresponding to endemic disease.

We separate individuals by vaccination status and assume that vaccine immunity does not wane. We scale time so that the recovery rate is 1. In our models, the vaccines are not assumed to be perfect: vaccinated individuals can become infected, albeit at a lower rate, contributing to the escape pressure. We split the (imperfect) transmission blocking from vaccines into two components, θ_S and θ_I , as defined in [4]. θ_S is the susceptibility reduction (VE against infection): $\theta_S = 0$ corresponds to perfect protection, while $\theta_S = 1$ corresponds to no protection. Similarly, θ_I is the infectivity reduction (VE against transmission), if an infection occurs. For mathematical convenience, the reduction in susceptibility from vaccination is polarised [16]: some vaccinated individuals are entirely immune to the infection, while the rest are as susceptible as the unvaccinated. Appendix A.2 shows that the same qualitative results hold with an alternative formulation for immunity.

In both scenarios, S_U and S_V are the vaccinated and unvaccinated proportions of the population susceptible to the disease; I_U and I_V , the vaccinated or unvaccinated proportions that are infected; R_U and R_V , the vaccinated or unvaccinated recovered proportions. R_0 is the basic reproduction number and c is the proportion of the population who are vaccinated.

Scenario (a): transient epidemic

In the transient scenario, recovered hosts have full permanent immunity and the system is

$$\dot{S}_U = -R_0 S_U \lambda(t) \qquad \dot{S}_V = -R_0 S_V \lambda(t) \qquad (2)$$

$$\dot{I}_U = R_0 S_U \lambda(t) - I_U \qquad \dot{I}_V = R_0 S_V \lambda(t) - I_V \qquad (3)$$

$$\dot{R}_U = I_U \qquad \dot{R}_V = I_V \qquad (4)$$

with $\lambda(t) = I_U + \theta_I I_V$. The initial conditions are $S_U = 1 - c$, $S_V = c\theta_S$, $R_U = 0 = R_V$, and an infinitesimal number of infected individuals. The remaining $c(1 - \theta_S)$ proportion of the population has full immunity from vaccination, so they do not appear in the compartments of the system. Appendix A.1 shows that for polarised vaccine immunity—and non-assortative mixing as here—the original proportion between vaccinated and unvaccinated susceptibles is maintained and extended to the other compartments:

$$(S_V, I_V, R_V) = \frac{c\theta_S}{1-c} (S_U, I_U, R_U) \qquad (5)$$

This relation (5) reduces the system to:

$$\dot{S}_U = -S_U I_U R_0 \left(1 + \frac{c\theta_S \theta_I}{1-c}\right) \quad \dot{I}_U = +S_U I_U R_0 \left(1 + \frac{c\theta_S \theta_I}{1-c}\right) - I_U \quad (6)$$

$$S_U(0) = 1 - c \quad I_U(0) = \epsilon \ll 1 \quad (7)$$

where ϵ accounts for the initial cases. (6)-(7) describe a standard SIR model, up to rescaling the effective reproduction number and the population size. The initial effective reproduction number is $R_e = R_0[1 - c + c\theta_S \theta_I]$. If the vaccination coverage is above $\tilde{c} = (1 - R_0^{-1})/(1 - \theta_S \theta_I)$, herd immunity prevents an outbreak and prevalence decreases exponentially. From (6)-(7), if $c < \tilde{c}$, the cumulative infections in unvaccinated are $N_U = (1 - c)(1 + W(-R_e e^{-R_e})/R_e)$, where W is the Lambert-W function [24] (see Appendix A.1). The breakthrough infections are $N_V = c\theta_S(1 + W(-R_e e^{-R_e})/R_e)$. Thus, the escape pressure (1) is

$$P_a = (1 - c + \theta_S \theta_E c) \left(1 + \frac{1}{R_e} W(-R_e e^{-R_e})\right) \text{ if } R_e > 1, \text{ else zero} \quad (8)$$

Scenario (b): endemic disease

To achieve an endemic scenario, we consider two possible extensions of the SIR model (2)-(4) that replace susceptibles. If starting from the same initial conditions as in the transient scenario, these extensions reach an endemic state if $R_e > 1$. In both cases, we find the proportions I_U^*, I_V^* of infected individuals at the endemic equilibrium and use them to evaluate the escape pressure P_b at endemicity. Both models lead the same expression P_b for the resulting escape pressure, up to a constant of proportionality:

$$P_b = I_U^* + \theta_E I_V^* \propto (1 - c(1 - \theta_S \theta_E)) \left(1 - \frac{1}{R_e}\right) \text{ if } R_e > 1, \text{ else zero} \quad (9)$$

In the first extension, we assume that immunity from infections wanes at a constant rate ω , instead of being permanent. We still assume that vaccine immunity is lifelong. The modified ODEs are

$$\dot{S}_U = -R_0 S_U \lambda(t) + \omega R_U \quad \dot{S}_V = -R_0 S_V \lambda(t) + \omega R_V \quad (10)$$

$$\dot{I}_U = R_0 S_U \lambda(t) - I_U \quad \dot{I}_V = R_0 S_V \lambda(t) - I_V \quad (11)$$

$$\dot{R}_U = I_U - \omega R_U \quad \dot{R}_V = I_V - \omega R_V \quad (12)$$

with λ as before. We use (5), which still holds (see Appendix A.1), to find the endemic state (setting to zero the time-derivatives). For $R_e > 1$, there is an equilibrium state with $I_U^* = (1 - c)(1 - R_e^{-1})/(1 + \omega^{-1}) = (1 - c)I_V^*/c\theta_S$. Hence, for $R_e > 1$, the escape pressure rate at the endemic equilibrium is $P_b = I_S^* + \theta_E I_U^* = (1 - c(1 - \theta_E \theta_S)) [1 - 1/R_e] / (1 + \omega^{-1})$.

In the second extension, we include births and deaths, at the same homogeneous per capita rate μ , so that the population size is conserved. A proportion c of the population is vaccinated at birth. We assume that neither infection nor immunity are maternally transmitted. Therefore, the births appear in \dot{S}_U and \dot{S}_V as $\mu(1 - c)$ and $\mu c\theta_S$. The number of individuals V with full vaccine immunity obeys $\dot{V} = \mu c(1 - \theta_S) - \mu V$ with $V(0) = c(1 - \theta_S)$, so $V(t) = c(1 - \theta_S)$ at all times. The new ODEs are

$$\dot{S}_U = \mu(1 - c) - R_0 S_U \lambda - \mu S_U \quad \dot{S}_V = \mu c\theta_S - R_0 S_V \lambda - \mu S_V \quad (13)$$

$$\dot{I}_U = R_0 S_U \lambda - I_U - \mu I_U \quad \dot{I}_V = R_0 S_V \lambda - I_V - \mu I_V \quad (14)$$

$$\dot{R}_U = I_U - \mu R_U \quad \dot{R}_V = I_V - \mu R_V \quad (15)$$

with λ as before. Condition (5) still holds (see Appendix A.1). For $R_e > 1 + \mu$, there is an endemic state $I_U^* = \mu(1 - c) [1/(1 + \mu) - 1/R_e] = (1 - c)I_V^*/c\theta_S$. Hence $P_b = I_S^* + \theta_E I_U^* = (1 - c(1 - \theta_E\theta_S))\mu [1/(1 + \mu) - 1/R_e]$. For $\mu \ll 1$, which corresponds to an infectious period much shorter than the host's life expectancy, $P_b = \mu(1 - c(1 - \theta_E\theta_S)) [1 - R_e^{-1}]$ to leading order in μ .

2.3 Output metric

To study the escape pressure as a function of the vaccination coverage, we normalise the expressions (8) and (9) by their value at $c = 0$ (in the absence of vaccination):

$$\hat{P}_a(c) = \frac{P_a(c)}{P_a(0)} = (1 - c(1 - \theta_S\theta_E)) \frac{1 + W(-R_e e^{-R_e}) R_e^{-1}}{1 + W(-R_0 e^{-R_0}) R_0^{-1}} \quad \text{if } R_e > 1, \text{ else zero} \quad (16)$$

$$\hat{P}_b(c) = \frac{P_b(c)}{P_b(0)} = (1 - c(1 - \theta_S\theta_E)) \frac{1 - R_e^{-1}}{1 - R_0^{-1}} \quad \text{if } R_e > 1, \text{ else zero} \quad (17)$$

If $\hat{P}_*(c) > 1$, the vaccination coverage c increases the escape pressure from no vaccination.

category	item	description
vaccine parameters	θ_S	susceptibility reduction (VE against infection)
	θ_I	infectivity reduction (VE against transmission)
	θ_E	relative escape contribution of breakthrough infections
epidemiological parameters	R_0	initial basic reproduction number
	c	vaccination coverage in the population
	R_e	initial effective basic reproduction number
	\tilde{c}	vaccination coverage threshold for $R_e < 1$
escape pressure	P	unscaled escape pressure, as per (1)
	\hat{P}_a	normalized, in the transient (polarised) scenario
	$\hat{P}_{a'}$	normalized, in the transient "leaky" scenario (see A.2)
	\hat{P}_b	normalized, in the endemic (polarised) scenario
output parameters	c_m	vaccination level that maximises the escape pressure
	c_m^∞	vaccination level that maximises breakthrough cases
	$\hat{\theta}_E$	threshold value of θ_E , defined by $dP/dc = 0$ at $c = 0$
	$\hat{\theta}_S$	threshold value of θ_S , defined by $dP/dc = 0$ at $c = 0$
	$\hat{\theta}_I$	threshold value of θ_I , defined by $dP/dc = 0$ at $c = 0$

Table 1: Parameters and functions from our models that appear in the results section.

3 Results

First, we focus on the effects of θ_E on the escape pressure in a fully vaccinated population. We vary θ_S , θ_I and θ_E , and fix the vaccination coverage at $c = 1$. If vaccination does not prevent the epidemic, i.e., $R_0\theta_S\theta_I < 1$:

$$\hat{P}_a(1) = \theta_S\theta_E \left(\frac{1 + \frac{1}{R_0\theta_S\theta_I} W(-R_0\theta_S\theta_I e^{-R_0\theta_S\theta_I})}{1 + \frac{1}{R_0} W(-R_0 e^{-R_0})} \right), \quad \hat{P}_b(1) = \theta_S\theta_E \left(\frac{1 - \frac{1}{R_0\theta_S\theta_I}}{1 - \frac{1}{R_0}} \right) \quad (18)$$

and $\hat{P}_*(1) = 0$ for $R_0\theta_S\theta_I \geq 1$. Both expressions in (18) depend only on R_0 , $\theta_S\theta_I$ and $\theta_S\theta_E$, but this dependence changes for "leaky" immunity (see Appendix A.2). Figures 1a, S1a show how the escape

pressure in a fully vaccinated population $\hat{P}_*(1)$ depends on the vaccine parameters θ_S, θ_I and θ_E . As expected, it is unchanged relative to no vaccination ($\hat{P}_*(1) = 1$) when $\theta_E = 1 = \theta_S = \theta_I$, because then vaccines have no effect in our model. For a fixed VE, the escape pressure increases with $\theta_S\theta_E$, because escape becomes more likely in the vaccinated. Similarly, for fixed $\theta_S\theta_E$, the escape pressure decreases as the vaccines become more effective, because fewer infections reduce the opportunities for escape mutations. Therefore, the escape pressure in a fully-vaccinated population is less than in an unvaccinated population ($\hat{P}_*(1) < 1$) if the population is close enough to herd-immunity. As a consequence, $\theta_S\theta_E < 1$ is not necessary for $\hat{P}_*(1) < 1$. It is sufficient, because if $\theta_E\theta_S < 1$, vaccination blocks the expected escape so $\hat{P}_*(1) < 1$ unsurprisingly. Our results shows that vaccines can reduce the total escape pressure even if $\theta_S\theta_E > 1$.




Figure 1: (a): Escape pressure $\hat{P}_b(1)$ in a fully vaccinated population relative to an unvaccinated population. White dotted curve ($\hat{P}_b(1) = 1$): vaccine parameters that give the same escape as in an unvaccinated population. White dashed line: VE that prevents an epidemic in a fully vaccinated population (the horizontal axis is scaled to above this value). $R_0 = 1.4$. (b): Escape pressure $\hat{P}_b(c)$ for variable vaccination coverage c and relative contribution to escape θ_E from the vaccinated, relative to an unvaccinated population. White solid curve: the vaccination coverage, c_m , at which the escape pressure peaks, as a function of θ_E . White dotted curve ($\hat{P}_b(c) = 1$): combination of c and θ_E that give the same escape in a population as in an unvaccinated population. Dashed line: vaccination coverage for herd-immunity threshold (the horizontal axis only shows values of c below this). $\theta_S, \theta_I = 0.6$, $R_0 = 1.4$.

Next, we consider how the escape pressure changes with a variable vaccination coverage c , and how this is affected by θ_E . We keep θ_S and θ_I fixed for simplicity. Figures 1b, S1b show the escape pressure \hat{P}_* as a function of c and θ_E , for fixed VEs. Intuitively, the escape pressure decreases to zero as the vaccination coverage approaches \tilde{c} , the herd immunity threshold ($\tilde{c} < 1$ here). When vaccination is slightly below \tilde{c} , the outbreak is small (a few infections are enough to build herd immunity) so there are fewer opportunities for escape. It is helpful to consider the vaccination coverage that maximises the escape pressure, which we call c_m , as it depends on θ_E . If θ_E is below a threshold $\hat{\theta}_E$, then $c_m = 0$. The escape pressure simply decreases with c , because reducing cases outweighs the escape pressure from breakthrough cases. If $\theta_E > \hat{\theta}_E$, then $c_m > 0$. Breakthrough infections can drive evolution and reducing cases may not suffice to reduce the escape pressure, which has a unimodal shape. We can characterise these results analytically. Here we refer to the endemic scenario (Appendix A.1 discusses the transient scenario). Since $P_b(c)$ is concave ($d^2P_b/dc^2 < 0$), by (17), the

condition for a peak of the escape pressure at a non-zero vaccination level ($c_m > 0$) is

$$0 < \left. \frac{dP_b}{dc} \right|_{c=0} = \frac{1}{R_0} ((\theta_E \theta_S - 1)(R_0 - 1) - (1 - \theta_S \theta_I)) \iff \frac{1}{\theta_S} \frac{R_0 - \theta_S \theta_I}{R_0 - 1} =: \hat{\theta}_E \quad (19)$$

Figure S5 plots $\hat{\theta}_E$, as given by (19). Interpreting $1 - \theta_S \theta_I$ as the transmission-blocking and $\theta_E \theta_S - 1$ as the escape boost provided by the vaccines, the condition (19) can be rewritten as (escape boost) \times ($R_0 - 1$) $>$ (transmission-blocking). If (19) is satisfied, we solve $dP_b/dc = 0$ for the maximiser of the escape pressure $c_m = (1 - \theta_S \theta_I)^{-1} \left[1 - \sqrt{1 + (1 - \theta_S \theta_I)/(\theta_S \theta_E - 1)} / \sqrt{R_0} \right]$. As expected, c_m continuously increases with θ_E from zero (at $\theta_E = \hat{\theta}_E$). In the limit $\theta_E \rightarrow \infty$ only breakthrough infections contribute to P_b , the analytical expression for c_m tends to $(1 - 1/\sqrt{R_0}) / (1 - \theta_S \theta_I) =: c_m^\infty$. This number c_m^∞ can be obtained directly as the vaccination coverage that maximises the total number of infections in vaccinated hosts. If herd-immunity from vaccination alone is not possible ($\tilde{c} > 1$), Figure S6 shows that a fully vaccinated population ($c = 1$) can have the largest escape pressure. However, this does not happen ($c_m^\infty < 1$) if and only if $\sqrt{R_0} \theta_S \theta_I < 1$, which holds if the threshold coverage for $R_e < 1$ is less than one ($\tilde{c} < 1$), because then $R_0 \theta_S \theta_I < 1 < R_0$. Therefore, unsurprisingly, the escape pressure cannot peak at $c = 1$ if $\tilde{c} < 1$ (since $c = 1$ prevents the epidemic).




Figure 2: Relative escape pressure $\hat{P}_b(c)$ at fixed $R_0 = 1.4$ and a high relative contribution to escape from the vaccinated, $\theta_E = 10$. (a) has fixed $\theta_I = 0.6$ and the susceptibility $VE \theta_S$ varies. (b) has fixed $\theta_I = 0.6$ and the infectiousness $VE \theta_I$ varies. On both (a) and (b), the white solid curves mark the vaccination coverage, c_m , at which the escape pressure peaks; the dashed lines give the herd-immunity threshold. Considered together, (a) and (b) show that there is more variation in the relative escape pressure if θ_S varies and θ_I is fixed.

Finally, we study how each of θ_S , θ_I affects the escape pressure as the vaccination coverage varies. When infections in the vaccinated drive selection (θ_E is large), θ_S contributes more to the escape pressure than θ_I . Both reduce the total number of cases at the same rate, through $R_e = R_0(1 - c(1 - \theta_S \theta_I))$. Additionally, θ_S (partially) protects the vaccinated from infections, so it lowers the proportion of the total cases in the vaccinated. Hence, if $\theta_E > 1$, a vaccine that reduces susceptibility has a quantitatively stronger effect on the escape pressure than a vaccine that reduces infectivity. Figures 2, S2, S3, S4 demonstrate that the escape pressure is more sensitive to θ_S than θ_I . In particular, Figures 2, S2 show that a sufficiently low θ_S can lead to the escape pressure peaking at no vaccination ($c_m = 0$), while for the same background parameter values the escape pressure peaks at intermediate vaccination levels ($c_m > 0$), regardless of θ_I . We can define threshold values $\hat{\theta}_*$ similar to $\hat{\theta}_E$ (see

Figure 1b, S5), such that $\theta_* < \hat{\theta}_*$ is the condition for the escape pressure to peak at no vaccination ($c_m = 0$). Figures 2a, S2a, S3a all have $\hat{\theta}_S \in (0, 1)$, while $\hat{\theta}_I \in (0, 1)$ only appears in Figure S3b. In the endemic scenario, (19) gives $\hat{\theta}_S = R_0 / (\theta_E(R_0 - 1) + \theta_I)$ and $\hat{\theta}_I = (1 - (\theta_E\theta_S - 1)(R_0 - 1)) / \theta_S$. Therefore, as in Figure 2b, $\hat{\theta}_I$ can be negative which means that the escape pressure peaks at intermediate levels even if the vaccines fully block transmission ($\theta_I = 0$). Vaccinated hosts are still able to generate escape strains, even if they are not able to transmit the original strain. On the contrary, $\hat{\theta}_S$ is always positive, so if vaccines fully prevent infection ($\theta_S = 0$), vaccination always reduces the escape pressure ($c_m = 0$).

4 Discussion

4.1 Summary of findings

Introducing a new parameter, θ_E , we have explored the effect of the contribution to immune escape of cases in vaccinated hosts, relative to the unvaccinated. Taking a parsimonious modelling approach we have studied how the population escape pressure depends on the vaccination coverage, the vaccine efficacies and θ_E . We have found that θ_E is critical to determine how the escape pressure depends on the vaccination coverage. If θ_E is low, escape decreases with vaccination. As θ_E increases, so that vaccinated hosts contribute relatively more to escape, escape is largest at intermediate vaccination levels. Therefore, our results show that models of immune escape should take into account the relative contribution to escape of cases in hosts with imperfect immunity. Our results also show mathematically that vaccines which are effective in reducing transmission can generally reduce the population escape pressure, even if they favour escape strains in individual hosts.

4.2 Implications for modelling evolution with imperfect immunity

Our results highlight not only the importance of θ_E itself, but also the role in reducing escape of θ_S , the susceptibility vaccine efficacy (VE), especially when vaccinated hosts contribute relatively more to escape ($\theta_E > 1$). Our results show mathematically the need to consider separately the VEs against infection (susceptibility) and onwards transmission (infectiousness). A purely epidemic model which does not consider immune-induced escape only needs the overall transmission-blocking VE [4]. However, our escape pressure is more sensitive to a reduction in susceptibility than infectivity. The susceptibility VE not only reduces infections but specifically protects the vaccinated, who may exert more selection pressure. Traditional models assume that partial immunity confers only reduced susceptibility or infectivity [16]. Often the dynamics are qualitatively the same [25], so the choice of VE is usually based on tractability [16]. Our work suggests that evolutionary models should split infections by immune status, so that θ_E can be used, and consider the susceptibility VE separately from the transmissibility VE, so that θ_S and θ_E can regulate together the escape pressure. These points may also apply in multi-strain models where partial cross-immunity drives evolution [16]. Furthermore, the contribution to escape of an infected host may change with different vaccine types or number of doses, so along different VEs, different θ_E s may be needed. Parameters akin to θ_E could also be appropriate for other phylodynamic models.

4.3 Relation to existing results

As in previous work [8], [3], in our models intermediate vaccination levels could lead to the largest selection pressure for antigenic escape. The same conclusion holds for a wide range of θ_S and θ_I

values [4] when dividing the population in two subgroups with different contact rates. This population-level result is analogous [4] to the within-host phylodynamic trade-off between viral load and selection pressure [22]. Our works shows that a population-level trade-off can appear without mixing heterogeneity, stochasticity, or detailed within-host dynamics. Past work implicitly uses fixed choices of θ_E or the VEs (or both). We instead specify the region of $(\theta_E, \theta_S, \theta_I)$ parameter space in which intermediate vaccination levels maximise the escape pressure. Beyond this region, vaccination reduces the escape pressure, as in [15].

4.4 Limitations and further work

Due to the simplicity of our approach, many epidemiological and evolutionary processes could modify our results. We assume that there are no changing mitigation measures, behavioural changes or seasonality effects. Vaccination comes through a single dose of a unique vaccine, administered before the outbreak or at birth. Further work could consider ongoing vaccination, as in [8]. We also neglect the waning of vaccine immunity and assume complete immunity after infection. Infections in individuals with partial or waned immunity may contribute with a different weight to the escape pressure [5]. Heterogeneity in mixing, susceptibility and transmissibility could all play a role in evolution. It may be best to vaccinate first those with more contacts [4], or immunocompromised hosts, who might contribute significantly more to the escape pressure [26]. Moreover, unlike in COVID-19 [21], both scenarios explained here assume that vaccination does not change the recovery rate and that the disease is not fatal. Disease-induced mortality, different infectious periods and hospitalisations would break the proportionality (5), since these effects would be larger in the unvaccinated. Similarly, lifting the well-mixing assumption and including any age or spatial structure could change these patterns.

Our work ignores the invasion dynamics of escape strains [6], but simply focuses on the pressure to generate one. An escape strain might be unable to grow if few individuals remain susceptible. In the transient scenario, this could occur if a strains appears late in the epidemic; in the endemic scenario, if the invasion fitness of the escape strain is not large enough for it to replace or coexist with the resident strain. Similarly, stochasticity could drive escape strains extinct shortly after their generation. For example, [8] finds that escape strains appearing in lockdowns are unlikely to survive stochastic extinction. Simple approaches to consider the outcome of the new variant [8], [6] could be incorporated in our models, but ultimately this is a complex process [3] so a full description will require significant further development of models.

Acknowledgements: MAG is supported by the Gates Cambridge Trust (grant OPP1144 from the Bill & Melinda Gates Foundation).

Declaration of interests: We declare we have no competing interests.

References

- [1] Telenti A, Arvin A, Corey L, Corti D, Diamond MS, García-Sastre A, et al. After the pandemic: perspectives on the future trajectory of COVID-19. *Nature*. 2021.
- [2] Saad-Roy CM, Wagner CE, Baker RE, Morris SE, Farrar J, Graham AL, et al. Immune life history, vaccination, and the dynamics of SARS-CoV-2 over the next 5 years. *Science*. 2020.
- [3] Morris DH, Petrova VN, Rossine FW, Parker E, Grenfell BT, Neher RA, et al. Asynchrony between virus diversity and antibody selection limits influenza virus evolution. *eLife*. 2020.
- [4] Gog JR, Hill EM, Danon L, Thompson RN. Vaccine escape in a heterogeneous population: insights for SARS-CoV-2 from a simple model. *Royal Soc Open Sci*. 2021.

- [5] Saad-Roy CM, Morris SE, Metcalf CJE, Mina MJ, Baker RE, Farrar J, et al. Epidemiological and evolutionary considerations of SARS-CoV-2 vaccine dosing regimes. *Science*. 2021.
- [6] McLean AR. Vaccination, evolution and changes in the efficacy of vaccines: a theoretical framework. *Proc Royal Soc B*. 1995.
- [7] Gandon S, Day T. The evolutionary epidemiology of vaccination. *J R Soc Interface*. 2007.
- [8] Rella SA, Kulikova YA, Dermitzakis ET, Kondrashov FA. Rates of SARS-CoV-2 transmission and vaccination impact the fate of vaccine-resistant strains. *Sci Rep*. 2021.
- [9] Yewdell JW. Antigenic drift: Understanding COVID-19. *Immunity*. 2021.
- [10] Parums DV. Editorial: Revised World Health Organization (WHO) Terminology for Variants of Concern and Variants of Interest of SARS-CoV-2. *Med Sci Monit*. 2021.
- [11] Alizon S, Sofonea MT. SARS-CoV-2 virulence evolution: Avirulence theory, immunity and trade-offs. *J Evol Biol*. 2021.
- [12] Hu J, Peng P, Cao X, Wu K, Chen J, Wang K, et al. Increased immune escape of the new SARS-CoV-2 variant of concern Omicron. *Cell Mol Immunol*. 2022.
- [13] Gandon S, Mackinnon MJ, Nee S, Read AF. Imperfect vaccines and the evolution of pathogen virulence. *Nature*. 2001.
- [14] Kennedy DA, Read AF. Monitor for COVID-19 vaccine resistance evolution during clinical trials. *PLOS Biol*. 2020.
- [15] Wen FT, Malani A, Cobey S. The Potential Beneficial Effects of Vaccination on Antigenically Evolving Pathogens. *Am Nat*. 2022.
- [16] Gog JR, Grenfell BT. Dynamics and selection of many-strain pathogens. *Proc Natl Acad Sci USA*. 2002.
- [17] Boni MF, Gog JR, Andreasen V, Feldman MW. Epidemic dynamics and antigenic evolution in a single season of influenza A. *Proc Royal Soc B*. 2006.
- [18] Hensley SE, Das SR, Bailey AL, Schmidt LM, Hickman HD, Jayaraman A, et al. Hemagglutinin receptor binding avidity drives influenza a virus antigenic drift. *Science*. 2009 10;326(5953):734-6.
- [19] Kelvin AA, Falzarano D. The influenza universe in an mRNA vaccine. *Science*. 2022 11;378(6622):827-8.
- [20] Arevalo CP, Bolton MJ, Le Sage V, Ye N, Furey C, Muramatsu H, et al. A multivalent nucleoside-modified mRNA vaccine against all known influenza virus subtypes. *Science*. 2022 11;378(6622):899-904.
- [21] Kissler SM, Fauver JR, Mack C, Tai CG, Breban MI, Watkins AE, et al. Viral Dynamics of SARS-CoV-2 Variants in Vaccinated and Unvaccinated Persons. *N Engl J Med*. 2021.
- [22] Grenfell BT, Pybus OG, Gog JR, Wood JLN, Daly JM, Mumford JA, et al. Unifying the Epidemiological and Evolutionary Dynamics of Pathogens. *Science*. 2004.
- [23] Garcia-Beltran WF, Lam EC, St Denis K, Nitido AD, Garcia ZH, Hauser BM, et al. Multiple SARS-CoV-2 variants escape neutralization by vaccine-induced humoral immunity. *Cell*. 2021.
- [24] Lehtonen J. The Lambert W function in ecological and evolutionary models. *Methods Ecol Evol*. 2016.
- [25] Ferguson NM, Andreasen V. The Influence of Different Forms of Cross-Protective Immunity on the Population Dynamics of Antigenically Diverse Pathogens. In: *Mathematical Approaches for Emerging and Reemerging Infectious Diseases: Models, Methods, and Theory. The IMA Volumes in Mathematics and its Applications.* vol. 126. Springer, New York, NY; 2002. p. 157-69.
- [26] Choi B, Choudhary MC, Regan J, Sparks JA, Padera RF, Qiu X, et al. Persistence and Evolution of SARS-CoV-2 in an Immuno-compromised Host. *N Engl J Med*. 2020.

A Mathematical proofs

A.1 Polarised vaccine immunity

Using the notation of the main text, define column vectors $\mathbf{u}(t)$ and $\mathbf{v}(t)$ as

$$\mathbf{u}^T = (S_U(t), I_U(t), R_U(t)) \text{ and } \mathbf{v}^T = (S_V(t), I_V(t), R_V(t)) \quad (\text{A1})$$

where \mathbf{x}^T denotes the transpose of a vector \mathbf{x} . Consider an ODE system

$$\dot{\mathbf{u}} = \mathbf{M}\mathbf{u} \quad \dot{\mathbf{v}} = \mathbf{M}\mathbf{v} \quad (\text{A2})$$

for a time-dependent matrix \mathbf{M} , which possibly depends on \mathbf{u} and \mathbf{v} . Suppose initial conditions

$$\mathbf{v}(0) = \alpha\mathbf{u}(0) \quad (\text{A3})$$

for some constant α . $\hat{\mathbf{u}}(t) := \alpha^{-1}\mathbf{v}(t)$ and $\mathbf{u}(t)$ obey the same first order differential equation (A2) and initial condition $\mathbf{u}(0) = \hat{\mathbf{u}}(0)$. Hence (by uniqueness of solutions), $\hat{\mathbf{u}}(t) = \mathbf{u}(t)$. Thus $\mathbf{v}(t) = \alpha\mathbf{u}(t)$.

Analytic results for the transient polarised model

The system (2)-(4) can be written in the form (A1)-(A3) with $\mathbf{M}(t) = \begin{pmatrix} -R_0\lambda(t) & 0 & 0 \\ R_0\lambda(t) & -1 & 0 \\ 0 & +1 & 0 \end{pmatrix}$ and $\alpha = c\theta_S/(1-c)$. Thus $(S_V(t), I_V(t), R_V(t)) = c\theta_S(1-c)^{-1}(S_U(t), I_U(t), R_U(t))$. This relation (5) allows

us to derive (6), which yields

$$\frac{\dot{I}_U}{\dot{S}_U} = \frac{dI_U}{dS_U} = -1 + \frac{1}{R_0 S_U} \left(1 + \frac{c\theta_S\theta_I}{1-c}\right)^{-1} \implies I_U = (1-c) - S_U - \left(1 + \frac{c\theta_S\theta_I}{1-c}\right)^{-1} \frac{1}{R_0} \log \frac{1-c}{S_U} \quad (\text{A4})$$

where the initial conditions (7) set the integration constant. The final size of the susceptible population, $S_U^\infty = \lim_{t \rightarrow \infty} S_U(t)$, is given implicitly by taking $t \rightarrow \infty$ and setting $0 = \lim_{t \rightarrow \infty} I_U(t)$:

$$S_U^\infty = (1-c) - \left(1 + \frac{c\theta_S\theta_I}{1-c}\right)^{-1} \frac{1}{R_0} \log \frac{1-c}{S_U^\infty} = -\frac{1-c}{R_e} W(-R_e e^{R_e}) \quad (\text{A5})$$

The second equality expresses S_U^∞ in terms of the Lambert-W function [24] and R_e . From S_U^∞ we obtain (8) for the escape pressure $P_a(c)$, which we study here. For simplicity, we set $W \equiv W(-R_e e^{-R})$ and $R \equiv R_e$.

$$\frac{dP_a}{dc} + (1 - \theta_E\theta_S) \left(1 + \frac{1}{R} W(-R_e e^{-R})\right) = -R_0(1 - \theta_S\theta_I)(1 - c(1 - \theta_S\theta_E)) \frac{d}{dR} \frac{W(-R_e e^{-R})}{R} \quad (\text{A6})$$

$$= R_0(1 - \theta_S\theta_I)(1 - c(1 - \theta_S\theta_E)) \left[\frac{W}{R^2} + \frac{1}{R} \frac{d}{dR} (-R_e e^{-R}) \frac{W}{(-R_e e^{-R})(1+W)} \right] \quad (\text{A7})$$

$$= WR_0(1 - \theta_S\theta_I)(1 - c(1 - \theta_S\theta_E)) \left[\frac{1}{R^2} + \frac{1}{R} \frac{-e^R + R e^{-R}}{R e^{-R}(1+W)} \right] \quad (\text{A8})$$

$$= WR_0(1 - \theta_S\theta_I)(1 - c(1 - \theta_S\theta_E)) \frac{1}{R^2} \frac{W+R}{1+W} \implies \quad (\text{A9})$$

$$\frac{dP_a}{dc} = (1 + W/R) \left[-(1 - \theta_E\theta_S) + (1 - \theta_S\theta_I) \frac{1 - c(1 - \theta_S\theta_E)}{1 - c(1 - \theta_S\theta_I)} \frac{W}{1+W} \right] \quad (\text{A10})$$

See, e.g., [24] for the derivative of the Lambert-W function used in (A7). We prove that there is either none or one root ($c = c_m$) solving $dP_a/dc = 0$ in $c \in [0, \tilde{c})$. $W \geq -1$, and hence $(1 + W/R) > 0$, because $R > 1$ for $c < c_m$. Thus it suffices to consider the sign and monotonicity of the bracketed (right-most) term in (A10). First suppose $(1 - \theta_S\theta_E) > 0$, which means $(1 - c(1 - \theta_S\theta_E)) > 0$. Since $W/(W+1) < 0$, $dP_a/dc < 0$. So if $\theta_E < 1/\theta_S$, vaccination reduces the escape pressure regardless of the coverage c . Now consider $\theta_E\theta_S > 1$. R decreases with c and $R > 1$ for $c < \tilde{c}$. $z := -R_e e^{-R} < 0$ increases with $R > 1$, so z decreases with $c \in (0, \tilde{c})$. The Lambert W function $W(z)$ increases with its argument $z \in (-1/e, 0)$, so it decreases with c . $W/(1+W)$ is an increasing function of $W \in (0, -1)$ and hence it is a negative, decreasing function of c . These imply

$$\frac{d}{dc} \left[\frac{1 + c(\theta_S\theta_E - 1)}{1 - c(1 - \theta_S\theta_I)} \frac{W}{1+W} \right] = - \left| \frac{W}{W+1} \right| \frac{d}{dc} \left(\frac{1 + c(\theta_S\theta_E - 1)}{1 - c(1 - \theta_S\theta_I)} \right) + \frac{1 + c(\theta_S\theta_E - 1)}{1 - c(1 - \theta_S\theta_I)} \frac{d}{dc} \left(\frac{W}{1+W} \right) \quad (\text{A11})$$

where the first derivative in the RHS is positive and the second is negative. Hence the LHS is negative. Thus dP_a/dc is a decreasing function of c , with at most one root, c_m . The if and only if condition for $c_m > 0$ is $dP_a/dc > 0$ at $c = 0$. Evaluating (A10) at $c = 0$, the condition for $c_m > 0$ becomes

$$\theta_E\theta_S - 1 > (1 - \theta_S\theta_I) \frac{-W_0}{W_0 + 1} =: \hat{\theta}_E\theta_S - 1 \quad \text{with } W_0 \equiv W(-R_0 e^{-R_0}) \quad (\text{A12})$$

(A12), as (19), can be written in the form (escape boost) $> g(R_0)$ (transmission blocking) for a function g . Here $g(R_0) = -W_0/(1+W_0)$; while in the endemic scenario, $g(R_0) = 1/(R_0 - 1)$.

Analytic results the endemic models

The SIRS system (10)-(12) can be written in the form (A1)-(A3) with $\alpha = \frac{c\theta_S}{1-c}$ and $\mathbf{M}(t) = \begin{pmatrix} -R_0\lambda(t) & 0 & \omega \\ R_0\lambda(t) & -1 & 0 \\ 0 & +1 & -\omega \end{pmatrix}$.

Thus, (5) holds. We write the system with natural dynamics (13)-(15) as $\dot{\mathbf{u}} = \mathbf{x}_U + \mathbf{M}(t)\mathbf{u}$, $\dot{\mathbf{v}} = \mathbf{x}_V + \mathbf{M}(t)\mathbf{v}$, with $\mathbf{M}(t) = \begin{pmatrix} -R_0\lambda(t) - \mu & 0 & 0 \\ R_0\lambda(t) & -1 - \mu & 0 \\ 0 & +1 & -\mu \end{pmatrix}$ and $\mathbf{x}_U^T = (\mu(1-c), 0, 0) = \frac{1}{\alpha}\mathbf{x}_V^T$, $\mathbf{v}(0) = \alpha\mathbf{u}(0)$ for $\alpha = c\theta_S/(1-c)$. Both $\mathbf{u}(t)$, $\hat{\mathbf{u}}(t) := \alpha^{-1}\mathbf{v}(t)$ solve the initial value problem $\dot{\mathbf{y}}(t) = \mathbf{F}(t, \mathbf{y}(t)) = \mathbf{M}(t)\mathbf{y}(t) + \mathbf{x}_U$, $\mathbf{y}(0) = \mathbf{u}(0)$. By uniqueness of solutions, $\mathbf{u}(t) = \hat{\mathbf{u}}(t)$, and hence (5) still holds.

A.2 Analytical results for leaky vaccine immunity

We modify the SIR model of Section 2.2 to have “leaky” immunity, instead of polarised [16]:

$$\dot{S}_U = -R_0 S_U \lambda(t) \qquad \dot{S}_V = -R_0 S_V \theta_S \lambda(t) \qquad (\text{A13})$$

$$\dot{I}_U = R_0 S_U \lambda(t) - I_U \qquad \dot{I}_V = R_0 S_V \theta_S \lambda(t) - I_V \qquad (\text{A14})$$

$$S_U(0) = 1 - c \gg I_U(0) \qquad S_V(0) = c \gg I_V(0) \qquad (\text{A15})$$

For $c = 1$ or $c = 0$, (A13)-(A15) become a standard SIR model: $\dot{S} = -R_0 S I$, $\dot{I} = R_0 S I - I$, $S(0) = 1 \gg I(0) > 0$ if $c = 0$, and the same but with $R_0 \rightarrow R_0 \theta_S \theta_I$ for $c = 1$. We assume $R_0 \theta_S \theta_I, R_0 > 1$ so that the outbreak grows in both situations. The final sizes are

$$N_U = 1 + \frac{1}{R_0} W(-R_0 e^{-R_0}) \qquad N_V = 1 + \frac{1}{R_0 \theta_S \theta_I} W(-R_0 \theta_S \theta_I e^{-R_0 \theta_S \theta_I}) \qquad (\text{A16})$$

respectively, where W is the Lambert-W function [24]. From (A16) and (18) we find that the relative escape pressure in a fully-vaccinated population under the “leaky” immunity assumption is $\hat{P}_{a'}(1) := \theta_E N_V / N_U = \theta_S^{-1} \hat{P}_a(1)$, proportional to the relative escape pressure under polarised immunity. Figures S1a, S1c are hence identical, except for the vertical variables, θ_E and $\theta_S \theta_E$.

For $c \in (0, 1)$, from (A13), $\frac{d}{dt} \log S_V = \theta_S \frac{d}{dt} \log S_U$ and so $S_V(t) = c(S_U(t)/(1-c))^{\theta_S}$, which deviates from (5). However, we can still study the system through $S_U(t)$ and $\lambda(t) = I_S + \theta_I I_V$:

$$\dot{S}_U = -R_0 S_U \lambda, \qquad \dot{\lambda} = R_0 [S_U + \theta_S \theta_I S_V] \lambda - \lambda = R_0 \left[S_U + \theta_S \theta_I \frac{c}{(1-c)^{\theta_S}} S_U^{\theta_S} \right] \lambda - \lambda \qquad (\text{A17})$$

Using the chain rule and the initial conditions (A15), $d\lambda/dS_U = -1 - S_U^{\theta_S-1} c \theta_S \theta_I / (1-c)^{\theta_S} + 1/R_0 S_U$

$$\implies \lambda(t) = 1 - c - S_U + \frac{c\theta_I}{(1-c)^{\theta_S}} [(1-c)^{\theta_S} - S_U^{\theta_S}] - \frac{1}{R_0} \log \frac{1-c}{S_U} \qquad (\text{A18})$$

Setting $\lim_{t \rightarrow \infty} \lambda(t) = 0$, (A19) defines $S_U^\infty = \lim_{t \rightarrow \infty} S_U < 1 - c$, the final size of the unvaccinated susceptible compartment:


$$S_U^\infty = 1 - c + \frac{c\theta_I}{(1-c)^{\theta_S}} [(1-c)^{\theta_S} - (S_U^\infty)^{\theta_S}] - \frac{1}{R_0} \log \frac{1-c}{S_U^\infty} \qquad (\text{A19})$$

$N_U(c) = 1 - c - S_U^\infty$ and $N_V(c) = c - \lim_{t \rightarrow \infty} S_V(t) = c \left[1 - (S_U^\infty / (1-c))^{\theta_S} \right]$ so, for $0 < c < 1$,


$$P_{a'}(c) = N_U(c) + \theta_E N_V(c) = \left[1 - c(1 - \theta_E) - S_U^\infty - c\theta_E \left(\frac{S_U^\infty}{1-c} \right)^{\theta_S} \right] \qquad (\text{A20})$$

and where S_U^∞ is the unique solution in $(0, 1 - c)$ of (A19) for $R_e > 1$. For $R_e \leq 1$, we set $P_{a'} = 0$. Figures S1b, S1d are the equivalent of Figure 1b for the “leaky” and polarised transient models. The effect of θ_E is qualitatively the same in all models, regardless of vaccination coverage.


S Supplementary Figures




(a) Relative escape pressure $\hat{P}_\alpha(1)$.



(b) Relative escape pressure $\hat{P}_\alpha(c)$.



(c) Relative escape pressure $\hat{P}_{\alpha'}(1)$.



(d) Relative escape pressure $\hat{P}_{\alpha'}(c)$.

Figure S1: As Figure 1 but for the transient scenario. The qualitative behaviour is the same as in the endemic scenario. Top row is with polarised immunity, bottom row is for leaky immunity. The escape pressure is larger with leaky immunity instead of polarised immunity (as seen in the different variables in the vertical axis of (c) or the maximum values of \hat{P}_* attained in (d)).




Figure S2: As Figure 2, but for the transient scenario with polarised immunity. The qualitative behaviour is the same as in the endemic scenario. The same behaviour appears in the transient scenario with leaky immunity (plots not shown).




Figure S3: As Figure 2, but with a lower relative contribution to escape from vaccinated hosts: $\theta_E = 5$ instead of $\theta_E = 10$. Here vaccination that fully blocks infection ($\theta_S = 0$) or onward transmission ($\theta_I = 0$) reduces the escape pressure at any vaccination coverage ($\hat{P}_b(c) < 1$ for $c > 0$). In other words, here $\hat{\theta}_I > 0$. However, the escape pressure is still more sensitive to θ_S than θ_I . The same behaviour appears in the transient scenario with leaky or polarised immunity (plots not shown).




Figure S4: Relative escape pressure $\hat{P}(0.1)$ as a function of the vaccine efficacies θ_S and θ_I for a fixed vaccination coverage $c = 0.1$. (a) is for the transient scenario and (b) is for the endemic scenario: the qualitative results are the same. As expected, reducing θ_S or θ_I decreases the escape pressure, because this corresponds to more effective vaccines. However, the contour lines that gives constant escape pressures are roughly vertical, meaning that the escape pressure decreases is much more sensitive to θ_S than θ_I . As in Figures 2, S2, $R_0 = 1.4$ and $\theta_E = 10$. The same behaviour appears in the transient scenario with leaky immunity (plots not shown).




Figure S5: $\hat{\theta}_E$ (the value of the relative contribution to escape from the vaccinated above which intermediate coverages maximise the escape pressure) in the endemic scenario, as given by (19). We only plot values of the susceptibility VE θ_S above 0.4, because $\hat{\theta}_E$ quickly grows without bounds as $\theta_S \rightarrow 0$ (the vaccines become better at preventing infection). In other words, $\hat{\theta}_E$ is more sensitive to θ_S than θ_I , as \hat{P} (see, e.g., Figure S4). $R_0 = 1.4$.




Figure S6: As Figures 1b, S1b, but with different R_0 such that vaccination alone cannot achieve herd-immunity ($\tilde{c} > 1$). (a), (b): $R_0 = 2.8$. The escape pressure is non-zero even in a fully vaccinated population ($c = 1$), because there is still transmission. Nevertheless, the vaccination coverage that maximises the escape pressure is always less than 1, so the escape pressure does not peak at a fully vaccinated population regardless of how much cases in vaccinated individuals contribute to the escape pressure ($c_m < 1$). $c_m^\infty < 1$ is the vertical asymptote of the white curve, the limit of c_m as $\theta_E \rightarrow \infty$. (c), (d): $R_0 = 9$. For large θ_E , the escape pressure increases monotonically with the vaccination coverage c and is largest at a fully-vaccinated population ($c = 1$). As with the other plots of this paper, the qualitative behaviour of the transient (left column) and endemic (right column) scenarios is the same. We do not show plots for the transient scenario with leaky immunity, but the qualitative behaviour is the same.