

Knowledge, Attitude and Practice (KAP) Survey about hepatitis B (HBV) and C (HCV) among migrant populations from Sub-Saharan Africa

Frhat M. A. Saaed¹

¹Department of Zoology, College of Arts and Sciences, Benghazi University, Al Kufra, Libya

Jerry E. Ongerth^{2*}

²Environmental Engineering, University of Wollongong, Wollongong, NSW, 2522, Australia
jongerth@uow.edu.au; +61 0408548427

Muna H. Ali³

³Department of Mathematics, Faculty of Arts & Science Kufrah, Benghazi University, Libya.

Abstract:

Background: Hepatitis B and C virus infections are serious blood borne diseases with global health concern particularly in developing countries. The association between survey participant's knowledge, attitude, and practices (KAP) about HBV and HCV infection is rarely studied, particularly among migrants seeking resettlement. Libya is a main transient station for migrants northward toward Europe and the flow is increasing into Al Kufra from neighboring countries that have endemic intermediate and high HBV and HCV prevalence. The purpose of this exploratory descriptive study in Al Kufra was to assess the level of participating migrant's KAP, regarding cause, transmission, prevention, and treatment of viral hepatitis due to HBV and HCV. **Method:** A cross-sectional study was conducted among 674 sub-Saharan African migrants in Al Kufra, Libya from January to October 2021. Migrant's KAP about HBV and HCV infection was assessed by interview questionnaire. Statistical and data analysis used SPSS version of 25. **Results:** Out of 700 questionnaires distributed, 674 were returned with a response rate of 96.3%. The study population included 580 (86.1%) males, mean age of 29.6 ± 7.2 SD years. A majority, 508 (75.4%) were illiterate and elementary level, 8 2.483+.232, respectively. Positive significant linear correlations were found between knowledge-attitude ($r = 0.524$, $p < 0.01$) knowledge-practice ($r = 0.123$, $p < 0.01$) and attitude-practice ($r = 0.278$, $p < 0.01$). Ethnicity and education variables were significantly associated with mean KAP. Gender identity and marital status were both significantly associated with mean knowledge and attitude. **Conclusion:** The migrant population from the neighboring and sub-Saharan African countries have inadequate KAP about HBV and HCV to limit infection.

Keywords: Knowledge, Attitude, Practice, hepatitis B, hepatitis C, Migrant, sub-Saharan

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Introduction: 37

Viral hepatitis B and C are blood borne viruses that infect millions of people worldwide. 38

Infections with hepatitis B virus (HBV) and C (HCV) virus are leading causes of chronic liver 39
disease, cirrhosis, liver cancer and associated morbidity and mortality worldwide and are a 40
primary indication for liver transplantation [19]. Chronic HBV and HCV infections are 41
responsible for nearly 57% of liver cirrhosis cases (HBV 30%, HCV 27%) and 78% of 42
hepatocellular carcinomas (HBV 53%, HCV 25%), the third most common cause of cancer 43
deaths worldwide [12, 24, 25]. 44

The endemicity of HBV is high in developing areas, including Africa, the Middle East and 45
Southeast Asia, where at least 8% of the population are HBV chronic carriers [39, 24]. The 46
largest burden of morbidity and mortality from chronic liver disease continues to be in nations 47
of the developing world. For example, 5–10% of the adult population in East Asia and sub- 48
Saharan Africa are estimated to have chronic HBV infection [28]. The highest reported 49
prevalence of chronic HCV is in Africa. Egypt has the highest prevalence of HCV, with some 50
studies reporting HCV antibody positive rates of up to 15%, with an estimated 10% with 51
chronic viremia [14, 11]. Viral hepatitis disproportionately affects sub- Saharan Africans 52
populations [10, 13, 31, 34, 41, 42] with high rates of migration from countries where 53
childhood HBV vaccination has only recently been implemented [15]. 54

The hepatitis B vaccine is the mainstay of HBV prevention [38]. No vaccine is available for 55
HCV, dictating the importance of education and precautions [38]. These viruses are blood- and 56
bodily fluid-borne, transmitted most commonly through sharing of injection equipment, reuse 57
or poorly sterilized of medical equipment, especially syringes and needles and sharp objects, 58
sexual contact, and mother to child transmission [4, 16, 21]. 59

Countries of high HBV & HCV prevalence are often economically disadvantaged and areas of 60
conflict that are a source of migrants seeking stable high-income regions including Europe and 61

North America. This poses challenges to the public health and immigration systems in the host nations [28] and countries along migration routes [26]. In 2020 the total of international migrants was 280.6 million, making this group the 5th most populous nation [38; 22]. Many migrants move into and remain part of ethnic minority groups maintaining traditional social and cultural behaviors that are reinforced and persist and may have adverse implications for exposure to HBV and HCV [30].

Countries such as Libya along migration routes from countries with high prevalence of viral hepatitis, e.g., sub-Saharan African countries have special problems. Migrants typically suffer from infectious diseases that are more prevalent in countries of origin [14, 26]. Migrants from neighboring and sub-Saharan African countries have been cited as one of the major factors spreading the chronic liver disease and hepatocellular carcinoma with risks of disease transmission to the local population [8, 29]. Lack of access to health care services, lack of basic knowledge, poor to nonexistent personal hygiene, and the inability to obtain information about the transmission and prevention all contribute to the importance of defining issues leading to improvement.

The knowledge, attitudes, and practices (KAP) about viral hepatitis transmission, prevention, and liver disease progression among immigrant populations from the neighboring and sub-Saharan African countries is likely to have significant bearing on transmission. Studies show that awareness of chronic HBV or HCV in most populations is low but that among migrant populations it tends to be even lower [30; 36]. A combination of factors contributes to disease propagation, including poor knowledge of the diseases, their risk factors and symptoms, lack of access to healthcare and health information, stigma associated with disease, as well as the lack of symptoms from the early stages of liver disease such as viral hepatitis [30; 33; 34]. General knowledge in immigrant populations is not adequate to avoid blood borne diseases. Social and cultural norms, unsafe sexual intercourse, exposure to the blood or body fluids of

an infected person, and intravenous drug use may increase the risk of HBV and HCV infection. 87
Furthermore, minimal education and resources, isolation, and stress contribute the refugees 88
tendency to engage in risky behavior, increasing the risk of these infections [28]. The purpose 89
of this study is to determine the state of knowledge, attitudes, and practices related to HBV and 90
HCV, transmission, consequences, treatment and prevention. 91

Methods:

 92

The study was conducted from January 1 to October 31, 2021 at the Libyan Red Crescent, Al 93
Kufra branch laboratory in a high migrant area. The study area is located in the southeast of 94
Libya having borders with Egypt to the East and with Sudan and Chad to the South. It is 1700 95
km from Tripoli, the capital city of Libya and 900 km South of the nearest Mediterranean port. 96
The study population of migrants attending at the Libyan Red Crescent Al Kufra branch 97
laboratory included 674 apparently healthy males and females, aged 21–52 years. Knowledge, 98
attitudes, and practices related to HBV and HCV infection were assessed by questionnaire 99
specially designed to permit literacy-independent answers. Questions concerned awareness, 100
beliefs, and attitudes focusing on the causes, transmission, prevention, and treatment of viral 101
hepatitis (Appendix 1-3). Subject categories included behaviors contributing to infection risk 102
with both HBV and HCV; sociodemographic data; and practices of the study population 103
regarding viral hepatitis. A pre-designed, structured questionnaire sheet was completed inside 104
the laboratory by direct personal interview with each migrant individually. The questionnaire 105
was developed and adapted through extensive literature review in English language [1; 5; 20; 106
31; 40]. The questions in knowledge section were further divided into 3 main sections (see 107
Appendix A): the first 19 knowledge questions were general on viral hepatitis diseases; seven 108
significant viral hepatitis transmission risk factors were included in question 20. Question 21 109
includes eight common complications of HBV and HCV infections. The attitude and practice 110

section of questionnaire consist of 13 and 12 questions, respectively. The demographic items 111
included age, gender, education level, and ethnicity. 112

Statistical Analysis: 113

Data were analyzed using Statistical Package for Social Sciences (SPSS) v. 25.0. The relationship 114
between independent categorical variables and the main outcomes of the study (knowledge, attitude, 115
and practice related to transmission and prevention of HBV and HCV infection) were tested using Chi- 116
square test. The p-value < 0.05 was considered to be significant. Descriptive statistics were used to 117
illustrate migrants' demographic characteristics. Categorical variables were measured as 118
percentages while continuous variables were expressed as mean \pm standard deviation. 119
Normality of data was tested using Kolmogorov-Smirnov test. Inferential statistics involving 120
Chi-square test, Mann-Whitney U test, Kruskal Wallis H test were used to assess the difference 121
while Spearman's rank correlation coefficient was used to evaluate the relationship between 122
the study variables. Cronbach's alpha was calculated as an indication of internal consistency. 123

Ethics Approval: 124

Written informed consent was obtained from the study participants after they have been clearly 125
briefed about the objective of the study. They were informed about their right to refuse to 126
participate in the study. 127

Results: 129

Descriptive statistics for each item in the questionnaire are given in Appendix Tables A1, A2, 130
and A3. A total of 674 African migrants from sub-Saharan Africa, 43.8% Eritrean, 26.6%, 131
Somalian, 21.7%, Ethiopian, and 8% Sudanese, participated in this study (Table 1). Among the 132
674 participants, 87.5% were married, 86.1% were male and 13.9% female. The age range was 133
21–52 years old with a mean age of 29.6 years. A majority of the participants, 75.4%, were 134

illiterate or had a basic education level, while 24.6% of participants had a high school level or above (Table 1).

Table 1. Sub-Saharan African migrant study population demography (n=674)

Migrants Characteristics	N (%)
Age (Mean ± SD)	29.6 ± 7.2
Gender identity	
Male	580 (86.1%)
Female	94 (13.9%)
Ethnicity	
Eritrean	295 (43.8%)
Somalian	179 (26.6%)
Ethiopian	146 (21.7%)
Sudanese	54 (8%)
Marital Status	
Single	84 (12.5%)
Married	590 (87.5%)
Education level	
Illiterate/Basic education	508 (75.4%)
High school and above	166 (24.6%)

Of the 700 distributed questionnaires, 674 were returned with a response rate of 96.3%. The Kolmogorov-Smirnov test revealed non normal distribution of the data. The Cronbach's alpha of the questionnaire was 0.76 for individual domains, maximum Cronbach's alpha was for attitude section followed by practices and knowledge section. The reason for low Cronbach's alpha for knowledge domain might be due to the number of questions as compared to attitude and practices sections.

Overall, knowledge regarding HBV and HCV was poor, Table 1a. more than 40% of participants answered Don't know to the 37 questions reflecting understanding. Most (34 of 37) questions had an answer either correct or not, of these, only 16% of migrants' answers were correct. For questions of overall indicative knowledge, more than 8 out of 10 migrants, 83.0%, had never heard of viral hepatitis. Less than 1 out of 10 migrants, 7.9%, could identify hepatitis B and C as viral infections (Table 2a). More than 90% of participants had no knowledge about the transmission of HBV and HCV and did not know that a simple test could identify HBV and HCV infections.

Table 2a. Knowledge segment, KAP study of migrant African populations regarding HBV and HCV infection in Al Kufra, Libya, 2021 (n = 674; 609 male (M), 94 female (F))

2a. Knowledge assessment questions	Item	Yes (%)	No (%)	% Correct	(%) Don't know
General questions:					Avg (33.8)
1. Have you ever heard about viral hepatitis?	1	115 (17.1%)	472(70.0%)	na	87(12.9%)
2. Are Hepatitis B and C viral infection diseases?	2	53 (7.9%)	500(74.2%)	7.9	121(18.0%)
3. Can hepatitis B or C be transmitted by contact with open wound/cut?	3	41 (6.1%)	490(72.7%)	6.1	143(21.2%)
4. Can hepatitis B and C virus be spread by someone who looks healthy?	4	58 (8.6%)	405(60.1%)	8.6	211(31.3%)
5. Is there a test that shows whether or not someone is hepatitis B or C positive?	5	62 (9.2%)	304(45.1%)	9.2	308(45.7%)
6. Do you think people with hepatitis B can be infected for life?	6	36 (5.3%)	411(61.0%)	5.3	227(33.7%)
7. Is there a vaccine for HBV?	7	37 (5.5%)	298(44.2%)	5.5	339(50.3%)
8. Can hepatitis B be cured?	8	202(30.0%)	91 (13.5%)	30.0	381(56.5%)
9. Can hepatitis C be cured?	9	130(19.3%)	319(47.3%)	19.3	225(33.4%)
10. Can HBV and HCV infection be passed from mother to child?	10	117 (17.4%)	205(30.4%)	17.4	352(52.2%)
11. Can HBV and HCV be spread through social contact (e.g. shaking hands, sharing food or sneezing)?	11	362(53.7%)	68(10.1%)	10.1	244(36.2%)
12. Can hepatitis infection cause death?	12	147 (21.8%)	196(29.1%)	21.8	331(49.1%)
13. Can people with Hepatitis B or C donate blood?	13	411 (61.0%)	166 (24.6%)	24.6	97 (14.4%)
14. Can HCV infection be prevented?	14	427 (63.4%)	126 (18.7%)	63.4	121(18.0%)
15. Can HBV infection be prevented?	15	234(34.7%)	247(36.6%)	59.1	193(28.6%)
16. Can Hepatitis B and C be transmitted through air and water?	16	398 (59.1%)	97 (14.4%)	14.4	179(26.6%)
17. Have you ever been screened for HBV and HVC?	17	115 (17.1%)	559 (82.9%)	na	0 (0.0%)
18. Have you been vaccinated against hepatitis B?	18	11 (1.6%)	663 (98.4%)	na	0 (0.0%)
19. Is a vaccine for Hepatitis C available?	19	413 (61.3%)	93 (13.8%)	13.8	168 (24.9%)
Significant risk factor questions:					Avg (48.0)
20. Can HBV and HCV be transmitted by un-sterilized syringes and needles?	20	66 (9.8%)	298 (44.2%)	9.8	310 (46.0%)
21. Are HBV and HCV passed through contact with infectious blood, blood products and body fluids?	21	39 (5.8%)	290 (43.0%)	5.8	345 (51.2%)
22. Can viral hepatitis B and C be transmitted by sharing needles for use drug?	22	70 (10.4%)	404 (59.9%)	10.4	200(29.7%)
23. Can viral hepatitis B and C be transmitted by tattooing or body piercing?	23	44 (6.5%)	294 (43.6%)	6.5	336 (49.9%)
24. Can viral hepatitis B and C be transmitted by sharing personal care items like toothbrush or razor?	24	60 (8.9%)	198 (29.4%)	8.9	416(61.7%)
25. Can hepatitis B and C be transmitted through sexual contact?	25	83 (12.3%)	229 (34.0%)	12.3	362 (53.7%)
26. Can HBV and HCV be transmitted by a lack of sanitation services like bathrooms?	26	257 (38.1%)	121(18.0%)	38.1	296 (43.9%)
Common complications questions:					Avg (53.3)
30. Can a person be a carrier of HBV and HCV and can infect others without knowing it?	30	88 (13.1%)	310 (46.0%)	13.1	276(40.9%)
31. Can HBV and HCV infections result in chronic hepatitis?	31	112(16.6%)	223 (33.1%)	16.6	339 (50.3%)
32. Can HBV and HCV infections result in liver cancer?	32	97 (14.4%)	174 (25.8%)	14.4	403 (59.8%)
33. Can HBV and HCV infections lead to liver cirrhosis?	33	78 (11.6%)	198 (29.4%)	11.6	398 (59.1%)
34. Can HBV and HCV infections lead to fulminant hepatitis?	34	57 (8.5%)	157 (23.3%)	8.5	460 (68.2%)
35. Can HBV and HCV infections cause yellow skin and eyes (jaundice) ?	35	72 (10.7%)	234 (34.7%)	10.7	368 (54.6%)
36. Can HBV and HCV infections cause liver failure?	36	112 (16.6%)	262 (38.9%)	16.6	300 (44.5%)
37. Can HBV and HCV infections cause death?	37	58 (8.6%)	287 (42.6%)	8.6	377 (55.8%)
Average %				16.4	Avg (42.0)

A majority, 86.5% falsely believed that hepatitis B could be cured (or didn't know), while nearly half, 47.3% believed that hepatitis C is not curable although a third (33.4%) 'did not know'. The vast majority, 98.4% did not know that an HBV vaccine existed and only 1.6% of the migrants had been vaccinated. Moreover, 82.9% had never been screened for HBV.

Knowledge of risk factors was also very poor. The overwhelming majority had no understanding that HBV and HCV could be transmitted by mechanisms including: by unsterilized syringes, 90.2%; by blood products and body fluids, 94.2%; and by sharing needles, 89.6%. Furthermore, similarly high majorities did not understand that both HBV & HCV could be transmitted by tattooing, 93.5%, sharing personal care items 91.0%, and through sexual contact, 87.7%, (Table 2). Among the 674 migrant participants, only ca. one in 10 understood that complications of chronic hepatitis include liver cancer, liver cirrhosis, fulminant hepatitis, and liver failure (Table 2a).

Results of attitude assessment indicated that nearly 90% of the migrants were not concerned about being infected with HBV or HCV and did not consider the infections a public health problem Table 2b. Only about 20% believed that they could get HBV or HCV infections and 15% were not worried about it. Only 15% agreed that migrants

Table 2b. Attitude assessment of the migrant population regarding HBV and HCV.

2b. Attitude assessment questions	Item no.	Definitely Agree	Moderately agree	Neutral	Moderately disagree
1. I have no concern for being infected with hepatitis B or C.	1	83(12.3%)	164(24.3%)	295(43.8%)	80(11.9%)
2. Hepatitis B vaccine is safe and effective.	2	23(3.4%)	36(5.3%)	369(54.7%)	176(26.1%)
3. All African migrant populations should be tested for hepatitis B or C before they receive health care.	3	38(5.6%)	74(11.0%)	259(38.4%)	198(29.4%)
4. I don't feel comfortable living with people with hepatitis B or C.	4	20(3.0%)	64(9.5%)	446(66.2%)	121(18.0%)
5. I believe that hepatitis infections can be transmitted through food.	5	49(7.3%)	182(27.0%)	340(50.4%)	78(11.6%)
6. I think hepatitis B is a curable disease.	6	87(12.9%)	167(24.8%)	369(54.7%)	27(4.0%)
7. I think that hepatitis B and C is a serious public health problem.	7	119(17.7%)	138(20.5%)	285(42.3%)	96(14.2%)
8. I can get hepatitis infection.	8	45(6.7%)	91(13.5%)	341(50.6%)	138(20.5%)
9. Viral hepatitis is something to worry about.	9	33(4.9%)	74(11.0%)	260(38.6%)	221(32.8%)
10. I believe that vaccination could prevent transmission.	10	49(7.3%)	111(16.5%)	244(36.2%)	197(29.2%)
11. I believe that healthy people need vaccination.	11	27(4.0%)	117(17.4%)	221(32.8%)	243(36.1%)
12. I believe that you need a vaccination at your age.	12	22(3.3%)	70(10.4%)	503(74.6%)	41(6.1%)
13. I believe that vaccine can be free or low cost through certain programs.	13	34(5.0%)	38(5.6%)	267(39.6%)	188(27.9%)

should be tested and less than 10% of participants agreed that the HBV vaccine is safe and effective (Table 2b). Similar to the knowledge response (Table 1a), less than 10% of participants believed or didn't know that HBV was not curable. Reflecting lack of concern, more than 90% of participants we felt comfortable living with people with hepatitis B or C, while 34.3% of the migrants believed that HBV and HCV infection could be transmitted by

food. More than two thirds of participants showed a negative attitude towards vaccination 179
(Table 2b). 180

Participating migrant responses to practice assessment questions indicate many unsafe 181
practices that may result in exposure to blood- and body fluid-borne diseases. Very few, 4.3%, 182
indicated that they would seek further diagnosis and treatment if they tested positive for HBV 183
or HCV, Table 2c. Very few indicated a ready source of information about hepatitis B or C, 184
social networks, 4.9%, or friends, 8.0%. Responses reflecting lack of knowledge or reliable 185
source of information, very high proportions of participants indicated high-risk practices for 186
transmission: 84% indicated shared use of injecting materials; 81% indicated shared intravenous 187
drug use; almost all, 98% indicated sharing shaving materials; and with a high rate of sexual 188
activity, ca. 90%, 189

Table 2c. Practice assessment of migrant the population regarding HBV and HCV 190

2c. Practice assessment questions	Item	Always	Frequently	Sometimes	Rarely
1. How often do you go for further investigation and treatment in case you are diagnosed with Hepatitis B or C?	1	11(0.02%)	7(0.01%)	12(0.02%)	9(0.01%)
2. How often are social networks on the internet your source of hepatitis B or C information?	2	15(0.02%)	2(0.003%)	5(0.007%)	11(0.02%)
3. How often are friends as your source of hepatitis B or C information?	3	13(0.02%)	10(0.02%)	17(0.03%)	14(0.02%)
4. How often do you use a used needle/syringe by more than one person?	4	147(21.8%)	155(23.0%)	187(27.7%)	108(16.0%)
5. How often do you share intravenous drug use?	5	134(19.9%)	144(21.4%)	173(25.7%)	127(18.8%)
6. How often are razors and shaving blades used by more than one person?	6	366(54.3%)	247(36.6%)	36 (5.3%)	14(2.1%)
7. How often do you screen for hepatitis B or C?	7	5(0.01%)	11(0.02%)	97(14.4%)	2(0.003%)
8. How often have you had sex within the last 12 months?	8	147(21.8%)	175(26.0%)	197(29.2%)	80(11.9%)
9. How often have you used condoms with non-steady/steady sexual partners within the last 12 months?	9	5(0.01%)	2(0.003%)	7(1.0%)	2(0.003%)
10. How often do you wash hands with tap water and soap?	10	71(10.5%)	93(13.8%)	142(21.1%)	161(23.9%)
11. How often do you refuse shaving with used instruments?	11	46(6.8%)	114(16.9%)	127(18.8%)	143(21.2%)
12. How often do you share eating utensils?	12	627(93.0%)	20(0.03%)	14(0.02%)	8(0.01%)

almost none indicated use of condoms. Reflecting lack of understanding the implications of 191
high-risk practices, very few, ca 15%, ever went for hepatitis B or C screening. 192

Inferential statistics, i.e., Mann-Whitney U and Kruskal Wallis H tests were applied to compare 193
scores of each domain with various demographic factors Table 3. Mean knowledge and attitude 194
scores of male and female differed for $p < 0.05$. Male showed higher knowledge and attitude 195

than females. Eritrean migrants had significantly higher mean knowledge and practice scores 196
than other demographic groups. The mean knowledge scores were significantly higher among 197
single than married migrants, while 198

**Table 3. Comparison of demographic characteristics for mean knowledge, attitude and 199
practice scores 200**

Demographic variables	K.score 0.794+.204 Mean rank	P- value	A.score 2.906+.380 Mean rank	P- value	P.score 2.483+.232 Mean rank	P- value
Gender identity**						
Male	355.27	0.001	343.47	0.047	335.08	0.418
Female	227.85		300.66		352.43	
Ethnicity*						
Eritrean	362.56	0.024	371.98	0.001	363.24	0.007
Somalian	320.64		288.41		317.97	
Ethiopian	309.08		305.26		331.08	
Sudanese	333.32		399.03		278.99	
Marital Status**						
Single	379.88	0.033	340.07	0.897	292.05	0.021
Married	331.47		337.13		343.97	
Education level**						
Illiterate/Basic education	291.61	0.001	305.40	0.001	352.02	0.001
High school and above	477.94		435.72		293.06	

* Kruskal Wallis Test, ** Mann Whitney Test, *P-value < 0.05 is statistically significant.* 202

the mean practice score of married migrants was significantly higher than among single 203
migrants. Finally, significantly higher differences were observed between mean knowledge and 204
attitude scores for migrants with high school education compared to those with illiterate/basic 205
education. However, those having high school education had significantly lower mean practice 206
scores. 207

A correlation between different domains of questionnaire was also assessed. The correlation 208
revealed significant positive linear correlations between knowledge-attitude ($r = 0.524$, $p <$ 209
 0.01) knowledge-practice ($r = 0.123$, $p < 0.01$) and attitude-practice ($r = 0.278$, $p < 0.01$), as 210
shown in Table 4. 211

Table 4 Correlation between knowledge, attitude, and practice scores 212

Variables	Correlation coefficient	P-value*
Knowledge, attitude	0.524	0.01
Knowledge, practice	0.123	0.01
Practice, Attitude	0.278	0.01

*Correlation significant at 0.01 level (2 tailed). 213

Using chi-square test, for knowledge scores of the study participants were divided into two 214
categories: good, and poor. Attitude and practice scores of the study participants were divided 215

into three categories: good, fair and poor. On this basis, knowledge was 'Poor', >85%, across all categories of gender, ethnicity, and marital status (Table 5). Attitudes could be considered 'Fair' for most categories. But practices were considered 'Poor' for > 2/3's of participants across all categories.

Table 5 Categorization of the study African migrant populations score on KAP domains

Variable	Knowledge					p-value	Attitude					p-value	Practices					p-value		
	Good		Poor		Good		Fair		Poor		Good		Fair		Poor					
	n	n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%			
Gender identity																				
Male	580	70	12.1%	510	87.9%	0.486	77	13.3%	418	72.1%	85	14.7%	0.649	5	0.9%	140	24.1%	435	75.0%	0.014
Female	94	9	9.6%	85	90.4%		10	10.6%	72	76.6%	12	12.8%		1	1.1%	34	36.2%	59	62.8%	
Ethnicity																				
Eritrean	295	26	8.8%	269	91.2%	0.188	16	5.4%	258	87.5%	21	7.1%	0.001	2	0.7%	93	31.5%	200	67.8%	0.036
Somalian	179	26	14.5%	153	85.5%		26	14.5%	111	62.0%	42	23.5%		1	0.6%	47	26.3%	131	73.2%	
Ethiopian	146	21	14.4%	125	85.6%		34	23.3%	88	60.3%	24	16.4%		1	0.7%	25	17.1%	120	82.2%	
Sudanese	54	6	11.1%	48	88.9%		11	20.4%	33	61.1%	10	18.5%		2	3.7%	9	16.7%	43	79.6%	
Marital Status																				
Single	84	14	16.7%	70	83.3%	0.132	11	13.1%	61	72.6%	12	14.3%	0.998	1	1.2%	11	13.1%	72	85.7%	0.017
Married	590	65	11.0%	525	89.0%		76	12.9%	429	72.7%	85	14.4%		5	0.8%	163	27.6%	422	71.5%	
Education Level																				
Basic	508	16	3.1%	492	96.9%	0.005	76	15.0%	429	84.4%	85	16.7%	0.001	-		146	28.7%	362	71.3%	0.024
High School	166	63	38.0%	103	62.0%		64	38.6%	86	51.8%	16	9.6%		6	3.6%	28	16.9%	132	79.5%	

P < 0.05 was set as statistically significant

Discussion:

Currently, the flow of immigrants into Libya from neighboring countries is increasing. However, there are no post-arrival screening mechanisms for hepatitis B and C viruses to assess migrant conditions or to support public health policy or actions. Accordingly, the purpose of this study was to assess the level of participating migrant's knowledge, attitude, and practices, regarding cause, transmission, prevention and treatment of viral hepatitis due to HBV and HCV. Libya public health professionals are concerned as the country as a whole is classified among geographical regions with low-intermediate endemicity for hepatitis B virus infection and an area of low endemicity for hepatitis C. The prevalence of hepatitis B surface antigen

(HBsAg) and anti-HCV antibodies in sub-Saharan African migrants has been found to exceed 233
30-50% indicating significant cause for concern (27). A plausible reason for this high 234
prevalence appears related to the lack of knowledge about HBV and HCV, their health 235
significance, their means of transmission, of actions for preventing infection. 236
The present study assessed the participant's knowledge, attitude, and practices about HBV and 237
HCV infection. Results clearly show poor KAP towards HBV and HCV among migrants at Al 238
Kufra. An overwhelming lack of knowledge about HBV and HCV, their mode of transmission, 239
and the possibility of prevention by vaccination. was observed among participating migrants 240
Most of these sub-Saharan participants did not know typical ways of transmission, prevention, 241
and development of liver disease. In addition, participants were not aware of the different types 242
of viral hepatitis and did not have access to information about their consequences. Only just 243
over 15% the migrants in our study had basic knowledge about HBV and HCV as a disease of 244
liver, consistent with a previous report of an Ethiopia study in which just 11.0% of the refugees 245
know the relationship between liver cancer and hepatitis B and C. Ethiopia has a population- 246
wide prevalence of 7.4 and 3.1% for HBV and HCV respectively [5]. 247
Most of the migrants in this study, 84.12%, did not consider HBV and HCV as a real risk for 248
themselves and for their family. Although this may explain their inadequate practices it almost 249
certainly contributes to high rates of HBV and HCV infection [27]. Migrants participating in 250
this study showed poorly supported attitudes and exceedingly poor practices towards HBV and 251
HCV reflecting their very low state of knowledge important to minimizing risk and infection. 252
A wide range of unsafe practices are evident in the survey responses, exposing the migrants to 253
infection by blood borne diseases. 254

Conclusions: 255

In this screening study, the level of knowledge and awareness of migrant African populations 256
transiting Libya was poor about the concepts of infectious risk associated HBV and HCV. Poor 257

knowledge and lack of awareness of the African immigrants about HBV and HCV must 258
contribute to the prevalence of these infections in developing countries compared with 259
developed ones. Survey results demonstrate that knowledge on transmission is lacking, and 260
awareness about HBV and HCV transmission to others is only fair. Lacking information about 261
transmission and prevention migrants are at risk of contracting HBV and HCV emphasizing 262
the need for education to limit the spread of the blood borne disease. Opportunities exist for 263
attention by Libyan health and immigration authorities to assist migrants seeking improved 264
conditions. Poor knowledge and lack of awareness about HBV and HCV among sub-Saharan 265
populations undoubtedly contributes to the spread of these infections in native countries. 266

References:

- 1- Alam M, Tariq WUZ: Knowledge, Attitudes and Practices about Hepatitis B and C among young 269
healthy males. *Pak J Pathol* 2006, 17:147–150. 270
- 2- Allain J-P, Candotti D, Soldan K, Sarkodie F, Phelps B, Giachetti C, et al. The risk of hepatitis B 271
virus infection by transfusion in Kumasi, Ghana. *Blood*. 2003;101(6):2419–25. 272
- 3- Ayele A, Abera D, Hailu M, Birhanu M and Desta K. Prevalence and associated risk factors for 273
Hepatitis B and C viruses among refugees in Gambella, Ethiopia. *BMC Public Health* (2020) 274
20:721. 275
- 4- Badawi M, Atif M, Mustafa Y. Systematic review and meta-analysis of HIV, HBV and HCV 276
infection prevalence in Sudan. *Virology* 2018;15(1):1–16. 277
- 5- Belyhun Y, Maier M, Mulu A, Diro E, Liebert UG. Hepatitis viruses in Ethiopia: a systematic 278
review and meta-analysis. *BMC Infect Dis*. 2016;16(1):761. 279
- 6- Burnett R, Francois G, Kew M, Leroux-Roels G, Meheus A, Hoosen A, et al. Hepatitis B virus and 280
human immunodeficiency virus co-infection in sub-Saharan Africa: a call for further investigation. 281
Liver Int. 2005;25(2):201–13. 282

- 7- Burnett R, Francois G, Kew M, Leroux-Roels G, Meheus A, Hoosen A, et al. Burgess C, Drucker 283
E. Epidemiology of chronic hepatitis C virus infection in sub-Saharan Africa. *Lancet Infect Dis.* 284
2002;2(5):293–302. 285
- 8- Daw MA, et al. Hepatitis C Virus in North Africa: An Emerging Threat. *The Scientific World 286*
Journal. Volume 2016, Article ID 7370524, 11 pages. <http://dx.doi.org/10.1155/2016/7370524>. 287
- 9- Daw MA, El-Bouzedi A. In association with Libyan Study Group of Hepatitis & HIV. Prevalence 288
of hepatitis B and hepatitis C infection in Libya: results from a national population based survey. 289
BMC Infect Dis 2014; 14: 17. 290
- 10- De Martel C, Ferlay J, Franceschi S, Vignat J, Bray F, Forman D, et al. Global burden of cancers 291
attributable to infections in 2008: a review and synthetic analysis. *Lancet Oncol.* (2012) 13:607– 292
15. doi: 10.1016/S1470-2045(12)70137-7 293
- 11- Frank C, Mohamed MK, Strickland GT, Lavanchy D, Arthur RR, Magder LS, et al. The role of 294
parenteral antischistosomal therapy in the spread of hepatitis C virus in Egypt. *Lancet* 295
2000;355:887–891. 296
- 12- Georgel P, Schuster C, Zeisel MB, Stoll-Keller F, Berg T, Bahram S. Virus-host interactions in 297
hepatitis C virus infection: implications for molecular pathogenesis and antiviral strategies. *Trends* 298
Mol Med. 2010;16(6):277–86. 299
- 13- Goldstein ST, Zhou F, Hadler SC, Bell BP, Mast EE, Margolis HS. A mathematical model to 300
estimate global hepatitis B disease burden and vaccination impact. *Int J Epidemiol.* (2005) 34:1329– 301
39. doi: 10.1093/ije/dyi206 302
- 14- Gushulak BD, MacPherson DW. Globalization of infectious diseases: the impact of migration. *Clin 303*
Infect Dis 2004 Jun 15; 38(12):1742e8. PMID: 15227621. 304
- 15- Hassan MA, Kim WR, Li R, Smith CI, Fried MW, Sterling RK, et al. Characteristics of US-born 305
versus foreign-born Americans of African descent with chronic hepatitis B. *Am J Epidemiol.* (2017) 306
186:356–66. doi: 101093/aje/kwx064. 307
- 16- Hauri AM, Armstrong GL, Hutin YJ. The global burden of disease attributable to contaminated 308
injections given in health care settings. *Int J STD AIDS.* 2004;15(1):7–16. 309

17- Howell J, Ladep NG, Lemoin M, Thursz MR, Taylor-Robinson SD. Hepatitis B in sub-saharan Africa. S Sudan Med J. 2014;7(3):59–61.	310 311
18- Kirbak ALS. Sero-prevalence for hepatitis B virus among pregnant women attending antenatal clinic in juba teaching hospital, republic of South Sudan. Pan Afr Med. 2017;26:72.	312 313
19- Lavanchy D. The global burden of hepatitis C. Liver Int. 2009;29 Suppl 1:74–81.	314
20- Mohamed EA, Giama NH, Shaleh HM, Kerandi L, Oseini AM, Ahmed Mohammed H, et al. Knowledge, attitudes, and behaviors of viral hepatitis among recent African immigrants in the United States: a community based participatory research qualitative study. Front Public Health. 2020;8:25.	315 316 317 318
21- Nelson PK, Mathers BM, Cowie B, Hagan H, Des Jarlais D, Horyniak D, et al. Global epidemiology of hepatitis B and hepatitis C in people who inject drugs: results of systematic reviews. Lancet. 2011;378(9791):571–83.	319 320 321
22- OECD. World Migration in Figures. < http://www.oecd.org/els/mig/World-Migration-in-Figures.pdf >.	322 323
23- Ott JJ, Stevens GA, Groeger J, Wiersma ST. Global epidemiology of hepatitis B virus infection: new estimates of age-specific HBsAg seroprevalence and endemicity. Vaccine. (2012) 30:2212–9. doi: 10.1016/j.vaccine.2011.12.116	324 325 326
24- Owiti JA, Greenhalgh T, Sweeney L, Foster GR, Bhui KS. Illness perceptions and explanatory models of viral hepatitis B & C among immigrants and refugees: a narrative systematic review. BMC Public Health. 2015; 15:151.	327 328 329
25- Perz JF, Armstrong GL, Farrington LA, Hutin YJ, Bell BP. The contributions of hepatitis B virus and hepatitis C virus infections to cirrhosis and primary liver cancer worldwide. J Hepatol. 2006;45(4):529–38.	330 331 332
26- Saeed, FMA, Ongerth, JE, 2019. Giardia and Cryptosporidium in children with diarrhea, Kufra Libya, a North African migration route city. International Journal of Hygiene and Environmental Health. 222(5):840-846. https://doi.org/jijheh.2019.04.006 .	333 334 335

- 27- Saaed, FMA, Ongerth, JE, 2022. Prevalence of hepatitis B and hepatitis C in migrants from sub-Saharan Africa newly arrived at Al Kufra in SE Libya in transit toward Europe. MEDRXIV/2022/27021. 336
337
338
- 28- Schilling T, Rauscher S, Menzel C, Reichenauer S, Müller-Schilling M, Schmid S, et al. Migrants and refugees in Europe: challenges, experiences and contributions. *Visc Med.* 2017;33(4):295–300. 339
340
- 29- Seeger C, Mason WS. Hepatitis B virus Biology. *Micro Mol Biol Rev* 2000; 64: 51-68. 341
- 30- Sharma S, Carballo M, Feld JJ, Janssen HLA. Immigration and viral hepatitis. *Journal of Hepatology.* 2015; 63: 515–522. 342
343
- 31- Shalaby S, Kabbash I, El Saleet G, Mansour N, Omar A, El Nawawy A. Hepatitis B and C viral infection: prevalence, knowledge, attitude and practice among barbers and clients in Gharbia governorate, Egypt. *East Mediterr Health J.* 2010;16:10–7. 344
345
346
- 32- Stanaway JD, Flaxman AD, Naghavi M, Fitzmaurice C, Vos T, Abubakar I, et al. The global burden of viral hepatitis from 1990 to 2013: findings from the Global Burden of Disease Study 2013. *Lancet.* (2016) 388:1081–8. doi: 10.1016/S0140-6736(16)30579-7 347
348
349
- 33- Strong C, Hur K, Kim F, Pan J, Tran S, Juon H-S. Sociodemographic characteristics, knowledge and prevalence of viral hepatitis infection among vietnamese Americans at community screenings. *J Immigr Minor Health* 2015;17:298–301. 350
351
352
- 34- Taylor VM, Jackson JC, Pineda M, Pham P, Fischer M, Yasui Y. Hepatitis B knowledge among Vietnamese immigrants: implications for prevention of hepatocellular carcinoma. *J Cancer Educ* 2000;15:51–55. 353
354
355
- 35- Tramuto F, Maida CM, Colomba GM, Di Carlo P, Mazzola G, Li Vecchi V, et al. Occult hepatitis B infection in the immigrant population of Sicily, Italy. *J Immigr Minor Health.* (2013) 15:725–31. doi: 10.1007/s10903-012-9699-y 356
357
358
- 36- Tugwell P, Pottie K, Welch V, Ueffing E, Chambers A, Feightner J, et al. Evaluation of evidence-based literature and formulation of recommendations for the clinical preventive guidelines for immigrants and refugees in Canada. *CMAJ* 2011;183:E933–E938. 359
360
361
- 37- UNHCR. Ethiopia refugee response plan; 2018. p. 1–68. 362

- 38- UN. UN International Migration Wallchart 2013. [Http://www.un.org/en/development/desa/population/publications/migration/migration-wallchart-2013.shtml](http://www.un.org/en/development/desa/population/publications/migration/migration-wallchart-2013.shtml). 363
364
- 39- Who.int. WHO | Hepatitis C [Internet]. 2015 [cited 16 October 2015]. Available from: <http://www.who.int/mediacentre/factsheets/fs164/en/>. 365
366
- 40- World Health Organization: Advocacy, communication and social mobilization for TB control: a guide to developing knowledge, attitude and practice surveys. http://whqlibdoc.who.int/publications/2008/9789241596176_eng.pdf. 367
368
369
- 41- World Health Organisation. Hepatitis: Fact sheets. Geneva: WHO. Available from: [www. http://www.who.int/topics/hepatitis/factsheets/en/](http://www.who.int/topics/hepatitis/factsheets/en/). Accessed on: 17 May 2014. 370
371
- 42- Yang JD, Roberts LR. Hepatocellular carcinoma: a global view. *Nat Rev Gastroenterol Hepatol*. (2010) 7:448–58. doi: 10.1038/nrgastro.2010.100 372
373
- 43- Yang JD, Gyedu A, Afihene MY, Duduyemi BM, Micah E, Kingham TP, et al. Hepatocellular carcinoma occurs at an earlier age in africans, particularly in association with chronic hepatitis B. *Am J Gastroenterol*. (2015) 110:1629–31. doi: 10.1038/ajg.2015.289 374
375
376
- 44- Yang JD, Mohamed EA, Aziz AO, Shousha HI, HashemMB, NabeelMM, et al. Characteristics, management, and outcomes of patients with hepatocellular carcinoma in Africa: a multicountry observational study from the Africa Liver Cancer Consortium. *Lancet Gastroenterol Hepatol*. (2017) 2:103–11. doi: 10.1016/S2468-1253(16)30161-3 377
378
379
380
381

Appendix

382

Table A1. Questions on Knowledge of study participating migrants about HBV and HCV

383

Knowledge Questions	Yes (%)	No (%)	Don't know (%)
General questions:			
1. Have you ever heard about viral hepatitis?	115 (17.1%)	472(70.0%)	87(12.9%)
2. Are Hepatitis B and C viral infection diseases?	53 (7.9%)	500(74.2%)	121(18.0%)
3. Can hepatitis B or C be transmitted by contact with open wound/cut?	41 (6.1%)	490(72.7%)	143(21.2%)
4. Can hepatitis B and C virus be spread by someone who looks healthy?	58 (8.6%)	405(60.1%)	211(31.3%)
5. Is there a test that shows whether or not someone is hepatitis B or C positive?	62 (9.2%)	304(45.1%)	308(45.7%)
6. Do you think people with hepatitis B can be infected for life?	36 (5.3%)	411(61.0%)	227(33.7%)
7. Is there a vaccine for HBV?	37 (5.5%)	298(44.2%)	339(50.3%)
8. Can hepatitis B be cured?	202(30.0%)	91 (13.5%)	381(56.5%)
9. Can hepatitis C be cured?	130(19.3%)	319(47.3%)	225(33.4%)
10. Can HBV and HCV infection be passed from mother to child?	117 (17.4%)	205(30.4%)	352(52.2%)
11. Can HBV and HCV be spread through social contact (e.g. shaking hands, sharing food or sneezing)?	362(53.7%)	68(10.1%)	244(36.2%)
12. Can hepatitis infection cause death?	147 (21.8%)	196(29.1%)	331(49.1%)
13. Can people with Hepatitis B or C donate blood?	411 (61.0%)	166 (24.6%)	97 (14.4%)
14. Can HCV infection be prevented?	427 (63.4%)	126 (18.7%)	121(18.0%)
15. Can HBV infection be prevented?	234(34.7%)	247(36.6%)	193(28.6%)
16. Can Hepatitis B and C be transmitted through air and water?	398 (59.1%)	97 (14.4%)	179(26.6%)
17. Have you ever been screened for HBV and HVC?	115 (17.1%)	559 (82.9%)	0 (00.00%)
18. Have you been vaccinated against hepatitis B?	11 (1.6%)	663 (98.4%)	0 (00.00%)
19. Is a vaccine for Hepatitis C available?	13 (61.3%)	93 (13.8%)	168 (24.9%)
Significant risk factor questions:			
20. Can HBV and HCV be transmitted by un-sterilized syringes and needles?	66 (9.8%)	298 (44.2%)	310 (46.0%)
21. Are HBV and HCV passed through contact with infectious blood, blood products and body fluids?	39 (5.8%)	290 (43.0%)	345 (51.2%)
22. Can viral hepatitis B and C be transmitted by sharing needles for use drug?	70 (10.4%)	404 (59.9%)	200(29.7%)
23. Can viral hepatitis B and C be transmitted by tattooing or body piercing?	44 (6.5%)	294 (43.6%)	336 (49.9%)
24. Can viral hepatitis B and C be transmitted by sharing personal care items like toothbrush or razor?	60 (8.9%)	198 (29.4%)	416(61.7%)
25. Can hepatitis B and C be transmitted through sexual contact?	83 (12.3%)	229 (34.0%)	362 (53.7%)
26. Can HBV and HCV be transmitted by a lack of sanitation services like bathrooms?	257 (38.1%)	121(18.0%)	296 (43.9%)
Common complications questions:			
27. Can a person be a carrier of HBV and HCV and can infect others without knowing it?	88 (13.1%)	310 (46.0%)	276(40.9%)
28. Can HBV and HCV infections result in chronic hepatitis?	112(16.6%)	223 (33.1%)	339 (50.3%)
29. Can HBV and HCV infections result in liver cancer?	97 (14.4%)	174 (25.8%)	403 (59.8%)
30. Can HBV and HCV infections lead to liver cirrhosis?	78 (11.6%)	198 (29.4%)	398 (59.1%)
31. Can HBV and HCV infections lead to fulminant hepatitis?	57 (8.5%)	157 (23.3%)	460 (68.2%)
32. Can HBV and HCV infections cause yellow skin and eyes (jaundice) ?	72 (10.7%)	234 (34.7%)	368 (54.6%)
33. Can HBV and HCV infections cause liver failure?	112 (16.6%)	262 (38.9%)	300 (44.5%)
34. Can HBV and HCV infections cause death?	58 (8.6%)	287 (42.6%)	329 (48.8%)

384

385

386

Table A2. Questions regarding Attitude of participating migrants about HBV and HCV

387

Attitude assessment questions	Definitely Agree	Moderately agree	Neutral	Moderately disagree	Definitely Disagree
1. I have no concern for being infected with Hepatitis B or C.	83(12.3%)	164(24.3%)	295(43.8%)	80(11.9%)	52(7.7%)
2. Hepatitis B vaccine is safe and effective.	23(3.4%)	36(5.3%)	369(54.7%)	176(26.1%)	70(10.4%)
3. All African migrant populations should be tested for hepatitis B or C before they receive health care.	38(5.6%)	74(11.0%)	259(38.4%)	198(29.4%)	105(15.6%)
4. I don't feel comfortable to live with people with Hepatitis B or C.	20(3.0%)	64(9.5%)	446(66.2%)	121(18.0%)	23(3.4%)
5. I believe that hepatitis infections can be transmitted through food.	49(7.3%)	182(27.0%)	340(50.4%)	78(11.6%)	25(3.7%)
6. I think hepatitis B is a curable disease.					
7. I think that hepatitis B and C is a serious public health problem.	87(12.9%)	167(24.8%)	369(54.7%)	27(4.0%)	24(3.6%)
8. I can get hepatitis infection.	119(17.7%)	138(20.5%)	285(42.3%)	96(14.2%)	36(5.3%)
9. Viral hepatitis is something to worry about.	45(6.7%)	91(13.5%)	341(50.6%)	138(20.5%)	59(8.8%)
10. I believe that vaccination could prevent transmission.	33(4.9%)	74(11.0%)	260(38.6%)	221(32.8%)	86(12.8%)
11. I believe that healthy people need vaccination.	49(7.3%)	111(16.5%)	244(36.2%)	197(29.2%)	73(10.8%)
12. I believe that you need a vaccination at your age.	27(4.0%)	117(17.4%)	221(32.8%)	243(36.1%)	66(9.8%)
13. I believe that vaccine can be free or low cost through certain programs.	22(3.3%)	70(10.4%)	503(74.6%)	41(6.1%)	38(5.6%)
	34(5.0%)	38(5.6%)	267(39.6%)	188(27.9%)	147(21.8%)

388

Table A3. Questions regarding Practices of participating migrants about HBV and HCV

389

Practice Assessment Questions	Always	Frequently	Sometimes	Rarely	Never
1. How often do you go for further investigation and treatment, if you are diagnosed with hepatitis B or C?	11(01.63%)	7(01.03%)	12(01.78%)	9(01.33%)	635(94.21%)
2. How often are social networks on the internet your source of hepatitis B or C information?	15(02.22%)	2(00.29%)	5(00.74%)	11(01.63%)	641(95.10%)
3. How often are friends as your source of hepatitis B or C information?	13(1.9%)	10(1.5%)	17(2.5%)	14(2.1%)	620(92.0%)
4. How often do you use a used needle/syringe by more than one person?	47(21.8%)	155(23.0%)	187(27.7%)	108(16.0%)	77(11.4%)
5. How often do you share intravenous drug use?	134(19.9%)	144(21.4%)	173(25.7%)	127(18.8%)	96(14.2%)
6. How often are razors and shaving blades used by more than one person?	366(54.3%)	247(36.6%)	36 (5.3%)	14(2.1%)	11(1.6%)
7. How often do you screen for hepatitis B or C?	5(7%)	11(1.6%)	97(14.4%)	2(.3%)	559(82.9%)
8. How often have you had sex within the last 12 months?	147(21.8%)	175(26.0%)	197(29.2%)	80(11.9%)	75(11.1%)
9. How often have you used condoms with non-steady/steady sexual partners within the last 12 months?	5(7%)	2(.3%)	7(1.0%)	2(.3%)	658(97.6%)
10. How often do you wash hands with tap water and soap?	71(10.5%)	93(13.8%)	142(21.1%)	161(23.9%)	207(30.7%)
11. How often do you refuse shaving with used instruments?	46(6.8%)	114(16.9%)	127(18.8%)	143(21.2%)	244(36.2%)
12. How often do you share eating utensils?	627(93.0%)	20(3.0%)	14(2.1%)	8(1.2%)	5(.7%)

390