Integrated fragmentomic profile and 5-Hydroxymethylcytosine of capture-based low-pass sequencing data enables pan-cancer detection via cfDNA

Zhidong Zhang[1,†], Xuenan Pi[1,†], Chang Gao[1], Jun Zhang[2], Lin Xia[1], Xiaoqin Yan[2], Xinlei Hu[1], Ziyue Yan[2], Shuxin Zhang[3], Ailin Wei[7], Yuer Guo[1], Jingfeng Liu[5], Ang Li[4], Xiaolong Liu[5], Wei Zhang[6], Yanhui Liu[3], Dan Xie1[*]

[1]National Frontier Center of Disease Molecular Network, State Key Laboratory of Biotherapy, West China Hospital, Sichuan University, No. 37 Guoxue Alley, Chengdu 610041, Sichuan Province, P. R. China.

[2]Tailai Inc., Shanghai 200233, P. R. China.

[3]Department of Neurosurgery, West China Hospital, Sichuan University, No. 37 Guoxue Alley, Chengdu 610041, Sichuan Province, P. R. China.

[4]Department of Pancreatic Surgery, West China Hospital, Sichuan University, No. 37 Guoxue Alley, Chengdu 610041, Sichuan Province, P. R. China.

[5]Mengchao Hepatobiliary Hospital of Fujian Medical University, Xihong Road 312, Fuzhou 350025, Fujian Province, P. R. China.

[6]Department of Respiratory and Critical Care Medicine, First Affiliated Hospital of the Second Military Medical University, Shanghai 200433, Shanghai, P. R. China.

[7]Guang'an People's Hospital, Guang'an, China

*Correspondence to: Dr Dan Xie, National Frontier Center of Disease

Molecular Network, State Key Laboratory of Biotherapy, West China Hospital,

Sichuan University, No. 37 Guoxue Alley, Chengdu 610041, Sichuan Province,

P. R. China. Tel: +86-28-8558-2944; Fax: +86-28-8558-2944; E-mail:

danxie@scu.edu.cn (D. Xie).

† These authors contributed equally to this work as joint first authors.

Abstract

Using epigenetic markers and fragmentomics of cell-free DNA for cancer

detection has been proven applicable. We further combine the two features

and explore the diagnostic potential of the features on pan-cancer detection.

We extracted cfDNA fragmentomic features from 191 whole-genome

sequencing data and investigated them in 396 low-pass 5hmC sequencing

data from four common cancer types and controls. We identified aberrant

ultra-long fragments (220-500bp) of cancer samples in 5hmC sequencing

data, both in size and coverage profile, and showed its dominant role in

cancer prediction. Since cfDNA hydroxymethylation and fragmentomic

markers can be detected simultaneously in low-pass 5hmC sequencing data,

we built an integrated model including 63 features of both fragmentomic

features and hydroxymethylation signatures for pan-cancer detection with

high sensitivity and specificity (88.52% and 82.35%, respectively). We

showed that fragmentomic information in 5hmC sequencing data is an ideal

marker for cancer detection and that it shows high performance in low-pass

sequencing data.

Introduction

Displaying a trend of increase, more than 19.3 million individuals were

diagnosed with cancer, and 10.0 million individuals died of cancer worldwide

in 2020 (1). Currently, tissue biopsy is the most widely used method for

cancer diagnosis and staging. However, owing to being an invasive

examination method, tissue biopsy has difficulties in tracking clonal evolution,

monitoring treatment response, and early detection of cancer and recurrence

(2). Therefore, the need for the development of accurate liquid biopsy

methods is urgent, with the potential to detect cancer and identify resistance

mechanisms early, to quantify minimal residual disease, and to monitor

treatment responses (2, 3).

Circulating cell-free tumor DNA (ctDNA) is a kind of tumor-derived material in

the blood of patients with cancer and is mainly derived from cancer cells via

apoptosis or necrosis (3, 4). ctDNA carries cancer-specific genetic mutations,

epigenetic alterations, and fragmentomic aberrations, allowing for cancer

detection and tissue-of-origin prediction in a non-invasive manner.

Nevertheless, as cancer is a highly heterogeneous disease, biomarkers

based on genetic and epigenetic aberrations often have limitations in practical

applications. Recent studies have shown that cell-free DNA (cfDNA)

fragmentation is a non-random process (5, 6), and it is thus possible to

develop a generic approach for non-invasive cancer detection using cfDNA

fragmentomics.

The fragmentomics analysis of cfDNA encompasses the sizes (7), coverage

(7), nucleosome footprints (8), end-points (9, 10) and end-motifs (11) of cfDNA.

For example, the size distribution of cfDNA fragments can indicate specific

cellular biological changes (12, 13). Coverage in cfDNA sequencing can be

used to analyze nucleosome positions (8, 14) and identify expressed genes

(14). Fragmentation patterns of cfDNA can be characterized by deferentially

phased fragment end signals, which were preferentially found in tissue-

specific open chromatin regions (15). Practically, Jiang et al. showed that

cfDNA fragments ending with particular genomic coordinates or motifs had

higher degrees of association with hepatocellular carcinoma (HCC) (10, 11).

Recently, the epigenetic marker 5-hydroxymethylcytosine (5hmC) has

attracted attention in the field of cancer research due to its regulatory role in

tissue-specific gene expression (16). The cfDNA captured by 5hmC comprises

a subset of all circulating cfDNA and, therefore, may contain fragmentomic

information. Integrative multi-omics analysis including the 5hmC profile of

cfDNA improves the sensitivity and specificity of cancer detection, as

demonstrated by recent studies (17, 18).

To highlight the potential of 5hmC sequencing data in cancer detection, we

obtained a large cohort of cancer samples, including HCC, pancreatic ductal

adenocarcinoma (PDAC), lung adenocarcinoma (LUAD), and glioblastoma

(GBM). We then established a pan-cancer preferred end map based on WGS

data from these four cancer types and checked their characteristics in low-

pass 5hmC sequencing data. We hypothesized that the fragmentomic

information would be consistent between WGS and 5hmC sequencing data

and tested it by comparison of each feature. We built classification models

and investigated the effect of combining 5hmC signatures with fragmentomic

information in pan-cancer screening.

Methods

Sample collection and study design

In total, low-pass (mean mapped reads: 38.32±8.02 million) 5hmC-seq data

from 85 healthy controls and 311 cancer patients were included in this study.

We obtained 5hmC sequencing data from 33 LUAD (18), 74 PDAC (37), 132

HCC (38), and 85 control (37) samples from three publications, and 72 GBM

samples from an unpublished paper.

Plasma samples from 15 healthy controls and 59 cancer patients were

collected and subjected to WGS (mean: 14×, range: 8.6-23.7×) to establish

the preferred end marker set, while the low-pass WGS data from 19 healthy

controls and 98 cancer patients (mean: 3×, range: 2.3-6.2×) were obtained to

cross-check with low-pass 5hmC sequencing data. 13 of the samples with

low-pass WGS sequencing had above 10x WGS sequencing data. The

demographics and clinical characteristics of the cohort are summarized in

Supplementary Tables 1 and 2. The design of the study is shown in Figure 1.

Plasma sample collection and cfDNA WGS sequencing

We collected plasma samples from 46 GBM patients, 29 PDAC patients and 34 controls from West China Hospital. Plasma samples of 37 HCC patients and 32 LUAD patients were collected from Mengchao Hepatobiliary Hospital of Fujian Medical University and the First Affiliated Hospital of the Second Military Medical University, respectively. For every subject, the QIAseq cfDNA All-in-one Kit (Qiagen) was applied on the 2 ml plasma sample for the cfDNA extraction and library construction. Pair-end 150 bp sequencing of the libraries was performed on the Illumina Novaseq 6000 platform.

cfDNA 5hmC profiling and sequencing

cfDNA extraction and 5hmC library construction was performed as previously described (28). Firstly, cfDNA was extracted from 2 ml plasma sample using QIAamp Circulating Nucleic Acid Kit (QIAGEN Inc., Valencia, CA, USA). Then, cfDNA (5–10 ng) ligated with sequencing adaptors was incubated in a 25 μL reaction solution containing HEPES buffer (50 mM, pH 8.0), 25 mM MgCl2, 60 μM N3-UDP-Glc (ActiveMotif, Carlsbad, CA, USA), and 12.5 U β-glucosyltransferase (NEB, Beverly, MA, USA) for 2 h at 37 °C. Next, 2.5 μL DBCO-PEG4-biotin (Sigma, Carlsbad, CA, USA) was directly added and incubated for 2 h at 37 °C. 10 μg sheared salmon sperm DNA (Life

Technologies, USA) was added, then the Micro Bio-Spin 30 Column (Bio-Rad,

Hercules, CA, USA) was used to purify the DNA following the instruction, and

the final volume was adjusted to 25μL. After that, the purified DNA was

incubated with 5 μL C1 streptavidin beads (Life Technologies, USA) in buffer 1

(5 mM Tris pH 7.5, 0.5 mM EDTA, 1 M NaCl and 0.2% Tween 20) for 30 min.

The beads were subsequently undergone three 5-min washes each with

buffer 1, buffer 2 (buffer 1 without NaCl), buffer 3 (buffer 1 with pH 9) and

buffer 4 (buffer 3 without NaCl). Then, the beads were resuspended in water

and amplified with 11 cycles of PCR amplification (initial denaturing at 98 °C

for 45 s, followed by 11 cycles of denaturing at 98 °C for 15 s, annealing at

60 °C for 30 s, extension at 72 °C for 30 s, and a final extension at 17 °C for 5

min). The amplified products were purified using 0.8 × AMPure XP beads

(Beck-man Coulter, Fullerton, CA, USA). Pair-end 150 bp sequencing was

performed on the Illumina Novaseq 6000 platform.

Fragmentomic profiling of cfDNA and cancer prediction analysis

The details of the preferred end, 5hmC signatures, size profile and coverage

profile calculation are described in the Supplementary Material. We used a

random forest model to distinguish healthy people from cancer patients using

fragmentomic features. To estimate the prediction error, we used five cross-

validations. Near-zero variance features were removed. The training,

validation, and test sets account for 60%, 20%, and 20% of the data,

respectively. The samples were selected randomly in a balanced way to keep the ratio of the number of cancer to non-cancer samples similar in the training, validation, and test subsets. Feature importance was calculated using the training data in each cross-validation run, and we sorted the features according to the mean value of feature importance. The number of features used in the final model was obtained by the highest AUC value in the validation set. To build an integrative model, we fitted the features selected by every feature-alone model into one random forest model, and performed feature selection, training, testing as described above. Random forest machine learning was implemented using the python package sklearn.ensemble.RandomForestClassifier with the following parameters: n_estimators = 100, criterion="gini".

Statistics

All statistical analyses were performed using R version 3.6.1 and python 3.7.0. All tests were two-sided, and P values < 0.05 were considered statistically significant. The ROC-AUC plot and AUC value were implemented using sklearn.metrics.plot_roc_curve.

Ethics

This study was approved by the Ethics Committee of Sichuan Cancer Hospital (SCCHEC-02-2016-005). The written informed consent was obtained from all

participants.

Results

Constructing the cfDNA preferred end map

cfDNA fragments ending at specific genomic coordinates that were statistically

more abundant than the Poisson distribution was an essential fragmentation

signature referred to as the preferred end (10). We utilized WGS data from

the four cancer types to build a preferred end map and checked the feature

using low-pass data. The basis of utilizing cfDNA preferred end coordinates to

indicate the occurrence and location of cancer was its ability to locate

nucleosomes, the position of which is related to cell identity (19). To confirm

this, we defined upstream (U) ends and downstream (D) ends by their

genomic coordinate order on the reference genome calculated via alignment.

We surveyed the distance distribution between adjacent U ends, D ends, and

nucleosome centers, and the result suggested the preferred ends were

enriched on the DNA linker (Supplementary Figure 1a-f). We further checked

the distribution of the predicted nucleosome location based on non-malignant

tissues at chr12: 34517269-34519122, and the preferred U and D ends of

WGS data from controls (n=15) and HCC patients (n=15). As expected, the

preferred U and D ends of the controls were enriched at the midpoint between

two nucleosomes, which is the location of the DNA linker (Figure 2a), and U

ends were found downstream of their nearest D ends. However, in HCC

patients, U ends were upstream of their nearest D ends, and only D ends

were found at one midpoint between two nucleosomes (Figure 2b), which

indicated inconsistent nucleosome positions between HCC patients with

controls. We observed the same abnormal preferred end distribution in GBM,

LUAD and PDAC patients (Supplementary Figure 1g-i).

By using WGS data, we constructed cfDNA fragment preferred end maps for

the four cancer types and control (Figure 2c-d). Preferred end enrichment was

observed in intron and open chromatin regions (Supplementary Figure 2).

Additionally, we noticed that for every cancer type, both U and D preferred

ends were more enriched in health-specific open chromatin regions than in

cancer-specific open chromatin regions (Supplementary Figure 2). One

explanation was the nucleosome pattern in the health-specific open chromatin

regions was unstable, as the enrichment of both U and D preferred ends in

healthy cfDNA were significantly lower than cancer patients (Mann–Whitney,

P<0.05). Another was the heterogeneous nature of cfDNA, as cfDNA can be

derived from various tissue sources. To search for preferred ends specific to

each of the four cancer types, we determined genome-wide cancer-specific

preferred U/D end coordinates and health-specific preferred U/D end

coordinates based on the Youden index. In total, we got 81,083 D and 80,505

U preferred ends. Intriguingly, compared with all repeatable ends (preferred

ends appeared in at least two samples in each cancer type), cancer-specific

preferred U/D ends were no longer enriched in health-specific open chromatin

regions, but they showed enrichment in promoter regions and cancer-specific

open chromatin regions (Figure 2e-f, Supplementary Figure 3-4). Furthermore,

health-specific preferred U/D ends showed enrichment in intergenic regions

and were not found in promoters, as cancer-specific preferred ends were

(Supplementary Figure 4). To cross check the cancer specificity of the

preferred ends in low-pass WGS data, we calculated a ratio, which was the

ratio of detected cancer-specific preferred end numbers to health-specific

preferred end numbers for each sample. The results showed significantly

higher ratios of cancer-specific to health-specific preferred U/D ends in the

three cancer types (except GBM) than in the control (Figure 2g,

Mann–Whitney, p value<0.001). In addition to the difference between cancer

and health, statistically significant differences in the ratios between cancer

types were observed (Supplementary Figure 5).

Fragmentomic features in 5hmC data

We checked the batch effect of the 5hmC sequencing data based on size

profile and coverage profile, and found no clustering of samples according to

cancer types (Supplementary Figure 6).

The cfDNA captured by 5hmC marks comprised a subset of DNA in whole

plasma. In theory, one sequenced fragment should contain at least one 5hmC.

Hence, we investigated the consistency of the size profile between WGS data

and 5hmC sequencing data. According to the results, fragment sizes of 5hmC

cfDNA in the four cancer samples were significantly shorter than those in the

control (Mann–Whitney test, p value<0.001, pan-cancer Median N50 = 174 bp,

control Median N50 = 210 bp), though the difference between cancer and

control groups in WGS data was only observed for HCC and LUAD (Figure

3a). Next, we confirmed that the cfDNA size profile of the 5hmC sequencing

data was consistent with that of the WGS data in the size range of 0-220 bp

(Figure 3b), with both data types displaying a dominant peak at 167 bp

(Supplementary Figure 7a). A difference between the cancer and control

samples in both short (100-150 bp) and long (151-220 bp) cfDNA fragments

(7) was found in the 5hmC data, as in the WGS data (Supplementary Figure

7b-c), though the 5hmC data showed a unique secondary peak at ~320 bp

(Figure 3c, Supplementary Figure 7a). In the WGS data, cfDNA fragments

ranging from 0-220 bp accounted for nearly 100% of fragments ranging from

0-800 bp, but the percent value of the same size interval in the 5hmC

sequencing data ranged from 31.8% to 99.6% (Figure 3f). Thus, we confirmed

the presence of ultra-long fragments (221-500 bp) in the 5hmC cfDNA. We

suspected that the existence of ultra-long fragments may have resulted from

the capture-based technique applied in 5hmC sequencing, and we

consequently assessed the relationship between fragment length and 5hmC

peak number. In ultra-long fragments, the percentage of fragments with more

than one peak was significantly higher than that in non-ultra-long fragments

(0-220 bp) (Mann–Whitney test, p value<0.001) (Figure 3d). Indeed,

enrichment analysis of the ultra-long fragments with more than one peak

revealed primary enrichment at CpG islands (Supplementary Figure 7d). This

indicated that the capture-based sequencing technique is more likely than

WGS to capture ultra-long cfDNA fragments. Another finding was that in all

cancer types, the percentage of ultra-long fragments was significantly lower

than that in the control (Supplementary Figure 7b). We examined the

percentage of ultra-long fragments with more than one 5hmC peak in the four

cancer types and the control and found that the value was significantly lower

in the former (Figure 3e). As reported by a recent study, the stability of

circulating DNA derives mostly from the nucleosome structure (21). We hence

evaluated the proportion of 5hmC cfDNA fragments at 146 bp, 166 bp, 312 bp,

and 332 bp, which indicates the length of one mono-nucleosome and one di-

nucleosome (plus the linker size). The result showed a higher proportion of

mono-nucleosome-sized cfDNA fragments and a lower proportion of di-

nucleosome-sized cfDNA fragments in the cancer group than in the control

(Supplementary Figure 7e).

We also explored the cancer specificity of the pan-cancer preferred end set

built upon WGS data in the 5hmC sequencing data by calculating the ratio of

cancer-specific to health-specific preferred U/D ends. The ratios were

significantly increased in cfDNA samples in three cancer types (except PDAC)

compared with healthy individuals (Figure 3g, Supplementary Figure 5c-d,

Mann–Whitney, p value<0.005). Coverage profiles in 5hmC sequencing data

also showed a higher variance than WGS data in samples from control and

cancer (Supplementary Figure 8). We explored the coverage patterns of short,

long, and ultra-long cfDNA fragments at the 5 Mb bins of the genome. Similar

to WGS data (Supplementary Material), cancer patients exhibited multiple

unstable genomic regions: coverage of short/long/ultra-long fragments was

inconsistent in cancer patients but consistent in controls (Figure 4a,

Supplementary Figure 9a). We analyzed the genome-wide coverage profile

correlation of cancer patients to controls in the 5hmC data and found that the

coverage profiles were consistent among controls and that the correlation

value in cancer patients was significantly lower (Wilcoxon rank-sum test, p

value<0.005). Importantly, the largest difference between cancer samples and

controls was found with respect to the coverage profile of ultra-long fragments

(Wilcoxon rank-sum test, p value<0.005) (Figure 4b).

Genomic distribution of 5hmC in cancer and control cohorts

To gain an understanding of the genomic regions associated with

hydroxymethylation in cfDNA, we first determined 5hmC-enriched loci in the

control and four types of cancer, which were detected as peaks via MACS2

(22). 5hmC-modified regions among samples were compared in 1 kb bins on

the reference genome, and thus a consensus list of the absence and

presence of 5hmC-modified peaks among the samples was obtained. 5hmC

signatures were enriched over genic features, most significantly in the

promoter, 5'UTRs, 3'UTRs, exons, and transcription end sites (TESs) (Figure

5a). Comparison of 5hmC signature enrichment in the four types of cancers

with the control revealed significant differences, whereby 5hmC peak

enrichment was lower in all cancer types than in the control over promoters,

5'UTRs, exons, and CpG islands (Wilcoxon rank-sum test, p value<0.001)

(Figure 5a).

Differential analysis of 5hmC-modified peaks between cancer patients and

controls by Fisher's exact test detected 1,010, 6,395 and 773 differentially

modified 5hmC peaks in GBM, HCC and PDAC, respectively. Moreover, this

pan-cancer 5hmC peak set was able to separate most cancer samples from

the control (Figure 5b). KEGG enrichment analysis of genes located within the

differentially modified 5hmC peaks revealed associated oncogenesis

pathways for each cancer type (Supplementary Figure 10), such as the

neurotrophin signaling pathway and platelet activation for GBM (23, 24), the

MAPK signaling pathway and focal adhesion for HCC (25, 26), and the FoxO

signaling pathway and insulin signaling pathway for PDAC (27).

Cancer detection by combining fragmentomic features and 5hmC signatures using

5hmC sequencing data

As 5hmC sequencing data retain both 5hmC signatures and fragmentomic

information, it is expected to theoretically show high sensitivity and specificity

in cancer detection. A receiver operator characteristic (ROC) curve was used

to evaluate the performance of the classifier. In total, 53 5hmC signatures

were selected, and the AUC of the classifier was 0.876 in the validation set

(sensitivity = 82.26%, specificity = 82.35%) and 0.872 in the test set

(sensitivity = 81.97%, specificity = 82.35%) (Figure 5d-e, Supplementary

Figure 11 c).

Overall, the fragmentomic information in the 5hmC sequencing data

performed well. Using the size profile as features, 40 were selected to build a

pan-cancer prediction model, among which 32 features were of the ultra-long

and longer size range (>= 220 bp) (Figure 5c). The AUC value was 0.981 in

the validation set (sensitivity = 93.55%, specificity = 94.12%) and 0.882 in the

test set (sensitivity = 71.43%, specificity = 88.24%) (Figure 5d-e). Using

preferred ends as features, 37 were selected to build a pan-cancer prediction

model. ROC analysis showed an AUC value of 0.940 in the validation set

(sensitivity = 83.87%, specificity = 94.12%) and 0.899 in the test set

(sensitivity = 73.02%, specificity = 94.12%) (Figure 5d-e, Supplementary

Figure 11a). Regarding the coverage profile of the 5hmC sequencing data, we

explored the genomic distribution of short, long, and ultra-long cfDNA

fragments on a 100 kb window of the genome. This model achieved an AUC

value of 0.946 in the validation set (sensitivity = 80.65%, specificity = 94.12%)

and 0.882 in the test set (sensitivity = 71.43%, specificity = 88.24%) (Figure

5d-e).

Finally, we constructed an integrated cancer screening model by combining

fragmentomic features and 5hmC signatures in 5hmC data. Sixty-three

features, including 10 5hmC signatures, 24 size profile features, 21 coverage

profile features, and 8 preferred end features, were selected to build the pan-

cancer prediction model. The AUC value was 0.927 in the validation set

(sensitivity = 93.44%, specificity = 88.24%) and 0.920 in the test set

(sensitivity = 88.52%, specificity = 82.35%) (Figure 5d-e). The performance of

single-cancer detection is depicted in Supplementary Figure 11b.

With data from the four cancer types, we further explored tissue-of-origin

prediction. The performance of the random forest model with 5hmC

fragmentomic features and the integrated model are shown in Supplementary

Tables 3-5.

Discussion

cfDNA 5hmC signatures are reported to have potential for cancer detection,

such as in lung cancer (28), hepatocellular carcinoma (28), colon cancer (29),

gastric cancer (29), and pancreatic cancer (17, 30). The large-scale cohort of

5hmC sequencing data utilized in this study additionally shows the potential of

cfDNA 5hmC signatures for cancer detection in glioblastoma and pan-cancer.

Besides, we further examined the classification effect using fragmentomic

features in tissue-of-origin prediction of cancer. Most importantly, we explored

cfDNA fragmentomic information in 5hmC sequencing data and found that it is

possible to detect cfDNA hydroxymethylation and fragmentomic markers

simultaneously. The size profile, preferred end, and coverage profile in 5hmC

sequencing data showed a large difference between cancer patients and

healthy individuals. The integrated model covering the 5hmC signature, size

profile, preferred end, and coverage profile contained more information than

the model with the 5hmC signature alone, as revealed by higher sensitivity

and specificity in pan-cancer detection.

Previous work reported a difference in the size profile between specific cancer

patients or pan-cancer patients and healthy controls (12, 13, 31), though the

analyses only focused on lengths less than 320 bp. Here, we characterized

ultra-long fragments in 5hmC sequencing data and identified their size and

coverage profile aberrations in cancer samples. Although it is commonly

acknowledged that ctDNA tends to be more fragmented than cfDNA in normal

tissue (12, 31), we found an even larger size difference in cfDNA between

cancer samples and controls in 5hmC data owing to ultra-long fragments.

Hence, we further expanded the size range to 800 bp by setting 80 adjacent,

non-overlapping bins to capture 10-bp interval signals (32). The coverage

profile is another major improvement we made with regard to previously

reported models based on fragmentation information (9). Indeed, only the

ratio of short to long cfDNA fragments has been analyzed in previously

reported models, with information on cfDNA ultra-long fragments being lost.

Our results suggest that the coverage distribution of size-selected cfDNA

fragments in ultra-long fragments has the strongest inconsistency between

control and cancer samples (Figure 4a). Of note, the classification models

selected the majority of features from ultra-long and longer size range

fragments in the coverage profile model (34/42), size profile model (32/40)

(Figure 5c, Supplementary Figure 12c) and integrated model (41/63),

suggesting that ultra-long features are a major contribution to the

classification model. Further research on the mechanism of the aberration of

the captured ultra-long fragments in cfDNA from cancer samples may provide

insights into the fundamental biological properties of plasma cfDNA from

patients with cancer.

In our work, we constructed a single-base resolution preferred end set based

on a set of WGS data. Our results indicated that the preferred end set had

good generalization ability because the differences between patients with

cancer and controls were also shown in other independent data sets, e.g.,

low-pass WGS data and 5hmC data. It has been reported that referred end

coordinates differ between HCC patients and healthy people at a whole-

genome scale (10). Apart from HCC, our results suggest the cfDNA preferred

end can also be used to distinguish patients with GBM, LUAD, and PDAC. We

herein discussed the distribution of cfDNA preferred ends in genomic features,

open chromatin regions, and nucleosome structure. The distribution of cfDNA

preferred ends can reflect nucleosome positioning, though there is no

apparent preference in genomic features and open chromatin regions,

consistent with previous work (15). However, informative cancer-specific

preferred ends were found to be significantly enriched in promoter and far

away from health-specific open chromatin regions (Figure 2e-f,

Supplementary Figure 4c-d, Supplementary Figure 5c-d).

In summary, our results indicate that cfDNA fragmentomic analysis with a

nonlinear classification algorithm using low-pass 5hmC sequencing data may

provide a simple, low-cost, and highly effective method for cancer detection.

The success of utilizing ultra-long fragments in 5hmC sequencing data as

biomarkers indicates the potential of other capture-based sequencing

approaches, such as cell-free methylated DNA immunoprecipitation and high-

throughput sequencing (cfMeDIP–seq) (33), in the diagnosis of cancer.

Notably, third-generation sequencing methods are expected to detect long

cfDNA molecules, which have been utilized in non-invasive prenatal testing

(34). As third-generation sequencing methods can detect sequence context

and epigenetic modification at the same time, they can provide simultaneous

analysis of genome-wide genetic, fragmentomic and epigenetic detection (35,

36). Moreover, standardized and effective workflows for analysis of cfDNA

fragments need to be developed.

Data availability

444　The 5hmC sequencing data for controls, LUAD, HCC and PDAC were publicly

445　available as described in the Method section. The 5hmC sequencing data for

446　GBM can be accessed at https://ngdc.cncb.ac.cn/gsa-human/s/15mfS98o.

Acknowledgments

This study was supported by the 1·3·5 project for disciplines of excellence,

West China Hospital, Sichuan University (ZYYC20006) to D. Xie; Thousand

Talents Program of the West China Hospital (0040205401F58) to D. Xie;

Sichuan Provincial Foundation of Science and Technology (2020YFS0051) to

D. Xie, (2017SZ0006) to Y. Liu; Clinical Research Innovation Project, West

China Hospital, Sichuan University (19HXCX009) to Y. Liu; the San Hang

Program of the Second Military Medical University, Medical basic research

project of the First Affiliated Hospital, the Second Military Medical University

(2021JCMS16) to W. Zhang; the Science and technology project of Sichuan

Province (2021YFS0109) to A. Li; Post-Doctor Research Project, West China

Hospital, Sichuan University (20HXBH035) to S. Zhang; the high quality

development of Guang 'an People's Hospital (21FZ003) to A. Wei.

Author Contributions

Conception and design: Dan Xie, Zhidong Zhang, Xuenan Pi and Lin Xia.

Collection and assembly of data: Jun Zhang, Xiaoqin Yan, Shuxing Zhang,

Ailin Wei, Jingfeng Liu, Ang Li, Xiaolong Liu, Wei Zhang and Yanhui Liu. Data

analysis and interpretation: Zhidong Zhang, Xuenan Pi, Chang Gao, Xinlei Hu,

Xiyue Yan and Yuer Guo. Manuscript writing: Zhidong Zhang and Xuenan Pi.

All authors read and approved the final version of the manuscript.

Competing interests

The authors declare no competing interests.

References

1.   Sung, H., et al. Global cancer statistics 2020: GLOBOCAN estimates of

incidence and mortality worldwide for 36 cancers in 185 countries. *CA*

Cancer. J. Clin. **0**, 1–41 (2021).

2.   Heitzer, E., Haque, I. S., Roberts, C. E. S. & Speicher, M. R. Current

and future perspectives of liquid biopsies in genomics-driven oncology.

Nat. Rev. Genet. **20**, 71–88 (2019).

3.   Siravegna, G., Marsoni, S., Siena, S. & Bardelli, A. Integrating liquid

biopsies into the management of cancer. *Nat. Rev. Clin. Oncol.* **14**,

531–48 (2017).

4.   Jahr, S., et al. DNA fragments in the blood plasma of cancer patients:

quantitations and evidence for their origin from apoptotic and necrotic

cells. *Cancer. Res.* **61**, 1659-65 (2001).

5.   Serpas, L., et al. Dnase1l3 deletion causes aberrations in length and

end-motif frequencies in plasma DNA. *Proc. Natl. Acad. Sci. U. S. A.*

116, 641-49 (2019).

6. Han, D. S. C. , et al. The Biology of Cell-free DNA Fragmentation and

the Roles of DNASE1, DNASE1L3, and DFFB. *Am. J. Hum. Genet.* **106**,

202-14 (2020).

7. Cristiano, S., et al. Genome-wide cell-free DNA fragmentation in

patients with cancer. *Nature.* **570**, 385-89 (2019).

8. Snyder, M. W., Kircher M., Hill A. J., Daza R. M. & Shendure J. Cell-free

DNA Comprises an In Vivo Nucleosome Footprint that Informs Its

Tissues-Of-Origin. *Cell.* **164**, 57-68 (2016).

9. Chan, K. C., et al. Second generation noninvasive fetal genome

analysis reveals de novo mutations, single-base parental inheritance,

and preferred DNA ends. *Proc. Natl. Acad. Sci. U. S. A.* **113**, E8159-68

(2016).

10. Jiang, P., et al. Preferred end coordinates and somatic variants as

signatures of circulating tumor DNA associated with hepatocellular

carcinoma. *Proc. Natl. Acad. Sci. U. S. A.* **115**, E10925-33 (2018).

11. Jiang, P., et al. Plasma DNA End-Motif Profiling as a Fragmentomic

Marker in Cancer, Pregnancy, and Transplantation. *Cancer. Discov.* **10**,

664-73 (2020).

12. Mouliere, F., et al. Enhanced detection of circulating tumor DNA by

fragment size analysis. *Sci. Transl. Med.* **10**, eaat4921 (2018).

13. Jiang, P., et al. Lengthening and shortening of plasma DNA in

hepatocellular carcinoma patients. *Proc. Natl. Acad. Sci. U. S. A.* **112**,

E1317-25 (2015).

14. Ulz, P., et al. Inferring expressed genes by whole-genome sequencing

of plasma DNA. *Nat. Genet.* **48**, 1273-8 (2016).

15. Sun, K., et al. Orientation-aware plasma cell-free DNA fragmentation

analysis in open chromatin regions informs tissue of origin. *Genome*

Res. **29**, 418-27 (2019).

16. He, B., et al. Tissue-specific 5-hydroxymethylcytosine landscape of the

human genome. *Nat. Commun.* **12**, 4249 (2021).

17. Chen, L., et al. Genome-scale profiling of circulating cell-free DNA

signatures for early detection of hepatocellular carcinoma in cirrhotic

patients. *Cell. Res.* **31**, 589-92 (2021).

18. [dataset]* Hu, X., et al. Integrated 5-hydroxymethylcytosine and

fragmentation signatures as enhanced biomarkers in lung cancer. *Clin.*

Epigenetics. **14**, 15 (2022).

19. Roadmap Epigenomics Consortium, et al. Integrative analysis of 111

reference human epigenomes. *Nature.* **518**, 317-30 (2015).

20. Gaffney, D. J., et al. Controls of nucleosome positioning in the human

genome. *PLoS. Genet.* **8**, e1003036 (2012).

21. Sanchez, C., et al. Circulating nuclear DNA structural features, origins,

and complete size profile revealed by fragmentomics. *JCI. Insight.* **6**,

e144561 (2021).

22. Zhang, Y., et al. Model-based analysis of ChIP-Seq (MACS). *Genome*

Biol. **9**, R137 (2008).

23. Lawn, S., et al. Neurotrophin signaling via TrkB and TrkC receptors

promotes the growth of brain tumor-initiating cells. *J. Biol. Chem.* **290**,

3814-24 (2015).

24. Nolte, I., Przibylla, H., Bostel, T., Groden, C. & Brockmann, M. A.

Tumor-platelet interactions: glioblastoma growth is accompanied by

increasing platelet counts. *Clin. Neurol. Neurosurg.* **110**, 339-42 (2008).

25. Moon, H. & Ro, S. W. MAPK/ERK Signaling Pathway in Hepatocellular

Carcinoma. *Cancers (Basel).* **13**, 3026 (2021).

26. Gnani, D., et al. Focal adhesion kinase depletion reduces human

hepatocellular carcinoma growth by repressing enhancer of zeste

homolog 2. *Cell. Death. Differ.* **24**, 889-902 (2017).

27. Li, J., et al. Knockdown of FOXO3a induces epithelial-mesenchymal

transition and promotes metastasis of pancreatic ductal

adenocarcinoma by activation of the β-catenin/TCF4 pathway through

SPRY2. *J. Exp. Clin. Cancer. Res.* **38**, 38 (2019).

28. Song, C. X., et al. 5-Hydroxymethylcytosine signatures in cell-free DNA

provide information about tumor types and stages. *Cell. Res.* **27**, 1231-

42 (2017).

29. Li, W., et al. 5-Hydroxymethylcytosine signatures in circulating cell-free

DNA as diagnostic biomarkers for human cancers. *Cell. Res.* **27**, 1243-

57 (2017).

30. Guler, G. D., et al. Detection of early stage pancreatic cancer using 5-

hydroxymethylcytosine signatures in circulating cell free DNA. *Nat.*

Commun. **11**, 5270 (2020).

31. Lam, W. K. J., et al. Sequencing-based counting and size profiling of

plasma Epstein-Barr virus DNA enhance population screening of

nasopharyngeal carcinoma. *Proc. Natl. Acad. Sci. U. S. A.* **115**, E5115-

24 (2018).

32. Lo, Y. M., et al. Maternal plasma DNA sequencing reveals the genome-

wide genetic and mutational profile of the fetus. *Sci. Transl. Med.* **2**,

61ra91 (2010).

33. Shen, S. Y., et al. Sensitive tumour detection and classification using

plasma cell-free DNA methylomes. *Nature.* **563**, 579-83 (2018).

34. Yu, S. C. Y., et al. Single-molecule sequencing reveals a large

population of long cell-free DNA molecules in maternal plasma. *Proc.*

Natl. Acad. Sci. U. S. A. **118**, e2114937118 (2021).

35. Tse, O. Y. O., et al. Genome-wide detection of cytosine methylation by

single molecule real-time sequencing. *Proc. Natl. Acad. Sci. U. S. A.*

118, e2019768118 (2021).

36. Liu, Q., et al. Detection of DNA base modifications by deep recurrent

neural network on Oxford Nanopore sequencing data. *Nat. Commun.* **10**,

2449 (2019).

37. [dataset]* Cao, F., et al. Integrated epigenetic biomarkers in circulating

cell-free DNA as a robust classifier for pancreatic cancer. *Clin.*

Epigenetics. **12**, 112 (2020).

38. [dataset]* Cai, Z., et al. Liquid biopsy by combining 5-

hydroxymethylcytosine signatures of plasma cell-free DNA and protein

biomarkers for diagnosis and prognosis of hepatocellular carcinoma.

ESMO Open. **6**, 100021 (2021).

Figure 1 Overview of the study design.

cfDNA from cancer patients and controls were sequenced with WGS.

601   Available 5hmC sequencing data from the same four cancer types and

602   controls were combined with WGS data to get the accurate fragmentomic

603   information and upon it a machine learning model was built to distinguish

604   healthy controls and cancer patients.

Figure 2 Genome-wide cfDNA fragment preferred end map construction.

cfDNA preferred end signals in a nucleosome array region (chr12: 34517269-
34519122) in healthy (a) and HCC (b) subjects. Brown dots at the bottom

represent the predicted nucleosome center loci reported in a previous study (8,

22). (c) Venn diagram showing the intersection of the preferred U and D ends

between the cancer cohort and the control cohort. (d) Circle diagram showing

the density of preferred ends in 1 Mb windows in four types of cancer and in

healthy controls (the order of sample sources from outer ring to inner ring are

control, GBM, HCC, LUAD and PDAC). (e) Enrichment of preferred U/D ends

in promoter regions. (f) Enrichment of preferred U/D ends in health-specific

open chromatin regions. (g) Boxplots showing the ratios of the cancer-specific

to health-specific preferred U/D ends in low-pass WGS data for the four

cancer types (Wilcoxon rank-sum test).

Figure 3 cfDNA fragmentomic information in 5hmC sequencing data.

(a) Violin plots showing fragment size profile comparison between cancer

samples and controls in terms of N50 (Mann–Whitney test). (b) tSNE plot

showing 5hmC sequencing data and WGS data by fragments below 220 bp.

(c) The cfDNA size profile of 5hmC sequencing data. (d) Comparison of the

percentage of fragments with more than one 5hmC peak between ultra-long

and non-ultra-long fragments in every sample. The dashed line indicates the

diagonal line. (e) Violin plots showing the percentage of ultra-long fragments

with more than one 5hmC peak in cancer samples and controls

(Mann–Whitney test). (f) Boxplots showing the percentage of fragments

630　ranging from 0-220 bp and 221-500 bp among fragments ranging from 0-800

631　bp in WGS data and 5hmC sequencing data (Wilcoxon rank-sum test). (g)

632　Boxplots showing the ratios of the cancer-specific to health-specific preferred

633　U/D ends in each of the four cancer types (Wilcoxon rank-sum test).

Figure 4 Size-selected coverage profile and correlation to median health of cfDNA fragments in 5hmC sequencing data.

(a) Genome-wide coverage profile of short, long, and ultra-long cfDNA fragments; color indicates sample-wise correlation to median health. (b)

Coverage profile correlation to median healthy in control and four types of cancer in short, long, and ultra-long cfNDA fragments (Wilcoxon rank-sum test, p value<0.005).

Figure 5 Performance of cancer detection combining fragmentomic

features and 5hmC signatures using 5hmC sequencing data.

(a) Enrichment of 5hmC in genomic features in control and four types of

cancer (Wilcoxon rank-sum test, p value<0.001). (b) Heatmap of differentially

modified 5hmC peaks. (c) Selected size profile features in the size profile

model. (d) ROC curves for the validation set and test set in pan-cancer

diagnosis. (e) Performance evaluation of validation set and test set in pan-

cancer diagnosis.