Genome-wide meta-analysis identifies novel maternal risk variants and enables polygenic prediction of preeclampsia and gestational hypertension

Michael C. Honigberg[1-3]*, Buu Truong[2,3]*, Raiyan R. Khan[4], Brenda Xiao[5], Laxmi Bhatta[6,7], Thi Ha Vy[8], Rafael F. Guerrero[9], Art Schuermans[2,3,10], Margaret Sunitha Selvaraj[2,3], Aniruddh P. Patel[1-3], Satoshi Koyama[2,3], So Mi Jemma Cho[2,3,11], Shamsudheen Karuthedath Vellarikkal[3,12], Mark Trinder[3,13], Sarah M. Urbut[1-3], Kathryn J. Gray[14,15], Ben M. Brumpton[6,7], Snehal Patil[16], Sebastian Zöllner[16], Mariah C. Antopia[17], Genes & Health Research Team, Estonian Biobank Research Team, nuMoM2b Research Team, Richa Saxena[3,15], Girish N. Nadkarni[8], Ron Do[8], Qi Yan[18], Itsik Pe'er[4], Shefali Setia Verma[5], Rajat M. Gupta[3,12], David M. Haas[19], Hilary C. Martin[20], David A. van Heel[21], Triin Laisk[22], Pradeep Natarajan[1-3,15]

[1]Cardiology Division, Department of Medicine, Massachusetts General Hospital, Boston, MA, USA
[2]Cardiovascular Research Center, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA
[3]Program in Medical and Population Genetics, Broad Institute of Harvard and MIT, Cambridge, MA, USA
[4]Department of Computer Science, Columbia University, New York, NY, USA
[5]University of Pennsylvania, Philadelphia, PA, USA
[6]K.G. Jebsen Center for Genetic Epidemiology, Department of Public Health and Nursing, NTNU, Norwegian University of Science and Technology, Trondheim, Norway
[7]HUNT Research Centre, Department of Public Health and Nursing, NTNU, Norwegian University of Science and Technology, Levanger, Norway
[8]The Charles Bronfman Institute for Personalized Medicine, Icahn School of Medicine at Mount Sinai, New York, NY, USA
[9]Department of Biological Sciences, North Carolina State University, Raleigh, NC, USA
[10]Faculty of Medicine, KU Leuven, Leuven, Belgium
[11]Integrative Research Center for Cerebrovascular and Cardiovascular Diseases, Seoul, Republic of Korea
[12]Cardiovascular Division, Department of Medicine, Brigham and Women's Hospital, Boston, MA, USA
[13]Centre for Heart Lung Innovation, University of British Columbia, Vancouver, BC, Canada
[14]Department of Obstetrics and Gynecology, Brigham and Women's Hospital, Boston, MA, USA
[15]Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, USA
[16]Department of Biostatistics and Center for Statistical Genetics, University of Michigan School of Public Health, Ann Arbor, MI, USA
[17]Department of Integrative Biology, University of Texas at San Antonio, San Antonio, TX, USA
[18]Department of Obstetrics and Gynecology, Columbia University, New York, NY, USA
[19]Indiana University School of Medicine, Indianapolis, IN, USA
[20]Department of Human Genetics, Wellcome Sanger Institute, Cambridge, UK
[21]Blizard Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, UK
[22]Estonian Genome Centre, Institute of Genomics, University of Tartu, Tartu, Estonia

*These authors contributed equally.

49   **Corresponding Authors:**
50
51   Michael C. Honigberg, MD MPP
52   Massachusetts General Hospital
53   185 Cambridge Street, CPZN 3.187
54   Boston, MA 02114
55   Phone: 617-726-1843
56   Fax: 617-726-2203
57   Email: mhonigberg@mgh.harvard.edu
58   Twitter: @mchonig
59
60   Pradeep Natarajan, MD MMSc
61   Massachusetts General Hospital
62   185 Cambridge Street, CPZN 3.184
63   Boston, MA 02114
64   Phone: 617-726-1843
65   Fax: 617-726-2203
66   Email: pnatarajan@mgh.harvard.edu
67   Twitter: @pnatarajanmd

Abstract

Preeclampsia and gestational hypertension are common pregnancy complications

associated with adverse maternal and offspring outcomes. Current tools for prediction,

prevention, and treatment are limited. We tested the association of maternal DNA sequence

variants with preeclampsia in 20,064 cases and 703,117 controls and with gestational

hypertension in 11,027 cases and 412,788 controls across discovery and follow-up cohorts

using multi-ancestry meta-analysis. Altogether, we identified 18 independent loci associated

with preeclampsia/eclampsia and/or gestational hypertension, 12 of which are novel (e.g.,

MTHFR-CLCN6, *WNT3A*, *NPR3*, *PGR*, and *RGL3*), including two loci (*PLCE1*, *FURIN*)

identified in multi-trait analysis. Identified loci highlight the role of natriuretic peptide signaling,

angiogenesis, renal glomerular function, trophoblast development, and immune dysregulation.

We derived genome-wide polygenic risk scores that predicted preeclampsia/eclampsia and

gestational hypertension in external datasets, independent of first trimester risk markers.

Collectively, these findings provide mechanistic insights into the hypertensive disorders of

pregnancy and advance pregnancy risk stratification.

The hypertensive disorders of pregnancy (HDPs) represent a leading global cause of

maternal and neonatal morbidity and mortality and account for approximately 14% of maternal

deaths worldwide[1,2]. Up to 15% of child-bearing women experience an HDP in at least one

pregnancy[3,4]. The HDPs include preeclampsia, defined as new-onset hypertension or worsening

hypertension after 20 weeks' gestation plus proteinuria or other evidence of end-organ

dysfunction; gestational hypertension, defined as new-onset hypertension without

accompanying features of preeclampsia; and eclampsia, defined as progression of

preeclampsia to maternal seizures[5,6]. In addition to short-term risks of end-organ failure and

death in the absence of prompt recognition and treatment, those who develop HDPs have

roughly two-fold long-term risk of cardiovascular disease events and premature cardiovascular

mortality compared with those who experience only normotensive pregnancies for reasons that

remain incompletely understood[7,8].

The pathophysiology of the HDPs is increasingly recognized to be heterogeneous with

both maternal and fetal contributions. In the contemporary model of preeclampsia

pathophysiology, defective trophoblast invasion in early placental development and incomplete

remodeling of the maternal spiral arteries lead to placental ischemia later in gestation[1,9,10]. The

distressed placenta secretes an excess of circulating anti-angiogenic proteins (e.g., soluble fms-

like tyrosine kinase receptor 1 [sFlt-1] and soluble endoglin) which induce the systemic maternal

endothelial dysfunction and vasoconstriction that drive the clinical manifestations of

preeclampsia (hypertension and proteinuria)[11]. In addition, maternal cardiometabolic risk factors

(e.g., pre-pregnancy chronic hypertension, diabetes, obesity) and pre-pregnancy kidney and

autoimmune disease strongly predict preeclampsia[1,12] and influence early placentation as well

as maternal vascular adaptation to pregnancy[10,13].

Genetic analysis may yield novel mechanistic insights into the pathophysiology of the

HDPs and may also help improve pregnancy risk prediction. An estimated 31-35% of

preeclampsia predisposition has been attributed to maternal genetics using familial aggregation-

based approaches[14,15]. However, few genetic loci linked to preeclampsia have been identified

and robustly validated to date. Several fetal variants near the *FLT1* gene, which encodes

placenta-derived sFlt-1, have been reported to associate with preeclampsia[16,17]. Steinthorsdottir

et al. recently performed the largest maternal GWAS of preeclampsia to date and identified

associations near *FTO* (the first reported obesity-associated locus)[18], *ZNF831*, and several

other blood pressure-associated genes (*MECOM*, *FGF5*, and *SH2B3*) in combined meta-

analysis of 12,150 cases and 164,098 controls[17]. In addition, several studies have now reported

that increased maternal polygenic risk of hypertension associates with risk of HDPs[17,19-22].

In this work, we performed expanded multi-ancestry GWAS meta-analysis for

preeclampsia/eclampsia and separately performed GWAS for gestational hypertension. We

then used these results to train and test polygenic risk scores (PRS) for each outcome in

independent datasets (**Extended Data Fig. 1**).

Results

Association of common genetic variants with preeclampsia/eclampsia and gestational

hypertension.

We tested the association of common variants (minor allele frequency [MAF] ≥1%) with

preeclampsia/eclampsia among 17,150 cases and 451,241 controls in discovery analysis

(78.0% European, 21.2% Asian, 0.5% admixed American, and 0.3% African ancestry;

Supplementary Table 1) using multi-ancestry fixed-effects meta-analysis in METAL[23]. Cases

were identified principally using *International Classification of Diseases* (ICD) codes and

phecodes corresponding to preeclampsia and, where available, eclampsia (**Supplementary**

Tables 1-3); controls were generally either women with exclusively normotensive pregnancies

or all women without codes corresponding to hypertension in pregnancy[17]. In discovery

analysis, we identified 12 independent loci at the commonly used statistical significance

threshold of $P<5\times10^{-8}$, including 6 previously nominated by Steinthorsdottir et al. in maternal or

fetal GWAS (*MECOM* [3q26], *FGF5* [4q21], *SH2B3* [12q22], *FLT1* [13q12], *FTO* [16q12], and

ZNF831 [20q13])[17] and 6 additional loci (*MTHFR-CLCN6* [1p36], *WNT3A* [1q42], *MICA* [6p21],

LINC00484 [9q22], *PGR* [11q22], and *RGL3* [19p13]; **Table 1, Supplementary Fig. 2a,**

Supplementary Table 4).

We pursued replication of these GWAS results in 4 follow-up cohorts that collectively

included 2,914 preeclampsia/eclampsia cases and 251,876 controls (96.7% European, 3.1%

African, and 0.3% admixed American ancestry). We replicated 7 of 12 associations from

discovery analysis with $P<0.05$ and consistent direction of effect, including the novel

associations at *MTHFR-CLCN6*, *PGR*, and *RGL3* (**Table 1**). Ten associations had consistent

direction of effect in follow-up cohorts, and 11 of 12 associated loci retained genome-wide

significance in combined meta-analysis of preeclampsia/eclampsia discovery and follow-up

cohorts. In meta-analysis of discovery and follow-up cohorts, 2 additional loci attained genome-

wide significance (*FGL1* [8p22] and *UPB1* [22q11]), yielding a total of 13 loci associated with

preeclampsia/eclampsia with genome-wide significance in combined meta-analysis (**Fig. 1a**).

We did not observe inflation in test statistics (lambda genomic inflation factor: 1.038,

Supplementary Fig. 1a). There was no discernible heterogeneity of these associations across

ancestries (**Supplementary Table 5**). Conditional analysis in GCTA-COJO[24] identified a second

independent association on chromosome 20 near *ZBTB46* (lead variant rs4809370, OR 1.08,

$P=1.4 \times 10^{-8}$).

We next tested the association of common variants with gestational hypertension among

8,961 cases and 184,925 controls in discovery analysis (91.3% European, 6.7% Asian, 0.7%

African, 1.3% admixed American) and among 2,066 cases and 227,863 controls in follow-up

cohorts (96.3% European, 3.4% African, 0.3% admixed American, **Supplementary Tables 1-2**).

Gestational hypertension cases were identified primarily based on qualifying ICD codes for

gestational hypertension and an absence of qualifying codes for preeclampsia/eclampsia. In

discovery analysis, we identified 7 independent genome-wide significant loci associated with

gestational hypertension, including 4 also associated with preeclampsia/eclampsia (*MECOM*,

FGF5, *RGL3*, and *ZNF831*) and 3 additional associations (*NPR3* [5p13], *TNS2-CSAD* [12q13],

and *PREX1* [20q13]; **Table 2**). Four of 7 significant associations replicated with *P*<0.05 in

follow-up cohorts (*FGF5*, *RGL3*, *PREX1*, and *ZNF831*), and all 7 loci had consistent direction of

effect in follow-up cohorts. In combined meta-analysis of discovery and follow-up cohorts, 6 of 7

loci retained genome-wide significance, and the *MTHFR-CLCN6* locus (1p36) additionally

reached genome-wide significance, yielding a total of 7 loci associated with gestational

hypertension in combined meta-analysis (**Fig. 1b**). As with preeclampsia/eclampsia, we did not

observe inflation in test statistics (lambda genomic inflation factor: 0.976, **Supplementary Fig.**

1b). Stratified analyses suggested potential heterogeneity of association by ancestry

($P_{\text{heterogeneity}}$=0.001) at the *MECOM* locus, with an inverse association with the lead risk variant

observed in those with admixed American ancestry (**Supplementary Table 5**).

Genetic correlation across hypertension-related phenotypes.

175   We used cross-trait LD score regression[25] to assess genetic correlations among

preeclampsia/eclampsia, gestational hypertension, systolic blood pressure (SBP), and diastolic

blood pressure (DBP)[26]. Preeclampsia/eclampsia and gestational hypertension were strongly

genetically correlated (r_g = 0.71, SE = 0.08). SBP demonstrated stronger genetic correlation with

gestational hypertension (r_g = 0.73, SE = 0.06) vs. preeclampsia/eclampsia (r_g = 0.52, SE =

0.05). Of note, correlation of SBP with gestational hypertension (r_g = 0.73) was larger than that

of SBP with DBP (r_g = 0.62, SE = 0.03, in the Million Veteran Program[26], with similar genetic

correlation between SBP and DBP observed previously in the UK Biobank[27]). Genetic

correlations with the HDPs were stronger for SBP vs. DBP (**Supplementary Table 6**).

 Multi-trait analysis of GWAS

Given the high degree of genetic correlation observed between preeclampsia/eclampsia

and gestational hypertension noted earlier, we used multi-trait analysis of genome-wide

association summary statistics (MTAG)[28] to boost power to identify additional associated

variants. Consistent with this high degree of correlation, MTAG yielded very similar results for

each trait (displayed for preeclampsia/eclampsia in **Extended Data Fig. 2**). MTAG identified 2

additional loci with genome-wide significance (**Supplementary Table 7**): *PLCE1* (10q23), a

blood pressure-associated gene involved in glomerular podocyte development[29], which also

narrowly missed statistical significance in combined meta-analysis of gestational hypertension

discovery and follow-up cohorts ($P=6.0\text{x}10^{-8}$); and *FURIN* (15q26), whose product is a protein

convertase involved in processing pro-natriuretic peptides[30] and whose expression is decreased

in preeclamptic placentas[31].

Gene prioritization at preeclampsia/eclampsia and gestational hypertension risk loci.

To prioritize causal genes, we first performed colocalization analysis with expression

quantitative trait loci (eQTLs) within ±500 kilobases of lead variants across 52 tissues in the

Genotype-Tissue Expression project (GTEx) (**Supplementary Tables 8-9**)[32]. Colocalization

analyses implicated *FGF5* and *NPR3* as causal genes at their respective loci. The lead variant

at the *MTHFR-CLCN6* locus colocalized with *CLCN6* eQTLs as well as expression of *NPPA*

(which encodes the precursor to atrial natriuretic peptide [ANP]). The lead

preeclampsia/eclampsia variant at the *ZNF831* locus colocalized with multiple genes but most

strongly with *ZBTB46* expression, including in arterial tissue. We also observed multiple

colocalizations with *WNT3A* (*WNT3A*, *GJC2,* and mitochondrial proteins *IBA57* and *MPRL55*),

MICA (*CLIC1* and psoriasis-associated genes *TCF19*, *CCHCR1*, and *PSORS1C1*), and *RGL3*

(*ZNF627* and *EPOR*). We observed no strong colocalizations with lead variants at *LINC00484*,

PGR, *SH2B3*, *FLT1*, *FTO*, or *TNS2-CSAD*.

Next, we queried variant-to-gene evidence in Open Targets Genetics v7

(**Supplementary Tables 10-11**)[33] and generated polygenic priority scores (PoPS,

Supplementary Table 12)[34] for lead variants. Both approaches nominated *MECOM*, *FGF5*,

SH2B3, *FTO*, and *NPR3* as the most likely causal gene and their respective loci. PoPS

prioritized *NPPA* as the most likely causal gene at the *MTHFR-CLCN6* locus and *TRPC6* as the

most likely causal gene at the *PGR* locus.

To further understand how identified genes might influence HDP risk, we queried lead

maternal variants in the fetal GWAS for maternal preeclampsia from Steinthorsdottir et al.[17]

(**Supplementary Table 13**). As published previously[16,17], the lead *FLT1* variant was strongly

associated with preeclampsia (P=3.9x10^{-11}); all other lead variants had P>10^{-4} in the fetal

GWAS. In addition, we queried nearest genes and those prioritized by colocalization, variant-to-

gene scores, and/or PoPS in a publicly available database of the human placental transcriptome

including preeclampsia cases and controls (**Supplementary Table 14**)[35]. Consistent with the

correlation of increased circulating placental sFlt-1 with preeclampsia incidence[11,36], *FLT1* gene

expression was increased in preeclamptic placentas (log$_2$(fold change) 0.39, false discovery

rate-adjusted P=0.003). Expression of *WNT3A,* which occurs almost exclusively in the

placenta[37], was increased in preeclamptic placentas vs. healthy controls (log$_2$(fold change) 0.21,

adjusted P=0.029), in line with another recent study[38]. *OBSCN,* which was most strongly

prioritized by PoPS at the *WNT3A* locus, was also overexpressed in preeclamptic vs. control

placentas (log$_2$(fold change) 0.18, adjusted P=0.037). Furthermore, preeclamptic placentas

demonstrated lower expression of *ARHGAP42*—which sits adjacent to *PGR* and encodes Rho

GTP-ase activating protein 42, a known regulator of vascular tone and blood pressure

expressed selectively in smooth muscle cells[39]—compared with controls (log$_2$(fold change)

-0.18, adjusted P=0.004).

We analyzed expression of prioritized genes in a dataset of single-nuclei RNA

sequencing (snRNA-seq) from non-atherosclerotic human aortic tissue. SnRNA-seq identified

subpopulations of vascular smooth muscle cells, fibromyocytes, fibroblasts, endothelial cells,

macrophages, natural killer T cells, and neuronal cells. The greatest enrichment was seen in the

two endothelial cell (EC) populations and in macrophages (**Extended Data Fig. 3**). The EC1

subpopulation is enriched for genes in angiogenesis and lipoprotein assembly and clearance,

while the EC2 subpopulation is enriched for genes in extracellular matrix production and integrin

expression[40]. Relative expression in ECs vs. other cell types was strongest for *FLT1* (EC1/EC2),

ZBTB46 (EC1), and *MECOM* (EC2).

Training and testing polygenic risk scores for preeclampsia/eclampsia and gestational

hypertension.

We used PRS-CS[41] to construct genome-wide PRS for preeclampsia/eclampsia

(PRS$_{preeclampsia}$) and gestational hypertension (PRS$_{GH}$) from our corresponding discovery GWAS

summary statistics. In addition, since blood pressure polygenic risk has previously been

associated with HDPs[17,19,20], we used PRS-CS to derive a PRS for SBP (PRS$_{SBP}$) using the SBP

GWAS from the Million Veteran Program[26] to compare prediction across scores and determine

whether a linear combination of each HDP PRS and PRS$_{SBP}$ improves performance.

We tuned polygenic scores among 423 women with a history of

preeclampsia/eclampsia, 482 women with a history of gestational hypertension only, and

211,129 parous women without a reported history of HDPs in the UK Biobank. The global

shrinkage parameter of 1×10^{-4} was chosen for PRS$_{preeclampsia}$ and PRS$_{SBP}$ and of 1×10^{-6} for

PRS$_{GH}$ as these values generated the highest R^2 (**Supplementary Table 15**). A linear

combination of PRS$_{preeclampsia}$ and PRS$_{SBP}$ (PRS$_{preeclampsia+SBP}$) improved performance vs. each

score individually for the outcome of preeclampsia (**Supplementary Table 16**).

The individual and combination PRS tuned in the UK Biobank were carried forward for

external validation in two complementary datasets: a Norwegian population-based cohort linked

to the Medical Birth Register of Norway (HUNT, preeclampsia/eclampsia only) and a

prospective U.S. pregnancy cohort (nuMoM2b). Among 25,582 Norwegian women in HUNT

(1,569 [6.1%] with preeclampsia/eclampsia), the prevalence of preeclampsia/eclampsia ranged

from ~4% among women in the bottom decile of $PRS_{preeclampsia+SBP}$ to ~10% among the top decile

of $PRS_{preeclampsia+SBP}$ (**Fig. 2a**). After adjustment for age at birth, age^2, and the first 10 principal

components (PC) of ancestry, the OR corresponding to the top 10%vs. bottom 90% of

$PRS_{preeclampsia+SBP}$ was 1.85 (95% CI 1.61-2.13, P=6.3x10^{-18}). $PRS_{preeclampsia+SBP}$ increased

Nagelkerke's R^2 by 28.5% compared with the $PRS_{preeclampsia}$ alone and by 79.3% compared with

PRS_{SBP} alone (**Supplementary Table 17**).

We next tested PRS performance in the prospective, multi-ancestry nuMoM2b cohort of

U.S. women recruited in the first trimester of their first pregnancy, including 481 (6.4%) who

developed preeclampsia, 1,319 (17.5%) who developed gestational hypertension, and 5,744

with normotensive pregnancies (overall 73.6% European, 16.5% African, and 1.0% admixed

American ancestry). Rates of preeclampsia/eclampsia ranged from ~4% among women in the

bottom decile of $PRS_{preeclampsia+SBP}$ to ~10% among the top decile of $PRS_{preeclampsia+SBP}$ (**Fig. 2b**).

Rates of gestational hypertension ranged from ~9% among women in the bottom decile of

PRS_{GH+SBP} to ~24% among the top decile of PRS_{GH+SBP} (**Fig. 2c**). As in HUNT, incorporating

SBP PRS in linear combination boosted PRS performance for HDPs, especially for gestational

hypertension, although PRS performance was better for European ancestry vs. other women

(**Supplementary Table 18**). After adjustment for age, PC 1-10, and self-reported race/ethnicity,

$PRS_{preeclampsia+SBP}$ and PRS_{GH+SBP} each predicted their respective outcomes

(preeclampsia/eclampsia: OR 1.78, 95% CI 1.35-2.31, for top 10% vs. bottom 90%

$PRS_{preeclampsia+SBP}$, P=2.6x10^{-5}; gestational hypertension: OR 1.52, 95% CI 1.26-1.84, for top 10%

vs. bottom 90% PRS_{GH+SBP}, P=1.0x10^{-5}). As pre-pregnancy hypertension and obesity are

established clinical predictors of HDPs, we performed further adjustment for first trimester SBP,

antihypertensive medication use (as a marker of chronic hypertension), and body mass index

(BMI). After this additional adjustment, the scores both remained predictive

(preeclampsia/eclampsia: OR 1.64, 95% CI 1.23-2.15, for top 10% vs. bottom 90%

$PRS_{preeclampsia+SBP}$, P=5.1x10^{-4}; gestational hypertension: OR 1.53, 95% CI 1.26-1.85, for top 10%

vs. bottom 90% PRS_{GH+SBP}, P=1.0x10^{-5}). Compared with a model including age, PC 1-10, self-

reported race/ethnicity, first trimester SBP, antihypertensive medication use, and first trimester

BMI, addition of $PRS_{preeclampsia+SBP}$ improved the C-statistic for preeclampsia/eclampsia from

0.690 to 0.701 (+0.011, 95% CI 0.001-0.021, Delong's P=3.7x10^{-2}). Similarly, addition of

PRS_{GH+SBP} improved the C-statistic for gestational hypertension from 0.649 to 0.659 (+0.010,

95% CI 0.003-0.018, Delong's P=5.7x10^{-3}).

Low-dose aspirin starting after 12 weeks' gestation represents an evidence-based but

underutilized[42] strategy to reduce risk of preeclampsia. To further probe the potential clinical

impact of incorporating PRS to guide aspirin allocation, we examined aspirin eligibility according

to current U.S. Preventive Service Task Force major criteria[43] with and without addition of

$PRS_{preeclampsia+SBP}$ as an additional eligibility criterion in the nuMoM2b cohort. Among singleton,

nulliparous women (i.e., the population enrolled in nuMoM2b), major criteria for aspirin eligibility

are chronic pre-pregnancy hypertension, pre-gestational diabetes, kidney disease, and

autoimmune disease.[43] Major risk factors identified only a small proportion of the cohort that

developed HDPs (**Table 3**), and aspirin would have been recommended in only 17.5% of

women who developed preeclampsia/eclampsia (i.e., sensitivity of major risk factors alone was

17.5%), with a positive predictive value of 12.8%. Incorporating the top 10% of $PRS_{preeclampsia+SBP}$

increased identification of the aspirin-eligible proportion to 30.4% of women with

preeclampsia/eclampsia (i.e., sensitivity 30.4% [95% CI 26.2-34.5%]) with specificity of 83.3%

(95% CI 82.5-84.2%), positive predictive value of 11.0% (95% CI 9.3-12.7%), and negative

predictive value of 94.6% (95% CI 94.1-95.2%; **Table 3**). Expanding aspirin eligibility further to

include the top 25% of $PRS_{preeclampsia+SBP}$ captured nearly half (47.0%) of women who developed

preeclampsia/eclampsia. Addition of high $PRS_{preeclampsia+SBP}$ to major risk factors to up-classify

risk of preeclampsia/eclampsia vs. other pregnancies yielded net reclassification of +1.8% (95%

CI -0.3 to +4.0%) for top 5% $PRS_{preeclampsia+SBP}$, +4.3% (95% CI 1.3-7.3%) for top 10%

$PRS_{preeclampsia+SBP}$, and +8.3% (95% CI 3.9-12.6%) for top 25% $PRS_{preeclampsia+SBP}$ (**Table 3**).

Phenome-wide associations with preeclampsia/eclampsia and gestational hypertension

polygenic risk.

We performed sex-stratified phenome-wide association analysis (pheWAS) for

$PRS_{preeclampsia}$ and PRS_{GH} across 1,445 phecode-based phenotypes in the UK Biobank.

$PRS_{preeclampsia}$ was significantly associated with 36 phenotypes in women and 37 phenotypes in

men with Bonferroni-corrected statistical significance ($P<0.05/1,445 = 3.5x10^{-5}$); PRS_{GH} was

significantly associated with 25 and 32 phenotypes in women and men, respectively (**Fig. 3**).

$PRS_{preeclampsia}$ and PRS_{GH} were most strongly associated with hypertension in both sexes

($PRS_{preeclampsia}$: OR_{women} 1.15 per SD, 95% CI 1.14-1.16, $P=1.4x10^{-175}$; OR_{men} 1.12 per SD, 95%

CI 1.11-1.13, $P=7.5x10^{-112}$; PRS_{GH}: OR_{women} 1.15 per SD, 95% CI 1.14-1.16, $P=5.1x10^{-184}$; OR_{men}

1.13 per SD, 95% CI 1.11-1.14, $P=1.3x10^{-133}$). Other strong phenotypic associations included

hypercholesterolemia, type 2 diabetes, obesity, and atherosclerotic cardiovascular disease

(**Supplementary Tables 19-20**). $PRS_{preeclampsia}$ predicted ischemic heart disease in women (OR

1.09 per SD, 95% CI 1.07-1.11, $P=4.4x10^{-27}$) and men (OR 1.09 per SD, 95% CI 1.07-1.10,

$P=2.6x10^{-40}$), as did PRS_{GH} (OR_{women} 1.07 per SD, 95% CI 1.05-1.08, $P=4.4x10^{-16}$; OR_{men} 1.08

per SD, 95% CI 1.07-1.09, $P=1.4x10^{-35}$). These similar associations between sexes suggest that

most genes identified are not pregnancy-specific, but rather that pregnancy likely unmasks

underlying risk. $PRS_{preeclampsia}$ was also associated with several autoimmune phenotypes,

including celiac disease, type 1 diabetes, hypothyroidism (in women), and a suggestive

association with rheumatoid arthritis in women ($P=5.7x10^{-6}$), whereas type 1 diabetes and celiac

disease were not significantly associated with PRS_{GH} in either sex.

Discussion

We present an expanded multi-ancestry maternal genome-wide association analysis of

preeclampsia/eclampsia and, to our knowledge, the first distinct GWAS of gestational

hypertension, two pregnancy complications on the spectrum of HDPs with substantial

associated public health impact. Altogether, we identified 18 independent genomic loci

associated with preeclampsia/eclampsia and/or gestational hypertension: 13 loci associated

with preeclampsia/eclampsia; 7 loci associated with gestational hypertension, of which 4 overlap

with preeclampsia/eclampsia; and 2 additional loci associated with both phenotypes in multi-trait

analysis. Identified loci highlight the role of angiogenesis and endothelial cell function (*FLT1*,

ZBTB46), natriuretic peptide signaling (*NPPA*, *NPR3*, *FURIN*), renal glomerular function

(*TRPC6*, *TNS2*, *PLCE1*), and immune dysregulation (*MICA*, *SH2B3*) in the pathogenesis of

these conditions, with some loci (*FLT1*[16,17,36], *WNT3A*[38,44]) previously described to influence risk

via the fetal genome. Furthermore, we demonstrated that an HDP PRS augmented by a systolic

blood pressure PRS stratified risk of HDP. Polygenic risk scores predicted HDP risk among

nulliparous women independent of first trimester blood pressure and BMI, indicating potential

clinical utility of these scores risk for pregnancy risk stratification. Collectively, these findings

may have implications for advancing HDP prediction, prevention, and treatment.

First, our findings provide insights into mechanisms of HDP pathogenesis and

underscore the causal role of blood pressure. As in previous GWAS of preeclampsia[17], most

genetic associations we identified were also associated with blood pressure[29], and genetic

correlation between blood pressure and the HDPs, especially gestational hypertension, was

high. These findings align with prior work demonstrating heightened polygenic BP risk in those

with HDPs[17,19,20]. Indeed, we observed that integrating SBP PRS with HDP PRS in linear

combination further improved predictive performance for both preeclampsia/eclampsia and

gestational hypertension. The recently published randomized CHAP trial of treatment for mild

chronic hypertension in pregnancy demonstrated that lowering blood pressure

pharmacologically reduced risk of progression to preeclampsia[45], supporting the notion that

elevated blood pressure is not merely a clinical manifestation of the HDPs but also plays a

causal role in disease pathogenesis.

Second, our GWAS findings suggest a key role of natriuretic peptide signaling in the

pathogenesis of the HDPs. The natriuretic peptides (e.g., ANP, B-type natriuretic peptide [BNP])

promote renal sodium excretion and counteract renin-angiotensin and sympathetic nervous

system activation[46]. The lead variant at the *MTHFR-CLCN6* locus colocalized with *NPPA*

expression in atrial tissue, its primary site of expression. ANP (the protein product of *NPPA*)

also plays a role in uterine decidualization and spiral artery remodeling[47], a process known to be

impaired in the early pathogenesis of preeclampsia[1,10]. Furthermore, ANP is cleared from the

circulation by the protein product of *NPR3*[48], which was identified in the GWAS of gestational

hypertension. Human data also support accelerated ANP clearance in the setting of

preeclampsia[49]. Of note, our lead risk variant at the *MTHFR-CLCN6* locus has been previously

associated with reduced levels of circulating N-terminal pro-BNP[50]. A recent analysis from the

nuMoM2b cohort found that first trimester levels of N-terminal pro-BNP were unexpectedly lower

among women who subsequently developed HDPs later in pregnancy after adjustment for race

and BMI[51]. Collectively, these findings suggest that a relative deficiency in endogenous

natriuretic peptide signaling may predispose to HDPs. Synthetic natriuretic peptides have been

developed previously (e.g., nesiritide), and the natriuretic peptides may represent a future

therapeutic target for direct or indirect modulation toward HDP prevention and/or treatment.

Third, our findings suggest several other potential novel mechanisms underlying HDP

risk. Although *ZNF831* has been previously associated with hypertension, the mechanisms by

which risk variants increase blood pressure have been unclear. In our conditional analysis at the

ZNF831 locus, we detected a second independent association with *ZBTB46*, which was also

strongly colocalized with the lead *ZNF831* variant in multiple tissues, including arterial tissue.

ZBTB46 is a transcription factor expressed in dendritic cells[52,53] and in vascular endothelial

cells, consistent with our observed expression in aortic endothelial cells, and *ZBTB46*

overexpression suppresses endothelial cell proliferation and angiogenesis *in vitro*[54].

Furthermore, ZBTB46 is sensitive to shear stress[54], which may have relevance to the

hyperdynamic hemodynamic state of pregnancy.

In addition, we identified and replicated a novel association in the intergenic region

between *PGR* and *TRPC6* (rs2508372), which has several plausible mechanistic links to the

HDPs. Along with other newly identified HDP-associated loci (*TNS2*[55] and *PLCE1*[29]), *TRPC6*

has also been linked to glomerular function. It has been implicated in focal segmental

glomerulosclerosis and diabetic nephropathy[56] and mediates the proteinuria and renal

dysfunction induced by exposure to hypertension and diabetes[57]. In addition, the progesterone

receptor mediates the physiologic response to progesterone, including pregnancy-associated

vasodilation[58]. As noted, *ARHGAP42* (adjacent to *PGR*), which was found to have reduced

expression in preeclamptic placentas, regulates vascular tone and blood pressure[39].

Furthermore, this same intergenic lead variant has been associated with reduced circulating

levels of matrix metalloproteinase-10[59], which mediates vascular endothelial growth factor

(VEGF)-induced angiogenesis[60]. Further research is necessary to clarify which of these

potential mechanisms primarily mediate the preeclampsia/eclampsia risk associated with the

PGR/TRPC6 locus.

Fourth, associations at *MICA* and *SH2B3* highlight the role of immune function in

preeclampsia pathogenesis. Differences in T-cell phenotypes and circulating pro- and anti-

inflammatory cytokines in preeclampsia are well described[9,10]. We detected an association near

MICA (encoding major histocompatibility complex class I polypeptide-related sequence A) within

the *HLA* region, which has also been linked to multiple autoimmune diseases and may reflect

the importance of maternal immune tolerance of fetal cells at the maternal-fetal interface in early

placentation[10]. Indeed, we found that preeclampsia/eclampsia PRS predicted type 1 diabetes,

celiac disease, and hypothyroidism. Subclinical thyroid dysfunction has been associated with

increased risk for preeclampsia[61], and previous Finnish data suggested term preeclampsia

predicts incident hypothyroidism later in life[62]. SH2B3 (also known as LNK) is expressed

primarily in endothelial and hematopoietic cells and negatively regulates cytokine signaling;

reduced SH2B3 function has been linked to atherosclerosis as well as several autoimmune

diseases[63]. The lead variant in our preeclampsia/eclampsia GWAS at *SH2B3* is in linkage

disequilibrium (D'=0.96, R^2=0.91)[64] with the well-described coronary artery disease risk allele at

this locus (rs3184504)[65]. Our lead *SH2B3* variant was also associated with increased leukocyte,

red cell blood, and platelet counts in the UK Biobank (nealelab.is/uk-biobank) and with

heightened levels of vascular cell adhesion protein 1, platelet glycoprotein IB alpha chain,

interleukin-2 receptor, and lymphocyte activation gene 3 protein, among other immune-related

proteins[50]. Recent data indicate that reduced SH2B3 function promotes neutrophil extracellular

trap formation, a process implicated in preeclampsia pathogenesis[66,67], and arterial

thrombosis[68]. This process was dependent on the presence of oxidized phospholipids[68], which

are increased in the setting of preeclampsia vs. normotensive pregnancy[69,70].

Fifth, we observed that polygenic risk scores for preeclampsia/eclampsia and gestational

hypertension were predictive in multiple external test cohorts and independent of first trimester

risk indicators (blood pressure and BMI), with women in the top PRS decile demonstrating 65-

85% greater risk of preeclampsia/eclampsia vs. other women. Predictive accuracy of clinical risk

factors for HDPs is modest[71,72]. Meta-analysis indicates that personal history of preeclampsia,

pre-pregnancy chronic hypertension, and pre-pregnancy diabetes are associated with relative

risks of 8.4, 5.1, and 3.7, respectively, for development of preeclampsia. However, nulliparity

carries the largest population attributable fraction of associated risk for preeclampsia

(approximately one-third)[12], and most affected women lack any overt pre-pregnancy risk factors

other than nulliparity[6]. Low-dose aspirin after 12 weeks' gestation represents one evidence-

based strategy to mitigate risk of preeclampsia and preterm birth in women at increased risk for

preeclampsia[43]. Improving pregnancy risk prediction, especially in the first pregnancy, therefore

remains a pressing clinical need to optimize HDP prevention[43,73]. First trimester screening

algorithms have been developed, with the UK Fetal Medicine Foundation combined prediction

model[74] incorporating clinical factors, mean arterial pressure, uterine artery pulsatility index, and

maternal serum pregnancy-associated plasma protein A and placental growth factor being most

extensively validated to date[75], although not currently endorsed by UK or US care guidelines[43].

Future studies are required to ascertain whether PRS may augment existing risk algorithms. In

contrast with markers measured during pregnancy, PRS can be calculated anytime from birth,

including preconception, and may therefore also inform preconception counseling and health

optimization.

Although our GWAS included substantially more non-European individuals than prior

GWAS, >80% of individuals were European, and as such, the preeclampsia/eclampsia and

gestational hypertension PRS generally performed better in European ancestry women vs.

others, consistent with many prior published PRS and a well-recognized challenge in

contemporary genetics[76]. Ongoing efforts to include accurate, detailed pregnancy and

reproductive history phenotypes in diverse genetic datasets and increase representation of non-

European individuals will be critical to improve genetic discovery and cross-ancestry polygenic

prediction and to achieve genomic equity[76,77].

This study should be considered in the context of other limitations. Due to HDP

phenotyping limitations in large datasets using ICD code-based ascertainment, some

participants may have had preeclampsia superimposed on chronic hypertension rather than *de*

novo preeclampsia, which may enrich genetic associations for hypertension predilection. In a

subset of cohorts, however, the control group included women with chronic hypertension in

pregnancy. Validation studies of ICD codes and registry diagnoses demonstrate that these

approaches have modest sensitivity but high specificity and positive predictive value (>80%) in

comparison with adjudicated HDP diagnoses[78-80] and that accuracy may be higher for severe vs.

mild preeclampsia[81]. We were unable to examine more granular HDP subtypes such as

preeclampsia with severe features, HELLP (hemolysis, elevated liver enzymes, and low

platelets) syndrome, preterm vs. term vs. postpartum onset, or HDP with vs. without

accompanying fetal growth restriction. The underlying pathophysiology of the HDPs is

heterogeneous and may vary across these subtypes; future adequately powered studies should

examine these subtypes separately as implications for pregnancy care and long-term maternal

health risk may differ[82]. In addition, we lacked paired maternal-fetal samples to condition

maternal risk variants on fetal genotype, although other complementary analyses such as

placental transcriptomics indicated particular variants more likely to be influencing risk via the

fetal genome. Finally, snRNA-seq analyses were performed in male aortic tissue; while findings

are consistent with current understanding of HDPs, future work is needed to verify that these

results are consistent in women and determine whether additional insights may be apparent in

women.

Overall, multi-ancestry genome-wide meta-analysis of preeclampsia/eclampsia and

gestational hypertension revealed novel distinct and overlapping risk loci and enabled polygenic

prediction of the HDPs, with implications for HDP prediction, prevention, and treatment.

Methods

Study cohorts, phenotypes, genotyping, and association analysis.

Preeclampsia/eclampsia and gestational hypertension case and control counts and

definitions for each cohort are summarized in Supplementary Tables 1-3. If an individual had

qualifying codes for both preeclampsia/eclampsia and gestational hypertension, she was

designated as having preeclampsia/eclampsia. In multi-ancestry cohorts, association analyses

were performed within each ancestry group separately and subsequently meta-analyzed.

FinnGen: Cases were identified using ICD-8, -9, and -10 codes; controls were women in

FinnGen without codes corresponding to hypertension and/or proteinuria during pregnancy

(ICD-10 O10-O16). Sample genotyping in FinnGen was performed using Illumina (Illumina Inc.,

San Diego, CA) and Affymetrix (Thermo Fisher Scientific, Santa Clara, CA, USA) arrays.

Genotype calls were made using GenCall or zCall for Illumina and AxiomGT1 algorithm for

Affymetrix data as described previously[83,84]. Individuals were removed for ambiguous sex,

genotype missingness >5%, heterozygosity >±4 SD, and non-Finnish ancestry. Variants were

removed for missingness >2%, Hardy-Weinberg equilibrium (HWE) $P<1\times10^{-6}$, and minor allele

count (MAC) <3. Pre-phasing was performed with Eagle v2.3.5 using 20,000 conditioning

haplotypes. Genotypes were imputed with Beagle 4.1 using the population-specific Sequencing

Initiative Suomi v3 imputation reference panel. Association analyses were performed using

SAIGE[85] with adjustment for age, genotyping batch, and PC 1-10.

Estonian Biobank: The Estonian Biobank is a population-based biobank with over

200,000 participants[86]. Preeclampsia/eclampsia cases were identified using ICD-10 O14-O15.

Gestational hypertension cases were defined using ICD-10 O13. Controls were defined using

ICD-10 O80-O84 and no codes for HDPs. All Estonian Biobank participants have been

genotyped at the Core Genotyping Lab of the Institute of Genomics, University of Tartu, using

the Illumina Global Screening Array v1.0, v2.0, and v2.0_EST arrays. Samples were genotyped

and PLINK format files were created using Illumina GenomeStudio v2.0.4. Individuals were

excluded from the analysis if their call-rate was <95% or if sex defined based on heterozygosity

of X chromosome did not match sex in phenotype data. Before imputation, variants were filtered

by call-rate <95%, HWE $P<1\times10^{-4}$, and MAF <1%. We also used the MAC filter --minMAC=5.

Variant positions were updated to genome build 37 and all variants were changed to be from

TOP strand using GSAMD-24v1-0_20011747_A1-b37.strand.RefAlt.zip files from

https://www.well.ox.ac.uk/~wrayner/strand/. Pre-phasing was performed with Eagle v2.3

software using 20,000 conditioning haplotypes, and imputation was done using Beagle

v.28Sep18.793 with effective population size n_e=20,000. Population specific imputation

reference of 2297 WGS samples was used. Analyses were carried out with SAIGE[85], adjusting

for year of birth and PC 1-10.

Genes & Health: Genes & Health is a cohort of British Pakistani and Bangladeshi

individuals recruited primarily in East London, England[87]. Cases and parous controls were

identified using qualifying ICD-10 and SNOMED codes (Supplementary Tables 1-2). Genotyping

was performed using the Illumina Infinium Global Screening Array v3.0, and quality control was

performed using Illumina GenomeStudio and PLINK v1.9. Individuals were removed who did not

have Pakistani or Bangladeshi ancestry defined as $>\pm3$ SD from the mean of PC1 and those

who self-reported another ethnicity. Variants with call rate <0.99, MAF <1%, and HWE $P<1\times10^{-6}$

were removed. Imputation was performed using the Michigan Imputation Server with the

GenomeAsia reference panel. Association analyses were performed using SAIGE[85] with

adjustment for age, age^2, and PC 1-10.

Michigan Genomics Initiative: The Michigan Genomics Initiative enrolls participants

receiving care at Michigan Medicine and links biospecimen data to electronic health record

(EHR) data. Preeclampsia cases were identified in the freeze 3 dataset using phecode 642.1

and its corresponding ICD codes[88], and women lacking this phecode served as controls.

Genotyping was performed using one of two versions of the Illumina Infinium CoreExome-24

bead array platform. Relatedness within the cohort was estimated using KING v2.1.3.

Individuals were removed for discordant, missing, or ambiguous sex, kinship coefficient >0.45

with another participant, call rate <99%, estimated contamination >2.5%, or missingness on any

chromosome >5%. Variants were excluded with poor intensity separation based on metrics from

GenomeStudio (GenTrain score <0.15 or Cluster Separation score <0.3), overall call rate <99%,

or HWE P-value $<1\times10^{-4}$. Genotypes were phased using EAGLE v2.4.1 and imputed using the

TOPMed reference panel. Association analysis was conducted using SAIGE[85] with adjustment

for age, genotype array, and PC 1-10.

Mass General Brigham Biobank: The Mass General Brigham Biobank is a health

system-based biobank linking genomic data to EHR data. Preeclampsia/eclampsia cases were

identified using ICD-10 O14-O15, gestational hypertension cases were identified using ICD-10

O13, and controls were identified using ICD-10 O80-O82. Variants with MAF <1%, missingness

per variant >1%, and HWE P-value <10^{-6} were removed. Imputation was performed using the

TOPMed reference panel. Association analysis was performed with variants filtered by MAC

≥50 and INFO score ≥0.6 using REGENIE[89] with adjustment for age, genotype batch, and PC 1-

10.

Biobank Japan: Biobank Japan is a biobank of approximately 200,000 Japanese adults.

Preeclampsia cases were identified using phecode 642 and its corresponding ICD codes, and

other women constituted the control group[90]. Genotyping was performed using the Illumina

HumanOmniExpressExome BeadChip or a combination of the Illumina HumanOmniExpress

and HumanExome BeadChip. Individuals with call rates <98% or closely related individuals

(PI_HAT >0.175 in PLINK) were excluded. Variants with call rate <99%, HWE P<1.0×10^{-6}, and

number of heterozygotes <5 were excluded. Genotype data were imputed with 1000 Genomes

Project Phase 3 v5 genotype data and Japanese whole-genome sequencing data. Association

analysis was performed using SAIGE[85] with adjustment for age, age^2, and PC 1-20.

BioMe: Bio*Me* is a health system-based biobank at the Icahn School of Medicine at

Mount Sinai in New York, NY, USA. Preeclampsia/eclampsia and gestational hypertension

cases were identified using ICD-10 codes O14-O15 and O13, respectively. Controls were other

women enrolled in Bio*Me* who indicated a history of childbirth on their intake questionnaire.

Genotyping was performed using the Illumina Global Screening Array. Individuals with ethnicity-

specific heterozygosity rate that surpassed ±3 SD of the population-specific mean, those with a

call rate of ≤95%, and those with discordance between EHR-recorded and genetic sex were

removed. For variant-level quality control, sites with a call rate below 95% and sites with HWE

P<1×10^{-8} were excluded. Imputation of variants was then performed with the Michigan

Imputation Server pipeline using the TOPMed reference panel. Association analysis was

performed separately in African, admixed American, and European ancestry women using

SAIGE[85] with adjustment for age, age^2, and PC 1-10.

InterPregGen consortium: We incorporated summary statistics for preeclampsia from the

discovery GWAS meta-analyses of European cohorts and Central Asian cohorts from the

InterPregGen consortium[17]. European-ancestry discovery cohorts included GOPEC (United

Kingdom), deCODE (Iceland), the Avon Longitudinal Study of Parents and Children (United

Kingdom), MoBa (Norway), SSI (Denmark), and FINRISK (Finland) (7,219 cases and 155,620

controls). Central Asian cohorts included two Kazakh cohorts and one Uzbek cohort (2,296

cases and 2,059 controls). Cohort-specific preeclampsia and control definitions are summarized

in Supplementary Table 3[17]. Fixed-effects inverse variance-weighted meta-analysis was

performed in METAL[23].

HUNT: The HUNT study is a population-based cohort study in Nord-Trøndelag County,

Norway. Genotyped, parous, European-ancestry women were included in the present analysis.

Preeclampsia/eclampsia was ascertained by linkage to the Medical Birth Registry of Norway,

which defines preeclampsia as systolic blood pressure ≥140 mmHg and/or diastolic blood

pressure ≥90 mmHg accompanied by proteinuria >0.3 g/24 h or >1+ on urine dipstick[91]. The

current analysis includes genetic data from approximately 90% of participants from HUNT2

(1995-1997) and HUNT3 (2006-2008) who were genotyped by genome-wide SNP arrays in

2015[92,93]. Genotyping, quality control metrics, and imputation have been described previously[92].

Briefly, one of three different Illumina HumanCoreExome arrays (HumanCoreExome12 v1.0,

HumanCoreExome12 v1.1, and UM HUNT Biobank v1.0) were used for genotyping the HUNT2

and HUNT3 samples[92]. Samples and variants with call rate <99% were excluded. Imputation

was performed using 2,201 HUNT samples with whole genome sequencing, the Haplotype

Reference Consortium, and TOPMed imputation panel (MAC >10). Association analysis was

performed using SAIGE[85] with adjustment for age, age^2, and PC 1-10.

UK Biobank: The UK Biobank is a population-based cohort study of adult residents of

the UK aged 40-69 years at the time of recruitment between 2006-2010.

Preeclampsia/eclampsia and gestational hypertension cases were identified using qualifying

ICD codes (Supplementary Table 1); controls were women self-reporting a history of at least

one live birth at the time of study enrollment without qualifying ICD codes for

preeclampsia/eclampsia or gestational hypertension or any self-reported history of HDPs.

Genotyping was performed using the UK BiLEVE Axiom Array or the UK Biobank Axiom Array

(both Affymetrix, Santa Clara, CA). Individuals with single nucleotide variant missingness ≥10%

were excluded. Imputation was performed centrally using the Haplotype Reference Consortium,

UK10K, and 1000 Genome reference panels[94]. Variants were required to pass the following

quality control filters: MAF ≥1%, single nucleotide variant missingness <10%, and HWE P≥10^{-15},

MAC ≥50, and INFO score ≥0.6. Association analysis was performed in European-ancestry

participants using REGENIE[89] with adjustment for age, genotyping array, and PC 1-10.

Penn Medicine Biobank (PMBB): PMBB is a health system-based biobank at the

University of Pennsylvania, Philadelphia, PA, USA. Preeclampsia/eclampsia cases were

identified using ICD-10 codes O14-O15 or phecode 642.1. Gestational hypertension cases were

identified using ICD-10 code O13. Controls were other women enrolled in PMBB. Genotyping

was performed using the Illumina Global Sequencing Array v2.0; genotype data were imputed to

the TOPMed reference panel using the Michigan Imputation Server. Variants with MAF<1%,

missing rate >10%, and HWE P<10^{-8} were filtered from the GWAS. Association analysis was

performed in REGENIE separately for African-ancestry and European-ancestry women with

adjustment for age, age^2, and PC 1-5.

nuMoM2b: The nuMoM2b study is a prospective U.S. pregnancy cohort of nulliparous

women enrolled in the first trimester of pregnancy between 2012-2015. HDPs were determined

by chart abstraction and adjudication according to previously published definitions[95].

Gestational hypertension was defined as SBP ≥140 mmHg or DBP ≥90 mmHg on two

occasions ≥6 hours apart or one occasion with subsequent antihypertensive therapy after 20

weeks' gestation, excluding blood pressures recorded during the second stage of labor, without

other qualifying features for preeclampsia or eclampsia. Preeclampsia was defined according to

the same blood pressure criteria plus proteinuria or other findings meeting criteria for severe

features, including HELLP syndrome[95]. Women documented as meeting these same blood

pressure criteria before 20 weeks' gestation were designated as having chronic hypertension.

All women were at risk for development of preeclampsia/eclampsia; only those without chronic

hypertension were at risk for the outcome of gestational hypertension. Blood pressure and BMI

were recorded at the first trimester study visit, which occurred at a mean (SD) 11.6 (1.5) weeks'

gestation. Genotyping was performed using the Illumina Infinium Multi-Ethnic Global D2

BeadChip. Individuals related within two degrees by the KING algorithm were removed. Variants

with MAF <1%, genotyping rate <95%, and HWE P<5x10^{-6} were removed[96]. After phasing with

EAGLE, imputation was performed for European, African, and admixed American ancestry

subjects using the TOPMed reference panel via the TOPMed Imputation Server. Association

analysis was performed in REGENIE separately by ancestry group (European, African, and

admixed American) with adjustment for age and PC 1-10. The same women in nuMoM2b with

imputed genotypes were used for external testing of optimized PRS (described below).

Genome-wide meta-analysis and replication.

Variants from GWAS summary statistics were matched by genome build 38 position and

alleles. GWAS summary statistics that were in genome build 37 were lifted to genome build 38

using UCSC liftOver (https://genome.ucsc.edu/cgi-bin/hgLiftOver). We used METAL[23] to

perform fixed-effect inverse variance-weighted meta-analysis. Correction for genomic inflation

factor was carried out prior to meta-analysis. Meta-analyses were conducted among discovery

cohorts, among follow-up cohorts, and across all cohorts. Given the potential overlap of 400

preeclampsia/eclampsia cases from FINRISK between the InterPregGen meta-analysis and

FinnGen (8.4% of FinnGen cases and 2.3% of overall discovery cases) and 7,805 controls

(5.7% of FinnGen controls and 1.7% of overall discovery controls), we conducted a sensitivity

analysis excluding FinnGen (Supplementary Table 4).

Lead variants for preeclampsia/eclampsia and gestational hypertension were

interrogated in multi-ancestry meta-analysis of follow-up cohorts using METAL: HUNT

(preeclampsia/eclampsia only), UK Biobank, PMBB (European- and African-ancestry women),

and nuMoM2b (European-, African-, and admixed American-ancestry women). $P<0.05$ in follow-

up cohorts, consistent direction of effect in follow-up cohorts, and genome-wide significance

($P<5\times10^{-8}$) in combined meta-analysis of discovery and follow-up cohorts indicated replication.

Conditional and joint analysis.

We conducted conditional analysis using GCTA-COJO[24] on the multi-ancestry meta-

analyses of preeclampsia/eclampsia and gestational hypertension to identify additional

association signals at the genome-wide significant loci. We used the European LD reference

from a randomly selected set of 10,000 unrelated individuals. The LD panel included variants

with MAF > 1% and INFO ≥0.3. The analysis was restricted to variants within ±1 Mb from lead

variants ($P<5\times10^{-8}$). In COJO, the lead variants were conditioned from each chromosome and

independent variants were iteratively included. All variants were then simultaneously fitted in the

joint analysis. Variants with $P<2\times10^{-7}$ were considered genome-wide significant. One additional

variant with conditional $P<2\times10^{-7}$ for association with preeclampsia/eclampsia was identified on

chromosome 20 ($P=1.4\times10^{-8}$).

Genetic correlation.

We used LD score regression (LDSC, https://github.com/bulik/ldsc)[25] with pre-computed

LD scores for 1.2 million HapMap3 variants after excluding MHC region in the European

population (https://data.broadinstitute.org/alkesgroup/LDSCORE/eur_w_ld_chr.tar.bz2) to

calculate genetic correlation between preeclampsia/eclampsia and gestational hypertension and

correlation of each HDP with systolic and diastolic blood pressure. In addition, the LDSC

intercept indicates potential confounding due to potential population stratification and cryptic

relatedness. In combined meta-analysis of discovery and follow-up cohorts, we observed

intercepts of 1.03 for preeclampsia/eclampsia and 0.95 for gestational hypertension.

Colocalization analysis.

We obtained the tissue-specific gene expression from GTEx data portal for 52 tissues[32].

We used marginal effect sizes, standard errors, and MAF for all SNPs within ±500 of lead

variants from discovery analysis ($P<5x10^{-8}$) as the input. We performed colocalization using the

coloc.abf() function in R package coloc v4. The H_4 test statistic estimates the posterior

probability of a shared causal variant between preeclampsia/eclampsia or gestational

hypertension and expression of a particular gene. $H_4 >0.7$ indicated strong evidence of

colocalization, H_4 0.5-<0.7 indicated weak evidence of colocalization, and $H_4 <0.5$ indicated no

colocalization.

Polygenic prioritization of causal genes.

We performed additional causal gene prioritization using the PoPS method[34]. Briefly,

PoPS integrates GWAS summary statistics with gene expression, biology pathways, and

predicted protein-protein interaction data to identify likely causal genes at genome-wide

significant loci. A linear model was trained to predict gene-level association scores and estimate

Z-scores indicating the confidence of the causal role at a given locus. In total, PoPS scores

were calculated for 18,000 genes. The top 5 available prioritized genes within 500 kilobases of

lead preeclampsia/eclampsia and gestational hypertension variants were extracted and

compared with the results of other *in silico* analyses.

Placental transcriptome data.

Genes nearest to lead variants, colocalization hits, and genes prioritized within the top 5

by either Open Targets variant-to-gene score or PoPS score were queried in a publicly available

database of placental gene expression (https://www.obgyn.cam.ac.uk/placentome/)[35]. Samples

were obtained from a prospective cohort of nulliparous women in Cambridge, United Kingdom.

Differential expression analysis for preeclampsia included 82 preeclampsia cases and 82

controls samples matched on presence of labor, Cesarean section, gestational age, fetal sex,

smoking status, maternal BMI, and maternal age. RNA sequencing was performed on placental

biopsy specimens with a median sequencing depth of 101 million reads per sample. Differential

expression (in $\log_2$-fold change) and corresponding P-values were generated using DESeq2; P-

values were then adjusted for multiple comparisons using the Benjamini-Hochberg method. We

report differentially expressed prioritized genes with adjusted P <0.05 (Supplementary Table

14).

Gene expression in human aortic tissue.

We queried expression of prioritized genes in subpopulations of vascular smooth muscle

cells, fibromyocytes, fibroblasts, endothelial cells, macrophages, natural killer T cells, and

neuronal cells in a dataset of single-nuclei RNA sequencing from human aortic tissue. In total,

1,114 unique molecular identifiers were obtained per cell. Prioritized genes were nearest genes,

genes with strong colocalization, genes with weak colocalization plus prioritization by another

method (top 5 Open Targets variant-to-gene score or PoPS), and genes prioritized by another

method plus statistically significant differential placental transcription in the human placental

transcriptome browser[35]; of these genes, 24 were available in the single-nuclei RNA

sequencing dataset (Extended Data Fig. 3). Single-nuclei RNA sequencing was performed on

non-atherosclerotic aortic root tissue from two individuals obtained during coronary artery

bypass graft surgery. Both individuals contributing aortic tissue specimens were European-

ancestry men, aged 49 and 51 years, with hypertension, hypercholesterolemia, and coronary

artery disease. Both were using aspirin and a statin preoperatively. A total of 4,537 nuclei were

obtained for downstream analysis. Cell types and subtypes were defined using top marker

genes and pathway enrichment scores. Raw *Cellranger* output data were filtered for removal of

ambient RNA using *cellbender* in 'full' running mode. The resultant filtered cell-gene matrix was

used for quality control and downstream analysis. All preliminary quality-control and clustering

were performed using *scanpy*. Any cells with fewer than 300 genes captured or greater than

0.1% mitochondrial reads were excluded from the analysis. Each sample was processed

with *Scrublet* to exclude doublets. The top 10,000 variable genes were used for

analysis. Relative expression of queried genes in each cell type against other cell types in

normal aortic tissue was quantified as Z-scores.

Derivation and testing of genome-wide polygenic risk scores.

742   We used PRS-CS to derive genome-wide PRS for preeclampsia/eclampsia and

gestational hypertension from the corresponding discovery GWAS summary statistics and for

systolic blood pressure from the Million Veteran Program GWAS summary statistics[26]. PRS-CS

is a Bayesian approach that estimates posterior effect sizes by placing a continuous shrinkage

prior on variant effect sizes[41]. The preeclampsia/eclampsia PRS included 1,087,033 HapMap3

variants, the gestational hypertension PRS included 1,087,916 HapMap3 variants, and the SBP

PRS included 1,064,898 HapMap3 variants. PRS were trained on the UK Biobank European LD

panel. Individual-level polygenic scores were generated in the tuning and test datasets as the

sum of genotypes*weights using PLINK. We used logistic regression to test the association of

each PRS with preeclampsia/eclampsia and gestational hypertension with adjustment for age,

age^2, and PC 1-10. PRS were tuned in the UK Biobank. Specifically, we used a small-scale grid

search of global shrinkage parameter $\square$ values (1, 1×10^{-2}, 1×10^{-4}, and 1×10^{-6}) for each PRS as

recommended to identify the $\square$ that produced the best predictive performance as measured by

R^2 in the tuning dataset. We then fitted a linear combination of optimized

preeclampsia/eclampsia PRS and SBP PRS for the outcome of preeclampsia/eclampsia and a

linear combination of gestational hypertension PRS and SBP PRS for the outcome of

gestational hypertension. The optimal linear combination derived for preeclampsia/eclampsia

was $0.1889*Z_{preeclampsia/eclampsia} + 0.1864*Z_{SBP}$, and the linear combination derived for gestational

hypertension was $0.1662*Z_{gestational\_hypertension} + 0.3050*Z_{SBP}$. We then carried these weighed

linear combination scores forward for final testing in nuMoM2b (European, African, and admixed

American ancestry groups) and HUNT (European ancestry). PRS performance was evaluated

using the OR for top decile vs. bottom 90% of PRS, the OR per SD of PRS, and Nagelkerke's

R^2.

We tested whether PRS correctly re-classified nuMoM2b participants with HDPs as

aspirin-eligible in comparison with the major criteria endorsed by the U.S. Preventive Services

Task Force[43]. Major criteria include: history of preeclampsia, which does not apply in nuMoM2b

as all women were nulliparous; multifetal gestation, which does not apply in nuMoM2b as all

women had singleton pregnancies; chronic hypertension, defined as a diagnosis of

hypertension before pregnancy or blood pressure ≥140/90 mmHg on two occasions at least 6

hours apart prior to 20 weeks' gestation; pre-gestational diabetes type 1 or type 2; any pre-

pregnancy kidney disease; and autoimmune disease, defined here as antiphospholipid antibody

syndrome, systemic lupus erythematosus, rheumatoid arthritis, inflammatory bowel disease

(Crohn's disease or ulcerative colitis), or "other collagen vascular or autoimmune disease." We

calculated sensitivity, specificity, positive predictive value, and negative predictive value for

major risk factors with or without different thresholds for $PRS_{preeclampsia+SBP}$ for the prediction of

preeclampsia/eclampsia, as well as net reclassification for composite HDPs vs. normotensive

pregnancy and for preeclampsia/eclampsia vs. all other pregnancies. Confidence intervals for

sensitivity, specificity, positive predictive value, and negative predictive value were calculated

using the normal approximation. Given use of PRS to up-classify risk, net reclassification was

calculated as P(up|case) - P(up|non-case). Bootstrap resampling performed 1,000 times was

used to estimate 95% confidence intervals for net reclassification.

Phenome-wide association analysis.

We tested the association of preeclampsia/eclampsia PRS and gestational hypertension

PRS with 1,445 phecode-based combined prevalent and incident phenotypes[97] in sex-stratified

fashion among genotyped UK Biobank participants with adjustment for age and PC 1-5 using

the PheWAS package[98] in R 3.6.0 (https://github.com/PheWAS/PheWAS). Bonferroni-corrected

$P<0.05/1,445 = 3.5 \times 10^{-5}$ indicated statistical significance.

Ethics approval.

FinnGen was approved by the Coordinating Ethics Committee of the Helsinki and Uusimaa

Hospital District. The Estonian Committee on Bioethics gave ethical approval for the work

conducted in the Estonian Biobank. The South East Research Ethics Committee gave ethical

approval for the work conducted in Genes & Health. The University of Michigan Medical School

Institutional Review Board gave ethical approval for the analyses conducted in the Michigan

Genomics Initiative. The Mass General Brigham Institutional Review Board gave ethical

approval for the work conducted in the Mass General Brigham Biobank. Biobank Japan

received ethics approval from the Institute of Medical Science, the University of Tokyo, the

RIKEN Yokohama Institute, and all participating hospitals. Bio*Me* received ethics approval from

the Icahn School of Medicine at Mt. Sinai Institutional Review Board. We used publicly available

summary statistics for discovery GWAS from the InterPregGen consortium; all contributing

studies received ethics approval as reported previously[17]. The work conducted in HUNT was

approved by the Regional Committee for Ethics in Medical Research, Norway (2018/2488). The

Penn Medicine Biobank received ethics approval from the University of Pennsylvania

Institutional Review Board. The North West Multi-centre Research Ethics Committee approved

the UK Biobank; the Mass General Brigham Institutional Review Board approved secondary

data analyses of the UK Biobank (application #7089). The nuMoM2b study was approved by the

institutional review boards of each participating site. All participants in all studies contributing

data for this study signed informed consent for participation and the use of data in research.

References

1.	Burton GJ, Redman CW, Roberts JM, Moffett A. Pre-eclampsia: pathophysiology and clinical implications. *BMJ*. **366**, l2381 (2019)

2.	Jiang L, et al. A global view of hypertensive disorders and diabetes mellitus during pregnancy. *Nat Rev Endocrinol*. **18**, 760-775 (2022).

3.	Garovic VD, et al. Incidence and Long-Term Outcomes of Hypertensive Disorders of Pregnancy. *J Am Coll Cardiol*. **18**, 2323-2334 (2020).

4.	Fraser A, et al. Associations of pregnancy complications with calculated cardiovascular disease risk and cardiovascular risk factors in middle age: the Avon Longitudinal Study of Parents and Children. *Circulation*. **125**, 1367-1380 (2012).

5.	Magee LA, et al. The 2021 International Society for the Study of Hypertension in Pregnancy classification, diagnosis & management recommendations for international practice. *Pregnancy Hypertens*. **27**, 148-169 (2022).

6.	ACOG Practice Bulletin No. 202: Gestational Hypertension and Preeclampsia. *Obstet Gynecol*. **133**, e1-e25 (2019).

7.	Honigberg MC, et al. Long-Term Cardiovascular Risk in Women With Hypertension During Pregnancy. *J Am Coll Cardiol*. **74**, 2743-2754 (2019).

8.	Wang YX, et al. Hypertensive Disorders of Pregnancy and Subsequent Risk of Premature Mortality. *J Am Coll Cardiol*. **77**, 1302-1312 (2021).

9.	Ives CW, Sinkey R, Rajapreyar I, Tita ATN, Oparil S. Preeclampsia-Pathophysiology and Clinical Presentations: JACC State-of-the-Art Review. *J Am Coll Cardiol*. **76**, 1690-1702 (2020).

10.	Rana S, Lemoine E, Granger J, Karumanchi SA. Preeclampsia. *Circ Res*. **124**, 1094-1112 (2019).

11.	Levine RJ, et al. Circulating angiogenic factors and the risk of preeclampsia. *N Engl J Med*. **350**, 672-683 (2004).

12.	Bartsch E, Medcalf KE, Park AL, Ray JG. Clinical risk factors for pre-eclampsia determined in early pregnancy: systematic review and meta-analysis of large cohort studies. *BMJ*. **353**, i1753 (2016).

13.	Jung E, et al. The etiology of preeclampsia. *Am J Obstet Gynecol*. **226**, S844-S866 (2022).

14.	Cnattingius S, Reilly M, Pawitan Y, Lichtenstein P. Maternal and fetal genetic factors account for most of familial aggregation of preeclampsia: a population-based Swedish cohort study. *Am J Med Genet A*. **130**, 365-371 (2004).

15.	Nilsson E, Salonen Ros H, Cnattingius S, Lichtenstein P. The importance of genetic and environmental effects for pre-eclampsia and gestational hypertension: a family study. *BJOG*. **111**, 200-206 (2004).

16.	McGinnis R, et al. Variants in the fetal genome near FLT1 are associated with risk of preeclampsia. *Nat Genet*. **49**, 1255-1260 (2017).

17.	Steinthorsdottir V, et al. Genetic predisposition to hypertension is associated with preeclampsia in European and Central Asian women. *Nat Commun*. **11**, 5976 (2020).

18.	Dina C, et al. Variation in FTO contributes to childhood obesity and severe adult obesity. *Nat Genet*. **39**, 724-726 (2007).

19.	Honigberg MC, et al. Genetic Variation in Cardiometabolic Traits and Medication Targets and the Risk of Hypertensive Disorders of Pregnancy. *Circulation*. **142**, 711-713 (2020).

20.	Gray KJ, et al. Risk of pre-eclampsia in patients with a maternal genetic predisposition to common medical conditions: a case-control study. *BJOG*. **128**, 55-65 (2021).

21.	O'Kelly AC, et al. Pregnancy and Reproductive Risk Factors for Cardiovascular Disease in Women. *Circ Res*. **130**, 652-672 (2022).

22.	Kivioja A, Toivonen E, Tyrmi J, et al. Increased Risk of Preeclampsia in Women With a Genetic Predisposition to Elevated Blood Pressure. *Hypertension*. **79**, 2008-2015 (2022).

23.   Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of genomewide
association scans. *Bioinformatics*. **26**, 2190-2191 (2010.
24.   Yang J, Lee SH, Goddard ME, Visscher PM. GCTA: a tool for genome-wide complex
trait analysis. *Am J Hum Genet*. **88**, 76-82 (2011).
25.   Bulik-Sullivan BK, et al. LD Score regression distinguishes confounding from
polygenicity in genome-wide association studies. *Nat Genet*. 47, 291-295, 2015.
26.   Giri A, et al. Trans-ethnic association study of blood pressure determinants in over
750,000 individuals. *Nat Genet*. **51**, 51-62 (2019).
27.   Bulik-Sullivan B, et al. An atlas of genetic correlations across human diseases and traits.
Nat Genet. **47**, 1236-1241 (2015).
28.   Turley P, et al. Multi-trait analysis of genome-wide association summary statistics using
MTAG. *Nat Genet*. **50**, 229-237 (2018).
29.   Padmanabhan S, Caulfield M, Dominiczak AF. Genetic and molecular aspects of
hypertension. *Circ Res*. **116**, 937-959 (2015).
30.   Rubattu S, Forte M, Marchitti S, Volpe M. Molecular Implications of Natriuretic Peptides
in the Protection from Hypertension and Target Organ Damage Development. *Int J Mol Sci*. **20**,
798 (2019).
31.   Ohwaki A, Nishizawa H, Kato A, et al. Altered serum soluble furin and prorenin receptor
levels in pregnancies with pre-eclampsia and fetal growth restriction. *J Gynecol Obstet Hum
Reprod*. **50**, 102198 (2021).
32.   Battle A, et al. Genetic effects on gene expression across human tissues. *Nature*. **550**,
204-213 (2017).
33.   Ghoussaini M, et al. Open Targets Genetics: systematic identification of trait-associated
genes using large-scale genetics and functional genomics. *Nucleic Acids Res*. **49**, D1311-
D1320 (2021).
34.   Weeks EM, et al. Leveraging polygenic enrichments of gene features to predict genes
underlying complex traits and diseases. Preprint at
https://www.medrxiv.org/content/10.1101/2020.09.08.20190561v1 (2020).
35.   Gong S, et al. The RNA landscape of the human placenta in health and disease. *Nat
Commun*. **12**, 2639 (2021).
36.   Maynard SE, et al. Excess placental soluble fms-like tyrosine kinase 1 (sFlt1) may
contribute to endothelial dysfunction, hypertension, and proteinuria in preeclampsia. *J Clin
Invest*. **111**, 649-658 (2003).
37.   Tekola-Ayele F, et al. Placental multi-omics integration identifies candidate functional
genes for birthweight. *Nat Commun*. **13**, 2384 (2022).
38.   Zhang Z, Wang X, Zhang L, Shi Y, Wang J, Yan H. Wnt/β-catenin signaling pathway in
trophoblasts and abnormal activation in preeclampsia (Review). *Mol Med Rep*. **16**, 1007-1013
(2017).
39.   Bai X, Lenhart KC, Bird KE, et al. The smooth muscle-selective RhoGAP GRAF3 is a
critical regulator of vascular tone and hypertension. *Nat Commun*. **4**, 2910 (2013).
40.   Kalluri AS, et al. Single-Cell Analysis of the Normal Mouse Aorta Reveals Functionally
Distinct Endothelial Cell Populations. *Circulation*. **140**, 147-163 (2019).
41.   Ge T, Chen CY, Ni Y, Feng YA, Smoller JW. Polygenic prediction via Bayesian
regression and continuous shrinkage priors. *Nat Commun*. **10**, 1776 (2019).
42.   Ray JG, Abdulaziz KE, Berger H. Aspirin Use for Preeclampsia Prevention Among
Women With Prepregnancy Diabetes, Obesity, and Hypertension. *JAMA*. **327**, 388-390 (2022).
43.   Davidson KW, et al. Aspirin Use to Prevent Preeclampsia and Related Morbidity and
Mortality: US Preventive Services Task Force Recommendation Statement. *JAMA*. **326**, 1186-
1191 (2021).
44.   Pollheimer J, et al. Activation of the canonical wingless/T-cell factor signaling pathway
promotes invasive differentiation of human trophoblast. *Am J Pathol*. **168**, 1134-1147 (2006).

45.   Tita AT, et al. Treatment for Mild Chronic Hypertension during Pregnancy. *N Engl J Med.*
386, 1781-1792 (2002).
46.   Daniels LB, Maisel AS. Natriuretic peptides. *J Am Coll Cardiol.* **50**, 2357-2368 (2007).
47.   Zhang W, et al. Atrial natriuretic peptide promotes uterine decidualization and a TRAIL-
dependent mechanism in spiral artery remodeling. *J Clin Invest.* **131**, e151053 (2021).
48.   Maack T, et al. Physiological role of silent receptors of atrial natriuretic factor. *Science.*
238, 675-678 (1987).
49.   Gu Y, et al. Aberrant pro-atrial natriuretic peptide/corin/natriuretic peptide receptor
signaling is present in maternal vascular endothelium in preeclampsia. *Pregnancy Hypertens.*
11, 1-6 (2018).
50.   Sun BB, et al. Genomic atlas of the human plasma proteome. *Nature.* **558**, 73-79 (2018).
51.   Hauspurg A, et al. Association of N-Terminal Pro-Brain Natriuretic Peptide Concentration
in Early Pregnancy With Development of Hypertensive Disorders of Pregnancy and Future
Hypertension. *JAMA Cardiol.* **7**, 268-276 (2022).
52.   Satpathy AT, et al. Zbtb46 expression distinguishes classical dendritic cells and their
committed progenitors from other immune lineages. *J Exp Med.* **209**, 1135-1152 (2012).
53.   Meredith MM, et al. Zinc finger transcription factor zDC is a negative regulator required
to prevent activation of classical dendritic cells in the steady state. *J Exp Med.* **209**, 1583-1593
(2012).
54.   Wang Y, Sun HY, Kumar S, Puerta MDM, Jo H, Rezvan A. ZBTB46 is a shear-sensitive
transcription factor inhibiting endothelial cell proliferation via gene expression regulation of cell
cycle proteins. *Lab Invest.* **99**, 305-318 (2019).
55.   Ashraf S, et al. Mutations in six nephrosis genes delineate a pathogenic pathway
amenable to treatment. *Nat Commun.* **9**, 1960 (2018).
56.   Hall G, Wang L, Spurney RF. TRPC Channels in Proteinuric Kidney Diseases. *Cells.* 9,
44 (2019).
57.   Wang Z, et al. Transient receptor potential cation channel 6 contributes to kidney injury
induced by diabetes and hypertension. *Am J Physiol Renal Physiol.* **322**, F76-F88 (2022).
58.   Pang Y, Dong J, Thomas P. Progesterone increases nitric oxide synthesis in human
vascular endothelial cells through activation of membrane progesterone receptor-α. *Am J*
Physiol Endocrinol Metab. **308**, E899-911 (2015).
59.   Folkersen L, et al. Genomic and drug target evaluation of 90 cardiovascular proteins in
30,931 individuals. *Nat Metab.* **2**, 1135-1148 (2020).
60.   Heo SH, Choi YJ, Ryoo HM, Cho JY. Expression profiling of ETS and MMP factors in
VEGF-activated endothelial cells: role of MMP-10 in VEGF-induced angiogenesis. *J Cell*
Physiol. **224**, 734-742 (2010).
61.   Toloza FJK, et al. Association between maternal thyroid function and risk of gestational
hypertension and pre-eclampsia: a systematic review and individual-participant data meta-
analysis. *Lancet Diabetes Endocrinol.* **10**, 243-252 (2022).
62.   Männistö T, et al. Preeclampsia, gestational hypertension and subsequent
hypothyroidism. *Pregnancy Hypertens.* **3**, 21-27 (2013).
63.   Wang W, et al. LNK/SH2B3 Loss of Function Promotes Atherosclerosis and Thrombosis.
Circ Res. **119**, e91-e103 (2016).
64.   Machiela MJ, Chanock SJ. LDlink: a web-based application for exploring population-
specific haplotype structure and linking correlated alleles of possible functional variants.
Bioinformatics. **31**, 3555-3557 (2015).
65.   Deloukas P, et al. Large-scale association analysis identifies new risk loci for coronary
artery disease. *Nat Genet.* **45**, 25-33 (2013).
66.   Gupta AK, Hasler P, Holzgreve W, Hahn S. Neutrophil NETs: a novel contributor to
preeclampsia-associated placental hypoxia? *Semin Immunopathol.* **29**, 163-167 (2007).

67.   Palomo M, et al. Differences and similarities in endothelial and angiogenic profiles of
preeclampsia and COVID-19 in pregnancy. *Am J Obstet Gynecol.* **227**, e1-e16 (2022).
68.   Dou H, et al. Oxidized Phospholipids Promote NETosis and Arterial Thrombosis in
LNK(SH2B3) Deficiency. *Circulation.* **144**, 1940-1954 (2021).
69.   Anand S, et al. Detection and confirmation of serum lipid biomarkers for preeclampsia
using direct infusion mass spectrometry. *J Lipid Res.* **57**, 687-696 (2016).
70.   He B, Liu Y, Maurya MR, et al. The maternal blood lipidome is indicative of the
pathogenesis of severe preeclampsia. *J Lipid Res.* **62**, 100118 (2021).
71.   Wright D, Syngelaki A, Akolekar R, Poon LC, Nicolaides KH. Competing risks model in
screening for preeclampsia by maternal characteristics and medical history. *Am J Obstet*
Gynecol. **213**, e1-e10 (2015).
72.   Al-Rubaie ZTA, et al. Prediction of pre-eclampsia in nulliparous women using routinely
collected maternal characteristics: a model development and validation study. *BMC Pregnancy*
Childbirth. **20**, 23 (2020).
73.   Myatt L. The prediction of preeclampsia: the way forward. *Am J Obstet Gynecol.* **226**,
S1102-S1107 (2022).
74.   Akolekar R, Syngelaki A, Poon L, Wright D, Nicolaides KH. Competing risks model in
early screening for preeclampsia by biophysical and biochemical markers. *Fetal Diagn Ther.* **33**,
8-15 (2013).
75.   Chaemsaithong P, Sahota DS, Poon LC. First trimester preeclampsia screening and
prediction. *Am J Obstet Gynecol.* **226**, S1071-S1097 (2022)
76.   Martin AR, Kanai M, Kamatani Y, Okada Y, Neale BM, Daly MJ. Clinical use of current
polygenic risk scores may exacerbate health disparities. *Nat Genet.* **51**, 584-591 (2019).
77.   Graham SE, Clarke SL, Wu KH, et al. The power of genetic diversity in genome-wide
association studies of lipids. *Nature.* **600**, 675-679 (2021).
78.   Angras K, Boyd VE, Gray C, Young AJ, Paglia MJ, Mackeen AD. Retrospective
application of algorithms to improve identification of pregnancy outcomes from the electronic
health record. *J Perinatol.* **43**, 10-14 (2023).
79.   Klungsøyr K, Harmon QE, Skard LB, et al. Validity of pre-eclampsia registration in the
medical birth registry of norway for women participating in the norwegian mother and child
cohort study, 1999-2010. *Paediatr Perinat Epidemiol.* **28**, 362-371 (2014).
80.   Klemmensen AK, Olsen SF, Osterdal ML, Tabor A. Validity of preeclampsia-related
diagnoses recorded in a national hospital registry and in a postpartum interview of the women.
Am J Epidemiol. **166**, 117-124 (2007).
81.   Geller SE, Ahmed S, Brown ML, Cox SM, Rosenberg D, Kilpatrick SJ. International
Classification of Diseases-9th revision coding for preeclampsia: how accurate is it? *Am J Obstet*
Gynecol. **190**, 1629-1633 (2004).
82.   Roberts JM, Rich-Edwards JW, McElrath TF, Garmire L, Myatt L. Subtypes of
Preeclampsia: Recognition and Determining Clinical Usefulness. **77**, 1430-1441 (2021).
83.   Kurki MI, et al. FinnGen provide genetic insights from a well-phenotyped isolated
population. *Nature.* **613**, 508-518 (2023).
84.   Sun BB, et al. Genetic associations of protein-coding variants in human disease. *Nature.*
603, 95-102 (2022).
85.   Zhou W, et al. Efficiently controlling for case-control imbalance and sample relatedness
in large-scale genetic association studies. *Nat Genet.* **50**, 1335-1341 (2018).
86.   Leitsalu L, et al. Cohort Profile: Estonian Biobank of the Estonian Genome Center,
University of Tartu. *Int J Epidemiol.* **44**, 1137-1147 (2015).
87.   Finer S, et al. Cohort Profile: East London Genes & Health (ELGH), a community-based
population genomics and health study in British Bangladeshi and British Pakistani people. *Int J*
Epidemiol. **49**, 20-21 (2020).

88. Wei WQ, et al. Evaluating phecodes, clinical classification software, and ICD-9-CM
codes for phenome-wide association studies in the electronic health record. *PLoS One.* **12**,
e0175508 (2017).
89. Mbatchou J, et al. Computationally efficient whole-genome regression for quantitative
and binary traits. *Nat Genet.* **53**, 1097-1103 (2021).
90. Sakaue S, et al. A cross-population atlas of genetic associations for 220 human
phenotypes. *Nat Genet.* **53**, 1415-1424 (2021).
91. Honigberg MC, et al. Heart Failure in Women With Hypertensive Disorders of
Pregnancy: Insights From the Cardiovascular Disease in Norway Project. *Hypertension.* **76**,
1506-1513 (2020).
92. Brumpton BM, et al. The HUNT study: A population-based cohort for genetic research.
Cell Genomics. **2**, 100193 (2022).
93. Åsvold BO, et al. Cohort Profile Update: The HUNT Study, Norway. *International Journal*
of Epidemiology. **52**, e80-e91 (2023).
94. Bycroft C, et al. The UK Biobank resource with deep phenotyping and genomic data.
Nature. **562**, 203-209 (2018).
95. Facco FL, et al. Association Between Sleep-Disordered Breathing and Hypertensive
Disorders of Pregnancy and Gestational Diabetes Mellitus. *Obstet Gynecol.* **129**, 31-41 (2017).
96. Guerrero RF, et al. Genetic Polymorphisms Associated with Adverse Pregnancy
Outcomes in Nulliparas. Preprint available at
https://www.medrxiv.org/content/10.1101/2022.02.28.22271641v1.
97. Denny JC, et al. PheWAS: demonstrating the feasibility of a phenome-wide scan to
discover gene-disease associations. *Bioinformatics.* **26**, 1205-1210 (2010).
98. Carroll RJ, Bastarache L, Denny JC. R PheWAS: data analysis and plotting tools for
phenome-wide association studies in the R environment. *Bioinformatics.* **30**, 2375-2376 (2014).
1037

Data availability

Upon publication, GWAS summary statistics for preeclampsia/eclampsia and gestational

hypertension will be made publicly available through the Broad Institute CVDi portal, and

corresponding genome-wide polygenic scores will be posted to the PGS Catalog. Summary

statistics used in this meta-analysis are publicly available for FinnGen r6

(https://www.finngen.fi/en/access_results) and for BioBank Japan

(https://pheweb.jp/pheno/PreEclampsia). Summary statistics from the InterPregGen consortium

are available at https://ega-archive.org. Placental transcriptome data are publicly available at

https://www.obgyn.cam.ac.uk/placentome/.

Code availability

The code used to conduct these analyses is available at https://github.com/buutrg/HDP.

Acknowledgements

This work was supported by grants from the U.S. National Heart Lung and Blood Institute

K08HL166687 (M.C.H.), K08HL146963 (K.J.G.), R01 HL163234 (R.S., K.J.G.), R01HL139865

(R.D.), R01HL155915 (R.D.), DP2HL152423 (R.M.G.), U01HL166060 (R.M.G.), R03HL148483

(R.M.G.), R01HL142711 (P.N.), R01HL127564 (P.N.), R01HL148050 (P.N.), R01HL151283

(P.N.), R01HL148565 (P.N.), R01HL135242 (P.N.), R01HL151152 (P.N.); the American Heart

Association 940166 (M.C.H.), 979465 (M.C.H.); the Korea Health Industry Development

Institute HI19C1330 (S.M.J.C.); Harvard Catalyst Medical Research Investigator Training

Program (A.P.P.); National Human Genome Research Institute U01HG011719 (A.P.P. and

P.N.); the Belgian American Educational Foundation (A.S.); the U.S. National Institute of

General Medical Sciences R35GM147197 (R.F.G.), R35GM124836 (R.D.); National Institute of

Diabetes and Digestive and Kidney Diseases R01DK125782 (P.N.); National Institute of Child

Health and Human Development R01HD101246 (D.M.H.); Preeclampsia Foundation (K.J.G.,

1064   R.S.); Fondation Leducq TNE-18CVD04 (P.N.); and the Massachusetts General Hospital Paul

1065   and Phyllis Fireman Endowed Chair in Vascular Medicine (P.N.).

1066

The work was conducted under UK Biobank application 7089. The Trøndelag Health Study

(HUNT) is a collaboration between HUNT Research Centre (Faculty of Medicine and Health

Sciences, NTNU, Norwegian University of Science and Technology), Trøndelag County Council,

Central Norway Regional Health Authority, and the Norwegian Institute of Public Health. The

genotyping in HUNT was financed by the National Institutes of Health; University of Michigan;

the Research Council of Norway; the Liaison Committee for Education, Research and

Innovation in Central Norway; and the Joint Research Committee between St Olavs hospital

and the Faculty of Medicine and Health Sciences, NTNU. The genetic investigations of the

HUNT Study is a collaboration between researchers from the K.G. Jebsen Center for Genetic

Epidemiology, NTNU and the University of Michigan Medical School and the University of

Michigan School of Public Health. The K.G. Jebsen Center for Genetic Epidemiology is financed

by Stiftelsen Kristian Gerhard Jebsen; Faculty of Medicine and Health Sciences, NTNU,

Norway. DNA extraction and genotyping of nuMoM2b was supported by the Indiana University

Precision Health Initiative. The authors thanks other members of the Estonian Biobank research

team: Lili Milani, Andres Metspalu, Tõnu Esko, Mari Nelis, Reedik Mägi, and Georgi Hudjashov.

Genes & Health is/has recently been core-funded by Wellcome (WT102627, WT210561), the

Medical Research Council (UK) (M009017, MR/X009777/1, MR/X009920/1), Higher Education

Funding Council for England Catalyst, Barts Charity (845/1796), Health Data Research UK (for

London substantive site), and research delivery support from the NHS National Institute for

Health Research Clinical Research Network (North Thames). Genes & Health is/has recently

been funded by Alnylam Pharmaceuticals, Genomics PLC; and a Life Sciences Industry

Consortium of Astra Zeneca PLC, Bristol-Myers Squibb Company, GlaxoSmithKline Research

and Development Limited, Maze Therapeutics Inc, Merck Sharp & Dohme LLC, Novo Nordisk

A/S, Pfizer Inc, Takeda Development Centre Americas Inc. The authors thank Social Action for

Health, Centre of The Cell, members of our Community Advisory Group, and staff who have

recruited and collected data from volunteers. We thank the NIHR National Biosample Centre

(UK Biocentre), the Social Genetic & Developmental Psychiatry Centre (King's College London),

Wellcome Sanger Institute, and Broad Institute for sample processing, genotyping, sequencing

and variant annotation. We thank: Barts Health NHS Trust, NHS Clinical Commissioning Groups

(City and Hackney, Waltham Forest, Tower Hamlets, Newham, Redbridge, Havering, Barking

and Dagenham), East London NHS Foundation Trust, Bradford Teaching Hospitals NHS

Foundation Trust, Public Health England (especially David Wyllie), Discovery Data

Service/Endeavour Health Charitable Trust (especially David Stables), NHS Digital for GDPR-

compliant data sharing backed by individual written informed consent. Most of all, we thank all

of the volunteers participating in Genes & Health. Current Genes & Health Research Team (in

alphabetical order by surname): Shaheen Akhtar, Mohammad Anwar, Elena Arciero, Omar

Asgar, Samina Ashraf, Saeed Bibi, Gerome Breen, Raymond Chung, Charles J Curtis, Shabana

Chaudhary, Maharun Chowdhury, Grainne Colligan, Panos Deloukas, Ceri Durham, Faiza

Durrani, Fabiola Eto, Sarah Finer, Joseph Gafton, Ana Angel Garcia, Chris Griffiths, Joanne

Harvey, Teng Heng, Sam Hodgson, Qin Qin Huang, Matt Hurles, Karen A Hunt, Shapna

Hussain, Kamrul Islam, Vivek Iyer, Ben Jacobs, Ahsan Khan, Cath Lavery, Sang Hyuck Lee,

Robin Lerner, Daniel MacArthur, Daniel Malawsky, Hilary Martin, Dan Mason, Rohini Mathur,

Mohammed Bodrul Mazid, John McDermott, Shefa Miah, Caroline Morton, Bill Newman,

Elizabeth Owor, Asma Qureshi, Samiha Rahman, Nishat Safa, Miriam Samuel, John Solly,

Daniel Stow, Farah Tahmasebi, Richard C Trembath, Karen Tricker, Nasir Uddin, David A van

Heel, Klaudia Walter, Caroline Winckley, John Wright, Julia Zollner.

1113

1114

Ethics declarations

M.C.H. reports consulting fees from CRISPR Therapeutics, advisory board service for Miga Health, and grant support from Genentech, all unrelated to this work. K.J.G. has served as a consultant for BillionToOne, Aetion, and Roche for projects unrelated to this work. R.S. is a co-founder of Magnet Biomedicine, unrelated to this work. R.D. reports receiving grants from AstraZeneca, grants and nonfinancial support from Goldfinch Bio, being a scientific co-founder, consultant, and equity holder for Pensieve Health (pending), and being a consultant for Variant Bio, all unrelated to this work. P.N. reports grant support from Amgen, Apple, AstraZeneca, Boston Scientific, and Novartis, spousal employment and equity at Vertex, consulting income from Apple, AstraZeneca, Novartis, Genentech / Roche, Blackstone Life Sciences, Foresite Labs, and TenSixteen Bio, and is a scientific advisor board member and shareholder of TenSixteen Bio and geneXwell, all unrelated to this work.

Table 1. Maternal sequence variants associated with preeclampsia/eclampsia.

Nearest gene	Lead variant	CHR	POS	Effect allele / other allele	Weighted average EAF	OR	P-value	P(het)*	OR	P-value	OR	P-value
					Discovery analysis (17,150 cases / 451,241 controls)				**Follow-up analysis (2,914 cases / 251,876 controls)**		**Combined discovery and follow-up meta-analysis (20,064 cases / 703,117 controls)**	
Loci with genome-wide significance in discovery analysis												
MTHFR / CLCN6	rs149764880	1	11820674	G / T	0.85	1.11	1.8e-8	0.30	1.17	6.8e-4	1.12	**8.8e-11**
WNT3A	rs708119	1	228015567	C / G	0.67	1.08	2.3e-8	0.92	1.05	0.10	1.08	**7.8e-9**
MECOM	rs9855086	3	169413542	A / T	0.53	1.08	5.7e-10	0.21	1.08	8.3e-3	1.08	**1.6e-11**
FGF5	rs16998073	4	80263187	T / A	0.32	1.11	3.1e-14	0.15	1.08	1.6e-2	1.11	**2.3e-15**
MICA	rs2442752	6	31383987	C / T	0.36	1.08	2.3e-9	0.57	0.99	0.84	1.07	**3.8e-8**
LINC00484	rs5899121	9	91145498	TG / T	0.85	1.12	4.6e-8	0.31	0.86	0.31	1.12	1.4e-7
PGR	rs2508372	11	101397592	A / G	0.79	1.10	1.3e-9	0.14	1.11	7.5e-3	1.11	**3.4e-11**
SH2B3	rs10774624	12	111395984	G / A	0.47	1.10	2.9e-12	0.40	1.03	0.29	1.09	**1.0e-11**
FLT1	rs7318880	13	28564148	T / C	0.48	1.09	6.7e-12	0.79	1.06	5.8e-2	1.09	**1.6e-12**
FTO	rs1421085	16	53767042	C / T	0.40	1.10	4.7e-12	0.61	1.07	1.8e-2	1.09	**3.3e-13**
RGL3	rs167479	19	11416089	G / T	0.55	1.10	7.6e-10	0.20	1.09	6.8e-3	1.10	**1.8e-11**
ZNF831	rs259983	20	59160402	C / A	0.16	1.13	1.0e-12	0.40	1.15	2.8e-4	1.14	**1.3e-15**
Additional loci identified in combined discovery and follow-up meta-analysis												
FGL1	rs2653414	8	17868560	A / C	0.02	1.27	1.0e-6	0.44	2.18	4.4e-9	1.36	**3.4e-11**
UPB1	rs17572606	22	24472204	T / C	0.02	1.45	1.9e-7	0.33	1.28	6.0e-2	1.41	**4.6e-8**

*P(het) indicates P-value for heterogeneity across discovery cohorts.

CHR indicates chromosome. POS indicates position (genome build 38). EAF indicates effect allele frequency. OR indicates odds ratio.

Table 2. Maternal sequence variants associated with gestational hypertension.

Nearest gene	Lead variant	CHR	POS	Effect allele / other allele	Weighted average EAF	Discovery analysis (8,961 cases / 184,925 controls)			Follow-up analysis (2,066 cases / 227,863 controls)		Combined discovery and follow-up meta-analysis (11,027 cases / 412,788 controls)	
						OR	P-value	P(het)*	OR	P-value	OR	P-value
Loci with genome-wide significance in discovery analysis												
MECOM	rs9855086	3	169413542	A / T	0.54	1.10	1.4e-9	0.51	1.04	0.28	1.08	6.6e-8
FGF5	rs16998073	4	80263187	T / A	0.32	1.12	4.6e-11	0.54	1.09	4.0e-2	1.12	**1.5e-11**
NPR3	rs13154066	5	32831564	C / T	0.58	1.11	6.1e-10	0.60	1.05	0.14	1.10	**4.5e-10**
TNS2 / CSAD	rs7139122	12	53173006	A / G	0.02	1.46	1.1e-9	0.13	1.05	0.65	1.37	**8.3e-9**
RGL3	rs167479	19	11416089	G / T	0.56	1.11	6.9e-9	0.76	1.12	4.6e-3	1.11	**2.5e-10**
PREX1	rs2208589	20	48791877	G / A	0.84	1.15	2.3e-8	0.17	1.11	2.8e-2	1.14	**2.9e-9**
ZNF831	rs260017	20	59141918	A / G	0.18	1.16	7.0e-13	0.32	1.14	5.4e-3	1.16	**1.2e-13**
Additional loci identified in combined discovery and follow-up meta-analysis												
MTHFR / CLCN6	rs17367504	1	11802721	A / G	0.85	1.13	1.7e-7	0.63	1.19	4.6e-4	1.14	**4.9e-10**

*P(het) indicates P-value for heterogeneity across discovery cohorts.

CHR indicates chromosome. POS indicates position (genome build 38). EAF indicates effect allele frequency. OR indicates odds ratio.

Table 3. Aspirin eligibility to prevent preeclampsia using major clinical risk factors and polygenic risk. This Table reports the proportion of participants in nuMoM2b (nulliparous women with singleton gestations recruited in the first trimester of pregnancy) who experienced each pregnancy outcome in the index pregnancy (normotensive pregnancy, gestational hypertension, or preeclampsia/eclampsia) who would have been identified as candidates for low-dose aspirin to prevent preeclampsia according to U.S. Preventive Services Task Force major criteria, with or without addition of preeclampsia/eclampsia and systolic blood pressure combination polygenic risk score (PRS$_{preeclampsia+SBP}$) as an additional criterion for aspirin eligibility. Net reclassification analyses reflect use of PRS$_{preeclampsia+SBP}$ to up-classify risk for composite hypertensive disorders of pregnancy (HDPs) vs. normotensive pregnancy and for preeclampsia/eclampsia vs. other pregnancies when added to major risk factors.

Total *N*=7,544	Normo-tensive pregnancy (*n*=5,744)	Gestational hypertension (*n*=1,319)	Preeclampsia/ eclampsia (*n*=481)	Net re-classification, HDPs vs. normotensive pregnancy (95% CI)	Net re-classification, preeclampsia vs. other pregnancy (95% CI)	Preeclampsia vs. other pregnancy			
						Sensitivity (95% CI)	Specificity (95% CI)	Positive predictive value (95% CI)	Negative predictive value (95% CI)
Major risk factors only (*n*=656 [8.7%])	476 (8.3%)	96 (7.3%)	84 (17.5%)	Ref.	Ref.	17.5% (14.1-20.9%)	91.9% (91.3-92.5%)	12.8% (10.2-15.4%)	94.2% (93.7-94.8%)
Major risk factors or top 5% PRS (*n*=997 [13.2%])	699 (12.2%)	184 (13.9%)	114 (23.7%)	+2.7% (1.4 to 4.0%)	+1.8% (-0.3 to 4.0%)	23.7% (19.9-27.5%)	87.5% (86.7-88.3%)	11.4% (9.5-13.4%)	94.4% (93.8-95.0%)
Major risk factors or top 10% PRS (*n*=1,324 [17.6%])	928 (16.2%)	250 (19.0%)	146 (30.4%)	+4.1% (2.5 to 5.8%)	+4.3% (1.3 to 7.3%)	30.4% (26.2-34.5%)	83.3% (82.5-84.2%)	11.0% (9.3-12.7%)	94.6% (94.1-95.2%)
Major risk factors or top 25% PRS (*n*=2,298 [30.5%])	1,612 (28.1%)	460 (34.9%)	226 (47.0%)	+8.3% (6.0 to 10.7%)	+8.3% (3.9 to 12.6%)	47.0% (42.5-51.4%)	70.7% (69.6-71.7%)	9.8% (8.6-11.1%)	95.1% (94.6-95.7%)

Figure 1. Manhattan plots of preeclampsia/eclampsia and gestational hypertension in combined discovery and follow-up meta-analysis.
Manhattan plots (chromosomal position on the X-axis and -log(10) of the P-value on the Y-axis) are displayed for **(A)** preeclampsia/eclampsia in 20,064 cases and 703,117 controls and **(B)** gestational hypertension in 11,027 cases and 412,788 controls. Analyses included multi-ancestry meta-analysis of common variants (minor allele frequency ≥1%). Loci are labeled by the gene nearest to the lead variant.

Figure 2. Polygenic prediction of preeclampsia/eclampsia and gestational hypertension in test cohorts. Polygenic risk scores (PRS) for preeclampsia/eclampsia and gestational hypertension were derived from our discovery genome-wide meta-analyses, tuned in the UK Biobank, and carried forward for testing in independent cohorts. Prevalence of preeclampsia/eclampsia vs. percentile of $PRS_{preeclampsia+SBP}$ in **(A)** HUNT and **(B)** nuMoM2b. **(C)** Prevalence of gestational hypertension vs. percentile of PRS_{GH+SBP} in nuMoM2b. Distribution of $PRS_{preeclampsia+SBP}$ by preeclampsia/eclampsia status in **(D)** HUNT and **(E)** nuMoM2b. **(F)** Distribution of PRS_{GH+SBP} by gestational hypertension status in nuMoM2b. Within each boxplot, horizontal lines reflect the median, top and bottom of the box reflect the interquartile range, and whiskers reflect the maximum and minimum value within each grouping.

Figure 3. Sex-stratified phenome-wide association study of preeclampsia/eclampsia and gestational hypertension polygenic risk in the UK Biobank. Preeclampsia/eclampsia polygenic risk **(A-B)** and gestational hypertension polygenic risk **(C-D)** were associated with 1,445 phenotypes among women (A, C) and men (B, D) in the UK Biobank. Associations with phenotypes were tested using logistic regression with adjustment for age and PC 1-5.

Extended Data Figure 1. Flow chart summarizing the study design and contributing cohorts.

Extended Data Figure 2. Results of multi-trait analysis of genome-wide summary statistics (MTAG) for preeclampsia/eclampsia from joint analysis of summary statistics for preeclampsia/eclampsia and gestational hypertension in discovery cohorts. The plot displays chromosomal position on the X-axis and -log(10) of the P-value on the Y-axis.

Extended Data Figure 3. Relative expression of prioritized genes in human aortic cells with single-nuclei RNA sequencing. We analyzed expression of genes prioritized by genome-wide meta-analysis of preeclampsia/eclampsia and gestational hypertension and secondary *in silico* analyses in a dataset of single-nuclei RNA sequencing from two normal human flash-frozen aortic specimens. Most prioritized genes were enriched in endothelial cell populations and/or macrophages.