Assessment of Environmental Factors Associated with Antibiotic Resistance Genes (ARGs) in the Yangtze Delta, China

Jiazheng Miao[1], Yikai Ling[1], Xiaoyuan Chen[1], Siyuan Wu[1], Sajid Umar[2], Shixin Xu[1,2,*], Benjamin D. Anderson[1,2,3,*]

[1] Division of Natural and Applied Science, Duke Kunshan University, Kunshan, Jiangsu, China

[2] Global Health Research Center, Duke Kunshan University, Kunshan, Jiangsu, China

[3] Department of Environmental and Global Health, College of Public Health and Health Professions, and Emerging Pathogens Institute, University of Florida, Gainesville, Florida, United States

* Shixin Xu and Benjamin D. Anderson serve as co-corresponding authors

*Corresponding Authors:

Benjamin D. Anderson, MPH, PhD, CPH

Assistant Professor

Department of Environmental and Global Health, College of Public Health and Health Professions, and Emerging Pathogens Institute, University of Florida, Gainesville, Florida, 32610 United States

Email: ander88@ufl.edu

Shixin Xu, PhD

Assistant Professor of Mathematics

Division of Natural and Applied Science, Duke Kunshan University, Kunshan,

Jiangsu, 215300, China

Global Health Research Center, Duke Kunshan University, Kunshan, Jiangsu,

215300, China

Email: shixin.xu@dukekunshan.edu.cn

Abstract

The emergence of antimicrobial resistance (AMR) is an urgent and complex public

health challenge worldwide. As a sub-problem of AMR, antibacterial resistance

(ABR) is of particular concern due to inadequacy of alternative medication. Earlier

studies have shown that ABR is not only impacted by antibiotics, but also affected by

the interactions between bacteria and their environments. Therefore, to combat ABR

in a specific region, local environmental conditions must be investigated to

comprehensively understand which environmental factors might contribute to ABR

and propose more tailored solutions. This study surveyed environmental contributors

of antibiotic resistance genes (ARGs), the parameter for measuring ABR, in the

Yangtze Delta. A high abundance of ARGs was detected, despite low antibiotic and

heavy metal concentrations. Phosphorus, chromium, manganese, calcium, and

strontium were identified as potential key contributors of ARGs. Suppression of

ARGs could be realized through decreasing the concentration of phosphorus in

surface water. Group 2A light metals (e.g., magnesium and calcium) could be

developed as eco-friendly reagents for controlling antibiotic resistance in the future.

Keywords: antimicrobial resistance, antibiotic resistance genes, antibiotic residues,

dissolved metal, surface water, environmental health

1. Introduction

Antimicrobial resistance (AMR) refers to the process by which bacteria, parasites, viruses, and fungi evolve mechanisms that protect them from drugs [1]. Due to the increasing prevalence, health effects, and economic impact of AMR, World Health Organization has characterized AMR as "a complex global public health challenge" (p. XIX) that requires urgent action [1]. Among the concerns about AMR, countermeasures against antibacterial resistance (ABR) are of particular urgency because of severe inadequacy of alternative therapeutic options [1]. In 2019, it is estimated that 4.95 million deaths were associated with ABR, and 1.27 million deaths were attributable to ABR [2]. In a recent study that estimates the economic burden of ABR in China, 27.45% of inpatient bacterial infection were antibiotic resistant, which is estimated to have caused a 1.5% increase in inpatient mortality rates and cost $77 billion in related societal expenses in 2017 [3]. What is worse, bacteria are even capable of spreading ABR at a high rate through exchanging ARGs in horizontal gene transfer (HGT) [4]. ARGs are a group of genes that encode the proteins responsible for resisting antibiotics and are frequently carried on mobile genetic elements (MGEs) like plasmids, transposons and integrons [4]. In HGT, one bacterial cell transfer its MGEs to other bacterial cells, facilitating environmental dispersal of the ARGs [4]. Thus, ARG has also been recognized as a new environmental pollutant that poses a threat to human health [4].

Earlier studies have attempted to suppress ABR by reducing antibiotic use to allow non-resistant bacteria to outcompete resistant bacteria [5-7], but reversibility only occurs at a slow rate. Later studies demonstrate a strong association between ABR and environmental factors [8]. In particular, heavy metals are reported to have a co-selection effect on ARGs [9-11]. Because bacteria resist antibiotics and heavy

metals through similar strategies (e.g., reduction in permeability, efflux,

sequestration, etc.), heavy metals may also pose a selective pressure on ARGs.

Furthermore, antibiotic resistant communities in urban lakes have also been

identified to be more competitive than non-resistant communities at higher

temperatures, but less competitive at higher concentration of magnesium or salinity

[8].

Therefore, to effectively understand and mitigate ABR in a specific region, local

environmental conditions must be investigated.

The Yangtze Delta is one of the most economically developed regions in China. It

consists of 26 cities including Shanghai City, southern part of Zhejiang Province, and

northern part of Jiangsu province [12]. In 2018, 154 million people were reported to

permanently resides in this area at a high population density of 720 persons/square

kilometre [12]. The hygiene of drinking water source in this area (e.g., Yangcheng

Lake in Suzhou and Huangpu River in Shanghai) is of vital importance since the

water sources provide drinking water for over 150 million residents. Due to its

economic and environmental importance, previous studies have investigated the

prevalence of ARGs and patterns of antibiotic use in the area.

In 2020, 64 out of 69 types of antibiotics monitored in the surface water in China

can be detected in Shanghai, indicating the widespread of antibiotics in the surface

water of Shanghai, with concentrations ranging from a few pg L^{-1} to hundreds ng L-1

[13-15]. While confirming the widespread distribution and persistence of antibiotics,

these studies estimate low environmental risks for most types of antibiotics [14, 16]

by calculating their hazard quotients[17]. However, the environmental risks of

enrofloxacin, amoxicillin and oxytetracyclines to aquatic organisms, represented by

the risk quotient method [18, 19], are distinctly higher than those in other countries

[14]. As for ARGs, tetracycline resistance genes (*tet*) and sulfonamide resistance

genes (*sul*) exist in most waterbodies in Yangtze Delta, and β-lactam resistance

genes also exist in some regions [20, 21]. For *sul* and *tet* genes, the detection

frequency of *sul*I, *sul*II, *tet*A, *tet*G, *tet*M and *tet*O were significantly higher than *sul*III,

which was still significantly higher than *tet*C [21]. It was observed that samples with

more *sul* genes tend to contain higher levels of total sulfonamides, and the same

relationship can be applied to *tet* genes [20].

While these studies focus on the link between ARGs and antibiotic detection.

Limited studies have investigated how ARGs are influenced by environmental

factors. In this study, we tested surface water samples for ARGs, antibiotics, and

dissolved elements to better understand what environmental factors might contribute

to the prevalence of ARGs in main water bodies in the Yangtze Delta.

2. Methods

2.1 Sampling Sites and Sample Collection

In 2021, surface water samples were collected from sampling sites located along

a watershed starting at Yangcheng Lake in Suzhou, running along the Wusong

River, passing Dianshan Lake in Shanghai, running along the Huangpu River in

Shanghai, and ending at an estuary at the Shanghai coastal area. The watershed

connects most of the main water bodies in Suzhou and Shanghai. The sampling

area contained four subregions: Dianshan Lake (Qingpu District, Shanghai),

Huangpu River (Songjiang District, Shanghai), estuary (Jinshan District, Shanghai),

and Kunshan (Suzhou).

Surface water samples were collected using brown HDPE bottles (NINGKE,

China) that were dipped into the top 10cm of the water body with the goal of

collecting 1000ml of water. Water temperature, salinity, and pH of the collected

samples were immediately measured and recorded using a salinity and pH meter

(BANTE, China) according to the manufacturer's instructions. Sample bottles were

transported on wet ice within 3 hours to Duke Kunshan University where 650ml of

each sample was filtered through a 0.45 µm glass microfiber filter membranes (Titan,

China). The filtrate and filter membranes were stored stored at a -80°C freezer.

2.2 Measurement of ARGs

Cells gathered on the filter membrane were first released in 10ml of ultrapure

water. A TaKaRa MiniBEST Bacteria Genomic DNA Extraction Kit was used to

extract 240µl of DNA from 10ml of stored sample, according to the manufacturer's

instructions. Quantitative PCR was used to screen extracts for ARGs using the

SYBR Green Real-Time PCR assay (TaKaRa Co. Dalian, China) on a Mic qPCR

Cycler (BioMolecular Systems, El Cajon, CA), according to the manufacturer's

instructions. One sample was randomly selected from each of the four subregions. A

total of 26 primers for ARGs were used to screen the four samples. The ARGs

include 15 tetracycline resistance genes (*tetA*, *tetC*, *tetE*, *tetK*, *tetL*, *tetA/P*, *tetG*,

tetM, *tetO*, *tetQ*, *tetS*, *tetT*, *tetW*, *tetBP*, and *tetX*), 4 β-lactam resistance genes

(*bla*$_{CTX-M}$, *bla*$_{TEM}$, *bla*$_{SHV}$, and *bla*$_{ampC}$), 3 sulfonamide resistance genes (*sul*I, *sul*II,

and *sul*III), 3 macrolides resistance genes (*ereA*, *ereB*, and *mphA*), and 1 beta-

lactam resistance gene. Primer sequences and cycling conditions for the quantitative

PCR experiments are listed in Table S1. Cultured environmental samples were used

as positive controls.

Based on the ARG screening results (Figure S1a and Figure S1b), the three most

abundant ARGs were selected for further quantitative PCR experiments: *tetBP*,

ampr, tetO. Each gene was commercially synthesized and used as an internal

standard to determine absolute abundance.

2.3 Measurement of Dissolved Elements Concentrations

After filtration, approximately 150ml of stored filtrate was acidified to pH < 2 by

(1+1) nitric acid and preserved at -80 °C freezer. The samples were sent to the

Nanjing Institute of Geography & Limnology, Chinese Academy of Sciences and

measured for Ca, K, Mg, Na, Si, Ag, Hg, Tl, Al, As, B, Ba, Be, Cd, Co, Cr, Cu, Fe,

Mn, Mo, Ni, P, Pb, Sb, Se, Sn, Sr, Ti, V, and Zn by ICP-MS 7900 (Agilent, US), as

described in [22].

2.4 Measurement of Antibiotic Concentration

Concentrations of amoxicillin and tetracycline were measured using 500 mL of

stored filtrate using the Amoxicillin ELISA kit (Shenzhen Lvshiyuan Biotechnology

Co., Ltd, China) and Tetracycline ELISA kit (Shenzhen Lvshiyuan Biotechnology Co.,

Ltd, China), according to the manufacturer's instructions. The procedures of the two

kits were technically the same. Briefly, samples and standard solutions with known

antigen (amoxicillin/tetracycline) concentration were separately mixed with

corresponding enzyme conjugates (enzyme-labelled antibodies) on a micro-well

plate where micro-well strips were pre-coated with coupling antigens. The antigens

in the sample and the coupling antigens would compete for the enzyme conjugates.

After removing the mixed solution and adding TMB substrate for coloration, the

enzyme conjugates bound with sample antigens were removed and only those

bound with pre-coated antigens remained, which led to higher optical density (OD)

value in the sample with lower antigen concentration. The OD values of the solutions

generated from samples and standard solutions were measured with the wavelength

of 450nm. All procedures above were performed in duplicate for each sample and

standard solution. A standard curve of absorption versus antigen concentration was

generated with the OD values from the standard solutions, and the OD values from

the samples were compared to the standard curve to get the antigen concentration.

2.5 Environmental Risk Assessment of Antibiotics

To assess the potential risk of antibiotics in the environment, a hazard quotient

(HQ) was used, which is the ratio between the measured environmental

concentration (MEC) of the antibiotic and the predicted no-effect concentration

(PNEC) [17]. HQ values >1.0 indicate a potential risk to the aquatic system [23]. This

study adopts PNEC values proposed by Bengtsson-Palme and Larsson, 2016 (0.25

µg/L for amoxicillin and 1 µg/L for tetracycline) [24].

2.6 Statistical Method

ARG screening data was calculated to relative abundance using the median Cq

value of *tetE* in Sample S7 as a reference with the following formula:

195   Relative Abundance = $2^{\wedge}(Cq_{reference} - Cq)$

TetBP, *ampr*, and *tetO* qPCR data was calculated to absolute abundance using a

standard curve derived by serially diluting (10-fold) synthetic genes at a range of 10^{12}

copies/mL to 10^2 copies/mL.

Concentrations of amoxicillin and tetracycline were calculated using a standard

curve derived from standard solutions provided in the commercial kits.

All figures were generated using R (version 4.0 and 4.1). Figure 1(a) used

country border data and river data from the rnaturalearth package. Population

density data was collected by the Ministry of Civil Affairs of the People's Republic of

China [25] and Statistical Bureau of Republic of China [26]. Figure 1(b) adopted the

Stamen map through the ggmap package.

Phosphorus, chromium, manganese, calcium, and strontium were used to predict

tetBP, *ampr*, and *tetO* abundance through ridge regression. Ridge regression is an

algorithm that produces stable estimation of the least-squares coefficients of

multiple-regression models with the condition of high collinearity [27]. Because *tetBP*

contained an outlier with an abundance several magnitudes higher than the others,

tetBP was re-scaled through the calculation of $\log_{10}(\text{Abd}+1)$.

3. Results

3.1 Abundance of Antibiotic Resistance Genes

The spatial distribution of the samples and absolute abundance of total ARG

(sum of *tetBP*, *ampr*, *tetO*) are shown in Figure 1. Notably, an ARG hotspot (S10)

was found in Yangcheng Lake, Kunshan. The hotspot contains an elevated amount

of *tetBP* (Figure S1).

The mean absolute abundance and composition of ARGs in the four regions are

shown in Figure 2. The ARG *ampr* accounts for the majority of detected

concentrations in all the regions. However, all the results shown a large standard

deviation, indicating high variability within each region.

3.2 Dissolved Elements & Water Quality

In general, all of the samples had a very low heavy metal concentration as shown

in Table 1 (according to the Environmental Quality Standard for Surface Water of

People's Republic of China [28]). However, Sample S8 collected from drinking water

reserve in Songjiang District, Shanghai had an iron concentration of 417.436 µg/L

that exceeds the standard limit of 300 µg/L (Table 2). Moreover, though the

concentration of phosphorus in many samples was relatively high (Table 1), the

values were still under the standard limits. One exception again was the Songjiang

sample (237.872 µg/L) which exceeds the phosphorus limit of 200 µg/L for drinking

water (Table 1).

3.3 Measurement of Antibiotic Concentration and Risk Assessment

Antibiotic concentrations are shown in Table 3. Unexpectedly, except for the

sample from Jinshan contains 0.13 µg/L of tetracycline, antibiotics were not detected

in any other samples. Also, the HQ value tetracycline concentration in Jinshan is

0.13, indicating no risk.

3.4 Correlation of ARGs and Environmental Risk Factors

Figure 3 shows the correlation of ARGs with salinity, temperature, and dissolved

elements. Water temperature and salinity show weak to little correlation with ARGs.

pH appears to have a weak negative correlation with *tetO* and *ampr* (r = -0.58, -

0.55). Interestingly, phosphorus, chromium, and manganese exhibit a strong positive

correlation with *tetO* (r = 0.79, 0.64, 0.63). Calcium has a strong negative correlation

with *ampr* (r = -0.70). And strontium has strong negative correlation with both *tetO*

and *ampr* (r = -0.71, -0.70). Linear regression of *tetO* and *ampr* with phosphorus,

chromium, manganese, calcium, and strontium is shown in Figure 4. These five

elements explain 71.2% variability of *ampr*, 79.1% variability of *tetO*, and 36.0%

variability of log-scaled *tetBP*, in a ridge regression model.

4. Discussion

This study aims to understand what environmental factors might contribute to the

prevalence of ARGs in main water bodies in the Yangtze Delta. In summary, a high

abundance of ARGs was detected and several dissolved elements (phosphorus,

chromium, manganese, calcium, and strontium) were potentially key contributors of

ARGs.

4.0 Interpretation of the abundance of ARGs

As shown in Figure 1 and Figure 2, a high abundance of ARGs was detected.

Compared with historical data of Huangpu River collected in July 2013 [29], the

ARGs abundance in Huangpu River detected in this study was several magnitudes

higher. Specifically, historical data shows that the abundance of tetracycline-resistant

genes was at the magnitude of 10 to 10^4 copies/mL [29], but in this study *tetO*

abundance in Huangpu River reached up to 10^6 copies/mL (Figure 2). The hotspot

sample from Yangcheng Lake, Kunshan, contained *tetBP* at the magnitude of 10^8

copies/mL (Figure 1). This finding suggests there could have been a surge of

antibiotic resistant bacteria in the Yangtze Delta during the past decade.

What is more, the hotspot and large standard deviations of ARG concentrations

imply that there might be point sources of ARG pollution, which elevated ARG

concentrations locally (e.g., hotspot of Sample S10), but could not disperse over a

wide area.

4.1 ARGs and Antibiotics

In contrast with the detection of ARGs, antibiotic measurement results indicate

that antibiotic persistence in the environment was limited, with ELISA results

showing that all samples contained less than 1 µg/L of amoxicillin and tetracycline

(Table 3). This finding is consistent with other studies that show antibiotic

concentrations in surface water in Shanghai is at the magnitude of ng/L [13-15, 29-

31]. This indicates, despite low antibiotic risk, ARGs remain persistent in the aquatic

system.

Notably, persistence of tetracycline resistance was also reported in coastal

regions of Southeast China, which is geologically close to this study's sampling

region [32]. Though other mechanisms may also contribute to persistence of

tetracycline resistance (e.g., compensatory evolution), existing evidence supports

co-selection between resistance genes is the most likely cause of persistence of

tetracycline resistance [6], which will be discussed in following parts.

Additionally, it is worth noting that overall antibiotic levels in the Yangtze Delta

were observed to vary in accordance to seasons, with the number of detected

antibiotics and their detection frequencies being significantly lower in summer than in

winter [14, 15, 33, 34], except for tetracyclines [34]. The high flow conditions in

summer might dilute the concentration of antibiotics in the surface water [35], and

faster degradation of antibiotics in summer could also contribute to the low

concentration of antibiotics detected [34, 36]. The results of this study do not

guarantee a risk of antibiotics during the winter.

Though the correlation coefficient of ARGs and antibiotics cannot be calculated,

considering the low level of antibiotic concentrations, it might be more prudent to

focus on direct remediation strategies for removing ARGs, instead of focusing only

on decreasing antibiotic contamination in surface water of the Yangtze Delta.

4.2 ARGs and Nonmetal Elements

As shown in Figure 3 and Figure 4, phosphorus exhibits a strong positive

correlation with *tetO*. One possible explanation for this correlation is that

phosphorus, as an essential biological element, enriches bacterial growth in

freshwater [37, 38]. The abundance of ARGs proliferate as the quantity of bacterial

cells increases. This positive connection between phosphorus and ARGs is also

reported by other studies in eutrophic environment [39, 40]. Though additional

assessment is needed to better understand the relationship between phosphorous

concentrations and ARG, the findings seem to suggest that regulating phosphorous

eutrophication could be a practical approach to manage ARG development in

surface water. One similar proposal is also given by Wang et al. 2020 [40].

4.3 ARGs and Heavy Metals

As shown in Figure 3 and Figure 4, chromium and manganese exhibit a strong

positive correlation with *tetO*. The positive correlation of chromium could be a result

of co-selection between tetracycline-resistant genes and chromium-resistant genes

though co-resistance mechanism. Co-resistance describes the spatial association of

HMRGs and ARGs when they are located on the same genetic elements, such as

plasmids and transposons. This association means either heavy metals or antibiotics

can be the selection pressure for both resistance genes. A previous study cured

plasmids from *Salmonella abortus equi* strains and found the plasmids were resistant

to ampicillin, arsenic, chromium, cadmium and mercury simultaneously [41].

However, co-selection of manganese and ARGs is rarely reported. Further research

is needed to confirm whether manganese truly has co-selective pressure on ARGs.

4.4 ARGs and Light Metals

A negative correlation was found between *ampr* and calcium, and strontium had

a strong negative correlation with both *tetO* and *ampr* (Figure 3 & Figure 4).

Interestingly, as noted, a previous study reported magnesium (Mg) could suppress

resistant bacterial community [8]. Magnesium, calcium, and strontium all belong to

Group 2A on the periodic table. This means the three elements have similar

chemical properties, so it is possible that the three elements have a potential

inhibition effect on ARG through similar mechanisms.

Although negative correlation of strontium may be explained by its cytotoxicity on

bacteria [42, 43], molecular biology studies have reported selective inhibition effects

of magnesium and calcium on antibiotic-resistant bacteria. Magnesium and calcium

may involve directly in the destabilization of the *Staphylococcus aureus* cell

membrane [44]. Calcium and magnesium can form a complex with Cardiolipin (CL),

one major component of the bacterial cell membrane, thus destabilizing the cell

membrane and eventually causing cell death [44]. In addition, calcium could

suppress bacterial growth through inhibiting peptidoglycan-layer (PG-layer) synthesis

or destabilizing the cell membrane. PG-layer synthesis involves transpeptidase (TP)

enzymes, including D,D-TP and L,D-TP. These TP enzymes are the target of many

antibiotic β-lactams [45]. Calcium hinders the binding of L,D-TP to PG-stem by

inducing the formation of L,D-TP dimer [45].

This finding reveals the possibility of developing Group 2A light metals as

reagents to control ABR. For example, as one of the major elements in nature,

calcium salts are of low toxicity and eco-friendly. Calcium-based reagent might be

capable of suppressing antibiotic resistant bacteria in natural environment if added to

natural surface water (e.g., exposure recreational water).

4.5 Limitations

Considering the reported seasonal variance in antibiotic concentrations [14, 15,

33, 34], the findings on antibiotic concentrations can only be generalizable to surface

water of the Yangtze Delta in summer. Also, impacted by the outbreak of COVID-19

Delta strain in China in 2021 Summer, the sampling stage of this study was

emergently terminated, ahead of schedule. The unexpected termination led to a

small sample size (15 samples). The small sample size decreases statistical power

and increases potential error in the results. Finally, ARG concentrations were of

large standard deviations in all four subregions. Though this could be explained as

indication of point sources of ARG pollution, one alternative explanation is there was

degradation of DNA during the experiments.

5. Conclusion

This study detected a high abundance of ARGs in surface water samples

collected from the Yangtze Delta, despite corresponding low antibiotic and low heavy

metal concentration. Phosphorus, chromium, manganese, calcium, and strontium

were identified as potential key contributors of ARGs in the Yangtze Delta. Reduction

of ARG abundance could be realized through decreasing the concentration of

phosphorus in surface water. In addition, Group 2A light metals (e.g., magnesium

and calcium) were identified to have a potential ARG suppression effect, though

future studies are still needed to further evaluate this association.

6. Acknowledgements

The authors thank Dr. Huansheng Cao for providing the positive controls for the quantitative PCR experiments and thank Dr. Yingxue Li for her help in measuring the antibiotic concentrations.

This work was funded by Interdisciplinary Health Research Fund at the Global Health Research Center, Duke Kunshan University, Kunshan, China.

References

1.   Antimicrobial Resistance: Global Report on Surveillance. Geneva: WHO Press, **2014**.

2.   Murray CJL, Ikuta KS, Swetschinski L, et al. Global burden of bacterial antimicrobial

resistance in 2019: a systematic analysis. The Lancet (British edition) **2022**;

399(10325): 629-55.

3.   Zhen X, Stålsby Lundborg C, Sun X, Zhu N, Gu S, Dong H. Economic burden of

antibiotic resistance in China: a national level estimate for inpatients. Antimicrob

Resist Infect Control **2021**; 10(1): 5.

4.   Tripathi V, Tripathi P. Antibiotic Resistance Genes: An Emerging Environmental

Pollutant. In: Kesari KK. Perspectives in Environmental Toxicology. Cham: Springer

International Publishing, **2017**:183-201.

5.   Andersson DI, Hughes D. Antibiotic resistance and its cost: is it possible to reverse

resistance? Nature Reviews Microbiology **2010**; 8(4): 260-71.

6.   Andersson DI, Hughes D. Persistence of antibiotic resistance in bacterial

populations. FEMS Microbiol Rev **2011**; 35(5): 901-11.

7.   Sundqvist M. Reversibility of antibiotic resistance. Ups J Med Sci **2014**; 119(2): 142-

8.

8.   Wang Q, Hamilton PB, Kang F, Zhu X, Zhang Y, Zhao H. Regional-scale

investigation for microbial competition-through-environment interactions modulating

antibiotic resistance. Sci Total Environ **2020**; 734: 139341.

9.   Baker-Austin C, Wright MS, Stepanauskas R, McArthur JV. Co-selection of antibiotic

and metal resistance. Trends Microbiol **2006**; 14(4): 176-82.

10.   Chen J, Li J, Zhang H, Shi W, Liu Y. Bacterial Heavy-Metal and Antibiotic Resistance

Genes in a Copper Tailing Dam Area in Northern China. Frontiers in Microbiology

2019; 10.

11.   Di Cesare A, Eckert E, Corno G. Co-selection of antibiotic and heavy metal

resistance in freshwater bacteria. Journal of Limnology **2016**; 75(s2).

12.   Xu Z, Yin Y. Regional Development Quality of Yangtze River Delta: From the

Perspective of Urban Population Agglomeration and Ecological Efficiency

Coordination. Sustainability **2021**; 13(22).

13.   Zhu F, Wang S, Liu Y, Wu M, Wang H, Xu G. Antibiotics in the surface water of

Shanghai, China: screening, distribution, and indicator selecting. Environmental

Science and Pollution Research **2021**; 28(8): 9836-48.

14.   Wu M, Que C, Tang L, et al. Distribution, fate, and risk assessment of antibiotics in

five wastewater treatment plants in Shanghai, China. Environmental Science and

Pollution Research **2016**; 23(18): 18055-63.

15.   Pan C, Bao Y, Xu B. Seasonal variation of antibiotics in surface water of Pudong

New Area of Shanghai, China and the occurrence in typical wastewater sources.

Chemosphere **2020**; 239: 124816.

16.   Wang H, Yang J, Yu X, et al. Exposure of Adults to Antibiotics in a Shanghai

Suburban Area and Health Risk Assessment: A Biomonitoring-Based Study.

Environmental Science & Technology **2018**; 52(23): 13942-50.

17.   Grung M, Källqvist T, Sakshaug S, Skurtveit S, Thomas KV. Environmental

assessment of Norwegian priority pharmaceuticals based on the EMEA guideline.

Ecotoxicology and Environmental Safety **2008**; 71(2): 328-40.

18.   Hu Y, Yan X, Shen Y, Di M, Wang J. Antibiotics in surface water and sediments from

Hanjiang River, Central China: Occurrence, behavior and risk assessment.

Ecotoxicology and Environmental Safety **2018**; 157: 150-8.

19.   Wu M-H, Que C-J, Xu G, et al. Occurrence, fate and interrelation of selected

antibiotics in sewage treatment plants and their receiving surface water.

Ecotoxicology and Environmental Safety **2016**; 132: 132-9.

20.   Jiang L, Hu X, Xu T, Zhang H, Sheng D, Yin D. Prevalence of antibiotic resistance

genes and their relationship with antibiotics in the Huangpu River and the drinking

water sources, Shanghai, China. Science of The Total Environment **2013**; 458-460:

267-72.

437 21. Yang Y, Liu W, Xu C, Wei B, Wang J. Antibiotic resistance genes in lakes from

438    middle and lower reaches of the Yangtze River, China: Effect of land use and

439    sediment characteristics. Chemosphere **2017**; 178: 19-25.

440 22. Zi XY, Zhang M, Gu XH, et al. Impact of Enclosure Culture on Heavy Metal Content

441    in Surface Sediments of Hongze Lake and Ecological Risk Assessment. Huanjing

442    Kexue/Environmental Science **2021**; 42(11): 5355-63.

443 23. López-Serna R, Postigo C, Blanco J, et al. Assessing the effects of tertiary treated

444    wastewater reuse on the presence emerging contaminants in a Mediterranean river

445    (Llobregat, NE Spain). Environmental Science and Pollution Research **2012**; 19(4):

446    1000-12.

447 24. Bengtsson-Palme J, Larsson DGJ. Concentrations of antibiotics predicted to select

448    for resistant bacteria: Proposed limits for environmental regulation. Environment

449    International **2016**; 86: 140-9.

450 25. Statistical list of administrative divisions of the People's Republic of China in 2020

451    [Chinese]. In: China MoCAotPsRo, **2020**.

452 26. Population density (end of year, persons per square kilometer). In: China SBoRo.

453    National Statistics, **2020**.

454 27. Hoerl AE, Kennard RW. Ridge Regression: Biased Estimation for Nonorthogonal

455    Problems. Technometrics **1970**; 12(1): 55-67.

456 28. Environmental quality standard for surface water [Chinese]. In: ENVIRONMENT

457    MOEA, CHINA TPSRO, **2002**.

458 29. Jiang L, Hu X, Xu T, Zhang H, Sheng D, Yin D. Prevalence of antibiotic resistance

459    genes and their relationship with antibiotics in the Huangpu River and the drinking

460    water sources, Shanghai, China. Sci Total Environ **2013**; 458-460: 267-72.

461 30. Chen K, Zhou JL. Occurrence and behavior of antibiotics in water and sediments

462    from the Huangpu River, Shanghai, China. Chemosphere **2014**; 95: 604-12.

463   31.   Jiang L, Hu X, Yin D, Zhang H, Yu Z. Occurrence, distribution and seasonal variation
464   of antibiotics in the Huangpu River, Shanghai, China. Chemosphere **2011**; 82(6):
465   822-8.

466   32.   Chen CQ, Zheng L, Zhou JL, Zhao H. Persistence and risk of antibiotic residues and
467   antibiotic resistance genes in major mariculture sites in Southeast China. Sci Total
468   Environ **2017**; 580: 1175-84.

469   33.   Zhu F, Wang S, Liu Y, Wu M, Wang H, Xu G. Antibiotics in the surface water of
470   Shanghai, China: screening, distribution, and indicator selecting. Environ Sci Pollut
471   Res Int **2021**; 28(8): 9836-48.

472   34.   Jiang L, Hu X, Yin D, Zhang H, Yu Z. Occurrence, distribution and seasonal variation
473   of antibiotics in the Huangpu River, Shanghai, China. Chemosphere (Oxford) **2011**;
474   82(6): 822-8.

475   35.   Hanamoto S, Nakada N, Jürgens MD, Johnson AC, Yamashita N, Tanaka H. The
476   different fate of antibiotics in the Thames River, UK, and the Katsura River, Japan.
477   Environmental Science and Pollution Research **2018**; 25(2): 1903-13.

478   36.   Karthikeyan KG, Meyer MT. Occurrence of antibiotics in wastewater treatment
479   facilities in Wisconsin, USA. Science of The Total Environment **2006**; 361(1): 196-
480   207.

481   37.   Coveney MF, Wetzel RG. Effects of nutrients on specific growth rate of
482   bacterioplankton in oligotrophic lake water cultures. Appl Environ Microbiol **1992**;
483   58(1): 150-6.

484   38.   Wang L, Miller TD, Priscu JC. Bacterioplankton nutrient deficiency in a eutrophic
485   lake. Archiv für Hydrobiologie **1992**; 125(4): 423-39.

486   39.   Ali L, Wang YQ, Zhang J, et al. Nutrient-induced antibiotic resistance in
487   Enterococcus faecalis in the eutrophic environment. J Glob Antimicrob Resist **2016**;
488   7: 78-83.

40.   Wang Z, Han M, Li E, et al. Distribution of antibiotic resistance genes in an

agriculturally disturbed lake in China: Their links with microbial communities,

antibiotics, and water quality. Journal of Hazardous Materials **2020**; 393: 122426.

41.   Ghosh A, Singh A, Ramteke PW, Singh VP. Characterization of large plasmids

encoding resistance to toxic heavy metals in Salmonella abortus equi. Biochem

Biophys Res Commun **2000**; 272(1): 6-11.

42.   Alshammari H, Neilands J, Svensäter G, Stavropoulos A. Antimicrobial Potential of

Strontium Hydroxide on Bacteria Associated with Peri-Implantitis. Antibiotics (Basel)

2021; 10(2): 150.

43.   Baheiraei N, Eyni H, Bakhshi B, Najafloo R, Rabiee N. Effects of strontium ions with

potential antibacterial activity on in vivo bone regeneration. Scientific Reports **2021**;

11(1): 8745.

44.   Xie Y, Yang L. Calcium and Magnesium Ions Are Membrane-Active against

Stationary-Phase Staphylococcus aureus with High Specificity. Sci Rep **2016**; 6:

20628.

45.   Gokulan K, Khare S, Cerniglia CE, Foley SL, Varughese KI. Structure and Inhibitor

Specificity of L,D-Transpeptidase (LdtMt2) from Mycobacterium tuberculosis and

Antibiotic Resistance: Calcium Binding Promotes Dimer Formation. AAPS J **2018**;

20(2): 44.

Table 1: Water Quality Rank (according to Environmental Quality Standard for Surface Water of PRC [28]) and Concentration (µg/L) of Dissolved Elements. Rank I is applicable for national nature reserves. Rank II is for first-class drinking water reserves. Rank III is for second-class drinking water reserves. Rank IV is for industry water and non-exposure entertainment water sources. Rank V is for agricultural water sources.

Sample	Area		P		Cu		Zn		As		Se		Hg		Cd		Pb		Cr
S1	Qingpu	III	128.006	I	0.174	I	ND	I	4.386	I	ND	I	0.047	I	0.018	I	2.922	I	0.973
S2	Qingpu	III	136.442	I	0.427	I	ND	I	4.345	I	ND	I	0.025	I	0.012	I	0.791	I	1.28
S3	Qingpu	II	94.828	I	0.433	I	7.425	I	4.288	I	ND	I	0.021	I	0.026	I	1.372	I	0.742
S4	Qingpu	II	79.516	I	0.056	I	1.762	I	5.011	I	ND	I	0.034	I	0.021	I	2.112	I	0.752
S5	Kunshan	II	37.645	I	ND	I	ND	I	2.329	I	ND	I	ND	I	0.017	I	1.914	I	0.522
S7	Jinshan	V	458.891	I	ND	I	5.135	I	3.743	I	ND	I	ND	I	0.004	I	0.668	I	2.074
S8*	Songjiang	IV	237.872	I	ND	I	2.272	I	3.626	I	0.535	I	ND	I	0.011	I	0.831	I	0.6
S9	Songjiang	III	180.972	I	0.529	I	3.414	I	3.276	I	ND	I	ND	I	0.023	I	1.205	I	1.33
S10	Kunshan	II	34.28	I	ND	I	ND	I	2.359	I	ND	I	ND	I	ND	I	0.457	I	ND
S11	Kunshan	II	30.058	I	ND	I	ND	I	2.442	I	ND	I	ND	I	0.005	I	0.642	I	ND
S12*	Kunshan	II	45.482	I	ND	I	9.226	I	2.708	I	ND	I	ND	I	ND	I	0.332	I	0.264
S13*	Kunshan	II	24.372	I	ND	I	ND	I	2.217	I	ND	I	ND	I	0.004	I	0.646	I	0.113
S14	Kunshan	IV	258.324	I	0.275	I	0.72	I	2.712	I	ND	I	ND	I	0.022	I	0.967	I	0.351
S15	Kunshan	III	114.272	I	2.07	I	11.143	I	2.309	I	ND	I	ND	I	0.031	I	2.364	I	1.125

509   *Labelled samples are from drinking water reserves. ND: Not detected.
510

Table 2: Additional requirements from Environmental Quality Standard for Surface Water of PRC for drinking water reserves.

Sample	Area	Mo		Co		Be		B		Sb		Ni		Ba		Ti		V		Tl		Fe	
S8	Songjiang	W	3.538	W	0.326	W	ND	W	99.268	W	0.751	W	0.548	W	ND	W	6.237	W	2.632	W	ND	E	417.436
S12	Kunshan	W	3.302	W	0.153	W	ND	W	119.726	W	1.407	W	1.998	W	ND	W	1.985	W	2.021	W	ND	W	153.924
S13	Kunshan	W	2.453	W	0.075	W	ND	W	115.066	W	0.742	W	1.312	W	ND	W	0.596	W	0.856	W	ND	W	61.913

511 "W" means concentration is within the limit for drinking water reserves, and "**E**" means exceeding the requirement; ND: Not
512 detected.

Table 3: Concentrations of amoxicillin and tetracycline (unit: μg/L).

Sample	Area	Amoxicillin	Tetracycline
S1	Qingpu	ND	ND
S2	Qingpu	ND	ND
S3	Qingpu	ND	ND
S4	Qingpu	ND	ND
S5	Kunshan	ND	ND
S6	Songjiang	ND	ND
S7	Jinshan	ND	ND
S8	Songjiang	ND	ND
S9	Songjiang	ND	ND
S10	Kunshan	ND	ND
S11	Kunshan	ND	ND
S12	Kunshan	ND	ND
S13	Kunshan	ND	ND
S14	Kunshan	ND	ND
S15	Kunshan	ND	ND

513   ND: Not detected. Detection limit: 1 μg/L.

 Figure Legend

Figure 1: Map of sampling area. (a) Map of China with population density in 2020.

Unit of population density: persons/square kilometre. The blue line represents the

Yangtze River. (b) Spatial distribution of samples and absolute abundance of total

ARGs in each sample. Unit of total ARG abundance: copies/mL. The black dash

represents the sampling lane.

Figure 2: Mean absolute abundance of ARGs in the four sampling regions. Unit of

ARG abundance: copies/mL.

Figure 3: Heatmap of correlation of ARG concentrations and environmental

parameters.

Figure 4: Linear regression of dissolved elements with ARGs (a) Calcium and *ampr*.

(b) Phosphorus and *tetO*. (c) Chromium and *tetO*. (d) Manganese and *tetO*. Unit:

tetO (copies/mL). Unit: ampr (copies/mL), tetO (copies/mL), calcium (mg/L),

phosphorus (µg/L), chromium (µg/L), manganese (µg/L), strontium (µg/L).