

1 **Healthy eating and physical activity: analysing Soweto's young adults' perspectives with**  
2 **an intersectionality lens**

3 **Authors:** Gudani Mukoma<sup>1\*</sup>, Edna N Bosire<sup>1,2</sup>, Sonja Klingberg<sup>1</sup> and Shane A Norris<sup>1,3</sup>

4

5 **Affiliations:**

6 <sup>1</sup>SAMRC/Wits Developmental Pathways for Health Research Unit, Department of Paediatrics,

7 Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

8 <sup>2</sup>Brain and Mind Institute, Aga Khan University

9 <sup>3</sup>School of Human Development and Health, Faculty of Medicine, University of Southampton,

10 Southampton, UK

11

12 **Corresponding author:** Gudani Mukoma

13 SAMRC/Wits Developmental Pathways for Health Research Unit

14 University of the Witwatersrand, School of Public Health

15 27 St Andrew's Road,

16 Parktown, Johannesburg, 2193

17 Phone: +27(0)79-903-7819

18 E-mail: [gudani.mukoma@wits.ac.za](mailto:gudani.mukoma@wits.ac.za)

19 **Abstract**

20 *Background and objectives:* Non-communicable diseases (NCDs) are taking a toll on Africa's  
21 youth at younger ages than in other regions. These are attributed to risk factors that usually  
22 advance in adolescence, such as unhealthy diets and reduced physical activity. Young adults in  
23 South Africa, particularly women, tend to be sedentary, consume energy-dense diets low in  
24 micronutrients, and are more likely to develop NCDs much earlier in life than those in high-  
25 income countries. With an intersectionality perspective, this study explored young adults'  
26 barriers and solutions to addressing these risk factors in Soweto.

27 *Setting:* Soweto, Johannesburg, South Africa, is one of the most well-known historically  
28 disadvantaged townships known for its established communities, and socioeconomic and  
29 cultural diversity.

30 *Design:* A qualitative investigation utilising focus group discussions (FGDs) with a topic guide.  
31 FGDs were transcribed verbatim and thematically analysed using a combination of deductive  
32 and inductive approaches.

33 *Participants:* Men and women 18-24 years of age living in Soweto (n =30).

34 *Results:* South African young adults have a basic understanding of the significance of nutrition,  
35 exercise, and their ties to health. However, numerous barriers to such behaviours were reported,  
36 arising from the participants' personal, domestic, social, and local community levels. Young  
37 women experienced sexism and had safety concerns while exercising in the streets, while  
38 young men tended to describe themselves as lazy.

39 *Conclusions:* Young adults face a multitude of intersecting barriers, making it difficult to adopt  
40 or sustain health-promoting behaviours. It is important that potential solutions focus on the  
41 intersections of barriers to healthy eating and physical activity in order to provide more realistic  
42 support for such behaviours.

43 **Keywords: Diet, physical activity, young adults, barriers, intersectionality**

## 44 **Introduction**

45 Non-communicable diseases (NCDs) account for 17 million deaths annually, 86% of which  
46 are premature and take place in low- and middle-income countries (1). NCDs like heart disease,  
47 stroke, and diabetes are becoming more prevalent causes of death and disability in sub-Saharan  
48 Africa (SSA). By 2035, NCDs are anticipated to surpass infectious diseases as the main cause  
49 of death and disability (2, 3). Particularly concerning is the fact that in Africa, NCDs are  
50 affecting people at a much younger age (10 or more years younger) than in wealthier regions  
51 (4). This rising burden of NCDs is the result of risk factors that typically begin in adolescence,  
52 such as unhealthy diets, low physical activity (PA), and sedentary behaviours, which are driven  
53 and exacerbated by urbanisation, globalisation, and commercial health determinants (5-8).  
54 Furthermore, evidence suggests that the health risks are transferred to the next generation as  
55 well as affecting young people themselves (9). To prevent the development of NCDs in future  
56 generations, the 2016 Lancet Commission on Adolescent Health and Wellbeing recommended  
57 investing in prevalent NCD-related health behaviours among adolescents (10).

58

59 Young adults (18-26 years) in South Africa, particularly women, tend to lead more  
60 sedentary lives compared to their male counterparts, consume energy dense but micronutrient  
61 poor diets, and are at a high risk of developing NCDs much earlier in life (11-14). Due to  
62 behavioural health risks, women were twice as likely as men to die from heart disease between  
63 2012 and 2016 (14.7% vs. 7.1% respectively) (15). A recent survey in urban South Africa  
64 found that 46.6% of young women aged 18-24 years were in the body mass index-based  
65 categories of overweight or obese, and 42.5% of them did not engage in regular physical  
66 activity (16). On the other hand, consumption of foods (fats, oils, sauces, dressings,  
67 condiments, sweets and savoury snacks) associated with excessive weight gain has increased  
68 by 30% over the past decade (17). While NCD risk factors are already high among young

69 women, they significantly increase with age (18, 19). It is therefore important to support young  
70 people to adopt and maintain behaviours that minimise NCD risks, setting them on healthier  
71 trajectories. In order to do so, we must first gain an understanding of the contextual factors  
72 influencing their health beliefs, diet and physical activity behaviour.

73

74 Individual characteristics, as well as, social and physical environment factors such as home,  
75 family, peers, neighbourhood, school, and workplace, have been shown in studies to influence  
76 health beliefs and behaviours (20-22). From previous studies in urban South Africa, we learned  
77 that perceived affordability, taste, availability, individual preferences and family were  
78 significant determinants of food choices (21, 23, 24), while physical activity opportunities are  
79 constrained by a lack of facilities and safety concerns (21). These barriers have been shown to  
80 vary by age and gender among children and adolescents from high-income countries (25);  
81 however, evidence of how they differ among young adults is scarce. Changes accompanying  
82 the transition from adolescence to adulthood may make individuals more susceptible to  
83 emotional and social influences, making them more likely to favour behavioural choices that  
84 will benefit them now rather than in later life (13, 16, 26, 27). When this is compounded by the  
85 complexity of overcoming a multitude of barriers that arise from the social and physical  
86 environment, engaging in healthy behaviours becomes much more difficult (28). In contrast to  
87 other studies that have used socio-ecological models to demonstrate the different levels of  
88 influence on health-related behaviours, we used an intersectional lens in order to explore how  
89 young people face multiple connected barriers to healthy behaviours (29-32). The aim of this  
90 study was therefore to explore young adults' barriers to healthy eating, physical activity, and  
91 the solutions to address these barriers with an intersectionality perspective.

92

93 **Methods**

94 *Setting and study design*

95 This study was conducted at the South African Medical Research Council (SAMRC)/Wits  
96 Developmental Pathways for Health Research Unit (DPHRU) at the Chris Hani Baragwanath  
97 Academic Hospital (CHBAH) in Soweto. CHBAH is the largest hospital in Africa and the third  
98 worldwide; it is also a public tertiary care institution that serves the low-income community of  
99 greater Soweto in south-western Johannesburg, South Africa (21). Soweto is one of the most  
100 well-known historically disadvantaged townships in South Africa known for its established  
101 communities, and socioeconomic and cultural diversity (33). However, while there is economic  
102 diversity in Soweto, poverty-related challenges are still a reality, including unemployment and  
103 food insecurity (34), and poor access to appropriate health services, especially for young adults  
104 (35).

105

106 Focus group discussions (FGDs) were utilised to collect data. We used a questionnaire to  
107 record the demographic information and anthropometric measures of participants. Participants  
108 were measured while wearing loose clothing and without shoes for all anthropometric  
109 measurements. A Seca 213 portable stadiometer was used to measure the participants' height  
110 (in cm), which was then translated to meters (m). Using a portable electronic bathroom scale,  
111 body weight was calculated to the nearest 0.1 kg. Weight in kilograms (kg) divided by height  
112 in meters (m<sup>2</sup>) was used to determine body mass index (BMI). These measurements were taken  
113 during the quantitative survey before the focus groups began.

114

115 *Participants and recruitment*

116 We purposively recruited and enrolled thirty black men and women between the ages of 18 and  
117 24 years from the Soweto household enumeration study database (36). The Soweto household  
118 enumeration database provided a sampling frame, which enabled identifying participants from


119 both low and high-income households hence ensuring participants' socioeconomic diversity  
120 (36, 37). During recruitment, research assistants contacted prospective participants using the  
121 contact information provided on the enumeration study database inviting them to participate.  
122 They outlined the purpose of the study, as well as provided contact information in case more  
123 information was required. Research assistants at DPHRU collated the information on  
124 prospective participants who were then called back to make appointments. The recruitment of  
125 these participants was to form part of a larger package of work by the Global Diet and Activity  
126 Research (GDAR) Network which started in the year 2020, explained in more detail elsewhere  
127 (37, 38).

128

### 129 *Theoretical framework*

130 The theoretical framework for this study is provided by an understanding of the  
131 Socio-Ecological model of health and the Theoretical Domains Framework (TDF) with an  
132 intersectionality perspective, which shows that health-related behaviours are influenced by  
133 physical, biological and psychological, sociological, economic and cultural barriers (39-42).  
134 The foundation of this paper is the assertion that various barriers intersect to affect how  
135 health-related behaviours turn out, which is supported by both the social-ecological model of  
136 health and the TDF (39-42). We applied the intersectionality framework to comprehend how  
137 context influences the emergence of health problems in complex ways rather than reducing  
138 perceptions and behaviours to a single characteristic (40). To be more precise, we assessed  
139 different factors (personal, domestic, societal, and neighbourhood) that affect people's  
140 decisions and beliefs and induce particular behaviours (39, 40, 42). The intersecting rings of  
141 the model (Figure 1) illustrate how factors at one level intersect with factors at another level  
142 and influence behavioural outcomes. For instance, easy access to unhealthy food may  
143 intersect with a person's taste preference, leading to higher consumption of unhealthy foods.

144 In recent years, there has been a strong push to develop environmental level health  
145 interventions such as school-based physical activity promotion to expand beyond programs  
146 that exclusively target individual level behaviours (43). Whilst research has extensively  
147 reported on the importance of households, parental feeding practices and family influence in  
148 shaping childhood health habits (44), the same cannot be said about adolescents and young  
149 adults (45). As young adults transition to adulthood, the personal, social, and environmental  
150 changes that transpire such as moving out of their home, studying, cohabitation with peers or  
151 partners, and starting a new job have a great influence on their lifestyle behaviour choices  
152 that are linked with poor diets and rapid weight gain (46). Therefore, interventions that target  
153 the environment may be more efficient and potentially more effective than individually  
154 targeted interventions because they are designed to change the context in which people live  
155 and work to create conditions supportive of healthy behavioural choices (41). In addition,  
156 although working with individuals to affect behaviour is difficult and resource consuming,  
157 interventions that influence policies and group-level behaviours can in turn affect individual-  
158 level behaviours among a much larger group of people and thus be more resource efficient  
159 (41, 47). Thus, health interventions targeting young men and women should take into account  
160 all intersecting elements [33].



161

162 Figure 1. A conceptual diagram showing different factors intersecting to influence young  
163 individuals health related behaviour.

164

#### 165 *Data collection*

166 The data were collected during May and June of 2020. The FGDs were conducted in English  
167 by two experienced multilingual qualitative researchers one facilitator and one note taker, with  
168 participants using vernacular where necessary. All FGDs were held remotely due to the  
169 COVID-19 pandemic and the necessity to ensure the safety of our participants. A day before  
170 the FGDs, all participants received a one-hundred-rand airtime recharge voucher to cover  
171 internet costs. Across all four FGDs, a topic guide (Additional file 1) was used to elicit  
172 discussions around perceived barriers and solutions to healthy eating and physical activity. A  
173 total of four FGDs were conducted, each comprised of 7-8 participants, and lasted between 45–  
174 90 minutes. The groups (G) were divided by gender, with two groups made up of young men  
175 and two groups made up of young women (all aged 18–24 years). All FGDs were audio-  
176 recorded with participants’ permission to record. The FGD sessions were meticulously  
177 documented, including observations. Audio recordings were transcribed verbatim and, when  
178 necessary, translated into English. All participants provided an electronically signed informed  
179 consent before taking part in the study. The Human Research Ethics Committee (HREC) at  
180 Witwatersrand University granted ethical clearance, with ethics number M190523.

181

#### 182 *Data analysis*

183 Data were analysed through thematic analysis (48, 49) with the help of the qualitative analysis  
184 software MAXQDA 2020 (version 20.4.2). Initially, data familiarisation was done by the first  
185 author, which involved checking transcripts and repeated reading of transcripts. Thereafter,  
186 coding, initial themes, reviewing themes, defining and naming themes, and finalising the



187 analysis followed (50). Other authors were asked to review the initial themes; the reviews were  
188 used to refine the key themes until an agreement was reached. The data were coded and  
189 analysed using a combination of deductive and inductive approaches. The deductive approach  
190 was based on pre-identified themes that focused on the research question, while the inductive  
191 approach was used for all themes generated from the transcripts and field notes (49). This  
192 process enabled an interpretation of young adults' perceptions of various barriers to healthy  
193 eating and physical activity, and specific solutions to address these barriers. The key identified  
194 domains in relation to barriers and solutions to healthy eating and physical activity are  
195 described in the results section.

196

## 197 **Results**

### 198 *Sample characteristics*

199 The analysis included a total of 30 participants. The average age of the men was 21.5 years,  
200 that of the women 20.5 years. Compared to the young women (46.7%), more than half of the  
201 young men (53.3%) had completed grade 12 or high school. Only 3.3% of the participants had  
202 a university degree (Additional Table 1). We present our findings from thematic analysis under  
203 the following four domains and nine main themes (in brackets): (1) Individual level factors  
204 (comprehension of health concepts and their interrelations; inability to apply nutrition  
205 knowledge to action; personal circumstances influence eating habits; not motivated enough to  
206 be active); (2) Influence at the household level (priority is for everyone to eat; immersed in  
207 household chores, no time for exercise); (3) Social influences (eating healthy means sickness)  
208 and (4) Influences by neighbourhood environment (junk food is inescapable, it's everywhere  
209 and cheap; no gyms, exercising outside is unsafe). Additional extracts of participants' views  
210 can be found in (Additional Table 2).

211

212 **Domain 1. Individual level factors**

213 ***Comprehension of health concepts and their interrelations***

214 In the various group discussions, young adults displayed a basic understanding of health  
215 concepts and their interrelationships. They described health as a daily process involving routine  
216 check-ups, sufficient sleep, a healthy and balanced diet, and physical activity to promote or  
217 maintain overall well-being. *“Health is how you take care of your body everyday like seeing  
218 the doctor to check BP [blood pressure] and with the kind of food that you eat like veggies and  
219 gym and the time you sleep. It is the wellbeing of your whole body.” (G2 men).*

220 When talking about nutrition, many young adults indicated that nutrition is about the  
221 foods people eat, which can be good or unhealthy. This in their opinion meant that nourishment  
222 could be both good and ill. Following a specific dietary plan that included eating fruit and foods  
223 high in protein and vitamins was generally recognised as good nutrition; *“[Nutrition is] a diet  
224 with proteins and vitamins that a person should follow like eating fruit and eating vegetables  
225 to stay healthy.” (G3 women).* Good nutrition is important for healthy growth, a robust immune  
226 system and a "sharp mind" [cleverness and quick thinking] according to participants' views;  
227 *“Food that helps our body in terms of growing strong and healthy, and also having a strong  
228 immune system and sharp mind”. (G1 women).* On the contrary, eating a modest amount of  
229 unhealthy food (referred to as "junk") was not considered a health risk by some young people,  
230 but eating too much of it was: *“I believe it is about how you eat the junk that is bad. If you  
231 have too much of junk food your health is going to be bad. But if watch what you eat then it's  
232 not bad.” (G1 women).*

233 In the discussions about physical activity, the young adults mostly defined it as  
234 participation in daily planned and organised exercise routines or sporting activity; *“Physical  
235 activity is part of our day-to-day life. It can be when you wake up in the morning take a walk  
236 or jog, play soccer or go to gym.” (G2 men).* Some of the reported benefits of physical activity

237 included strengthening muscles, maintaining good posture and preventing disease, especially  
238 heart disease; *“Physically activity, we also need that because that strengthens the muscles  
239 which ensures healthy posture and prevents diseases.” (G3 women)*. However, the young  
240 adults did not see unstructured and unplanned tasks such as cleaning, cooking or laundry as  
241 physical activities, but rather as barriers, that takes up a large part of their daytime and leaves  
242 them no time for exercise; *“I’m always busy like cooking and cleaning you know house duties  
243 and get tired so where do I get time to exercise.” (G3 women)*.

244 In many cases, young adults mentioned that physical inactivity was associated with  
245 obesity burden and increased risk of NCDs. *“[high obesity risk is] when you are a couch potato  
246 [always sited] and don’t exercise you gain fat..... you easily get sick with BP [high blood  
247 pressure] and heart problem when fat.” (G4 men)*. Physical activity was recognised as the only  
248 option for maintaining good health. From participants' narratives, physical activity is a more  
249 effective means of controlling obesity and preventing diseases than diet; *“Even if you eat junk  
250 foods, you can be healthy and safe from obesity and diseases if you get exercise.” (G4 men)*.  
251 In several instances, participants also acknowledged that genetics is a factor contributing to the  
252 burden of obesity *“Sometimes the obesity is taken from other people in your family.” (G4 men)*  
253 demonstrating a knowledge of risk factors outside their control.

254 Further, in the discussions, participants generally agreed that obesity meant a large  
255 amount of fat in the body that causes NCDs and could lead to death: *“[Obesity is] A drastic  
256 amount of fat in a human body. It can actually kill you as it brings heart problems and high  
257 BP.” (G3 women)*. Undoubtedly, young adults recognised obesity as a significant burden in  
258 their communities, but they believed that obesity had been normalised, as even those affected  
259 perceived it as a signal of contentment rather than a health risk. *“Yes, obesity is a problem  
260 where I live, but many people that are fat are fine with that and say it shows they don’t have  
261 problems [social and financial].” (G3 women)*.

262

263 ***Inability to apply nutrition knowledge in practice***

264 Across all FGDs, the majority of participants expressed that they did not know how to  
265 translate their knowledge about nutrition and physical activity into healthy behaviour; *“The*  
266 *thing is we don’t know a way to use this knowledge of how healthy foods are good to our body,*  
267 *at the same time our places [home and neighbourhood] make it hard to do exercise.” (G2*  
268 *men)*. One proposed solution to address this barrier was to provide health education in local  
269 languages, covering topics such as nutrition and physical activity through community TV and  
270 radio stations, social media platforms (WhatsApp, Facebook and Twitter) and integrated into  
271 school lessons, especially at the primary level (grades R-3). This solution, in the participants’  
272 opinion, is beneficial as it targets more than one barrier and takes into account language,  
273 outreach and accessibility issues that they have already encountered in other educational  
274 programmes:

275 *“We can start using TV or radio programs in the community to teach about*  
276 *healthy food and physical activity. Also periods in primary classes like grade R*  
277 *or all lower classes can also teach what is obesity in our language and what*  
278 *are the effects of obesity and what to do to avoid it instead of just saying we*  
279 *need to eat healthy and exercise.” (G4 men).*

280

281 ***Personal circumstances influence eating habits***

282 From participant discussions, it appeared that their basic understanding of health  
283 concepts, especially nutrition and physical activity, did not translate or influence an intent to  
284 engage in healthy behaviours. In all FGDs, participants admitted that their personal  
285 preferences, such as the taste of healthy foods, addiction to junk food, price and convenience  
286 of unhealthy foods were the main drivers to unhealthy eating patterns. In other cases, young

287 people mentioned that they were able to eat both healthy and unhealthy foods, depending on  
288 circumstances such as affordability, taste and weather. For example, some reported that certain  
289 foods, such as those high in fibre, were of great benefit in cold weather. *“I do eat fruit and veg.*  
290 *I also eat Kota [A quarter loaf of bread stuffed with foods such as fried chips, pork sausage,*  
291 *cheese, polony, tomato sauce and mango atchaar] when it’s cold days [because it is] high in*  
292 *fibre to keep warm.” (G1 women).* To overcome these barriers, participants suggested  
293 practising and promoting family meal sharing, as they recognised that time as an excellent  
294 opportunity for parents or adults in the family to teach the young one’s about healthy eating.  
295 *“We all eat together in my family, we eat the same food. I am forced to eat the greens I mean*  
296 *veggies like cabbage and spinach, [that] mom cooks a lot.” (G2 men).*

297

### 298 ***Not motivated enough to be active***

299 Further in the discussions, participants cited personal feelings such as laziness,  
300 discomfort and the monotony of exercise as the main reasons that prevent them from engaging  
301 in physical activity. *“I do know about exercising just that I just don’t want to exercise, it is*  
302 *boring and the pain, no.” (G2 men).* Some participants expressed interests in PA engagement;  
303 however, a lack of motivation was a barrier to their participation. *“In your back yard you are*  
304 *required to do the exercise alone and you are discouraged.” (G3 women).* Getting external  
305 motivators was proposed as a solution that could influence physical activity participation.  
306 According to participants, exercising with family or friends, or even joining a community club,  
307 might provide good motivation and promote regular participation in physical activity. *“I*  
308 *sometimes exercise with my friend in the morning and with my two older brothers in evenings.*  
309 *They make it fun and safe to exercise, I never miss.” (G1 women).*

310

### 311 **Domain 2. Influence at the household level**

312 ***Priority is for everyone to eat***

313 Some young adults' expressed a desire to lead a healthy lifestyle, but family and  
314 domestic barriers such as the inability to make decisions on what food to eat at home, limited  
315 food options due to household size, poverty and lack of family support to eat healthily were  
316 the main barriers that always put a halt to such a thought; "*The household where I live is poor,*  
317 *only mom works, I think I like to live healthy but they can't afford to buy healthy foods just for*  
318 *me alone.*" (G1 women). At this point, they declared that there was nothing else to do; moving  
319 away from home to become independent was the only solution to overcome these barriers. For  
320 them, independence meant taking control of their lives and deciding for themselves what they  
321 consume; "*Like when you start having your own place, you start buying groceries for*  
322 *yourself.*" (G4 men).

323

324 While being independent was one of the proposed solution to unhealthy eating,  
325 discussants argued that young children who are heavily dependent on their parents or relatives  
326 are a greater concern as they cannot make this choice of moving out of the family. They  
327 commented on how providing unhealthy foods to young children with the notion that it is a  
328 good way to pamper them increased their likelihood of health issues. Participants noted that  
329 many young children grow up believing that eating unhealthy foods is a form of self-indulgence  
330 because they have learned it at home; "*Small children are growing with obesity because of the*  
331 *junk like KFC they [caregivers/parents] provide to them, they believe in spoiling [ a nice treat]*  
332 *them with that KFC.*" (G1 women). To overcome this barrier, the young adults indicated that  
333 practising and teaching healthy eating at home is a much more viable strategy that can reduce  
334 the risk of premature mortality and diseases caused by eating poor quality meals from an early  
335 age. "*We should teach the young ones at home first about how do we eat healthy so they won't*  
336 *fall in the trap of eating junk food that causes diseases that can kill them later.*" (G2 men).

337

338 ***Immersed in household chores, no time for exercise***

339 Discussions around physical activities in the context of other everyday activities at  
340 home were also extensively noted. Our findings show that men and women face different  
341 challenges in regard to physical activities within the home. In all FGDs, young adult women  
342 reported that they found it difficult to find time to exercise because of other household  
343 responsibilities such as washing or cleaning, helping to cook or caregiving roles. “I’m always  
344 busy like cooking and cleaning you know house duties and get tired.” (G3 women). On the  
345 other hand, participants noted that men who could not exercise were just lazy, as they were  
346 not expected to do most of the household chores. “*I do know about exercising just that I’m*  
347 *lazy and just don’t want to exercise, it is boring.*” (G2 men). Becoming independent was their  
348 proposed solution to overcome this barrier because staying alone meant less chores. “*Staying*  
349 *alone is better, less duties and more time for exercising.*” (G3 women).

350

351 **Domain 3. Social influence**

352 ***Eating healthy means sickness***

353 Participants expressed their opinions on how peers, community members, and people in their  
354 communities have a general negative perception that eating healthy food is a sign of sickness.  
355 In particular, eating vegetables and losing weight are seen as indicators of illnesses like  
356 HIV/AIDS or those struggling with NCDs like diabetes and hypertension. As such, gaining  
357 weight and consuming unhealthy foods like "Kotas" seemed ideal because it meant no  
358 judgment from others; “*You know you start to lose fat when you eat veggies right. The majority*  
359 *of them [peers/community members] believe if you eat healthy stuff like veggies right, it means*  
360 *that you are sick with things like AIDS, so I rather not eat.*” (G4 men). This was another way  
361 on in which consumption of unhealthy food was normalised. Participants proposed that

362 nutrition education that can be delivered through group discussions in community can be a  
363 solution to create awareness and address this barrier. Accordingly, the discussions would  
364 encourage learning from one another, improving their understanding of the concept of healthy  
365 eating and associated benefits; “[*The solution is*]....*make a group discussion with people and*  
366 *educate each other on nutrition and the benefits so they don’t judge you.*” (G3 women)

367

#### 368 **Domain 4. Influence by neighbourhood environment**

##### 369 ***Junk food is all over and cheap***

370 Discussants reported that most street vendors and tuck shop sellers primarily sell  
371 unhealthy foods (commonly referred to as "junk") at lower prices. In addition, they noted that  
372 the high visibility of advertisements for unhealthy foods on main roads and neighbourhood  
373 streets exacerbates unhealthy eating habits. Based on their accounts, such visualizations,  
374 advertisements and proximity to junk foods have influenced unhealthy eating especially  
375 amongst young people:

376 *“When it comes to changing the situation and eat healthy you can’t because of*  
377 *the neighbourhood that you live around, we are exposed to junk food all over*  
378 *and it’s cheap at the spaza [neighbourhood convenient tuck shop]. As you walk*  
379 *down the road all you see is a big billboard with coke, burger and chips so you*  
380 *always think of eating them.*” (G2 men).

381 These challenges led participants to propose setting up community gardens as a solution. They  
382 also mentioned a need to have more healthier food advertisements and awareness campaigns  
383 by health professionals about the dangers of unhealthy foods. Others argued that unhealthy  
384 eating should be considered as a pandemic, meaning government regulations on unhealthy food  
385 must be imposed in order to address these neighbourhood-level barriers; “[*The government must*  
386 *address the nation like with Covid, address with a couple of food that is unhealthy and show*



387 *healthy one, do same like hand sanitizers when you go to the shop, you know it is law.” (G3*  
388 *women).*

389

### 390 ***No gyms! Exercising outside is unsafe***

391 While discussing about the environmental barriers to physical activity, it was noted that  
392 structural barriers, such as a lack of access to gyms or facilities for physical activity were  
393 among the most prevalent challenges that young adults’ experienced; *“The fact is that we don’t*  
394 *have places that we can go maybe to exercise after eating that Kota”.* (G4 men). Participants’  
395 narratives also revealed gender disparities, with women generally fearing for their safety and  
396 encountering sexist remarks while exercising around the neighbourhood. *“In my area I can’t*  
397 *even jog outside, there are always boys sitting at the corner saying embarrassing [sexist*  
398 *remarks] things about my body and their presence makes me feel uncomfortable, I don’t feel*  
399 *safe.”* (G1 women). To address these barriers, participants proposed that young women should  
400 be encouraged to use their household backyards for physical activity. In addition, participants  
401 noted a need for the government to construct more facilities for physical activity in the  
402 community as well as employing security officers in community parks, or having a community  
403 patrol to man security. This was said to be beneficial to all community members and may  
404 influence many young people to engage in physical activity.

405

### 406 **Discussion**

407 We set out to explore young adults' barriers to healthy eating, physical activity, and the  
408 solutions to address these barriers with an intersectionality perspective. Findings from the  
409 FGDs suggested that young adults in this urban South African township understood the  
410 significance of a healthy diet and physical activity and their relationship with health. However,  
411 they felt unable to put their knowledge into practice. Young adults in this study reported several


412 barriers to healthy eating including taste and preference; a lack of autonomy on food decisions  
413 at home; negative beliefs about healthy foods; street vendors, advertisements and convenience  
414 stores that primarily sell unhealthy foods at lower prices. The commonly discussed barriers to  
415 physical activity were laziness; lack of facilities; and safety concerns when engaging in PA in  
416 their neighbourhoods.

417 Furthermore, we found that barriers to physical activity differed by gender. On one hand,  
418 young women found it more difficult to exercise because of housework chores, experienced  
419 sexist remarks or they had safety concerns. On the other hand, laziness, discomfort and the  
420 monotony of exercise and lack of exercise facilities demotivated young men to engage in  
421 physical activity. That said, young adults proposed a number of solutions to addressing these  
422 barriers and among these were: the practice and promoting family meal sharing; providing  
423 health education in local languages; becoming independent; getting external motivators;  
424 practising and teaching healthy eating at home by adults; creating community gardens; more  
425 advertising of healthier foods; conducting more education campaigns by health professionals  
426 on the dangers of unhealthy foods; establishing a community patrol forum; and building  
427 accessible physical activity infrastructure by the government.

428

429 Our findings are consistent with those reported in studies conducted with adolescents,  
430 young women, and adults [19, 21, 22, 41-44]. According to Ware et al.,[22] in a South African  
431 study of urban young women, those who lived with their parents believed they had no influence  
432 over what was cooked and consumed in the home, and they perceived unhealthy foods to be  
433 cheaper, more accessible, and convenient because they were readily available. Among adults,  
434 Bosire et al. reported that the stigma associated with being thin, which links it to conditions  
435 such as HIV/AIDS, has a negative impact on people's risk perceptions of being fat or  
436 overweight and makes them want to gain weight in an effort to escape this stigma. Wrottesley

437 et al. [19] also reported on how adolescent girls were discouraged from exercising because they  
438 were concerned about losing weight or becoming too thin. Our findings, as well as those of  
439 previous studies, support our hypothesis that the young adults in this study are confronted with  
440 multiple barriers arising from the macro and micro levels of society [41] (individual, family,  
441 community-level, grassroots institutions, and policies ) that intersect to complicate their ability  
442 to engage in healthy behaviours, as shown in the conceptual diagram below (Figure 2), which  
443 illustrates how individual-level factors intersect with household-, neighbourhood-, and  
444 societal-level factors to influence behaviour. There is a need to understand the complex nature  
445 of these health inequities especially among young adults. Intersectionality is becoming more  
446 widely recognised as an important theoretical approach for studying health-related inequalities  
447 by highlighting the intersections of individuals' multiple identities within social power systems  
448 that exacerbate and reinforce experiences of poor health behaviours [42]. To the authors'  
449 knowledge, this is the first study carried out in South Africa that analyses the barriers that  
450 hinder young adults from engaging in physical activity and eating healthily as well as the  
451 solutions to overcome these barriers using an intersectionality framework.



452  
453 Figure 2. Conceptual diagram showing the intersectionality between different multilevel  
454 barriers to healthy eating and physical activity influencing young adults' health related  
455 behaviour.

456            Since the 1970s, neoliberal ideologies have grown in popularity and provide little  
457 support for addressing social determinants of health and wellbeing (51, 52). These ideologies  
458 emphasize individual attributes and choice as major drivers of health (53) and have led to a  
459 health promotion practice where interventions are implemented without taking contextual  
460 factors into account [49]. Intersectionality goes beyond looking at individual factors like  
461 biology, socioeconomic status, sex, gender, and race. It focuses on how these factors relate to  
462 one another and interact with one another at various societal levels to understand how health is  
463 shaped across demographic groups and geographical contexts (54). In this study,  
464 intersectionality is employed to explain how individual, household, societal, and  
465 neighbourhood barriers to healthy eating and physical activity participation interact, as well as  
466 their proposed solutions. As Louis Althusser pointed out, people (in this case, young adults)  
467 are conditioned to not only hear messages but also obey, comply, and live up to them (55). It  
468 is these context-based generated messages that address social ideas of race, class, gender,  
469 sexual orientation, age, and even physical characteristics often associated with disabilities or  
470 abilities in young adults (55). This means that healthy eating and physical activity perceptions  
471 are more strongly influenced by the messages and meanings transmitted by meaning providers  
472 through social conditioning. Parents, friends, family and community members are among the  
473 people who can provide meaning, as do groups like those involved in school sports, churches  
474 or other religious institutions, and the media (visual and audio). Consequently, what young  
475 adults encounter in their families (lack of family support to eat healthily), neighbourhood or  
476 communities (easy access to unhealthy foods), and through media like as TV and music  
477 (advertisements of unhealthy foods) influence what they perceive as important and what they  
478 ultimately prioritise in a given society. At this point, intersectionality is when all of these  
479 categories cross paths, revealing how inequities are shaped by interactions between different  
480 sites and spheres of power, which allows us to observe how these are all interconnected on a

481 larger scale. Considering the intersectionality of social constructs, identities, and histories, we  
482 can see that there is even more connection between barriers to healthy eating and physical  
483 activity at the individual, household, social, and neighbourhood levels. To fully understand  
484 one, we need to understand the others.

485

486 Furthermore, applying an intersectionality lens in this study contributes to highlighting  
487 significant differences between young men and women that are often overlooked when  
488 addressing barriers to healthy eating and physical activity. In particular, it offers an explanation  
489 to how young men and women from lower socioeconomic groups might decide to exercise  
490 outside because they lack access to gyms, but only the young women would encounter sexism  
491 and concerns about safety. Because of this, young men have a slight gender advantage over  
492 young women when choosing to exercise on neighbourhood streets. Furthermore, this may also  
493 explain why some health education interventions are unsuccessful despite being recognised by  
494 research as a crucial component in the management and prevention of NCDs and their  
495 associated risk factors [49]. For example, South Africa is a country with a wide range of  
496 linguistic and cultural diversity, yet health systems and public health interventions frequently  
497 use English as the main language for imparting health education, either orally or in handouts  
498 [51]. This is despite the evidence that shows there is a lack of health education among South  
499 Africans [14, 50, 51]. In this case, it becomes apparent that English-speaking people from  
500 lower socioeconomic groups may have a greater advantage in accessing health services and  
501 comprehend how health education pertaining to nutrition and physical activity can be applied  
502 than Africans who do not speak and understand English but have similar backgrounds. All this  
503 various aspects at different contextual levels interact and mutually inform one another and  
504 should be taken into account when implementing interventions or solutions that aim to  
505 influence healthy behaviour choices.

506           There are some limitations on the current study. The FGDs were held in the midst of  
507 the COVID-19 pandemic using online video conferencing software rather than where  
508 participants lived and went about their daily lives, so we were unable to supplement the data  
509 obtained through the FGDs with additional sources of information like field observations.  
510 Some conversations had to be halted due to unstable network conditions. The young men and  
511 women who formed part of the discussions did not practice healthy eating or engage in regular  
512 physical activity; however, holding discussions with those who have adopted health related  
513 behaviours might have been of interest.

514

## 515 **Conclusion**

516 This study is important because it highlights the barriers to healthy eating and physical activity  
517 among young adults in Soweto and the proposed solutions. While some solutions have been  
518 proposed, young adults believe that they are not able to put them into practice. Our findings  
519 have also shown that young adults face a multitude of barriers that occur simultaneously at the  
520 individual, household, neighbourhood and societal levels, making it difficult for them to make  
521 healthy behavioural choices. We advocate for young people to be involved in developing or  
522 designing solutions to promote health or influence healthy lifestyles. By focusing on the  
523 intersection or overlap of different factors, findings of this study could be used to influence  
524 health-related programmes in Soweto and other similar contexts.

525

## 526 **Declarations**

527 *Ethics approval and consent to participate* – The Human Research Ethics Committee (HREC)  
528 of the Witwatersrand University, under ethics number M190523, granted ethical approval.

529 *Consent for publication* – All participants gave written informed consent prior to taking part in  
530 the study.

531 *Availability of data and materials* – The dataset used and/or analysed during the current study  
532 are available from the corresponding author on reasonable request.

533 *Competing interests* – No potential conflict of interest was reported by the authors.

534 *Funding* – This research was funded by the National Institute for Health Research (NIHR)  
535 (GHR: 16/137/34) using UK aid from the UK Government to support global health research.

536 The views expressed in this publication are those of the authors and not necessarily those of  
537 the NIHR or the UK Department of Health and Social Care. GM is supported by the DSI-NRF  
538 Centre of Excellence in Human Development at the University of the Witwatersrand,  
539 Johannesburg, South Africa.

540 *Authors' contributions* – G.M. collected data, conceptualised the paper, analysed data and  
541 wrote the manuscript; S.A.N. conceptualised the paper and commented on the manuscript;  
542 E.N.B. and SK commented on the manuscript.

543 *Acknowledgements* – The authors thank the participants who took part in this research and the  
544 Soweto GDAR research team whose work they represent here.

545

## 546 **References**

547 1. World Health Organisation (WHO). Deaths from noncommunicable diseases on the  
548 rise in Africa <https://www.afro.who.int/news/deaths-noncommunicable-diseases-rise-africa>  
549 Accra/Brazzaville: WHO Regional Office for Africa; 2022 [updated 11 April 2022. Available  
550 from: <https://www.afro.who.int/news/deaths-noncommunicable-diseases-rise-africa>.

551 2. Nyirenda MJ. Non-communicable diseases in sub-Saharan Africa: understanding the  
552 drivers of the epidemic to inform intervention strategies. *International Health*. 2016;8(3):157-

553 8.

- 554 3. Girum T, Mesfin D, Bedewi J, Shewangizaw MJJjocd. The burden of  
555 noncommunicable diseases in Ethiopia, 2000–2016: analysis of evidence from global burden  
556 of disease study 2016 and global health estimates 2016. 2020;2020.
- 557 4. Gouda HN, Charlson F, Sorsdahl K, Ahmadzada S, Ferrari AJ, Erskine H, et al. Burden  
558 of non-communicable diseases in sub-Saharan Africa, 1990-2017: results from the Global  
559 Burden of Disease Study 2017. *The Lancet Global health*. 2019;7(10):e1375-e87.
- 560 5. Hay P, Mitchison D, Collado AEL, González-Chica DA, Stocks N, Touyz S. Burden  
561 and health-related quality of life of eating disorders, including Avoidant/Restrictive Food  
562 Intake Disorder (ARFID), in the Australian population. *Journal of eating disorders*. 2017;5:21.
- 563 6. World Health Organization. Noncommunicable diseases country profiles 2018.  
564 Geneva; 2018.
- 565 7. Mialon M. An overview of the commercial determinants of health. *Globalization and*  
566 *Health*. 2020;16(1):74.
- 567 8. World Health Organisation (WHO). Commercial determinants of health  
568 <https://www.who.int/news-room/fact-sheets/detail/commercial-determinants-of-health>:  
569 WHO; 2021 [updated 5 November 2021; cited 2022 22 June]. Available from:  
570 <https://www.who.int/news-room/fact-sheets/detail/commercial-determinants-of-health>.
- 571 9. Organization WH. Global status report on noncommunicable diseases 2014: World  
572 Health Organization; 2014.
- 573 10. Patton GC, Sawyer SM, Santelli JS, Ross DA, Afifi R, Allen NB, et al. Our future: a  
574 Lancet commission on adolescent health and wellbeing. 2016;387(10036):2423-78.
- 575 11. Shisana O, Labadarios D, Rehle T, Simbayi L, Zuma K, Dhansay A, et al. South African  
576 National Health and Nutrition Examination Survey (SANHANES-1). Cape Town: HSRC  
577 Press. 2013.



- 578 12. Guthold R, Stevens GA, Riley LM, Bull FC. Worldwide trends in insufficient physical  
579 activity from 2001 to 2016: a pooled analysis of 358 population-based surveys with 1·9 million  
580 participants. *The Lancet Global health*. 2018;6(10):e1077-e86.
- 581 13. Pisa PT, Pedro TM, Kahn K, Tollman SM, Pettifor JM, Norris SA. Nutrient patterns  
582 and their association with socio-demographic, lifestyle factors and obesity risk in rural South  
583 African adolescents. *Nutrients*. 2015;7(5):3464-82.
- 584 14. National Department of Health (NDoH), Statistics South Africa (Stats SA), South  
585 African Medical Research Council (SAMRC), ICF. South Africa Demographic and Health  
586 Survey 2016. Pretoria, South Africa, and Rockville, Maryland, USA: NDoH, Stats SA,  
587 SAMRC, and ICF; 2019.
- 588 15. De Wet-Billings N. Preventable deaths among youth in South Africa: Measuring life  
589 expectancy in the absence of non-communicable diseases and its implications for the healthcare  
590 system. *S Afr Med J*. 2021;111(4):361-4.
- 591 16. Mukoma G, Wrottesley SV, Kagura J, Oni T, Micklesfield L, Norris SA. The  
592 relationships between socioeconomic status, dietary knowledge and patterns, and physical  
593 activity with adiposity in urban South African women. *South African Journal of Clinical  
594 Nutrition*. 2022:1-7.
- 595 17. Ronquest-Ross L-C, Vink N, Sigge GO. Food consumption changes in South Africa  
596 since 1994 %J *South African Journal of Science*. 2015;111:01-12.
- 597 18. Shisana O, Labadarios D, Rehle T, Simbayi L, Zuma K, Dhansay A, et al. The South  
598 African National Health and Nutrition Examination Survey, 2012: SANHANES-1: the health  
599 and nutritional status of the nation. 2014.
- 600 19. NDoH SS, SAMRC IJSSA, South African Medical Research Council , ICF. South  
601 Africa Demographic and Health Survey 2016. Pretoria, South Africa, and Rockville, Maryland,  
602 USA: National Department of Health (NDoH). 2019.

- 603 20. Seguin R, Connor L, Nelson M, LaCroix A, Eldridge G. Understanding barriers and  
604 facilitators to healthy eating and active living in rural communities. *J Nutr Metab.*  
605 2014;2014:146502.
- 606 21. Wrottesley SV, Bosire EN, Mukoma G, Motlathledi M, Mabena G, Barker M, et al.  
607 Age and gender influence healthy eating and physical activity behaviours in South African  
608 adolescents and their caregivers: Transforming Adolescent Lives through Nutrition Initiative  
609 (TALENT). *Public Health Nutr.* 2021;24(16):5187-206.
- 610 22. Sallis JF, Floyd MF, Rodríguez DA, Saelens BE. Role of built environments in physical  
611 activity, obesity, and cardiovascular disease. *Circulation.* 2012;125(5):729-37.
- 612 23. Bosire EN, Stacey N, Mukoma G, Tugendhaft A, Hofman K, Norris SA. Attitudes and  
613 perceptions among urban South Africans towards sugar-sweetened beverages and taxation.  
614 *Public Health Nutr.* 2020;23(2):374-83.
- 615 24. Ware LJ, Prioreshi A, Bosire E, Cohen E, Draper CE, Lye SJ, et al. Environmental,  
616 Social, and Structural Constraints for Health Behavior: Perceptions of Young Urban Black  
617 Women During the Preconception Period-A Healthy Life Trajectories Initiative. *J Nutr Educ*  
618 *Behav.* 2019;51(8):946-57.
- 619 25. Leech RM, McNaughton SA, Timperio A. The clustering of diet, physical activity and  
620 sedentary behavior in children and adolescents: a review. *The international journal of*  
621 *behavioral nutrition and physical activity.* 2014;11:4.
- 622 26. National Academies of Sciences E, Medicine J Birth Settings in America: Outcomes  
623 Q, Access,, Choice. Health and Medicine Division; Division of Behavioral and Social Sciences  
624 and Education; Board on Children, Youth, and Families; Committee on the Neurobiological  
625 and Socio-behavioral Science of Adolescent Development and Its Applications; Backes EP,  
626 Bonnie RJ, editors. *The Promise of Adolescence: Realizing Opportunity for All Youth.* 2019.

- 627 27. Micklesfield LK, Munthali RJ, Prioreshi A, Said-Mohamed R, van Heerden A,  
628 Tollman S, et al. Understanding the Relationship between Socio-Economic Status, Physical  
629 Activity and Sedentary Behaviour, and Adiposity in Young Adult South African Women Using  
630 Structural Equation Modelling. *Int J Environ Res Public Health*. 2017;14(10).
- 631 28. Koshedo SA, Paul-Ebhohimhen VA, Jepson RG, Watson MCJBph. Understanding the  
632 complex interplay of barriers to physical activity amongst black and minority ethnic groups in  
633 the United Kingdom: a qualitative synthesis using meta-ethnography. 2015;15(1):1-16.
- 634 29. Uchendu C, Windle R, Blake H. Perceived facilitators and barriers to nigerian nurses'  
635 engagement in health promoting behaviors: a socio-ecological model approach. *International*  
636 *journal of environmental research and public health*. 2020;17(4):1314.
- 637 30. Upreti YR, Bastien S, Bjonness B, Devkota B. The socio-ecological model as a  
638 framework for understanding junk food consumption among schoolchildren in Nepal. *Nutrition*  
639 *and Health*. 2021;27(3):337-46.
- 640 31. Ngwenya N, Nkosi B, Mchunu LS, Ferguson J, Seeley J, Doyle AMJPo. Behavioural  
641 and socio-ecological factors that influence access and utilisation of health services by young  
642 people living in rural KwaZulu-Natal, South Africa: Implications for intervention.  
643 2020;15(4):e0231080.
- 644 32. Wang C-W, Chen D-R, Chan C-C, Yeh Y-P, Chen H-HJPHN. 'I Knew I Should Stop,  
645 but I Couldn't Control Myself': a qualitative study to explore the factors influencing  
646 adolescents' consumption of sugar-sweetened beverages and sugary snacks from a socio-  
647 ecological perspective. 2022;25(9):2465-74.
- 648 33. Alexander P, Ceruti C, Motseke K, Phadi M, Wale K. *Class in Soweto: University of*  
649 *KwaZulu-Natal Press*; 2013.
- 650 34. Battersby J, McLachlan M. Urban food insecurity: a neglected public health challenge.  
651 *S Afr Med J*. 2013;103(10):716-7.

- 652 35. Miller CL, Nkala B, Closson K, Chia J, Cui Z, Palmer A, et al. The Botsha Bophelo  
653 Adolescent Health Study: A profile of adolescents in Soweto, South Africa. *Southern African*  
654 *journal of HIV medicine*. 2017;18(1):731.
- 655 36. Prioreshi A, Wrottesley SV, Norris SA. Physical Activity Levels, Food Insecurity and  
656 Dietary Behaviours in Women from Soweto, South Africa. *J Community Health*.  
657 2021;46(1):156-64.
- 658 37. Odunitan-Wayas FA, Wadende P, Mogo ERI, Brugulat-Panes A, Micklesfield LK,  
659 Govia I, et al. Adolescent Levers for a Diet and Physical Activity Intervention Across  
660 Socioecological Levels in Kenya, South Africa, Cameroon, and Jamaica: Mixed Methods  
661 Study Protocol. *JMIR Res Protoc*. 2021;10(7):e26739.
- 662 38. Oni T, Assah F, Erzse A, Foley L, Govia I, Hofman KJ, et al. The global diet and  
663 activity research (GDAR) network: a global public health partnership to address upstream NCD  
664 risk factors in urban low and middle-income contexts. *Globalization and Health*.  
665 2020;16(1):100.
- 666 39. Etherington C, Rodrigues IB, Giangregorio L, Graham ID, Hoens AM, Kasperavicius  
667 D, et al. Applying an intersectionality lens to the theoretical domains framework: a tool for  
668 thinking about how intersecting social identities and structures of power influence behaviour.  
669 *BMC Medical Research Methodology*. 2020;20(1):169.
- 670 40. Sallis JF, Owen N, Fisher EB. Ecological models of health behavior. *Health behavior*  
671 *and health education: Theory, research, and practice*, 4th ed. San Francisco, CA, US: Jossey-  
672 Bass; 2008. p. 465-85.
- 673 41. Eldredge LKB, Markham CM, Ruiters RA, Fernández ME, Kok G, Parcel GS. *Planning*  
674 *health promotion programs: an intervention mapping approach*: John Wiley & Sons; 2016.
- 675 42. Stokols D. Translating social ecological theory into guidelines for community health  
676 promotion. *Am J Health Promot*. 1996;10(4):282-98.

- 677 43. Long KZ, Beckmann J, Lang C, Seelig H, Nqweniso S, Probst-Hensch N, et al. Impact  
678 of a school-based health intervention program on body composition among South African  
679 primary schoolchildren: results from the KaziAfyra cluster-randomized controlled trial. *BMC*  
680 *Medicine*. 2022;20(1):27.
- 681 44. Finnane JM, Jansen E, Mallan KM, Daniels LAJ. Behavior. Mealtime structure and  
682 responsive feeding practices are associated with less food fussiness and more food enjoyment  
683 in children. 2017;49(1):11-8. e1.
- 684 45. Li K-K, Concepcion RY, Lee H, Cardinal BJ, Ebbeck V, Woekel E, et al. An  
685 examination of sex differences in relation to the eating habits and nutrient intakes of university  
686 students. 2012;44(3):246-50.
- 687 46. Winpenny EM, van Sluijs EMF, White M, Klepp K-I, Wold B, Lien N. Changes in diet  
688 through adolescence and early adulthood: longitudinal trajectories and association with key life  
689 transitions. *International Journal of Behavioral Nutrition and Physical Activity*. 2018;15(1):86.
- 690 47. Bauer KW, Yang YW, Austin SB. “How can we stay healthy when you’re throwing all  
691 of this in front of us?” Findings from focus groups and interviews in middle schools on  
692 environmental influences on nutrition and physical activity. *Health Education & Behavior*.  
693 2004;31(1):34-46.
- 694 48. Braun V, Clarke V. Using thematic analysis in psychology. *Qualitative Research in*  
695 *Psychology*. 2006;3(2):77-101.
- 696 49. Braun V, Clarke V. Can I use TA? Should I use TA? Should I not use TA? Comparing  
697 reflexive thematic analysis and other pattern-based qualitative analytic approaches.  
698 *Counselling and Psychotherapy Research*. 2020;21(1):37-47.
- 699 50. Clarke V, Braun V. *Successful Qualitative Research: A Practical Guide for Beginners*.  
700 London: Sage; 2013.

- 701 51. Hankivsky O, Christoffersen A. Intersectionality and the determinants of health: a  
702 Canadian perspective. *Critical Public Health*. 2008;18(3):271-83.
- 703 52. Lindsay J. Healthy living guidelines and the disconnect with everyday life. *Critical*  
704 *Public Health*. 2010;20(4):475-87.
- 705 53. Baum F, Fisher M. Why behavioural health promotion endures despite its failure to  
706 reduce health inequities. *Sociol Health Illn*. 2014;36(2):213-25.
- 707 54. Kapilashrami A, Hankivsky O. Intersectionality and why it matters to global health.  
708 *Lancet*. 2018;391(10140):2589-91.
- 709 55. Althusser L. Ideology and ideological state apparatuses (notes towards an  
710 investigation). *The anthropology of the state: A reader*. 92006. p. 86-98.
- 711