Distinct neurofunctional alterations during motivational and hedonic processing of natural and monetary rewards in depression – a neuroimaging meta-analysis

Mercy Chepngetich Bore[1,2], Xiqin Liu[1,2], Xianyang Gan[1,2], Lan Wang[1,2], Ting Xu[1,2], Stefania Ferraro[1,2], Liyuan Li[1], Bo Zhou[1], Jie Zhang[3,4], Deniz Vatansever[3,4], Bharat Biswal[5], Benjamin Klugah-Brown[1,2*], Benjamin Becker[1,2*]

[1]Center of Psychosomatic Medicine, Sichuan Provincial Center for Mental Health, Sichuan Provincial People's Hospital, University of Electronic Science and Technology of China, Chengdu, China

[2]MOE Key Laboratory for Neuroinformation, School of Life Science and Technology, University of Electronic Science and Technology of China, Chengdu, China

[3]Institute of Science and Technology for Brain-Inspired Intelligence, Fudan University, Shanghai, China

[4]Key Laboratory of Computational Neuroscience and Brain Inspired Intelligence, Fudan University, Ministry of Education, China

[5]Department of Biomedical Engineering, New Jersey Institute of Technology, Newark, NJ, USA

*Corresponding authors

Correspondence:

Benjamin Becker (ben_becker@gmx.de) or Benjamin Klugah-Brown (bklugah@gmail.com)

University of Electronic Science and Technology of China

Xiyuan Ave 2006, 611731 Chengdu, China

Abstract

Reward processing alterations have been suggested as a candidate mechanism underlying anhedonia and apathy in depression. Neuroimaging studies have documented that neurofunctional alterations in mesocorticolimbic circuits may neurally mediate reward processing deficits in depression. However, common and distinct neurofunctional alterations during motivational and hedonic evaluation of monetary (extrinsic) and natural (intrinsic) rewards in depression have not been systematically examined. Here, we capitalized on a series of pre-registered neuroimaging meta-analyses to (1) establish general reward-related neural alterations in depression, (2) determine common and distinct alterations during anticipation of monetary rewards, receipt of monetary rewards, and receipt of natural rewards, and, (3) characterize the differences on the behavioral, network and molecular level. The coordinate-based meta-analysis included a total of 633 depressed patients and 644 healthy controls and revealed generally decreased subgenual anterior cingulate cortex (ACC) and striatal reactivity towards rewards in depression. Subsequent quantitative comparison analysis indicated that monetary rewards led to decreased hedonic reactivity in the right ventral caudate while natural rewards led to decreased reactivity in the bilateral putamen. These regions exhibited distinguishable profiles on the behavioral, network and receptor level. Further analyses demonstrated that the right thalamus and left putamen showed decreased activation during the anticipation of monetary reward. The present results indicate that distinguishable neurofunctional alterations may neurally mediate reward-processing alterations in depression, particularly with respect to monetary and natural rewards. Given that natural rewards prevail in everyday life, our findings suggest that reward-type specific interventions are warranted and challenge the generalizability of

experimental tasks employing monetary incentives to capture reward dysregulations in everyday life.

Keywords: Reward; Depression; Striatum; Functional magnetic resonance imaging; Meta-analysis

Introduction

Pleasure and rewards constitute vital attractors that not only shape our momentary decisions, but also motivate future behavior. Deficits in reward processing have been proposed as a candidate mechanism underlying anhedonia, which describes a consistently diminished pleasure and interest in almost all daily life activities (1), (see also, integration in the Research Domain Criteria framework: positive valence system) (2,3). Anhedonia is a cardinal symptom of Major Depressive Disorder (MDD) and it represents a transdiagnostic symptom of several debilitating neurological and mental health disorders (4) . With currently over 320 million people living with depression, MDD has become a leading cause of disability worldwide (5). While the established interventions can alleviate some of the symptoms of depression, anhedonia is not only inefficiently targeted by the available treatments, but it is also highly resistant (6–11). Understanding the specific dysfunctions of behavioral and neural reward processing in depression may thus provide better treatment and management options (12).

Reward processing encompasses an entire array of subprocesses that can be defined based on different perspectives including the class of external reinforcers as well as the stage and function of the rewarding process. With respect to the nature of the reinforcers, intrinsic rewards referring to non-conditioned and inherently pleasurable stimuli such as music or food, can be differentiated from extrinsic rewards (e.g., money) which are not inherently pleasurable but exert their motivational value via learned (conditioned) associations with intrinsic rewards (13,14). Moreover, the anticipation and receipt stages of rewards are commonly differentiated and considered to reflect motivational and hedonic subprocesses, respectively (15–17).

On the neural level, reward processing has been closely linked to activation in mesocorticolimbic systems, including the striatum and amygdala as well as medial prefrontal

and anterior cingulate regions (13,18). While some evidence suggests that the specific neural systems vary as a function of the reinforcers and the stage of the reward process (16,19,20), the ventral striatum has been consistently engaged during the anticipation and receipt of monetary rewards (21,22), as well as the receipt of different intrinsic (natural) rewards such as food or music (23,24). In accordance with overarching frameworks (25,26), these meta-analytic findings may support the notion of a "common neural currency" for reward (27). However, see also (28), who suggests phylogenetically different classes of rewards.

Previous meta-analyses have provided convergent evidence for reward processing deficits in depression (1,29), with functional neuroimaging studies suggesting altered neural activity during reward processing (30–32). Alterations have been observed along the mesocorticolimbic reward circuit, including prefrontal regions such as the orbitofrontal cortex (OFC), medial prefrontal cortex (mPFC) and the anterior cingulate cortex (ACC), as well as subcortical regions including the striatum, amygdala, and hippocampus (30,33–35). Neurofunctional alterations in the striatum and ACC during monetary reward processing have been consistently observed during different stages of depression, including subthreshold depression (36), major depressive episodes (30,37,38) or remitted depression (39), and may thus represent a common marker for depression-associated reward deficits. However, findings with respect to depression-related alterations during natural reward processing have revealed less consistent results (12,40,41), and an accumulating number of recent electroencephalography (EEG) studies that directly compared processing of monetary and natural rewards in depression suggest not only common but also domain-specific neurofunctional alterations (42,43). Against this background, we examined if depression is characterized by a dysregulation of the 'common neural currency' for rewards by capitalizing on a robust fMRI meta-analytic strategy that allowed us to determine common and

separable neurofunctional alterations during natural versus monetary reward processing and during anticipation versus receipt of reward in depression.

We conducted a preregistered coordinate-based meta-analysis according to the latest guidelines (44,45) and included case-control fMRI studies examining the anticipation and receipt of monetary and natural rewards in depression. To facilitate a robust determination, we employed Seed-based d Mapping with Permutation of Subject Images (SDM-PSI), a novel method of performing neuroimaging meta-analyses (46). SDM-PSI allows (nearly) unbiased estimation of effect sizes and generation of neurofunctional maps based on both positive and negative differences and hence can produce signed differential effect size maps. In line with the goals of the present study, we employed a three-step approach to meta-analytically determine common and domain-specific reward dysfunctions in depression, including a pooled analysis determining non-specific reward alterations, as well as separate meta-analyses determining common and domain-specific hedonic dysregulations during the receipt of monetary versus natural rewards or the anticipation and receipt of monetary rewards, respectively. We hypothesized domain-general alterations in the ventral striatal reward processing system (reflecting dysregulations in the 'common neural currency'), but domain-specific alterations in prefrontal systems involved in more complex value and evaluation processes. To further characterize common and separable alterations, we next employed a series of connectivity and meta-analytic strategies to distinguish the identified regions on the behavioral, network, as well as molecular genetics and receptor levels.

Method

Data collection procedures, inclusion criteria and data extraction

The current pre-registered meta-analysis adhered to the guidelines of conducting coordinate-based neuroimaging meta-analyses (44). Procedures, hypotheses and analyses

were pre-registered on the OSF-repository (https://osf.io/ay3r9). A comprehensive literature search was performed based on the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (45). The search aimed to identify case-control fMRI studies examining reward processing in patients with depression in comparison to healthy controls. For the literature search, biomedical and life science databases including PubMed (https://www.ncbi.nlm.nih.gov/pubmed) and Web of Science (https://www.webofscience.com/wos/alldb/basic-search) were utilized.

Original case-control fMRI studies examining monetary - (47), (48), (49), (50), (51), (39), (52), (53), (54), (55), (56), (41), (57), (58), (59), (60), (37) and natural - (12), (61), (62), (63), (64), (65), (66), (67), (68), (41) reward processing in depression were extracted and suitable studies from the reference lists of review articles were additionally included. The literature was screened based on titles and abstracts returned by search results of the search terms: "functional magnetic resonance imaging OR fMRI" AND "depression OR major depressive disorder OR unipolar depression OR sub-clinical depression OR at-risk of depression" in combination with either "monetary reward" OR "monetary incentive delay" OR "natural reward" OR "social reward". Studies written in English, published between 2001-2021 and reporting whole brain results in standard stereotactic space (Talairach or Montreal Neurological Institute) were included. Only peer-reviewed, original case-control studies comparing patients and healthy controls were included.

Additional exclusion criteria were: (a) studies focusing entirely on participants <18 years and >60 years of age, (b) studies focusing on other mood disorders, including bipolar disorder and postpartum depression, (c) studies reporting region-of-interest (ROI) results only, (d) studies focusing on other reward-related processes such as probabilistic reward learning, prediction error etc., (e) studies reporting findings from an identical dataset as already included studies. The systematic literature review identified 26 suitable neuroimaging

studies according to the inclusion criteria (see Figure 1), 17 examined monetary reward and

10 examined natural reward (such as exposure to positive images, positive social feedback,

or chocolate) processing in depression.

Aims and coordinate-based meta-analytic implementation

The present case-control neuroimaging meta-analysis aimed to segregate neurofunctional

reward alterations in depression, in particular to disentangle general from domain-specific

dysregulations in the domains of (a) monetary versus natural reward processing as well as in

the domains of (b) anticipation versus receipt processing of rewards. In an initial step,

general reward processing alterations were examined in an overarching meta-analysis that

pooled the data from all studies (analysis 1). Next, our primary meta-analysis determined

domain-specific alterations in hedonic processing by means of two separate meta-analyses

comparing alterations during the receipt of monetary rewards versus natural rewards in

depression (analysis 2). Finally, domain-specific alterations with respect to anticipation

versus receipt processing of monetary rewards in depression were disentangled by another

meta-analysis (analysis 3). All analyses were implemented in Seed-based d Mapping with

Permutation of Subject Images (SDM-PSI) version 6.21 (https://www.sdmproject.com/) a

novel and highly robust technique for performing neuroimaging meta-analyses (46).

The SDM-PSI approach followed the following steps: (a) Peak coordinates of between-group

differences (patients versus healthy subjects) and their effect sizes in terms of either t/z

values were extracted in accordance with the SDM inclusion criteria. Z-values representing

coordinates from whole-brain between-group differences of depressed patients and healthy

controls were transformed into t-values using the statistical converter

(https://www.sdmproject.com/utilities/?show=Coordinates). (b) For studies reporting peak

coordinates, the initial preprocessing step (Anisotropic full width half maximum (FWHM) =

20mm and voxel size = 2mm) estimated the lower and upper bounds (Hedges' g) of the most

probable effect size images (69). (c) The mean is analyzed by the maximum-likelihood estimation (MLE) and meta-analysis of non-statistically significant unreported effects (MetaNSUE) algorithm. This not only estimates the most likely effect size and the standard error, but it also creates numerous imputations based on the estimates that are within the bounds (70). (d) Next, imputed study images are recreated. (e) Finally, the permutation test evaluated the combined meta-analysis images for statistical significance.

To ensure the independence of all studies and to evade any possible bias in our meta-analyses, the coordinates of each study was put in a separate text file according to the guidelines of conducting meta-analysis using SDM-PSI. All analyses were thresholded at p ≤ 0.0025 uncorrected, k ≥ 10 voxels. The generated maps were visualized using multi-image analysis GUI (MANGO; http://ric.uthscsa.edu/mango). Furthermore, inter-study heterogeneity for each cluster was examined by the I^2 index which represents the proportion of the total variation caused by study heterogeneity (71). Generally, I^2 values have been categorized into three groups namely low (25%), moderate (50%) and high (70%) (20). Quantitative comparison analyses between monetary and natural reward outcomes were performed using Statistical Parametric Mapping (SPM12) software package (https://www.fil.ion.ucl.ac.uk/spm/).

Characterization of domain-specific hedonic dysregulations on the network, behavioral, genetic and receptor level

Given that our primary meta-analysis revealed distinct neurofunctional alterations during processing of monetary and natural rewards, we further characterized the identified signatures on the functional and neurobiological levels. First, meta-analytic network analyses were utilized to determine whether the identified regions represent nodes of separable networks. Functional decoding was conducted using the Neurosynth database

(72) (https://www.neurosynth.org/). Peak coordinates of the identified signatures were used

to generate region-specific unthresholded resting-state functional connectivity maps, meta-

analytic co-activation maps and their respective conjunction maps. The resultant resting-

state functional connectivity maps were thresholded at $r > 0.2$. The thresholded map was

transformed into z-scores. Next, conjunction maps between the thresholded resting state

functional connectivity and meta-analytic co-activation maps were generated for each ROI

signature using SPM and the resultant regions were identified with the Anatomy toolbox

(73). The functional connectivity maps represent brain regions co-activated across the

resting-state fMRI time series with the seed regions. Meta-analytic co-activation maps

symbolize co-activation of brain regions across all fMRI studies in the Neurosynth database.

The combination of functional connectivity and meta-analytic co-activation maps allows for

analyzing both task-independent and task-driven functional networks emerging from the

seed region. Additional seed to whole brain functional connectivity analyses on the voxel

level were conducted to provide a more fine-grained mapping of the common and separable

intrinsic network organization of the identified striatal subregions (74).

To this end, data from an original independent study was included. To conduct functional

connectivity analyses, data was collected from the Open Access Series of Imaging Studies

(OASIS)-3 dataset (https://central.xnat.org) including n = 100 healthy subjects. Preprocessing

was performed using Data Processing Assistant for Resting-State fMRI

(http://rfmri.org/DPARSF). Briefly, functional imaging preprocessing procedures consisted of

the following steps: (a) Deletion of the initial 5 volumes of the time-series, (b) Motion

correction using rigid body translation and rotation and exclusion of subjects with maximum

motion > 2 mm or 2°, (c) Two-step procedure of normalization into standard stereotactic

space including co-registration to anatomical images and application of the corresponding

segmentation matrix to the functional time series. 6mm radius spheres centered at the

corresponding peak coordinates (caudate: x=10, y=10, z=4 and putamen: x=22, y=8, z=-4)

served as seed regions for the seed-to-whole-brain voxel-wise analyses implemented in Data

Processing Assistant for Resting-State fMRI (http://rfmri.org/DPARSF).

Second, distinct behavioral functions of the regions were identified through meta-analytic

topic mapping using peak coordinates of the identified brain regions by means of the Brain

annotation toolbox (BAT) (75). The top twenty behavioral terms together with their p-values

values were obtained (permutation p values arranged in descending order, where $p_{perm} <$

0.05 is significant). Third, to examine separable genetic underpinnings of the identified

signatures (structurally defined by the Automated Anatomical Labeling, AAL2, atlas), we

examined genetic expressions of these regions by means of the Brain annotation toolbox

(BAT). The aim was to determine which genes had the highest densities and separable

expressions in the identified regions. Here, the top ten genes were acquired in terms of their

correlation values. Using PubMed gene (https://www.ncbi.nlm.nih.gov/gene), we

determined the identified genes' main functions.

Fourth, given the proposed role of serotonin and dopamine in both reward processing and

emotional dysregulations including depression (20,75), we determined the density of the

corresponding receptors in the identified regions. To begin, the Positron Emission

Tomography (PET) atlases of Serotonin (5-HT$_{1B}$) (https://xtra.nru.dk/FS5ht-atlas/) and

Dopamine (DAT) (https://www.nitrc.org/projects/spmtemplates) were obtained. Next, the

reference regions, i.e. cerebellum and occipital areas of 5-HT$_{1B}$ and DAT respectively, were

removed using WFU PickAtlas (v2.4) (76). Conjunction between the two no-reference region

masks and their respective receptor maps was performed in SPM12. Lastly, the final

conjunction maps were normalized using PyCharm platform (Edition 2021.2.2), based on

Python (v 3.7.2). The resultant receptor map was then obtained. Additionally, 6mm sphere-

defined ROIs of caudate and putamen masks were generated by MarsBaR ROI toolbox

(http://marsbar.sourceforge.net) and the Cognitive and Affective Neuroscience Lab (CANlab)

Core toolbox (https://canlabweb.colorado.edu/) was used to acquire the neurotransmitter density values.

Exploratory analyses of potential confounders - Linear Model analysis

Given the critical effects of anti-depressive medication on reward-related brain activity, we conducted a linear model analysis with medication status as a variable of interest to determine the potential effects of medication on neurofunctional alterations in the two primary domains (monetary and natural outcome). The analysis was run on SDM-PSI (Linear Model) where medicated studies and non-medicated studies in both datasets were coded 1 and 0, respectively. Moreover, we repeated the meta-analysis including only unmedicated studies in SDM-PSI to confirm the stability of the results, i.e. whether the results remain stable without medication or treatment. Moreover, given that previous studies have shown that age and gender distinctively regulate the brain's activity when processing monetary and natural rewards (77–80), additional meta-regressions were performed to explore the influences of other potential confounders such as age and gender.

Results

The comprehensive literature search resulted in 26 suitable original fMRI studies on reward processing in depression, with a total of n = 1 277 participants (n = 633 patients and n = 644 healthy controls). The database included patient data from monetary reward studies (n=391, mean age=31.47, SD=8.41) and natural reward studies (n=242, mean age=35.04, SD=10.26). There were no significant differences in age (p = .21, t = 2.07) and gender (p = .97, t = 2.10). Healthy controls data from monetary reward studies (n=393, mean age=30.95, SD=7.89) and natural reward studies (n=251, mean age=31.9, SD=7.98) had no significant differences in age (p = .69, t = 2.08) and gender (p = .53, t = 2.26). **Fig. 1** shows the flow diagram of the selection process. Demographic information of the included studies is displayed in **Table 1**

(see also **Supplementary Table 1** for the characteristics of included studies and

Supplementary Table 3 for the PRISMA checklist).

Neurofunctional reward alterations in depression

Analysis 1: General reward processing alterations in depression

Meta-analytic results obtained from the pooled studies of monetary and natural reward

outcome revealed decreased activation in the right striatum (peak MNI coordinates = 14, 4, -

8, z = -4.615, p < 0.0025, 294 voxels), with the cluster encompassing mainly the putamen and

extending into the caudate, and the subgenual anterior cingulate (peak MNI coordinates = 0,

2, -8, z = -3.740, p < 0.0025, 35 voxels) in depression (**Fig. 2**). There was low between-study

heterogeneity (I^2= 2.55% and 6.76%). Funnel plots for both clusters were asymmetrical and

the tests for publication bias failed to reach statistical significance (right putamen, p = 0.55

and anterior cingulate, p = 0.854) (to check examples of funnel plots generated in our meta-

analysis, see **Supplementary Figure 1**).

Analysis 2: Domain-specific (monetary versus natural) reward processing alterations in depression

During monetary reward outcome, depressed patients exhibited decreased activation in the

right caudate (peak MNI coordinates = 10, 10, 4, z = -4.260, p < 0.0025, 87 voxels) compared

to healthy controls (**Fig. 2B**). Low heterogeneity was observed for this analysis (I^2 = 11.75%),

with funnel plots showing asymmetry. There was no statistically significant publication bias

(p = 0.57).

During natural reward outcome, depressed patients exhibited decreased activation in more

dorsal parts of the striatum with the clusters being specifically located in the bilateral

putamen (right putamen, peak MNI coordinates = 22, 8, -4, z = -4.826, p < 0.0025, 223

voxels; left putamen, peak MNI coordinates = -14, 8, -6, z = -4.164, p < 0.0025, 143 voxels)

(**Fig. 2C**). There was low between-study heterogeneity (I^2 = 0.7% and 2.08%). Funnel plots illustrated asymmetry for these two clusters. Testing for potential publication bias was not statistically significant (left putamen, p = 0.89 and right putamen, p = 0.95).

Analysis 3: Stage-specific (anticipation versus outcome) reward processing alterations in depression

Examining stage-specific alterations during the anticipation and outcome of monetary reward showed decreased activation in the right thalamus (peak MNI coordinates = 18, -26, 12, z = -3.214, *p* < 0.0025, 18 voxels) and left putamen (peak MNI coordinates = -30, 4, -2, z = 3.466, *p* < 0.0025, 12 voxels) during anticipation (**Fig. 2D**) in depression relative to controls. Low heterogeneity was recorded for both clusters with asymmetric funnel plots (I^2 = 4.17% and 18.63% for right thalamus and left putamen, respectively). Publication bias tests were not statistically significant (right thalamus, p = 0.61 and left putamen, p = 0.99). For monetary reward outcome, as discussed in analysis 2, patients displayed decreased activation in the right caudate (peak MNI coordinates = 10, 10, 4, z = -4.260, *p* < 0.0025, 87 voxels). Similarly, low heterogeneity was witnessed in this step (I^2 = 11.75%), with funnel plots showing asymmetry. Publication bias failed to reach statistical significance (p = 0.57). There were no regions of increased activation between depressed patients and healthy controls across all four meta-analyses.

Domain-specific (natural vs monetary) reward dysregulations: regional quantitative, network level and behavioral characterization

In line with the main aim of the present study, we followed up the reward domain-specific neurofunctional alterations in depression. A quantitative comparison analysis was implemented to further determine common versus separable regions of altered monetary vs natural reward outcome processing in depression. The direct comparison did not reveal an overlap between neural alterations in the two domains. It confirmed that decreased

monetary outcome activity was located in the right caudate. In contrast, decreased natural

outcome alterations were located in more dorsal regions of the striatum, i.e., the bilateral

putamen (see **Fig. 3A**, monetary reward alterations, M-N depicted in green and natural

reward alterations, N-M depicted in blue).

Network-level and co-activation characterization of the identified striatal regions for

monetary (**Fig. 3B**) and natural reward dysregulations (**Fig. 3C**) revealed functional

connectivity, meta-analytic co-activation, and voxel-wise functional connectivity maps. The

caudate region was identified for monetary reward outcome dysfunctions coupled with core

regions of the mesocorticolimbic pathways, including bilateral ventral striatal regions,

ventral tegmental area (VTA) and cortical midline structures, while the putamen region was

identified for natural reward outcome dysfunctions coupled stronger with bilateral dorsal

striatal regions as well as lateral frontal and insular regions. For further details, see the

conjunction of the meta-analytic maps in **Supplementary Figure 2**.

Behavioral characterization of the identified regions (caudate and putamen) was conducted

using the Brain annotation toolbox. Examination of the top 20 behavioral terms revealed

that both regions were characterized by reward, motivation and emotional arousal

associated terms, while the caudate region was strongly related to decision making and

compulsive behavior and the putamen region was stronger involved in motor and behavioral

control (see also **Fig. 4A** and **Fig. 4B,** taking note that a stronger association is depicted as

smaller p-values, which indicates more significance).

Genetic and receptor level characterization

Genetic expression analyses revealed that S-antigen visual arrestin (SAG) was the most

expressed gene in both regions of interest (caudate and putamen). Also, the top ten genes

were generally highly similar. The summary of the top ten genes acquired for each region is

displayed in **Fig. 4C**. Comparison of the mean neurotransmitter density values of DAT and

5HT neurotransmitter in the caudate (t = 10.2335, p < 0.001, df = 244) and putamen (t= 10.1430, p < 0.001, df = 244) is displayed in **Fig. 4D** (see also **Supplementary Figure 3** for the normalized receptor maps of 5HT and DAT).

Exploratory analyses

Meta-regression analyses revealed that the results obtained from the main meta-analysis were not influenced by age or gender (**Supplementary Figure S4**). Linear model analyses further revealed that medication had no significant effect on the alterations observed during natural reward outcome (left and right putamen). However, the linear model for medication effects during monetary outcome overlapped with the meta-analytic effects of depression identified in the right caudate (**Supplementary Figure S5**). To further determine a potential confounding effect of medication, we recomputed the analyses after excluding data from patients with medication and differences between patients and healthy controls in terms of decreased activity during monetary reward outcome remained stable in the right caudate (retaining data from 7/11 studies on monetary reward outcome, *p* < 0.0025).

Discussion

Depressive mood and anhedonia represent key symptoms of major depression and have been associated with neurobiological dysregulations in reward and motivational processes. A growing body of studies have therefore utilized case-control fMRI designs to map the neurofunctional basis of reward dysregulations in individuals with depression. Several of these studies revealed altered neural activity in the mesocorticolimbic reward circuits in depression. However, it currently remains unclear whether the neurofunctional alterations vary as a function of the reward domain, i.e., natural or intrinsic rewards versus monetary or extrinsic rewards. We hereby conducted a pre-registered neuroimaging meta-analysis to systematically determine common and separable reward dysfunctions in depression.

Quantitative analyses of 26 suitable neuroimaging studies focusing on reward-related

processes in 633 patients and 644 healthy controls revealed that the right striatum and

subgenual ACC exhibit generally reduced reward reactivity in patients with depression, with

subsequent quantitative comparison analysis demonstrating that more ventral parts of the

right striatum (caudate) show reduced reactivity during monetary reward outcome while

more dorsal and bilateral parts of the striatum (putamen) show reduced reactivity during

the receipt of natural rewards. Additional meta-analytic characterization and an

independent fMRI dataset revealed that the two identified regions exhibit common as well

as separable characteristics. On the network level, the caudate interacted with ventral

striatum, ventral tegmental area (VTA), and cortical midline structures while the putamen

exhibited stronger connectivity with more dorsal striatal and lateral fronto-insular regions.

On the behavioral and molecular level, both regions were characterized by similar genetics,

high dopaminergic neurotransmitter density and an involvement in reward-related

processes. However, the putamen showed a stronger involvement in cognitive and motor

control related processes plus a higher serotonergic receptor density. Finally, an exploratory

meta-analytic examination of stage-specific reward alterations yielded decreased activation

in the right thalamus and left putamen during monetary reward anticipation in depression.

Common alterations - general hedonic reward processing alterations in depression

The current meta-analysis revealed that the subgenual ACC and right putamen extending to

the caudate, exhibited decreased reactivity in depression during general processing of

reward. Our results align with several original studies that have demonstrated altered brain

activity in these regions during reward processing in depression. The striatum has been

associated with both, monetary and natural reward processing in healthy individuals (51,81–

83) and dysfunctional reward processing in this region has been reported in several mental disorders, including depression, schizophrenia and addiction (51,84–87).

The subgenual ACC has been consistently involved in reward processing (88), as well as regulatory control in cognitive as well as emotional domains (89–91), and may serve an integrative function to evaluate rewards and guide the most appropriate choices between rewards (92). Recent neuropathological models suggest a key role of the subgenual ACC in mood disorders including depression (93,94). In fact, it has been described as a region that lies between the cognitive and emotional networks of the brain, which is critical in the conveyance of emotional information from the limbic system to higher cognitive systems of the brain (95,96). A previous meta-analysis has also shown decreased volume of the subgenual ACC in mood disorders (97). Reduced activation in this region is distinctively associated with depression as it cuts across behavioral, emotional and cognitive aspects.

Distinct neurofunctional alterations during hedonic processing of natural and monetary rewards

While our findings on the neurofunctional basis of general reward processing alterations in depression align with previous meta-analyses, the key aim of the present meta-analysis was to further segregate neurobiological dysregulations during hedonic processing of different rewards. We hypothesized a common ventral striatal deficit across natural and monetary rewards in depression, however we observed non-overlapping alterations such that ventral caudal regions displayed reduced reactivity during monetary rewards while more dorsal striatal regions showed decreased reactivity towards natural rewards. Both regions have been involved in reward processing across domains (98) and general reward processing alterations in depression (30,51,99,100). However, the striatum is organized along a ventral to dorsal axis with respect to different behavioral domains and network-level interactions. The ventral parts of the caudate have been primarily involved in reward, reinforcement and

incentive salience processing while the more dorsal parts of the striatum mediate sensorimotor processes and some executive functions such as inhibitory control (101–103). In line with this functional specialization of the striatum subregions, the identified caudate region exhibited strong connections with ventral parts of the striatum, the VTA and cortical midline structures strongly involved in reward-related processing while the putamen region coupled with dorsal striatal regions as well as lateral frontal and insular regions involved in interoceptive processes, pain empathy and regulatory control (101,104,105). The conceptual and network level differentiation was additionally mirrored on the behavioral level such that the meta-analytic behavioral characterization indicated that both regions contribute to reward processes, but that the more ventral region was characterized by an additional involvement in compulsivity, learning and motivational processes while the more dorsal region additionally showed an involvement in motor and behavioral control processes. Together, the neurofunctional, network and behavioral characterization may indicate that, (1) different neurobiological mechanisms underlie depression-related reward dysregulations in the hedonic processing of monetary and natural rewards, and that, (2) the generalization and ecological validity of the prevailing reward paradigms incorporating small amounts of monetary incentives to situations in everyday life that engage primarily natural rewards is limited.

On the molecular level, both regions exhibited a similar genetic profile such that S-antigen visual arrestin (SAG), SLC35D3 and MME showed high expressions. SAG has been associated with the G-protein-coupled receptor reactivity and thus sensitivity to several neurohormones or neurotransmitters. The SLC35D3 gene has been associated with mid-brain and striatal dopamine signaling (106) while MME has been associated with the inactivation of several neuropeptides such as oxytocin (107) (for a detailed characterization, see also Supplementary Table 2). Dopamine and oxytocin have been associated with network-level modulation of striatal circuits and reward processing across natural and

monetary reward domains (74,108,109) and, in interaction with the respective stimuli may shape context-dependent reward dysregulations in depression. In terms of neurotransmitter expression, both regions exhibited a high expression of DA while the dorsal striatal region identified for natural reward processing deficits exhibited a considerably higher 5-HT expression. Both neurotransmitter systems have been strongly associated with the pathophysiology of depression (110,111), but see also, (112). Dopamine has been described as the common currency by which animals pursue goals and therefore alterations in the dopaminergic system could underlie symptoms including anhedonia (113). Serotonin has been found to play a role in emotional and reward processes, in particular (dysregulated) natural reward processing (114,115). Within the context of the present results differences in the serotonin system may reflect a stronger engagement of serotonergic signaling in natural reward processing deficits in depression.

Stage-specific neurofunctional reward processing alterations in depression – anticipation vs outcome

While the outcome of rewards is commonly considered as a hedonic process, the anticipation of rewarding outcomes is commonly considered as a motivational process. Dysregulated motivational and approach tendencies towards pleasant events have long been associated with depression (116). However, previous findings in depression have remained largely inconsistent with respect to the anticipation of rewards. For instance, depressed patients have demonstrated elevated activation during monetary reward anticipation in the anterior cingulate (37,117), while other studies reported a relatively decreased anticipatory signal in striatal regions including the bilateral caudate, bilateral putamen, ventral striatum and globus pallidus (30,36,38). Partly resembling these previous findings, we observed decreased anticipatory activity during monetary rewards in the thalamus and putamen of depressed patients. The thalamus represents an early detection

node for salient stimuli including rewards and its connections with the striatum are involved in orchestrating behavioral and attentional reactions in response to different environmental cues (118). Decreased striatal activity during the anticipation of reward has been reported in depression (30,51) as well as other mental disorders characterized by motivational dysregulations, such as schizophrenia and addiction (119,120) and thus may represent a transdiagnostic marker for deficient motivation and engagement.

The present findings have to be interpreted in the context of some limitations inherent to the meta-analytic approach and the available original studies. First, the exploratory linear model analysis revealed potential confounding effects of treatment on the identified region during monetary outcome. Previous studies have demonstrated effects of anti-depressive treatment on brain structure and function (121). Although natural reward processing was not affected and both results remained stable after excluding studies with medication, future original studies are required to explore effects of treatment on reward-related neural dysregulations in depression. Second, some studies mixed unmedicated and medicated patient groups when performing experiments in a single study and as pointed out by (117), this increases the chances of type II errors. Future studies may recruit patients who have a uniform medication status in order to promote more reliable and cohesive results in original studies and meta-analyses. Third, the number of studies for the anticipation of natural rewards was insufficient to explore differences between natural and monetary reward anticipation. Fourth, the number of original studies was moderate, for example, only 10 original studies were eligible for inclusion of natural reward outcome category. Finally, while SDM-PSI allows to incorporate original studies with non-reported effect sizes due to non-significant effects, some of the included studies did not explicitly report the lack of significant results for all contrast directions which is common practice in fMRI studies but may affect meta-analytic results (70,122).

In summary, the present meta-analysis supports the notion of general reward processing deficits in depression. However, the underlying neurofunctional dysregulations vary according to reward type (e.g., monetary and natural rewards) and sub-processes (e.g., anticipation and outcome). Previous work has largely focused on extrinsic (monetary) rewards to map disrupted reward processing in mental disorders such as depression. However, this may have overshadowed deficits in the processing of natural rewards. Natural rewards prevail in everyday life and shape our social interactions and thus dysregulations in this domain may lead to marked impairments in everyday life. The present meta-analysis has revealed distinguishable neurofunctional reward processing deficits in depression, suggesting that natural and monetary reward processing alterations may be rooted in distinct neurobiological dysfunctions.

Funding

The present study was supported by the National Natural Science Foundation of China (NSFC 82271583; 32250610208) and the China Brain Project (MOST2030, Grant No. 2022ZD0208500).

Conflict of interest

The authors declare no conflict of interest.

Data availability

Studies included in the meta-analyses have been cited in the manuscript. Coordinates will be made available upon request.

References

1. Halahakoon DC, Kieslich K, O'Driscoll C, Nair A, Lewis G, Roiser JP. Reward-Processing Behavior in Depressed Participants Relative to Healthy Volunteers: A Systematic Review and Meta-analysis. Vol. 77, JAMA Psychiatry. American Medical Association; 2020. p. 1286–95.

2. Bedwell JS, Gooding DC, Chan CC, Trachik BJ. Anhedonia in the age of RDoC. Vol. 160, Schizophrenia Research. Elsevier B.V.; 2014. p. 226–7.

3. Insel T, Cuthbert B, Garvey M, Heinssen R, Pine DS, Quinn K, et al. Research Domain Criteria (RDoC): Toward a New Classification Framework for Research on Mental Disorders. Vol. 167, The American Journal of Psychiatry. 2010. p. 748–51.

4. Husain M, Roiser JP. Neuroscience of apathy and anhedonia: A transdiagnostic approach. Vol. 19, Nature Reviews Neuroscience. Nature Publishing Group; 2018. p. 470–84.

5. Friedrich MJ. Depression Is the Leading Cause of Disability Around the World. JAMA. 2017;317(15):1517–1517.

6. Kendler KS. The phenomenology of major depression and the representativeness and nature of DSM criteria. Vol. 173, American Journal of Psychiatry. American Psychiatric Association; 2016. p. 771–80.

7. Admon R, Pizzagalli DA. Dysfunctional reward processing in depression. Vol. 4, Current Opinion in Psychology. Elsevier; 2015. p. 114–8.

8. Alloy LB, Olino T, Freed RD, Nusslock R. Role of Reward Sensitivity and Processing in Major Depressive and Bipolar Spectrum Disorders. Behav Ther. 2016 Sep 1;47(5):600–21.

9. Treadway MT, Zald DH. Reconsidering anhedonia in depression: Lessons from translational neuroscience. Vol. 35, Neuroscience and Biobehavioral Reviews. 2011. p. 537–55.

10. Robbins TW. Illuminating anhedonia: Optogenetics and fMRI reveal the brain circuitry of anhedonia. Vol. 351, Science. American Association for the Advancement of Science; 2016. p. 24–5.

11. Becker B. Ketamine as a Potential Transdiagnostic Treatment for Anhedonia? Vol. 7, Biological Psychiatry: Cognitive Neuroscience and Neuroimaging. Elsevier Inc.; 2022. p. 241–2.

12. Oh H, Lee J, Patriquin MA, Oldham J, Salas R. Reward Processing in Psychiatric Inpatients With Depression. Biol Psychiatry Cogn Neurosci Neuroimaging. 2022;

13. Schultz W. Neuronal Reward and Decision Signals: From Theories to Data. Physiol Rev [Internet]. 2015 Jul 1 [cited 2022 Oct 21];95(3):853. Available from: /pmc/articles/PMC4491543/

14. Perry DC, Kramer JH. Reward processing in neurodegenerative disease. Neurocase. 2015 Jan 2;21(1):120–33.

15. Berridge KC, Robinson TE. Parsing reward. Vol. 26, Trends in Neurosciences. Elsevier Ltd; 2003. p. 507–13.

16. Oldham S, Murawski C, Fornito A, Youssef G, Yücel M, Lorenzetti V. The anticipation and outcome phases of reward and loss processing: A neuroimaging meta-analysis of the monetary incentive delay task. Hum Brain Mapp. 2018 Aug 1;39(8):3398–418.

17. Schultz W. Dopamine reward prediction error coding. Dialogues Clin Neurosci. 2016 Mar 1;18(1):23–32.

18. Rizvi SJ, Pizzagalli DA, Sproule BA, Kennedy SH. Assessing anhedonia in depression: Potentials and pitfalls. Vol. 65, Neuroscience and Biobehavioral Reviews. Elsevier Ltd; 2016. p. 21–35.

19. Yang X, Liu X, Zeng Y, Wu R, Zhao W, Xin F, et al. Secondary rewards acquire enhanced incentive motivation via increasing anticipatory activity of the lateral orbitofrontal cortex. Brain Struct Funct. 2021 Sep 1;226(7):2339–55.

20. Martins D, Rademacher L, Gabay AS, Taylor R, Richey JA, Smith D v., et al. Mapping social reward and punishment processing in the human brain: A voxel-based meta-analysis of neuroimaging findings using the social incentive delay task. Vol. 122, Neuroscience and Biobehavioral Reviews. Elsevier Ltd; 2021. p. 1–17.

21. Jauhar S, Fortea L, Solanes A, Albajes-Eizagirre A, McKenna PJ, Radua J. Brain activations associated with anticipation and delivery of monetary reward: A systematic review and meta-analysis of fMRI studies. Vol. 16, PLoS ONE. Public Library of Science; 2021.

22. Wilson RP, Colizzi M, Bossong MG, Allen P, Kempton M, Abe N, et al. The Neural Substrate of Reward Anticipation in Health: A Meta-Analysis of fMRI Findings in the Monetary Incentive Delay Task. Neuropsychol Rev. 2018 Dec 15;28(4):496–506.

23. Mas-Herrero E, Maini L, Sescousse G, Zatorre RJ. Common and distinct neural correlates of music and food-induced pleasure: A coordinate-based meta-analysis of neuroimaging studies. Vol. 123, Neuroscience and Biobehavioral Reviews. Elsevier Ltd; 2021. p. 61–71.

24. Sescousse G, Caldú X, Segura B, Dreher JC. Processing of primary and secondary rewards: A quantitative meta-analysis and review of human functional neuroimaging studies. Vol. 37, Neuroscience and Biobehavioral Reviews. 2013. p. 681–96.

25. Fehr E, Camerer CF. Social neuroeconomics: the neural circuitry of social preferences. Vol. 11, Trends in Cognitive Sciences. 2007. p. 419–27.

26. Izuma K. Social Reward. In: Brain Mapping: An Encyclopedic Reference. Elsevier Inc.; 2015. p. 21–3.

27. Ruff CC, Fehr E. The neurobiology of rewards and values in social decision making. Vol. 15, Nature Reviews Neuroscience. Nature Publishing Group; 2014. p. 549–62.

28. Knutson B, Bossaerts P. Neural antecedents of financial decisions. Vol. 27, Journal of Neuroscience. 2007. p. 8174–7.

29. Katz BA, Matanky K, Aviram G, Yovel I. Reinforcement sensitivity, depression and anxiety: A meta-analysis and meta-analytic structural equation model. Vol. 77, Clinical Psychology Review. Elsevier Inc.; 2020.

30. Keren H, O'Callaghan G, Vidal-Ribas P, Buzzell GA, Brotman MA, Leibenluft E, et al. Reward processing in depression: A conceptual and meta-analytic review across fMRI and EEG studies. American Journal of Psychiatry. 2018 Nov 1;175(11):1111–20.

31. Yang X, Su Y, Yang F, Song Y, Yan J, Luo Y, et al. Neurofunctional mapping of reward anticipation and outcome for major depressive disorder: a voxel-based meta-analysis. Psychol Med. 2022 Sep 1;1–14.

32. Zhang WN, Chang SH, Guo LY, Zhang KL, Wang J. The neural correlates of reward-related processing in major depressive disorder: A meta-analysis of functional magnetic resonance imaging studies. J Affect Disord. 2013 Nov;151(2):531–9.

33. Geugies H, Mocking RJT, Figueroa CA, Groot PFC, Marsman JBC, Servaas MN, et al. Impaired reward-related learning signals in remitted unmedicated patients with recurrent depression. Brain. 2019 Aug 1;142(8):2510–22.

34.   Ng TH, Alloy LB, Smith D v. Meta-analysis of reward processing in major depressive disorder reveals distinct abnormalities within the reward circuit. Transl Psychiatry. 2019 Dec 1;9(1).

35.   Russo SJ, Nestler EJ. The brain reward circuitry in mood disorders. Vol. 14, Nature Reviews Neuroscience. 2013. p. 609–25.

36.   Stringaris A, Belil PVR, Artiges E, Lemaitre H, Gollier-Briant F, Wolke S, et al. The brain s response to reward anticipation and depression in adolescence: Dimensionality, specificity, and longitudinal predictions in a community-based sample. American Journal of Psychiatry. 2015 Dec 1;172(12):1215–23.

37.   Knutson B, Bhanji JP, Cooney RE, Atlas LY, Gotlib IH. Neural Responses to Monetary Incentives in Major Depression. Biol Psychiatry. 2008 Apr 1;63(7):686–92.

38.   Takamura M, Okamoto Y, Okada G, Toki S, Yamamoto T, Ichikawa N, et al. Patients with major depressive disorder exhibit reduced reward size coding in the striatum. Prog Neuropsychopharmacol Biol Psychiatry. 2017 Oct 3;79:317–23.

39.   Dichter GS, Kozink R v., McClernon FJ, Smoski MJ. Remitted major depression is characterized by reward network hyperactivation during reward anticipation and hypoactivation during reward outcomes. J Affect Disord. 2012 Feb;136(3):1126–34.

40.   Jenkins LM, Skerrett KA, DelDonno SR, Patrón VG, Meyers KK, Peltier S, et al. Individuals with more severe depression fail to sustain nucleus accumbens activity to preferred music over time. Psychiatry Res Neuroimaging. 2018 May 30;275:21–7.

41.   Smoski MJ, Rittenberg A, Dichter GS. Major depressive disorder is characterized by greater reward network activation to monetary than pleasant image rewards. Psychiatry Res Neuroimaging. 2011 Dec 30;194(3):263–70.

42.   Ait Oumeziane B, Jones O, Foti D. Neural Sensitivity to Social and Monetary Reward in

        Depression: Clarifying General and Domain-Specific Deficits. Front Behav Neurosci.

        2019 Oct 9;13.

43.   Nelson BD, Jarcho JM. Neural response to monetary and social feedback

        demonstrates differential associations with depression and social anxiety. Soc Cogn

        Affect Neurosci. 2021 Oct 1;16(10):1048–56.

44.   Müller VI, Cieslik EC, Laird AR, Fox PT, Radua J, Mataix-Cols D, et al. Ten simple rules

        for neuroimaging meta-analysis. Vol. 84, Neuroscience and Biobehavioral Reviews.

        Elsevier Ltd; 2018. p. 151–61.

45.   Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The

        PRISMA 2020 statement: An updated guideline for reporting systematic reviews. Vol.

        372, The BMJ. BMJ Publishing Group; 2021.

46.   Albajes-Eizagirre A, Solanes A, Fullana MA, Ioannidis JPA, Fusar-Poli P, Torrent C, et al.

        Meta-analysis of voxel-based neuroimaging studies using seed-based d mapping with

        permutation of subject images (Sdm-psi). Journal of Visualized Experiments. 2019

        Nov 1;2019(153).

47.   Fischer AS, Ellwood-Lowe ME, Colich NL, Cichocki A, Ho TC, Gotlib IH. Reward-circuit

        biomarkers of risk and resilience in adolescent depression. J Affect Disord. 2019 Mar

        1;246:902–9.

48.   Mori A, Okamoto Y, Okada G, Takagaki K, Jinnin R, Takamura M, et al. Behavioral

        activation can normalize neural hypoactivation in subthreshold depression during a

        monetary incentive delay task. J Affect Disord. 2016 Jan 1;189:254–62.

49.   Ubl B, Kuehner C, Kirsch P, Ruttorf M, Diener C, Flor H. Altered neural reward and loss

        processing and prediction error signalling in depression. Soc Cogn Affect Neurosci.

        2014 Oct 3;10(8):1102–12.

50.   Martin-Soelch C, Guillod M, Gaillard C, Recabarren RE, Federspiel A, Mueller-Pfeiffer

        C, et al. Increased Reward-Related Activation in the Ventral Striatum During Stress

        Exposure Associated With Positive Affect in the Daily Life of Young Adults With a

        Family History of Depression. Preliminary Findings. Front Psychiatry. 2021 Jan 18;11.

51.   Pizzagalli DA, Holmes AJ, Dillon DG, Goetz EL, Birk JL, Bogdan R, et al. Reduced

        Caudate and Nucleus Accumbens Response to Rewards in Unmedicated Subjects with

        Major Depressive Disorder. Am J Psychiatry. 2009 Jun;166(6):702.

52.   Hall GBC, Milne AMB, MacQueen GM. An fMRI study of reward circuitry in patients

        with minimal or extensive history of major depression. Eur Arch Psychiatry Clin

        Neurosci. 2014;264(3):187–98.

53.   Arrondo G, Segarra N, Metastasio A, Ziauddeen H, Spencer J, Reinders NR, et al.

        Reduction in ventral striatal activity when anticipating a reward in depression and

        schizophrenia: A replicated cross-diagnostic finding. Front Psychol. 2015 Aug 26;6.

54.   Chase HW, Nusslock R, Almeida JR, Forbes EE, Labarbara EJ, Phillips ML. Dissociable

        patterns of abnormal frontal cortical activation during anticipation of an uncertain

        reward or loss in bipolar versus major depression. Bipolar Disord. 2013;15(8):839–54.

55.   Burrows K, Stewart JL, Kuplicki R, Figueroa-Hall L, Spechler PA, Zheng H, et al.

        Elevated peripheral inflammation is associated with attenuated striatal reward

        anticipation in major depressive disorder. Brain Behav Immun. 2021 Mar 1;93:214–

        25.

56.   Smoski MJ, Felder J, Bizzell J, Green SR, Ernst M, Lynch TR, et al. fMRI of alterations in reward selection, anticipation, and feedback in major depressive disorder. J Affect Disord. 2009 Nov;118(1–3):69–78.

57.   Segarra N, Metastasio A, Ziauddeen H, Spencer J, Reinders NR, Dudas RB, et al. Abnormal Frontostriatal Activity During Unexpected Reward Receipt in Depression and Schizophrenia: Relationship to Anhedonia. Neuropsychopharmacology. 2016 Jul 1;41(8):2001–10.

58.   Admon R, Nickerson LD, Dillon DG, Holmes AJ, Bogdan R, Kumar P, et al. Dissociable cortico-striatal connectivity abnormalities in major depression in response to monetary gains and penalties. Psychol Med. 2015 Jan 12;45(1):121–31.

59.   DelDonno SR, Mickey BJ, Pruitt PJ, Stange JP, Hsu DT, Weldon AL, et al. Influence of childhood adversity, approach motivation traits, and depression on individual differences in brain activation during reward anticipation. Biol Psychol. 2019 Sep 1;146.

60.   Gorka SM, Huggins AA, Fitzgerald DA, Nelson BD, Phan KL, Shankman SA. Neural response to reward anticipation in those with depression with and without panic disorder. J Affect Disord. 2014 Aug 1;164:50–6.

61.   Epstein J, Pan H, Kocsis JH, Yang Y, Butler T, Chusid J, et al. Lack of Ventral Striatal Response to Positive Stimuli in Depressed Versus Normal Subjects. Vol. 163, Am J Psychiatry. 2006.

62.   Gradin VB, Pérez A, MacFarlane JA, Cavin I, Waiter G, Engelmann J, et al. Abnormal brain responses to social fairness in depression: An fMRI study using the Ultimatum Game. Psychol Med. 2015 Apr 18;45(6):1241–51.

63.   Keedwell PA, Andrew C, Williams SCR, Brammer MJ, Phillips ML. A double dissociation of ventromedial prefrontal cortical responses to sad and happy stimuli in depressed and healthy individuals. Biol Psychiatry. 2005 Sep 15;58(6):495–503.

64.   McCabe C, Cowen PJ, Harmer CJ. Neural representation of reward in recovered depressed patients. Psychopharmacology (Berl). 2009 Sep;205(4):667–77.

65.   Kumari V, Mitterschiffthaler MT, Teasdale JD, Malhi GS, Brown RG, Giampietro V, et al. Neural Abnormalities during Cognitive Generation of Affect in Treatment-Resistant Depression. Biol Psychiatry. 2003;54(8):777–91.

66.   Canli T, Sivers H, Thomason ME, Whit¢eld-Gabrieli S, Gabrieli JDE, Gotlib IH. Brain activation to emotional words in depressed vs healthy subjects. Neuroreport. 2004;15(17):2585–8.

67.   Fournier JC, Keener MT, Mullin BC, Hafeman DM, LaBarbara EJ, Stiffler RS, et al. Heterogeneity of amygdala response in major depressive disorder: The impact of lifetime subthreshold mania. Psychol Med. 2013 Feb;43(2):293–302.

68.   Gotlib IH, Sivers H, Gabrieli JDE, Whit¢eld-Gabrieli S, Goldin P, Minor KL, et al. Subgenual anterior cingulate activation to valenced emotional stimuli in major depression. Neuroreport. 2005;16(16):1731–4.

69.   Radua J, Rubia K, Canales-Rodríguez EJ, Pomarol-Clotet E, Fusar-Poli P, Mataix-Cols D. Anisotropic kernels for coordinate-based meta-analyses of neuroimaging studies. Front Psychiatry. 2014;5(FEB).

70.   Albajes-Eizagirre A, Solanes A, Radua J. Meta-analysis of non-statistically significant unreported effects. Stat Methods Med Res. 2019 Dec 1;28(12):3741–54.

71.   Higgins JPT, Thompson SG. Quantifying heterogeneity in a meta-analysis. Stat Med. 2002 Jun 15;21(11):1539–58.

72. Yarkoni T, Poldrack RA, Nichols TE, van Essen DC, Wager TD. Large-scale automated synthesis of human functional neuroimaging data. Nat Methods. 2011 Aug;8(8):665–70.

73. Eickhoff SB, Stephan KE, Mohlberg H, Grefkes C, Fink GR, Amunts K, et al. A new SPM toolbox for combining probabilistic cytoarchitectonic maps and functional imaging data. Neuroimage. 2005 May 1;25(4):1325–35.

74. Zhao Z, Ma X, Geng Y, Zhao W, Zhou F, Wang J, et al. Oxytocin differentially modulates specific dorsal and ventral striatal functional connections with frontal and cerebellar regions. Neuroimage. 2019 Jan 1;184:781–9.

75. Liu Z, Rolls ET, Liu Z, Zhang K, Yang M, Du J, et al. Brain annotation toolbox: Exploring the functional and genetic associations of neuroimaging results. Bioinformatics. 2019 Oct 1;35(19):3771–8.

76. Maldjian JA, Laurienti PJ, Kraft RA, Burdette JH. An automated method for neuroanatomic and cytoarchitectonic atlas-based interrogation of fMRI data sets. Neuroimage. 2003 Jul 1;19(3):1233–9.

77. Rademacher L, Salama A, Gründer G, Spreckelmeyer KN. Differential patterns of nucleus accumbens activation during anticipation of monetary and social reward in young and older adults. Soc Cogn Affect Neurosci. 2013;9(6):825–31.

78. Spreckelmeyer KN, Krach S, Kohls G, Rademacher L, Irmak A, Konrad K, et al. Anticipation of monetary and social reward differently activates mesolimbic brain structures in men and women. Soc Cogn Affect Neurosci. 2009;4(2):158–65.

79. Wang D, Liu T, Shi J. Development of Monetary and Social Reward Processes. Sci Rep. 2017 Dec 1;7(1).

80. Greimel E, Bakos S, Landes I, Töllner T, Bartling J, Kohls G, et al. Sex differences in the neural underpinnings of social and monetary incentive processing during adolescence. Cogn Affect Behav Neurosci. 2018 Apr 1;18(2):296–312.

81. Nieuwenhuis S, Heslenfeld DJ, von Geusau NJA, Mars RB, Holroyd CB, Yeung N. Activity in human reward-sensitive brain areas is strongly context dependent. Neuroimage. 2005;25(4):1302–9.

82. Breiter HC, Aharon I, Kahneman D, Dale A, Shizgal P. Functional Imaging of Neural Responses to Expectancy and Experience of Monetary Gains and Losses. Neuron. 2001 May 1;30(2):619–39.

83. Knutson B, Adams CM, Fong GW, Hommer D, Varner J, Kerich M, et al. Anticipation of Increasing Monetary Reward Selectively Recruits Nucleus Accumbens. Vol. 21, The Journal of Neuroscience. 2001.

84. Surguladze S, Brammer MJ, Keedwell P, Giampietro V, Young AW, Travis MJ, et al. A differential pattern of neural response toward sad versus happy facial expressions in major depressive disorder. Biol Psychiatry. 2005 Feb 1;57(3):201–9.

85. Zimmermann K, Kendrick KM, Scheele D, Dau W, Banger M, Maier W, et al. Altered striatal reward processing in abstinent dependent cannabis users: Social context matters. European Neuropsychopharmacology. 2019 Mar 1;29(3):356–64.

86. Klugah-Brown B, Di X, Zweerings J, Mathiak K, Becker B, Biswal B. Common and separable neural alterations in substance use disorders: A coordinate-based meta-analyses of functional neuroimaging studies in humans. Hum Brain Mapp. 2020 Nov 1;41(16):4459–77.

87.   Chase HW, Loriemi P, Wensing T, Eickhoff SB, Nickl-Jockschat T. Meta-analytic

        evidence for altered mesolimbic responses to reward in schizophrenia. Hum Brain

        Mapp. 2018 Jul 1;39(7):2917–28.

88.   Rogers RD, Ramnani N, Mackay C, Wilson JL, Jezzard P, Carter CS, et al. Distinct

        portions of anterior cingulate cortex and medial prefrontal cortex are activated by

        reward processing in separable phases of decision-making cognition. Biol Psychiatry.

        2004 Mar 15;55(6):594–602.

89.   Drevets WC, Savitz J, Trimble M. The subgenual anterior cingulate cortex in mood

        disorders. CNS Spectr. 2008;13(8):663–81.

90.   Rushworth MFS, Walton ME, Kennerley SW, Bannerman DM. Action sets and

        decisions in the medial frontal cortex. Vol. 8, Trends in Cognitive Sciences. 2004. p.

        410–7.

91.   Critchley HD. Neural mechanisms of autonomic, affective, and cognitive integration.

        In: Journal of Comparative Neurology. 2005. p. 154–66.

92.   Holroyd CB, Yeung N. Motivation of extended behaviors by anterior cingulate cortex.

        Vol. 16, Trends in Cognitive Sciences. 2012. p. 122–8.

93.   Botteron KN, Raichle ME, Drevets WC, Heath AC, Todd RD. Volumetric Reduction in

        Left Subgenual Prefrontal Cortex in Early Onset Depression. Vol. 51, Biol Psychiatry.

        2002.

94.   Drevets WC, Price JL, Furey ML. Brain structural and functional abnormalities in mood

        disorders: Implications for neurocircuitry models of depression. Vol. 213, Brain

        Structure and Function. 2008. p. 93–118.

95.   Disner SG, Beevers CG, Haigh EAP, Beck AT. Neural mechanisms of the cognitive

        model of depression. Vol. 12, Nature Reviews Neuroscience. 2011. p. 467–77.

96.   Greicius MD, Flores BH, Menon V, Glover GH, Solvason HB, Kenna H, et al. Resting-State Functional Connectivity in Major Depression: Abnormally Increased Contributions from Subgenual Cingulate Cortex and Thalamus. Biol Psychiatry. 2007 Sep 1;62(5):429–37.

97.   Hajek T, Kozeny J, Kopecek M, Alda M, Höschl C, - A, et al. Reduced subgenual cingulate volumes in mood disorders: a meta-analysis. Vol. 33, J Psychiatry Neurosci. 2008.

98.   Arsalidou M, Vijayarajah S, Sharaev M. Basal ganglia lateralization in different types of reward. Brain Imaging Behav. 2020 Dec 1;14(6):2618–46.

99.   Connolly ME, Gollan JK, Cobia D, Wang X. Reduced striatal activation in females with major depression during the processing of affective stimuli. J Psychiatr Res. 2015 Sep 1;68:384–91.

100.   Foell J, Klawohn J, Bruchnak A, Brush CJ, Patrick CJ, Hajcak G. Ventral striatal activation during reward differs between major depression with and without impaired mood reactivity. Psychiatry Res Neuroimaging. 2021 Jul 30;313.

101.   Zhuang Q, Xu L, Zhou F, Yao S, Zheng X, Zhou X, et al. Segregating domain-general from emotional context-specific inhibitory control systems - ventral striatum and orbitofrontal cortex serve as emotion-cognition integration hubs. Neuroimage. 2021 Sep 1;238.

102.   Haber SN. Corticostriatal circuitry. Dialogues Clin Neurosci. 2016 Mar 31;18(1):7–21.

103.   Robbins TW, Gillan CM, Smith DG, de Wit S, Ersche KD. Neurocognitive endophenotypes of impulsivity and compulsivity: Towards dimensional psychiatry. Vol. 16, Trends in Cognitive Sciences. 2012. p. 81–91.

104. Craig AD. How do you feel — now? The anterior insula and human awareness. Nat Rev Neurosci. 2009;10:59–70.

105. Zhou F, Zimmermann K, Xin F, Scheele D, Dau W, Banger M, et al. Shifted balance of dorsal versus ventral striatal communication with frontal reward and regulatory regions in cannabis-dependent males. Hum Brain Mapp. 2018 Dec 1;39(12):5062–73.

106. Zhang Z, Hao CJ, Li CG, Zang DJ, Zhao J, Li XN, et al. Mutation of SLC35D3 Causes Metabolic Syndrome by Impairing Dopamine Signaling in Striatal D1 Neurons. PLoS Genet. 2014 Feb;10(2).

107. Sterchi EE, Naim HY, Lentze MJ, Hauri HP, Fransens JAM. N-Benzoyk-tyrosyl-p-aminobenzoic Acid Hydrolase: A Metalloendopeptidase of the Human intestinal Microvillus Membrane Which Degrades Biologically Active Peptides. Vol. 265, ARCHIVES OF BIOCHEMISTRY AND BIOPHYSICS. 1988.

108. Scheele D, Wille A, Kendrick KM, Stoffel-Wagner B, Becker B, Güntürkün O, et al. Oxytocin enhances brain reward system responses in men viewing the face of their female partner. Proc Natl Acad Sci U S A. 2013 Dec 10;110(50):20308–13.

109. Grimm O, Kopfer V, Küpper-Tetzel L, Deppert V, Kuhn M, de Greck M, et al. Amisulpride and l-DOPA modulate subcortical brain nuclei connectivity in resting-state pharmacologic magnetic resonance imaging. Hum Brain Mapp. 2020 May 1;41(7):1806–18.

110. Schultz W, Tremblay L, Hollerman JR. Reward processing in primate orbitofrontal cortex and basal ganglia. Cerebral Cortex. 2000;10(3):272–84.

111. Nemeroff CB, Owens MJ. The role of serotonin in the pathophysiology of depression: As important as ever. Vol. 55, Clinical Chemistry. 2009. p. 1578–9.

112. Moncrieff J, Cooper RE, Stockmann T, Amendola S, Hengartner MP, Horowitz MA. The serotonin theory of depression: a systematic umbrella review of the evidence. Molecular Psychiatry. Springer Nature; 2022.

113. Belujon P, Grace AA. Dopamine system dysregulation in major depressive disorders. Vol. 20, International Journal of Neuropsychopharmacology. Oxford University Press; 2017. p. 1036–46.

114. Watson KK, Ghodasra JH, Platt ML. Serotonin transporter genotype modulates social reward and punishment in rhesus macaques. PLoS One. 2009 Jan 14;4(1).

115. Dölen G, Darvishzadeh A, Huang KW, Malenka RC. Social reward requires coordinated activity of nucleus accumbens oxytocin and serotonin. Nature. 2013;501(7466):179–84.

116. Davidson RJ. Affective Style and Affective Disorders: Perspectives from Affective Neuroscience. Cogn Emot. 1998;12(3):307–30.

117. Dichter GS, Felder JN, Petty C, Bizzell J, Ernst M, Smoski MJ. The Effects of Psychotherapy on Neural Responses to Rewards in Major Depression. Biol Psychiatry. 2009 Nov 1;66(9):886–97.

118. Cho YT, Fromm S, Guyer AE, Detloff A, Pine DS, Fudge JL, et al. Nucleus accumbens, thalamus and insula connectivity during incentive anticipation in typical adults and adolescents. Neuroimage. 2013 Feb 1;66:508–21.

119. Leroy A, Amad A, D'Hondt F, Pins D, Jaafari N, Thomas P, et al. Reward anticipation in schizophrenia: A coordinate-based meta-analysis. Schizophr Res. 2020 Apr 1;218:2–6.

120. Luijten M, Schellekens AF, Kühn S, MacHielse MWJ, Sescousse G. Disruption of reward processing in addiction: An image-based meta-analysis of functional magnetic resonance imaging studies. JAMA Psychiatry. 2017 Apr 1;74(4):387–98.

121. Delaveau P, Jabourian M, Lemogne C, Guionnet S, Bergouignan L, Fossati P. Brain

      effects of antidepressants in major depression: A meta-analysis of emotional

      processing studies. J Affect Disord. 2011 Apr;130(1–2):66–74.

122. Albajes-Eizagirre A, Solanes A, Vieta E, Radua J. Voxel-based meta-analysis via

      permutation of subject images (PSI): Theory and implementation for SDM.

      Neuroimage. 2019 Feb 1;186:174–84.

Fig. 1 Flow diagram of the systematic literature search and identification of suitable original studies.

Fig. 2 Results of the primary meta-analyses on dysfunctional reward processing in depression. **A** Display of regions that showed generally reduced reactivity during reward processing in depression as compared to healthy controls. Subsequent meta-analyses revealed regions that are specifically dysfunctional during the hedonic processing of **B** monetary and **C** natural rewards. **D** shows region with altered activation in depression during the anticipation of monetary rewards (p < 0.0025 uncorrected, k ≥ 10 voxels)

Fig. 3 Regional and network level characterization of the monetary and natural reward outcome alterations in depression. **A** Quantitative comparison analysis analyses of monetary versus natural outcome alterations in depression. Meta-analytic and voxel-wise mapping of the identified regions using Neurosynth and independent resting state fMRI data. **B** Brain networks of the identified caudate region of monetary reward outcome alterations. **C** Brain networks of the identified putamen region of natural reward outcome alterations. MAFC, meta-analytic functional connectivity; MACo, meta-analytic co-activation; VWFC, voxel-wise functional connectivity.

Fig. 4 Behavioral and molecular characterization of monetary and natural reward alterations in depression. **A** Meta-analytic topic mapping of the right caudate. **B** Meta-analytic topic mapping of the right putamen. The terms are arranged and read from top in clockwise direction according to p-values. **C** Genetic expression of the caudate (left side) and putamen (right side). The plot shows the various genetic symbols according to their respective correlation values. **D** Bar chart illustrating the mean density values of 5-HT and DAT.

Table 1. Demographic information of included studies

Monetary rewards	Patients N/F	Mean_p	SD	Medication (%)	Controls N/F	Mean_c	SD
Fischer et al.	32/NA	17.87	2.58	0	18/NA	19.09	2.93
Mori et al.	15/9	18.5	0.6	0	15/7	19.1	0.7
Ubl et al.	30/16	46	11.85	0	28/15	43.96	2.18
Martin-Soelch et al.	16/12	24.31	4.08	0	16/12	25.19	4.79
Pizzagalli et al.	26/15	43.17	12.98	0	31/13	38.8	14.48
Dichter et al.	19/15	23.6	4.09	0	19/12	27.9	6.3
Hall et al.	29/16	37.01	8.48	51.72	29/16	37.38	9.78
Arrondo et al.	24/7	33.08	9.15	54	21/4	34.33	10.11
Chase et al.	40/31	31.04	8.04	NA	37/25	33.09	6.23
Burrows et al.	88/61	34.34	11.06	70.45	44/25	30.91	10.15
Smoski_a et al.	14/7	34.8	14.3	0	15/9	30.8	9.7
*Smoski_b 2011 et al.	9/NA	34.4	15.1	44.44	13/NA	26.2	6.3
Segarra et al.	24/7	33.08	9.15	54.16	21/4	34.33	10.11
Admon et al.	26/NA	42.66	11.72	0	29/NA	37.75	14.05
DelDonno et al.	23/16	25.09	3.32	0	27/23	29.15	9
Gorka et al.	9/6	25.4	7.7	11.11	18/13	29.5	13.1

Knutson et al.	14/NA	30.71	8.8	0	12/NA	28.67	4.25
Natural rewards							
Oh et al.	108/55	31.98	13.09	0	108/86	29.86	10.58
Epstein et al.	10/9	35.6	NA	0	12/7	32	NA
Gradin et al.	25/17	24.48	5.52	0	25/17	25.44	5.02
Keedwell et al.	12/8	43	9.8	91.67	12/8	36	14.6
Cabe et al.	13/11	NA	NA	46.15	14/9	NA	NA
Kumari et al.	6/6	44	NA	0	6/6	43.5	NA
Canli et al.	15/12	35.1	NA	46.7	15/12	30.7	NA
Fournier et al.	26/18	30.6	7.8	26.92	28/16	32.6	3.4
Gotlib et al.	18/13	35.2	10.262	50	18/13	30.8	NA
*Smoski_b et al.	9/NA	34.4	15.1	44.44	13	26.2	6.3

N/F Total number of participants/Total number of female participants, Mean_p mean age of patients, Mean_c mean age of controls, SD standard deviation, NA not available. *The coordinates of this study were merged into one text file when performing the general reward meta-analysis to preserve its independence.

A General reward

y = 4   z = -8

R Putamen

R Caudate   Subgenual
                   ACC

B Monetary reward outcome

y = 10   z = 5

R Caudate

C Natural reward outcome

y = 6   z = -4

R Putamen   L Putamen

D Monetary reward anticipation

y = 16   z = 4

L Putamen

Brain Orientation: Left | Right   **Brain Orientation: Right | Left**

A

B

C

D