Bio-psycho-social factors' associations with brain age: a large-scale UK Biobank diffusion study of 35,749 participants

Max Korbmacher[1,2,3,*], Tiril P. Gurholt[2], Ann Marie de Lange[2,4,5], Dennis van der Meer[2,6], Dani Beck[2,7,8], Eli Eikefjord[1,3], Arvid Lundervold[1,3,9,10], Ole A. Andreassen[2,11], Lars T. Westlye[2,8,11], Ivan I. Maximov[1,2,*]

[1] Department of Health and Functioning, Western Norway University of Applied Sciences, Bergen, Norway
[2] Norwegian Centre for Mental Disorder Research (NORMENT), Division of Mental Health and Addiction, University of Oslo and Oslo University Hospital, Oslo, Norway
[3] Mohn Medical Imaging and Visualisation Center (MMIV), Bergen, Norway
[4] Department of Psychiatry, University of Oxford, Oxford, UK
[5] LREN, Centre for Research in Neurosciences - Department of Clinical Neurosciences, CHUV and University of Lausanne, Lausanne, Switzerland
[6] Faculty of Health, Medicine and Life Sciences, Maastricht University, Maastricht, Netherlands
[7] Department of Psychiatric Research, Diakonhjemmet Hospital, Oslo, Norway
[8] Department of Psychology, University of Oslo, Oslo, Norway
[9] Department of Radiology, Haukeland University Hospital, Bergen, Norway
[10] Department of Biomedicine, University of Bergen, Bergen, Norway
[11] KG Jebsen Centre for Neurodevelopmental Disorders, University of Oslo, Oslo, Norway

* Correspondence:
Max Korbmacher, MSc & Ivan I. Maximov, PhD
Email: max.korbmacher@hvl.no, ivan.maximov@hvl.no

Keywords: brain age, age prediction, magnetic resonance imaging, diffusion MRI, health, cognition, brain variability.

Abstract

Brain age refers to age predicted by brain features. Brain age has previously been associated with various health and disease outcomes and suggested as a potential biomarker of general health. Few previous studies have systematically assessed brain age variability derived from single and multi-shell diffusion magnetic resonance imaging data. Here, we present multivariate models of brain age derived from various diffusion approaches and how they relate to bio-psycho-social variables within the domains of sociodemographic, cognitive, life-satisfaction, as well as health and lifestyle factors in midlife to old age (N = 35,749, 44.6 to 82.8 years of age). Bio-psycho-social factors could uniquely explain only a small proportion of the brain age variance. We observed a similar pattern across diffusion approaches. We consistently identified bio-psycho-social associations of interest across models. Waist-to-hip ratio, diabetes, hypertension, smoking, matrix puzzles solving, and job and health satisfaction and perception were associated with brain age. At the same time, sex and ethnicity showed large variability in their predictor strengths. As expected, chronological age contributed the strongest to explaining variability in brain age. Our results show that brain age cannot be sufficiently explained by bio-psycho-social variables alone. However, the observed associations suggest to adjust for sex, ethnicity, cognitive factors, as well as health and lifestyle factors.

Number of words: 7322
Number of figures and tables: 9

1   Introduction

Developmental trajectories of brain morphology are informative signaling markers of health. For example, significant deviations from normative morphology values can signify the presence or development of disease (Marquand et al., 2019; Remiszewski et al., 2022). Based on the idea that a general normative pattern could describe brain trajectories, the concept of brain age has been introduced. Here, different brain features are used to predict individuals' age. The difference between such predicted age and chronological age, the brain age gap (BAG), has the potential as a general health biomarker, sensitive to various neurological, neuropsychiatric, and neurodegenerative disorders (Kaufmann et al., 2019; Cole, 2020; Rokicki et al., 2021). Brain age can be derived using different imaging modalities. Structural and diffusion MRI (dMRI) have shown high prediction accuracy (e.g., Beck et al., 2022b; Chen et al., 2022; Cole, 2020; Leonardsen et al., 2022; Sone et al., 2022). Different dMRI-derived parameters allow one to describe multiple changes in WM micro-structure using various diffusion-weighted approaches. Such dMRI measures provide invaluable information about WM architecture at the micrometer scale and can be associated with macroscopic outcomes. The most popular dMRI approach, diffusion tensor imaging (DTI), is often used to describe WM organization (Basser et al., 1994). However, methodological advances and newer diffusion approaches may provide more meaningful bio-physical information (Novikov et al., 2019), thereby increasing the power of cross-validation of findings and their comparability with other clinical markers (Beck et al., 2021; Billiet et al., 2015; Kamagata et al., 2020; Wood et al., 2022).

The bio-psycho-social model (Engel, 1977) strives for a holistic perspective on medical research to understand health and disease by integrating information on biological, psychological, and social factors (e.g., Ghaemi, 2018; Wade & Halligan, 2017). Brain age can be utilized in this context as an indicator of general health (Kaufmann et al., 2019), using the different levels of observation (bio-psycho-social) to describe brain age relationships with different phenotypes. While there are some studies providing evidence for brain age associations with bio-psycho-social factors, including demographic, biomedical, lifestyle, cognitive, and behavioral factors (Beck et al., 2022b; Cole, 2020; de Lange et al., 2021; Leonardsen et al., 2022; Sone et al., 2022), it remains unclear whether brain age derived from different diffusion approaches relates differentially to sociodemographic, health, life-satisfaction, and cognitive factors (**Figure 1**), and what the qualities of such relationships are.

--Figure 1 around here--

It is necessary to increase our understanding and the interpretability of brain age by observing the associations of common phenotypes with brain age. There are large differences in the usage of underlying data and machine learning approaches applied to the data for brain age predictions (Franke & Gaser, 2019). Practical effects of such differences, for example, on phenotype associations, have yet to be systematized to better interpret brain age and BAG. The bio-psycho-social approach (Engel, 1977) lends itself to categorizing phenotypes into concrete groups. Within the groups phenotype associations with brain age in general can be considered in addition to differences in underlying data used to calculate brain age. Here we limit our investigations to dMRI-derived brain age.

DMRI can describe various biological processes by providing markers of brain tissue changes across the lifespan (Beck et al., 2021). These markers are not only heritable (Elliott et al., 2018) but also indicative of health, for example, by being associated with psychiatric and neurological disorders, addiction, stroke (Le Bihan & Iima 2015), or cardiovascular health (Beck et al., 2022b). Various diffusion metrics that have previously been related to cognitive and mental health traits have also shared genetic underpinnings with cognitive and mental health phenotypes (Zhao et al., 2021). The

biological underpinnings of dMRI markers become particularly apparent in WM abnormalities
observed in severe mental disorders, including schizophrenia (Cetin-Karayumak et al., 2020) or
bipolar disorder (Houenou et al., 2007). Furthermore, dMRI-derived brain age is higher in people
showing accumulations of cardiometabolic risk and markers of adipose tissue distribution (Beck et
al., 2022a, 2022b). Such associations between phenotypes and brain age can also be observed when
comparing high to low socioeconomic status (SES) groups, where low SES individuals have lower
WM integrity (Pavlakis et al., 2014; Shaked et al., 2019).

Furthermore, dMRI offers both single and multi-shell approaches and various meaningful metrics
describing white matter microstructure (Fieremans et al., 2011; Jensen et al., 2005; Kaden et al.,
2016a, 2016b; Reisert et al., 2017), which serve as a good basis for brain age estimations (Beck et al.,
2021; Korbmacher et al., 2022). Differences in brain age-phenotype relationships can be expected
when varying dMRI approaches, as varying underlying dMRI approaches will also produce
variability in brain age predictions (see Beck et al., 2021; Korbmacher et al., 2022), potentially due to
measuring different bio-physical processes (Fieremans et al., 2011; Jensen et al., 2005; Kaden et al.,
2016a, 2016b; Reisert et al., 2017). These potential differences become important when attempting to
generalize findings on brain age across the literature and setting standards for brain age predictions.
However, to what extent age predictions based on single and multi-shell dMRI approaches relate
differentially to phenotypes requires further investigation. Hence, comparing dMRI-based brain age
predictions can be fruitful, not only when expanding current efforts of examining brain age
associations with phenotypes but also by investigating whether differences in the underlying data can
influence relationships of brain age with bio-psycho-social variables.

State-of-the-art conceptualizations of health, such as the bio-psycho-social model (Engel, 1977),
recommend considering various domains or levels of explanation when assessing health outcomes,
such as brain age. To date, brain age has usually been described from single variables while
controlling for sex and age (e.g., Cole, 2020; Leonardsen et al., 2022). However, cumulative and
synergy effects can be expected to partly explain health, which has, for example, been shown for
cardiometabolic risk factors explaining brain age (see Beck et al., 2022a, 2022b). Hence, we group
available phenotypes that have previously been found influential for health (**Figure 1**) into health and
lifestyle factors, representing the biological dimension of the bio-psycho-social model, respectively.
Life satisfaction factors and cognitive factors represent the psychological dimension, and
sociodemographic factors the social dimension of the bio-psycho-social model, respectively.

Generally, explaining brain age variance is required to further our understanding of brain age and its
multivariate relationship with different phenotypes influencing physiology directly or indirectly. We,
therefore, extend previous work by explaining variance in brain age by combining sets of bio-psycho-
social variables into domains of sociodemographic, health, life satisfaction, and cognitive factors
(**Figure 1**) to assess their associations with brain age. In addition to exploring associations of bio-
psycho-social variables with dMRI-based brain age, we differentiate between diffusion approaches
used for brain age predictions. Previous findings revealed weak associations of various phenotypes
with brain age in the UK Biobank (e.g., Cole, 2020; Smith et al., 2019). Hence, we expect only small
proportions of the variance in brain age to be predicted by bio-psycho-social variables, whereby
factors directly representing or impacting physiology are more predictive of brain age than those
which impact physiology only indirectly. Thus, health factors are presumed to be more predictive of
brain age than sociodemographic, cognitive, and life satisfaction factors. Finally, we expect some
variability in these associations to be due to the underlying diffusion approach, as different WM
properties are also expected to be differentially related to phenotypes. We may move brain age closer
to the clinical utility by furthering our understanding of brain age.

136 2 Methods

137 2.1 Sample characteristics

The sample used has been described elsewhere (Korbmacher et al., 2022). In brief, the UK Biobank
(UKB) (Sudlow et al., 2015) diffusion MRI data consisted of N = 42,208 participants. We excluded
subjects who withdrew their informed consent (up to 22[nd] of February 2022) or with an ICD-10
diagnosis from categories F, G, I, or stroke from the general health assessment (Field 42006;
excluded: N = 3,521). We also excluded data that did not pass our quality control (N = 2,938) using
the YTTRIUM method (Maximov et al. 2021). Our final sample consisted of 35,749 healthy adults.
For an overview of demographics and the bio-psycho-social variables included in this study and their
relationship with brain age, see **Table 1**.

146 2.2 MRI acquisition, diffusion post-processing, and TBSS analysis

UKB MRI data acquisition procedures are described elsewhere (Alfaro-Almagro et al., 2018; Miller
et al., 2016; Sudlow et al., 2015). Briefly, single and multi-shell data were acquired at four different
locations using identical scanners: 3T Siemens Skyra, with a standard 32-channel head coil and key
diffusion parameters being MB = 3, R = 1, TE/TR = 92/3600 ms, PF 6/8, fat sat, b = 0 s/mm^2 (5x +
3× phase-encoding reversed), b = 1,000 s/mm^2 (50×), b = 2,000 s/mm^2 (50×) (Alfaro-Almagro et al.,
2018).

We obtained access to the raw diffusion data and pre-processed the data using an optimized pipeline
as described by Maximov et al. (2019). The pipeline includes corrections for noise (Veraart et al.,
2016), Gibbs ringing (Kellner et al., 2017), susceptibility-induced and motion distortions, and eddy
current artifacts (Andersson & Sotiropoulos, 2016). Isotropic 1 mm^3 Gaussian smoothing was carried
out using FSL's (Smith et al., 2004; Jenkins et al., 2012) *fslmaths*. Employing the multi-shell data,
Diffusion Tensor Imaging (DTI), Diffusion Kurtosis Imaging (DKI) (Jensen et al., 2005) and White
Matter Tract Integrity (WMTI) (Fieremans et al., 2011) metrics were estimated using Matlab 2017b
code (https://github.com/NYU-DiffusionMRI/DESIGNER). Spherical mean technique SMT (Kaden
et al., 2016b), and multi-compartment spherical mean technique (mcSMT) (Kaden et al., 2016a)
metrics were estimated using original code (https://github.com/ekaden/smt) (Kaden et al., 2016a,
2016b). Estimates from the Bayesian Rotational Invariant Approach (BRIA) were evaluated by the
original Matlab code (https://bitbucket.org/reisert/baydiff/src/master/) (Reisert et al., 2017).

Previous advances observing age-dependent WM changes have largely focused on single-shell
diffusion, such as DTI with DTI-derived metrics being fractional anisotropy (FA), and axial (AD),
mean (MD), and radial (RD) diffusivity, all being highly sensitive to age (Beck et al., 2021; Cox,
2016; Westlye et al., 2010). More recently developed multi-shell diffusion approaches which extend
the space of derivable diffusion metrics appear more sensitive to brain changes and sex differences
(Lawrence et al., 2021), and at the same time less sensitive to motion artefacts than single-shell
models (Pines et al., 2020). Newer approaches are 1) BRIA, as an alternative to not rely on fiber
orientation but rotation invariant feature (Reisert et al., 2017), 2) DKI, a method tackling the problem
of non-Gaussian diffusion (Jensen et al., 2005); 3) WMTI, which extends DKI by calculating inter
and extra-axonal features (Fieremans et al., 2011); and 4) SMT (Kaden et al., 2016b) and 5) mcSMT,
which factor out neurite orientation to give a better estimate of microscopic diffusion anisotropy
(Kaden et al., 2016a).

In total, we obtained 28 metrics (**Supplementary Table 1**) from six diffusion modeling approaches
(DTI, DKI, WMTI, SMT, mcSMT, and BRIA). To normalize all metrics, we used tract-based spatial
statistics (TBSS) (Smith et al., 2006) as part of FSL (Smith et al., 2004; Jenkins et al., 2012). In brief,

initially, all FSL BET-extracted (Smith, 2002) FA images were aligned to MNI space using non-
linear transformation (FNIRT) (Jenkins et al., 2012). Subsequently, we derived the mean FA image
and the related mean FA skeleton. Each diffusion scalar map was projected onto the mean FA
skeleton using the standard TBSS procedure. To provide a quantitative description of diffusion
metrics we evaluated averaged values over the skeleton and two WM atlases, namely the John
Hopkins University (JHU) atlas (Mori et al., 2005) and the JHU tractography atlas (Hua et al., 2008;
see **Supplementary Table 4** for an overview). Finally, we obtained 20 WM tracts and 48 regions of
interest (ROIs) based on a probabilistic WM atlas (JHU) (Hua et al., 2008) for each of the 28 metrics,
including the mean skeleton values. Altogether, we derived 1,932 features per individual (28 metrics
* (48 ROIs + 1 skeleton mean + 20 tracts)); see **Table 2** for number of dMRI features**,**
Supplementary Table 1 for metrics, **Supplementary Table 4** for regions and tracts.

2.3 Brain Age Predictions

Brain age was predicted using the XGBoost algorithm (Chen & Guestrin, 2016) implemented in
Python (v3.7.1). From the total included sample (N = 35,749), we used 10% (N = 3,575) for
parameter tuning on a data set containing data from all diffusion approaches using 5-fold cross-
validation. The remaining 90% were used as validation data (N = 32,174) to which we applied 10-
fold cross-validation (hyperparameters: learning rate = 0.05, max layers/depth = 3 and the number of
trees = 750; see Korbmacher et al., 2022). All presented statistics were calculated from the test data
set (N = 32,174).

2.4 Statistical Analyses

All statistical analyses were carried out using Python, version 3.7.1 and R, version 4.2.0 (www.r-
project.org/). These analyses focused on the associations between brain age and (1) demographics,
(2) social factors, (3) cognitive test scores, (4) life satisfaction, and (5) health and lifestyle factors
(with weight on cardiometabolic factors). For detailed information on how variables were extracted
and coded see **Supplementary Table 9**. We adjusted *p*-values for multiple comparison using
Bonferroni correction, dividing the alpha-level by eight, the number of brain age models used.

Initially, we used linear mixed effects models with the random intercepts at the level of scanner site
to explain changes in brain age from socio-demographics, cognitive test scores, life satisfaction (self-
assessment), and health and lifestyle factors. For an overview of the statistical models used see
Figure 2. Brain ages were derived from each of the six diffusion approach's metrics, whole-brain
averaged metrics for all approaches (mean multimodal model) and from all data combined (full
multimodal model), resulting in eight brain age models (see **Table 2** for models and **Supplementary**
Table 1 for dMRI approach-specific metrics). Each of these brain ages were used in the analyses.

213   --Figure 2 around here--

We established the following models to compare:

(1) A baseline model capturing the relationship of age, sex, the age-sex interaction, and scanner site
with brain age. This baseline model was selected as predicted age is expected to be largely reflected
by chronological age. However, also sex (e.g., Rokicki et al., 2021), and scanner site (here, Bristol,
Cheadle, Newcastle, Reading) and prediction bias (e.g., Jirsaraie et al., 2022) have been shown to be
influential for brain age. Using a baseline model and additional models for comparison had the goal
to estimate added variance explained by the bio-psycho-social models above and beyond the baseline
model, and to observe predictors within these bio-psycho-social models.

$$brainage = sex + age + sex * age + site$$

(2) A sociodemographic model additionally included ethnic ancestry (binary yes/no self-reported
white European; for additional information sample groupings by ethnicity see **Supplementary Table**
13), average annual total household income before tax (coded as continuous variable 1-5, with low
<£18,000 to high income >£100,000), and higher education (binary yes/no self-report of having
obtained higher education) relative to the baseline model.

$$brainage = sex + age + sex * age + ethnicity + income + education + site$$

(3) A cognitive model testing how non-verbal cognitive abilities add to the baseline model
(overview: Fawns-Ritchie & Deary, 2020). Namely, the number of matrix puzzles solved (matrixS)
testing non-verbal reasoning using COGNITO Matrices, tower arranging correctly solved (towerS)
testing executive function using the Delis-Kaplan Executive Function System Tower Test,
prospective memory (memory) assessed with the Rivermead Behavioural Memory Test, fluid
intelligence (intel) from the UKB own Fluid IQ test, digits remembered (digits) from the Symbol
Digit Modalities Tests, and the mean number of incorrect pair matches (IPM) across trail A and B
assessing visual declarative memory using the Wechsler Memory Scale IV Designs I and Designs II.
Correlations were small to moderate (r_{max} = 0.41) with the variance inflation factor (VIF) indicating
low levels of multicollinearity (**Supplementary Figure 2**)

$$brainage = sex + age + sex * age + matrixS + towerS + memory + intel + digits + IPM + site$$

(4) A life satisfaction model that additionally included job satisfaction (jobS), financial satisfaction
(financeS), overall health rating (healthR), health satisfaction (healthS), family relation satisfaction
(famS), happiness, friend relationship satisfaction (friendS) relative to the baseline model. Some of
the model features were highly correlated (r_{max} = 0.65), yet VIF values indicated low levels of
multicollinearity (**Supplementary Figure 3**).

$$brainage = sex + age + sex * age + jobS + financeS + healthR + healthS + famS + friendS + happiness + site$$

(5) A health and lifestyle model testing how body mass index (BMI), pulse pressure (Ppressure: the
difference between systolic and diastolic blood pressure), waist-to-hip-ratio (WHR), binary smoking
status, binary diabetes diagnosis (both type I and II), binary high cholesterol (chol), binary diagnosed
vascular problem (DVP), birth weight (Bweight), sleeping hours, and daily coffee intake (coffee) add
to the baseline model, with only BMI and WHR showing a moderate correlation r = 0.42, but all
other correlations being small rs < 0.16, with VIF values indicating only low levels of
multicollinearity (**Supplementary Figure 4**).

$$brainage = sex + age + sex * age + BMI + WHR + Ppressure + diabetes + smoking + chol + DVP + Bweight + coffee + site$$
254
255
3   Results

3.1   Overview of brain age gap and bio-psycho-social factors

Table 1 gives an overview of brain age calculated from combined single and multi-shell diffusion
data in relation to the observed bio-psycho-social factors. Strongest associations when adding single
factors to a model explaining brain age from age were found for WHR (β = 4.86, $p < .001$), overall
health rating (β = -0.36, $p < .001$), and health satisfaction (β = 0.26, $p < .001$). Strongest brain age
group differences were found for sex (β = -1.07, $p = .001$), diabetes (β = 0.99, $p < .001$), and
hypertension (β = 0.86, $p < .001$).

264   --Table 1 around here--

3.2   Brain age predictions

Brain age predictions from data derived separately for each diffusion approach (BRIA, DKI, DTI,
SMT, mcSMT, WMTI), their combination in a full data set (full multimodal), or global mean metrics
(mean multimodal) can be found in **Table 2**. Combining single and multi-shell diffusion approaches
led to the highest-performing age predictions, whereas using only whole brain average scores led to
the least accurate predictions. Differences in age predictions across single diffusion approaches were
small, with WMTI predicting age slightly better than the other approaches and mcSMT theleast.

272   --Table 2 around here--

3.3   Linear mixed effect models explaining brain age gap from bio-psycho-social factors

274 **Figure 3** shows the marginal variance in brain age explained from the different models from (1)
275 baseline, (2) social factors, (3) cognitive test scores, (4) life satisfaction, and (5) health and lifestyle
276 factors (with weight on cardiometabolic factors). For a full overview of model metrics see
277 **Supplementary Table 10**.

278   --Figure 3 about here--

Only a small proportion of either marginal or conditional variance in brain age could be uniquely
explained by bio-psycho-social variables (marginal and conditional $R^2 < .03$; **Figure 3**,
Supplementary Table 10). Differences between this uniquely explained variance were small across
diffusion approaches. Model comparisons showed that bio-psycho-social models explained
significantly more variance in brain age than the baseline model (with age, sex, and age-by-sex
interaction as fixed and scanner site as random effect), irrespective of the diffusion approach used to
calculate brain age ($ps < .01$; **Supplementary Table 2**). Across models, age was used as a control
variable to correct for the mere reflection of age by brain age producing stable associations across
models (1-5) for multimodal brain age (**Figure 4-7**). However, the other diffusion approaches' brain
ages were negatively associated with brain age, giving another indication of overall poor model fit.
Even more so, the effect of sex was dependent on the model, producing mixed effects with large
uncertainty surrounding β-values. Finally, the age-by-sex interaction's associations with brain age
were stable across models and approaches (**Figure 4-7**).

3.3.1 Sociodemographic factors' associations with brain age

In the model including sociodemographic factors explaining brain age (see **Figure 4** for the
predictors), results were mixed for the significant predictors. Sex was a significant predictor for mean
DKI, DTI, and WMTI ($ps_{Bonferroni} < .05$), the age-by-sex interaction only for BRIA ($p_{Bonferroni} = .045$),
and ethnicity only for DKI ($p_{Bonferroni} = .012$; **Figure 4**). Overall, only 95% confidence intervals of β-
values for age and ethnicity were not consistently overlapping, indicating differential effects of these
variables on brain age based on the underlying data. All other 95% confidence intervals surrounding
coefficients' β-values were overlapping across diffusion approaches, with expected strong age
contributions predicting brain age.

301   --Figure 4 about here--

3.3.2 Health and lifestyle factors' associations with brain age

Similarly, in the model including health and lifestyle factors explaining brain age (see **Figure 5** for
the predictors), significant health factors leading to higher brain age were WHR ($ps_{Bonferroni}$ < .001),
pulse pressure ($ps_{Bonferroni}$ < .001), and hypertension ($ps_{Bonferroni}$ < .001). Evidence across diffusion
approaches was mixed for the other predictors with smoking predicting brain age derived from
BRIA, DTI, mean scores, SNT, and WMTI ($ps_{Bonferroni}$ < .05), diabetes diagnosis for all models except
DKI and DTI(ps < .05), the diagnosis of at least one vascular disease for BRIA, mean scores, and
mcSMT ($ps_{Bonferroni}$ < .02), and average daily cups of coffee for brain age estimates except the one
based on BRIA ($ps_{Bonferroni}$ < .01), and the age-by-sex interaction for BRIA and mcSMT ($ps_{Bonferroni}$
< .04).

Interestingly, WHR was a stronger predictor of brain age in males than in females (**Supplementary**
Figure 1). Practically, a WHR $β$-value of, for example $β$ = 4 would mean that for every 0.1 step
change in WHR, the brain age can be expected to increase by 0.4 years. Importantly, this association
was controlled for age, as age is correlated with WHR at r = 0.14 and brain age at r = 0.80. Mean
population values for WHR were found to be WHR < 1 (Molarius et al., 1999), with our sample
corresponding with these estimates (M_{WHR} = 0.871±0.088, min = 0.534, max = 1.472) with males
having a higher WHR (M_{WHR} = 0.923±0.064) than females (M_{WHR} = 0.817±0.069).

BMI was potentially non-significant due to the model construction as the highly correlated WHR
(**Supplementary Figure 4**) was a significant predictor of brain age, and BMI alone being a
significant predictor of brain age (**Table 1**). Finally, higher birth weight was associated with lower
brain age estimated from full and mean models, as well as BRIA and WMTI ($ps_{Bonferroni}$ < .02).

Generally, 95% confidence intervals around coefficients' $β$-values were overlapping across models
indicating no significant differences in $β$-values across diffusion approaches. As a control, we ran the
same model without WHR as predictor, due to its high correlation with BMI, rendering BMI as
significant predictor across diffusion approaches' brain ages except the mean model ($βs$ > 0.01
$ps_{Bonferroni}$ < .004), also showing now clearer evidence for higher brain age when smoking ($ps_{Bonferroni}$
< .05), with other predictors unchanged (**Supplementary Figure 5**). Furthermore, leaving out
hypertension, being a substrate of blood pressure, did not lead to changes in the model
(**Supplementary Figure 6**). For both models, variance explained is slightly reduced compared to the
models including the respective variables, making the reduced models significantly different (ps
< .001) from the full health models (**Supplementary Table 7-8**).

333   --Figure 5 about here--

3.3.3 Life satisfaction factors' associations with brain age

When modeling brain age from life satisfaction (see **Figure 6** for the predictors), self-rated health
was a significant predictor of all brain age estimates except for DKI brain age ($ps_{Bonferroni}$ < .05) and
health satisfaction for all brain age estimates except the mean model's brain age ($ps_{Bonferroni}$ < .02).
Only the 95% confidence intervals of $β$-values for age do not overlap across models (with the mean
model having the largest $β$ and full model the smallest $β$-value for age). All other 95% confidence
intervals around coefficients' $β$-values overlap across models indicating no significant differences in
$β$-values across diffusion approaches.

Perceived health is moderately correlated with health satisfaction and was left out in a control model
resulting in a slightly stronger effect of health satisfaction and significantly worse performing model
(ps < .001; **Supplementary Figure 7, Supplementary Table 7-8**). Differently, when leaving out

happiness as being correlated with several variables the model remains unaffected (*ps* > .23;
Supplementary Figure 8, Supplementary Table 7-8).

347   --Figure 6 about here--

3.3.4 Cognitive factors' associations with brain age

The only cognitive factor explaining brain age across all models was symbol digit substitution
(*ps*$_{Bonferroni}$ < .001; **Figure 7**). Matrix puzzles solved was only a significant predictor for the full
multimodal brain age (*p*$_{Bonferroni}$ = .014), and sex only for DTI and WMTI (*ps*$_{Bonferroni}$ < .02). Confidence
intervals around coefficients' *β*-values are overlapping across models indicating no significant
differences in *β*-values across diffusion approaches. Fluid intelligence and matrix puzzles are highly
correlated and hence, matrix puzzles were left out in a quality control model, not significantly
affecting the structure of most models (**Supplementary Figure 9, Supplementary Tables 7-8**).

356   --Figure 7 about here--

3.3.5 Follow-up and quality control analyses

Due to the strong variability in sex *β*-values across models (**Figures 4-7**), we also ran the described
analyses separately for males and females (for the predictions from the models presented in **Table 2**)
showing some differences in model performance. For example, bio-psycho-social models explained a
differential of between 1%-4% of conditional variance for males (**Supplementary Table 3**) and
differences in contributions of the different models' predictors, predictors' *β*-values being generally
higher for males (**Supplementary Figure 1**). Finally, as a quality control step, we ran the proposed
five baseline and bio-psycho-social models with the first principal component of the numeric
predictors from each of the models showing an even smaller proportion of the variance in brain age
uniquely explained by the principal components (R^2 < 1%; **Supplementary Table 5**), with
differences between these models and respective baseline models yet being highly significant
(**Supplementary Table 6**). Overall, quality checks show small levels of multicollinearity, and that
each predictor contributes individual to the models (**Supplementary Figures 2-9**, **Supplementary**
Tables 7-8), supporting assumptions about the robustness of the utilized models, as well as that
simply adding all variables together saturates the model leading to lower model performance than at
baseline across brain ages based on different diffusion approaches with a differential in marginal R^2 =
3.38%.

4   Discussion

We assessed the influence of various bio-psycho-social variables on brain age estimated from
different diffusion approaches (and their combinations). Linear mixed effects models showed that
bio-psycho-social variables uniquely explain a small proportion of brain age variability and
predictors' *β*-values' confidence intervals overlap for most predictors. As predicted, bio-psycho-
social domain-specific brain age variance explained variability was small, with health and lifestyle
factors being most indicative of brain age. However, differences in brain age variance explained
between bio-psycho-social models were small. Furthermore, differences between predictor strength
were small, indicating small group differences in brain age and small association differences between
phenotypes and brain ages derived from different diffusion approaches. Significant predictors of
brain age were job satisfaction, health satisfaction, WHR (and to a lesser extent BMI when excluding
WHR as a predictor), diabetes, hypertension, any vascular diagnosis, daily coffee consumption,

smoking, birth weight, matrix puzzles, and symbol digit substitution performance. Our findings
indicate that brain age estimates derived from different diffusion approaches will relate similarly to
the examined bio-psycho-social factors. The bio-psycho-social models were however limited in
explaining brain age and more impacting predictors and predictor combinations are still to be
explored.

4.1   Explaining brain age from bio-psycho-social factors

Recent research has made a strong case for the conjunction effects of various bio-psycho-social
factors in explaining general health (Lehman et al., 2017). Applied to brain age, for example,
cardiometabolic effects have been shown to influence brain age (Beck et al., 2022a, 2022b).
However, assessments of how much of the variance explained in brain age above and beyond age,
sex, age-by-sex interaction, and scanner site have not been described in the literature. We find close-
to-zero added brain age variance explained by models including single bio-psycho-social variables
(**Table 1**). Comparably more, yet still a small proportion of brain age variance ($R^2 < 4\%$) is uniquely
explained by bio-psycho-social variable groups of socio-demographics, health and lifestyle, life
satisfaction, and cognitive ability (**Figure 3**, **Supplementary Table 10**). Importantly, the added brain
age variance explained is not just an effect of adding predictors randomly to the model, which rather
decreases the variance explained, as shown when adding all bio-psycho-social variables to the model.
Hence, it seems more sensible to employ models incorporating several compared to single domain-
specific variables to explain brain age. However, our results also indicate that a large part of the
variance in brain age cannot be explained by our proposed bio-psycho-social models. Whether this
unexplained variance is due to actual biologically founded individual differences, or the
characteristics of brain age, for example, how the metric is being estimated (de Lange et al., 2022),
remains unclear. BAG might also be rather static and indicated by constants such as genetic
architecture and birth weight (Vidal-Pineiro et al., 2021). This would explain the smaller influence of
more variable bio-psycho-social variables. Strong deviations from the norm, for example, due to
atrophy will also have a strong influence on brain age (Kaufmann et al., 2019). Hence, for diseases
impacting brain structure, brain age can be a useful indicator of health status (Kaufmann et al., 2019).
Potentially, the health and lifestyle factors which are most likely to impact brain structure are
therefore also more predictive of brain age than other bio-psycho-social variables (**Figure 3-6**).
While our models failed to explain larger proportions of the variance of brain age, there are various
interesting phenotype associations within these models which will be discussed in the following.

4.1.1 *The importance of age, sex, and ethnicity*

Usually, age, sex, and at times, scanner site, are used as covariates for brain age-phenotype
associations as they are expected to influence various phenotypes (Jirsaraie et al., 2022). As brain age
reflects chronological age, age also explains most of the brain age variance (**Table 3**). We also find
that the effects of sex and ethnicity were highly variable across models predicting brain age with sex
and ethnicity being mostly non-significant predictor across diffusion models (**Figure 3-6**). Yet,
models were more predictive of bio-psycho-social factors in males than females (**Supplementary
Table 3, Supplementary Figure 1**). Where the influence of sex changes based on the model
construction, while potentially also influencing the model (**Table 2-5**; **Supplementary Figure 4-7**).
Some of the observed sex differences might be based on anatomical features, such as higher
intracranial volume in males and different sex-specific aging (Eikenes et al., 2022). Brain age was
differentially sensitive to ethnicity dependent on the approach it was calculated on (**Figure 4-7**), with
these differences being influenced by sex (**Supplementary Figure 1**). A previous study showed that
being a UK immigrant might influence brain age estimates (Leonardsen et al., 2022). Potentially,
genetic contributions to brain age both estimated from T1-weighted (Ning et al., 2020; Vidal-Pineiro

et al., 2021) and dMRI data (Salih et al., 2021) also have a connection with the mentioned brain age
differences by ethnicity. However, the causal structure of sex and ethnicity differences in brain age
estimates requires further investigation.

Previous research has shown the effects of sex on metrics derived from conventional and advanced
diffusion approaches, such as BRIA, DKI, DTI, NODDI, RSI, SMT, SMT mc, and WMTI (Beck et
al., 2021; Eikenes et al., 2022). While a systematic assessment of sex-related effects on diffusion
metrics from both conventional and advanced dMRI approaches from voxel-to-whole-brain averages
over the lifespan is yet to be established, different studies presented sex-related developmental
trajectories in the structural connectome in children (Ingalhalikar et al., 2013), and sex related WM
changes during aging (Hsu et al., 2008). Furthermore, sex differences in aging reflected in WM
microstructure can be expected due to menopause and cascading biological processes, affecting both
brain and body systems in various ways (Barth & de Lange, 2020; Lohner et al., 2022; Mosconi et
al., 2021). Hence, developmental trajectories differing between males and females can be expected
which makes sex-separated analyses useful to providing important additional information (e.g., as in
Subramaniapillai et al., 2022). To which extend this applies to ethnicity requires further research.
Hence, further research is required to delineate the underlying causal structure of sex and ethnicity to
explain their highly variable associations with brain age.

4.1.2 *Health and lifestyle factors*

Interestingly, while the health and lifestyle factors models explained only a small proportion of the
brain age variance, most of its predictors were significant. Furthermore, these predictors are generally
only weakly correlated (**Supplementary Figure 4**), but when added explaining more variability in
brain age than on their own (**Table 1**). To a certain degree, this is not surprising, due to dependencies
between these predictors. For example, WHR, being the strongest predictor of brain age (see **Figure
5**), shows a clear relationship with pulse pressure (**Supplementary Figure 4**). For the extreme cases,
this is expressed in a well-established relationship between obesity and hypertension (Kotsis et al.,
2010) or any vascular diagnosis (Mathew et al., 2008). This is reflected in brain age, where minimum
and maximum values show that there is an expected difference of up to 4 years in brain age between
those with lowest compared to highest WHR, or a 2.4-years brain age difference between mean and
maximum WHR. Interestingly, blood pressure is expected to increase with age, and higher blood
pressure is positively associated with BAG (Cherbuin et al., 2021). However, these effects were not
exclusively driven by hypertension but across the spectrum of measured blood pressure values
(Cherbuin et al., 2021). This was supported by our findings showing both an effect of pulse pressure
and hypertension on brain age. These effects are not surprising, as hypertension has been suggested
as one of the most important risk factors for various cerebrovascular complications such as cerebral
small vessel disease and resulting cognitive impairments (Forte et al., 2020; Meissner, 2016).

Another aspect of high WHR and BMI is obesity increasing diabetes risk (Kahn et al., 2006). While
the evidence for the direction of the effect of diabetes is mixed (Cole et al., 2018; Franke et al., 2013;
Sone et al., 2022), we find participants with diabetes to show higher brain age than those without
diabetes (**Table 2, Figure 5**). Several complications within the central nervous system have been
associated with diabetes, including morphological, electrophysiological, and cognitive changes, often
in the hippocampus (Wrighten et al., 2009), just as WM lesions and altered metabolite ratios
(Biessels & Reijmer, 2014, van der Harten et al., 2006), supporting the idea of higher brain age
among those with diabetes. But also generally, the increase in risk of cardiovascular disease by WHR
is mediated by BMI, systolic blood pressure, diabetes, lipids, and smoking (Gill et al., 2021). In
relation to the brain, higher WHR has been generally associated with lower gray matter volume
(Gurholt et al., 2021; Hamer & Batty, 2019), and higher WM brain age (Beck et al., 2022a, 2022b;

Subramaniapillai et al., 2022). Hence, to which extent high WHR accelerates brain aging requires
further investigation, which might be particularly informative when observed in combination with
other health and lifestyle variables (Hamer & Batty, 2019) and sex (Subramaniapillai et al., 2022).

For example, negative health consequences of smoking (Erhardt, 2009) are reflected in smoker's
cortex being thinner (Gurholt et al., 2021), and smokers' brains being 1.5 years older on average than
non-smokers' brains (**Table 1**). Smoking is a known risk factor for cardiovascular health significantly
increasing its mortality and inducing various negative downstream effects on health (Erhardt, 2009),
with negative impacts on the reward system (Le Foll et al., 2022), repeatedly shown in rats (e.g., Cao
et al., 2013; Gozzi et al., 2006; Kenny & Markou, 2006). It can hence be expected that both general
and brain health are influenced by smoking, making it an important control variable in assessing
brain age.

The findings for coffee on the other hand are mixed, suggesting coffee consumption to be generally
positive for cardiovascular health and decreasing the risk of Parkinson's disease, stroke, and
Alzheimer's (Nehling, 2016). The consumption of higher doses of caffeine is, however, associated
with smaller brain volume and an increased risk of dementia (Pham et al., 2021). Practically, the
direct effect of the number of daily cups of coffee consumed is small in our study. It would require
on average 10 cups of coffee daily for an increase of 0.6 years of brain age, fitting the observations
made by Pham et al. (2021). It also remains unclear whether the effect of coffee consumption on
brain age is rather mediated by third variables such as poor sleep and mental health downstream
effects which show direct negative effects on health (Distelberg et al., 2017). Additionally, there are
vulnerable groups in which caffeine can cause adverse effects such as people with hypertension
(Higdon & Frei, 2007). We conclude that health and lifestyle factors function in synergy in
influencing brain age.

4.1.3 *Health perception and satisfaction, and job satisfaction*

We find significant assignations of self-rated health, friendship/relationship satisfaction, and job
satisfaction with brain age. Self-assessments and self-rated scores are some of the fastest and easiest
assessments. Yet, their reliability is under constant scrutiny, particularly when assessing health
outcomes (e.g., Crossley & Kennedy, 2002; Reychav et al., 2019). In our study, self-rated overall
health was a significant predictor of brain age, suggesting that asking participants about their health
can be a useful preliminary assessment of different aspects of health. Self-rated health was
additionally moderately correlated with health perception (**Supplementary Figure 3**), indicating
both variables measure, to a certain degree, the same underlying phenomenon. However, self-rated
health brain age associations were stronger and more variable across diffusion approaches' brain ages
(**Figure 6**). These associations support the idea of brain age is not only indicative of brain health, but
also overall health (Kaufmann et al., 2019).

Lastly, there was a trend of individuals' job satisfaction being associated with brain age (**Figure 6**).
Conceptually, this would not be surprising as associations between wealth and health (e.g., Adler &
Ostrove, 1999) as well as job (e.g., Faragher et al., 2013) and financial satisfaction and health (e.g.,
Hsieh, 2001) have already been investigated. However, in the case of our study, higher job
satisfaction was also indicative of higher brain age. Potential reasons are speculative but might reflect
the tendency of people engaged in their jobs to work long hours which has previously been related
with various negative mental and physical health outcomes (Bannai & Akiko, 2014; Lim et al.,
2011). Nevertheless, the underlying mechanisms of the associations between these single items in
their relationship with brain age require further investigation.

4.1.4 *Cognitive scores*

Cognitive scores' impact on brain age might be small in the current study, yet still important in
general (**Table 1**). This might be due to the selection of the observed cognitive test scores, with many
more possible tests to be included which are potentially more indicative of brain age, such as IQ
(Elliott et al., 2021). Another opportunity lies in assessing associations of cognitive performance and
brain age in clinical groups. For example, brain age has been found to be explanatory of symbol digit
modality test scores in multiple sclerosis suggesting brain age as a biomarker for cognitive
dysfunction (Denissen et al., 2022). Similar to such findings, we find a similarly sized effect of
symbol digit substitution test scores in our healthy aging data (**Figure 7**). Associations of cognitive
performance and brain age are also sensitive to sex. For example, the number of solved matrix
puzzles showing an effect when analyzing males and females data together seemed to be a predictor
of brain age only in females when analyzing females from males data separately (**Supplementary**
Figure 1). The quality of these differences requires further investigation.

4.2   Variability in brain age-phenotype relationships

Imaging phenotypes derived from diffusion UKB data contributes to a small additional proportion of
the variability in the obtained results. For example, SMT mc as an extension of SMT produces brain
age estimates closely related to SMT brain age (Korbmacher et al., 2022). This is clearly reflected in
the associations of the brain ages with different bio-psycho-social variables or group differences
(**Figures 4-7**) for strongly correlated BRIA, SMT, and SMT mc brain age estimates, and strongly
correlated brain ages of the full multimodal model (combining diffusion approaches' data), DTI, and
WMTI. However, differences between diffusion approaches are generally small (**Figures 4-7**).

The used brain age estimation model might also introduce variability in brain age phenotype
associations. However, model evaluation metrics such as R^2, MAE, or RMSE depend additionally on
cohort- and study-specific data characteristics making brain age model comparison not
straightforward (de Lange et al., 2022). However, there are differences between models trained on
voxel-level compared to region-wise data. Deep learning models using voxel-level data reach age
predictions errors as low as MAE = 2.14 years in midlife to late adulthood (Peng et al., 2021) or
MAE = 3.90 years across the lifespan (Leonardsen et al., 2022) while explaining large proportions of
variance in age ($R^2 > 0.90$), whereas models trained on regional and global average measures predict
age usually with larger error, MAE > 3.6 years, and/or lower variances explained $R^2 < 0.75$ (Beck et
al., 2021, 2022; de Lange et al., 2020a, 2020b; Korbmacher et al., 2022; Rokicki et al., 2021).
However, Niu et al. (2019) showed that with different shallow and deep machine learning algorithms
(ridge regression, support vector regressor, Gaussian process regressor, deep neural networks) high
prediction accuracies ($R^2 > 0.75$, MAE < 1.43) could be reached when using multimodal regional
average data using a young sample with narrow age range. When comparing brain age retrieved from
the same database (UK Biobank), using different samples, data, and different methods to calculate
brain age, similar patterns of associations between brain age and phenotypes can be observed in this
study. For example, diabetes diagnosis, diagnosed vascular problems or place of birth (see Figure 4
in Leonardsen et al., 2022), hip circumference, diabetes, trail-making tasks (here symbol digit
substitution), and matrix pattern completion were significantly associated with brain age (see Table 5
in Cole, 2020). Nevertheless, it remains unclear whether the differences in the findings are due to
analysis degrees of freedom, sample characteristics, or actual bio-physical manifestations. For
example, the underlying data used to estimate brain age, being dMRI in this study were T1-weighted
images in Cole (2020) and Leonardsen et al. (2022). Furthermore, phenotype data processing, such as
dummy coding, differed, as well as models to predict brain age. Here we used four mixed models
grouping a) demographics, b) cognitive, c) life satisfaction, and d) health and lifestyle variables to

predict brain age. In contrast, Cole (2020) predicted bias-adjusted brain age from simple linear
models with sex, age, and age^2 as covariates, and Leonardsen et al. (2022) observed similar
associations for uncorrected brain age predicted from the respective phenotype and age and sex as
covariates. However, bio-psycho-social variables are likely to interact forming a complex pattern
when explaining variables such as brain age. When adding only single bio-psycho-social variables to
a baseline model and then comparing the two models, the differences in variance explained are
smaller than when adding blocks of meaningfully related variables (compare **Table 1** and
Supplementary Table 10). The composition of variables forming such blocks/models requires
further investigation. In sum, there are various sources introducing variability in brain age phenotype
associations encompassing not only the underlying data but also researchers' degrees of freedom
such as data selection, processing, and analysis.

4.3   Conclusion and future directions

Microstructural WM measures derived from conventional and advanced diffusion MRI approaches
produce brain age estimates that are related to various bio-psycho-social factors. In that context, the
spatial specificity of the dMRI metrics plays a significant role in brain age estimations and their
interpretation, for example, in their associations with bio-psycho-social factors. The influence of sex
and ethnicity is largely variable suggesting the usage of sensible control mechanisms, such as
separate analyses or exclusions in case of strongly imbalanced samples. Health and lifestyle factors,
particularly physiological measures, WHR, birth weight, diabetes, hypertension, and related
diagnoses, as well as smoking status and coffee consumption, were more predictive of brain age than
cognitive and life satisfaction measures. Apart from health satisfaction and self-ratings, we found
relationships of life satisfaction variables with brain age to be non-significant. Digit substitution task
performance on the other hand, might be relevant in samples from midlife to old age. We recommend
future study designs taking observable interactions between the different bio-psycho-social effects
into account. A potentially helpful guiding principle in the search for bio-psycho-social variables
affecting brain age could be to focus on measures which are directly or indirectly related to or reflect
physiology and health.

5   References

Adler, N. E., & Ostrove, J. M. (1999). Socioeconomic status and health: what we know and what we don't. *Annals of the New York academy of Sciences*, *896*(1), 3-15. https://doi.org/10.1111/j.1749-6632.1999.tb08101.x

Alfaro-Almagro, F., Jenkinson, M., Bangerter, N. K., Andersson, J. L., Griffanti, L., Douaud, G., ... & Smith, S. M. (2018). Image processing and Quality Control for the first 10,000 brain imaging datasets from UK Biobank. *Neuroimage*, *166*, 400-424. https://doi.org/10.1016/j.neuroimage.2017.10.034

Andersson, J. L., & Sotiropoulos, S. N. (2016). An integrated approach to correction for off-resonance effects and subject movement in diffusion MR imaging. *Neuroimage*, *125*, 1063-1078. https://doi.org/10.1016/j.neuroimage.2015.10.019

Bannai, A., & Tamakoshi, A. (2014). The association between long working hours and health: a systematic review of epidemiological evidence. *Scandinavian journal of work, environment & health*, 5-18. https://doi.org/10.5271/sjweh.3388

Barth, C., & de Lange, A. M. G. (2020). Towards an understanding of women's brain aging: the immunology of pregnancy and menopause. *Frontiers in Neuroendocrinology*, *58*, 100850. https://doi.org/10.1016/j.yfrne.2020.100850

Beck, D., de Lange, A. M. G., Maximov, I. I., Richard, G., Andreassen, O. A., Nordvik, J. E., & Westlye, L. T. (2021). White matter microstructure across the adult lifespan: A mixed longitudinal and cross-sectional study using advanced diffusion models and brain-age prediction. *NeuroImage*, *224*, 117441. https://doi.org/10.1016/j.neuroimage.2020.117441

Beck, D., de Lange, A. M. G., Alnæs, D., Maximov, I. I., Pedersen, M. L., Leinhard, O. D., ... & Westlye, L. T. (2022a). Adipose tissue distribution from body MRI is associated with cross-sectional and longitudinal brain age in adults. *NeuroImage: Clinical*, *33*, 102949. https://doi.org/10.1016/j.nicl.2022.102949

Beck, D., de Lange, A. M. G., Pedersen, M. L., Alnæs, D., Maximov, I. I., Voldsbekk, I., ... & Westlye, L. T. (2022b). Cardiometabolic risk factors associated with brain age and accelerate brain ageing. *Human brain mapping*, *43*(2), 700-720. https://doi.org/10.1002/hbm.25680

Bentler, P. M. (1980). Multivariate analysis with latent variables: Causal modeling. *Annual review of psychology*, *31*(1), 419-456.

Biessels, G. J., & Reijmer, Y. D. (2014). Brain changes underlying cognitive dysfunction in diabetes: what can we learn from MRI?. *Diabetes*, *63*(7), 2244-2252. https://doi.org/10.2337/db14-0348

Billiet, T., Vandenbulcke, M., Mädler, B., Peeters, R., Dhollander, T., Zhang, H., ... & Emsell, L. (2015). Age-related microstructural differences quantified using myelin water imaging and advanced diffusion MRI. *Neurobiology of aging*, *36*(6), 2107-2121. https://doi.org/10.1016/j.neurobiolaging.2015.02.029

Cao, J., Wang, J., Dwyer, J. B., Gautier, N. M., Wang, S., Leslie, F. M., & Li, M. D. (2013). Gestational nicotine exposure modifies myelin gene expression in the brains of adolescent rats with sex differences. *Translational psychiatry*, *3*(4), e247-e247. https://doi.org/10.1038/tp.2013.21

Chen, T., & Guestrin, C. (2016, August). Xgboost: A scalable tree boosting system. In *Proceedings of the 22nd acm sigkdd international conference on knowledge discovery and data mining* (pp. 785-794). https://doi.org/10.1145/2939672.2939785

Chen, C. L., Kuo, M. C., Chen, P. Y., Tung, Y. H., Hsu, Y. C., Huang, C. W. C., ... & Tseng, W. Y. I. (2022). Validation of neuroimaging-based brain age gap as a mediator between modifiable

risk factors and cognition. *Neurobiology of Aging, 114,* 61-72.
https://doi.org/10.1016/j.neurobiolaging.2022.03.006
Cherbuin, N., Walsh, E. I., Shaw, M., Luders, E., Anstey, K. J., Sachdev, P. S., ... & Gaser, C.
(2021). Optimal blood pressure keeps our brains younger. *Frontiers in Aging Neuroscience,*
529. https://doi.org/10.3389/fnagi.2021.694982
Cetin-Karayumak, S., Di Biase, M. A., Chunga, N., Reid, B., Somes, N., Lyall, A. E., ... & Kubicki,
M. (2020). White matter abnormalities across the lifespan of schizophrenia: a harmonized
multi-site diffusion MRI study. *Molecular psychiatry, 25*(12), 3208-3219.
https://doi.org/10.1038/s41380-019-0509-y
Cole, J. H. (2020). Multimodality neuroimaging brain-age in UK biobank: relationship to biomedical,
lifestyle, and cognitive factors. *Neurobiology of aging, 92,* 34-42.
https://doi.org/10.1016/j.neurobiolaging.2020.03.014
Cole, J. H., & Franke, K. (2017). Predicting age using neuroimaging: innovative brain ageing
biomarkers. *Trends in neurosciences, 40*(12), 681-690.
https://doi.org/10.1016/j.tins.2017.10.001
Cole, J. H., Ritchie, S. J., Bastin, M. E., Hernández, V., Muñoz Maniega, S., Royle, N., ... & Deary,
I. J. (2018). Brain age predicts mortality. *Molecular psychiatry, 23*(5), 1385-1392.
https://doi.org/10.1038/mp.2017.62
Cox, S. R., Ritchie, S. J., Tucker-Drob, E. M., Liewald, D. C., Hagenaars, S. P., Davies, G., ... &
Deary, I. J. (2016). Ageing and brain white matter structure in 3,513 UK Biobank
participants. *Nature communications, 7*(1), 1-13. https://doi.org/10.1038/ncomms13629
Crossley, T. F., & Kennedy, S. (2002). The reliability of self-assessed health status. *Journal of health*
economics, 21(4), 643-658. https://doi.org/10.1016/S0167-6296(02)00007-3
de Lange, A. M. G., Anatürk, M., Suri, S., Kaufmann, T., Cole, J. H., Griffanti, L., ... & Ebmeier, K.
P. (2020a). Multimodal brain-age prediction and cardiovascular risk: The Whitehall II MRI
sub-study. *NeuroImage, 222,* 117292. https://doi.org/10.1016/j.neuroimage.2020.117292
de Lange, A. M. G., Barth, C., Kaufmann, T., Maximov, I. I., van der Meer, D., Agartz, I., &
Westlye, L. T. (2020b). Women's brain aging: Effects of sex-hormone exposure, pregnancies,
and genetic risk for Alzheimer's disease. *Human brain mapping, 41*(18), 5141-5150.
https://doi.org/10.1002/hbm.25180
de Lange, A. M. G., & Cole, J. H. (2020). Commentary: Correction procedures in brain-age
prediction. *NeuroImage: Clinical, 26.* https://doi.org/10.1016%2Fj.nicl.2020.102229
de Lange, A. M. G., Kaufmann, T., Quintana, D. S., Winterton, A., Andreassen, O. A., Westlye, L.
T., & Ebmeier, K. P. (2021). Prominent health problems, socioeconomic deprivation, and
higher brain age in lonely and isolated individuals: A population-based study. *Behavioural*
Brain Research, 414, 113510. https://doi.org/10.1016/j.bbr.2021.113510
de Lange, A. M. G., Anatürk, M., Rokicki, J., Han, L. K., Franke, K., Alnæs, D., ... & Cole, J. H.
(2022). Mind the gap: Performance metric evaluation in brain-age prediction. *Human Brain*
Mapping. https://doi.org/10.1002/hbm.25837
Denissen, S., Engemann, D. A., De Cock, A., Costers, L., Baijot, J., Laton, J., ... & Nagels, G.
(2022). Brain age as a surrogate marker for cognitive performance in multiple sclerosis.
European journal of neurology, 29(10), 3039-3049. https://doi.org/10.1111/ene.15473
Distelberg, B. J., Staack, A., Elsen, K. D. D., & Sabaté, J. (2017). The effect of coffee and caffeine
on mood, sleep, and health-related quality of life. *Journal of Caffeine Research, 7*(2), 59-70.
https://doi.org/10.1089/jcr.2016.0023
Eikenes, L., Visser, E., Vangberg, T., & Håberg, A. K. (2022). Both brain size and biological sex
contribute to variation in white matter microstructure in middle-aged healthy adults. *Human*
Brain Mapping. https://doi.org/10.1002/hbm.26093

Elliott, L. T., Sharp, K., Alfaro-Almagro, F., Shi, S., Miller, K. L., Douaud, G., ... & Smith, S. M. (2018). Genome-wide association studies of brain imaging phenotypes in UK Biobank. *Nature, 562*(7726), 210-216. https://doi.org/10.1038/s41586-018-0571-7

Elliott, M. L., Belsky, D. W., Knodt, A. R., Ireland, D., Melzer, T. R., Poulton, R., ... & Hariri, A. R. (2021). Brain-age in midlife is associated with accelerated biological aging and cognitive decline in a longitudinal birth cohort. *Molecular psychiatry, 26*(8), 3829-3838. https://doi.org/10.1038/s41380-019-0626-7

Engel, G. L. (1977). The need for a new medical model: a challenge for biomedicine. *Science, 196*(4286), 129-136. https://doi.org/10.1126/science.847460

Erhardt, L. (2009). Cigarette smoking: an undertreated risk factor for cardiovascular disease. *Atherosclerosis, 205*(1), 23-32. https://doi.org/10.1016/j.atherosclerosis.2009.01.007

Faragher, E. B., Cass, M., & Cooper, C. L. (2013). The relationship between job satisfaction and health: a meta-analysis. *From stress to wellbeing Volume 1*, 254-271.

Fieremans, E., Jensen, J. H., & Helpern, J. A. (2011). White matter characterization with diffusional kurtosis imaging. *Neuroimage, 58*(1), 177-188. https://doi.org/10.1016/j.neuroimage.2011.06.006

Kenny, P. J., & Markou, A. (2006). Nicotine self-administration acutely activates brain reward systems and induces a long-lasting increase in reward sensitivity. *Neuropsychopharmacology, 31*(6), 1203-1211. https://doi.org/10.1038/sj.npp.1300905

Le Foll, B., Piper, M. E., Fowler, C. D., Tonstad, S., Bierut, L., Lu, L., ... & Hall, W. D. (2022). Tobacco and nicotine use. *Nature Reviews Disease Primers, 8*(1), 1-16. https://doi.org/10.1038/s41572-022-00346-w

Fawns-Ritchie, C., & Deary, I. J. (2020). Reliability and validity of the UK Biobank cognitive tests. *PloS one, 15*(4), e0231627. https://doi.org/10.1371/journal.pone.0231627

Forte, G., De Pascalis, V., Favieri, F., & Casagrande, M. (2019). Effects of blood pressure on cognitive performance: A systematic review. *Journal of clinical medicine, 9*(1), 34. https://doi.org/10.3390/jcm9010034

Franke, K., Gaser, C., Manor, B., & Novak, V. (2013). Advanced BrainAGE in older adults with type 2 diabetes mellitus. *Frontiers in aging neuroscience, 5*, 90. https://doi.org/10.3389/fnagi.2013.00090

Ghaemi, S. N. (2009). The rise and fall of the biopsychosocial model. *The British Journal of Psychiatry, 195*(1), 3-4. https://doi.org/10.1192/bjp.bp.109.063859

Gill, D., Zuber, V., Dawson, J., Pearson-Stuttard, J., Carter, A. R., Sanderson, E., ... & Elliott, P. (2021). Risk factors mediating the effect of body mass index and waist-to-hip ratio on cardiovascular outcomes: Mendelian randomization analysis. *International Journal of Obesity, 45*(7), 1428-1438. https://doi.org/10.1038/s41366-021-00807-4

Gozzi, A., Schwarz, A., Reese, T., Bertani, S., Crestan, V., & Bifone, A. (2006). Region-specific effects of nicotine on brain activity: a pharmacological MRI study in the drug-naive rat. *Neuropsychopharmacology, 31*(8), 1690-1703. https://doi.org/10.1038/sj.npp.1300955

Gurholt, T. P., Kaufmann, T., Frei, O., Alnæs, D., Haukvik, U. K., van der Meer, D., ... & Andreassen, O. A. (2021). Population-based body–brain mapping links brain morphology with anthropometrics and body composition. *Translational psychiatry, 11*(1), 1-12. https://doi.org/10.1038/s41398-021-01414-7

Hamer, M., & Batty, G. D. (2019). Association of body mass index and waist-to-hip ratio with brain structure: UK Biobank study. *Neurology, 92*(6), e594-e600. https://doi.org/10.1212/WNL.0000000000006879

Higdon, J. V., & Frei, B. (2006). Coffee and health: a review of recent human research. *Critical reviews in food science and nutrition, 46*(2), 101-123. https://doi.org/10.1080/10408390500400009

Houenou, J., Wessa, M., Douaud, G., Leboyer, M., Chanraud, S., Perrin, M., ... & Paillere-Martinot, M. L. (2007). Increased white matter connectivity in euthymic bipolar patients: diffusion tensor tractography between the subgenual cingulate and the amygdalo-hippocampal complex. *Molecular psychiatry*, *12*(11), 1001-1010. https://doi.org/10.1038/sj.mp.4002010

Hsieh, C. M. (2001). Correlates of financial satisfaction. *The International Journal of Aging and Human Development*, *52*(2), 135-153. https://doi.org/10.2190/9YDE-46PA-MV9C-2JRB

Ingalhalikar, M., Smith, A., Parker, D., Satterthwaite, T. D., Elliott, M. A., Ruparel, K., ... & Verma, R. (2014). Sex differences in the structural connectome of the human brain. *Proceedings of the National Academy of Sciences*, *111*(2), 823-828. https://doi.org/10.1073/pnas.1316909110

Jenkinson, M., Beckmann, C. F., Behrens, T. E., Woolrich, M. W., & Smith, S. M. (2012). Fsl. *Neuroimage*, *62*(2), 782-790. https://doi.org/10.1016/j.neuroimage.2011.09.015

Jensen, J. H., Helpern, J. A., Ramani, A., Lu, H., & Kaczynski, K. (2005). Diffusional kurtosis imaging: the quantification of non-gaussian water diffusion by means of magnetic resonance imaging. *Magnetic Resonance in Medicine: An Official Journal of the International Society for Magnetic Resonance in Medicine*, *53*(6), 1432-1440. https://doi.org/10.1002/mrm.20508

Kaden, E., Kelm, N. D., Carson, R. P., Does, M. D., & Alexander, D. C. (2016a). Multi-compartment microscopic diffusion imaging. *NeuroImage*, *139*, 346-359. https://doi.org/10.1016/j.neuroimage.2016.06.002

Jirsaraie, R. J., Kaufmann, T., Bashyam, V., Erus, G., Luby, J. L., Westlye, L. T., ... & Sotiras, A. (2022). Benchmarking the generalizability of brain age models: Challenges posed by scanner variance and prediction bias. *Human Brain Mapping*. https://doi.org/10.1002/hbm.26144

Kahn, S. E., Hull, R. L., & Utzschneider, K. M. (2006). Mechanisms linking obesity to insulin resistance and type 2 diabetes. *Nature*, *444*(7121), 840-846. https://doi.org/10.1038/nature05482

Kaden, E., Kruggel, F., & Alexander, D. C. (2016b). Quantitative mapping of the per-axon diffusion coefficients in brain white matter. *Magnetic resonance in medicine*, *75*(4), 1752-1763. https://doi.org/10.1002/mrm.25734

Kamagata, K., Andica, C., Hatano, T., Ogawa, T., Takeshige-Amano, H., Ogaki, K., ... & Aoki, S. (2020). Advanced diffusion magnetic resonance imaging in patients with Alzheimer's and Parkinson's diseases. *Neural regeneration research*, *15*(9), 1590. https://doi.org/10.4103/1673-5374.276326

Kaufmann, T., van der Meer, D., Doan, N. T., Schwarz, E., Lund, M. J., Agartz, I., ... & Westlye, L. T. (2019). Common brain disorders are associated with heritable patterns of apparent aging of the brain. *Nature neuroscience*, *22*(10), 1617-1623. https://doi.org/10.1038/s41593-019-0471-7

Kellner, E., Dhital, B., Kiselev, V. G., & Reisert, M. (2016). Gibbs-ringing artifact removal based on local subvoxel-shifts. *Magnetic resonance in medicine*, *76*(5), 1574-1581. https://doi.org/10.1002/mrm.26054

Korbmacher, M., de Lange, A. M., van der Meer, D., Beck, D., Eikefjord, E. N., Lundervold, A., ... & Maximov, I. I. (2022). Brain-wide associations between white matter and age highlight the role of fornix microstructure in brain age. *bioRxiv*. https://doi.org/10.1101/2022.09.29.510029

Kotsis, V., Stabouli, S., Papakatsika, S., Rizos, Z., & Parati, G. (2010). Mechanisms of obesity-induced hypertension. *Hypertension research*, *33*(5), 386-393. https://doi.org/10.1038/hr.2010.9

Lawrence, K. E., Nabulsi, L., Santhalingam, V., Abaryan, Z., Villalon-Reina, J. E., Nir, T. M., ... & Thompson, P. M. (2021). Age and sex effects on advanced white matter microstructure measures in 15,628 older adults: a UK biobank study. *Brain imaging and behavior*, *15*(6), 2813-2823. https://doi.org/10.1007/s11682-021-00548-y

787   Lehman, B. J., David, D. M., & Gruber, J. A. (2017). Rethinking the biopsychosocial model of
788   health: Understanding health as a dynamic system. *Social and personality psychology*
789   *compass*, *11*(8), e12328. https://doi.org/10.1111/spc3.12328
790   Leonardsen, E. H., Peng, H., Kaufmann, T., Agartz, I., Andreassen, O. A., Celius, E. G., ... & Wang,
791   Y. (2022). Deep neural networks learn general and clinically relevant representations of the
792   ageing brain. *NeuroImage*, *256*, 119210. https://doi.org/10.1016/j.neuroimage.2022.119210
793   Liem, F., Varoquaux, G., Kynast, J., Beyer, F., Masouleh, S. K., Huntenburg, J. M., ... & Margulies,
794   D. S. (2017). Predicting brain-age from multimodal imaging data captures cognitive
795   impairment. *Neuroimage, 148*, 179-188. https://doi.org/10.1016/j.neuroimage.2016.11.005
796   Lim, N., Kim, E. K., Kim, H., Yang, E., & Lee, S. M. (2010). Individual and work-related factors
797   influencing burnout of mental health professionals: A meta-analysis. *Journal of Employment*
798   *Counseling, 47*(2), 86-96. https://doi.org/10.1002/j.2161-1920.2010.tb00093.x
799   Lohner, V., Pehlivan, G., Sanroma, G., Miloschewski, A., Schirmer, M. D., Stöcker, T., ... &
800   Breteler, M. M. (2022). Relation Between Sex, Menopause, and White Matter
801   Hyperintensities: The Rhineland Study. *Neurology, 99*(9), e935-e943.
802   https://doi.org/10.1212/WNL.0000000000200782
803   Marquand, A. F., Kia, S. M., Zabihi, M., Wolfers, T., Buitelaar, J. K., & Beckmann, C. F. (2019).
804   Conceptualizing mental disorders as deviations from normative functioning. *Molecular*
805   *psychiatry*, *24*(10), 1415-1424. https://doi.org/10.1038/s41380-019-0441-1
806   Mathew, B., Francis, L., Kayalar, A., & Cone, J. (2008). Obesity: effects on cardiovascular disease
807   and its diagnosis. *The Journal of the American Board of Family Medicine, 21*(6), 562-568.
808   https://doi.org/10.3122/jabfm.2008.06.080080
809   Maximov, I. I., Alnæs, D., & Westlye, L. T. (2019). Towards an optimised processing pipeline for
810   diffusion magnetic resonance imaging data: Effects of artefact corrections on diffusion
811   metrics and their age associations in UK Biobank. *Human Brain Mapping*, *40*(14), 4146-
812   4162. https://doi.org/10.1002/hbm.24691
813   Maximov, I. I., van Der Meer, D., de Lange, A. M. G., Kaufmann, T., Shadrin, A., Frei, O., ... &
814   Westlye, L. T. (2021). Fast qualitY conTrol meThod foR derIved diffUsion Metrics
815   (YTTRIUM) in big data analysis: UK Biobank 18,608 example. *Human brain mapping,*
816   *42*(10), 3141-3155. https://doi.org/10.1002/hbm.25424
817   Meissner, A. (2016). Hypertension and the brain: a risk factor for more than heart disease.
818   *Cerebrovascular diseases*, *42*(3-4), 255-262. https://doi.org/10.1159/000446082
819   Miller, K. L., Alfaro-Almagro, F., Bangerter, N. K., Thomas, D. L., Yacoub, E., Xu, J., ... & Smith,
820   S. M. (2016). Multimodal population brain imaging in the UK Biobank prospective
821   epidemiological study. *Nature neuroscience*, *19*(11), 1523-1536.
822   https://doi.org/10.1038/nn.4393
823   Molarius, A., Seidell, J. C., Sans, S., Tuomilehto, J., & Kuulasmaa, K. (1999). Waist and hip
824   circumferences, and waist-hip ratio in 19 populations of the WHO MONICA Project.
825   *International journal of obesity and related metabolic disorders : journal of the International*
826   *Association for the Study of Obesity*, *23*(2), 116–125. https://doi.org/10.1038/sj.ijo.0800772
827   Mori, S., Wakana, S., Van Zijl, P. C., & Nagae-Poetscher, L. M. (2005). *MRI atlas of human white*
828   *matter*. Elsevier.
829   Mosconi, L., Berti, V., Dyke, J., Schelbaum, E., Jett, S., Loughlin, L., ... & Brinton, R. D. (2021).
830   Menopause impacts human brain structure, connectivity, energy metabolism, and amyloid-
831   beta deposition. *Scientific reports*, *11*(1), 1-16. https://doi.org/10.1038/s41598-021-90084-y
832   Nehlig, A. (2016). Effects of coffee/caffeine on brain health and disease: What should I tell my
833   patients?. *Practical neurology, 16*(2), 89-95. http://dx.doi.org/10.1136/practneurol-2015-
834   001162

Ning, K., Zhao, L., Matloff, W., Sun, F., & Toga, A. W. (2020). Association of relative brain age
with tobacco smoking, alcohol consumption, and genetic variants. *Scientific reports*, *10*(1), 1-
10. https://doi.org/10.1038/s41598-019-56089-4
Niu, X., Zhang, F., Kounios, J., & Liang, H. (2019). Improved prediction of brain age using
multimodal neuroimaging data. *Human brain mapping*, *41*(6), 1626-1643.
https://doi.org/10.1002/hbm.24899
Nowok, B., Raab, G. M., & Dibben, C. (2016). synthpop: Bespoke creation of synthetic data in R.
Journal of statistical software, *74*, 1-26. https://doi.org/10.18637/jss.v074.i11
Pavlakis, A. E., Noble, K., Pavlakis, S. G., Ali, N., & Frank, Y. (2015). Brain imaging and
electrophysiology biomarkers: is there a role in poverty and education outcome research?.
Pediatric Neurology, *52*(4), 383-388. https://doi.org/10.1016/j.pediatrneurol.2014.11.005
Pham, K., Mulugeta, A., Zhou, A., O'Brien, J. T., Llewellyn, D. J., & Hyppönen, E. (2022). High
coffee consumption, brain volume and risk of dementia and stroke. *Nutritional Neuroscience*,
25(10), 2111-2122. https://doi.org/10.1080/1028415X.2021.1945858
Pines, A. R., Cieslak, M., Larsen, B., Baum, G. L., Cook, P. A., Adebimpe, A., ... & Satterthwaite, T.
D. (2020). Leveraging multi-shell diffusion for studies of brain development in youth and
young adulthood. *Developmental cognitive neuroscience*, *43*, 100788.
https://doi.org/10.1016/j.dcn.2020.100788
Peng, H., Gong, W., Beckmann, C. F., Vedaldi, A., & Smith, S. M. (2021). Accurate brain age
prediction with lightweight deep neural networks. *Medical image analysis*, *68*, 101871.
https://doi.org/10.1016/j.media.2020.101871
Reisert, M., Kellner, E., Dhital, B., Hennig, J., & Kiselev, V. G. (2017). Disentangling micro from
mesostructure by diffusion MRI: a Bayesian approach. *Neuroimage*, *147*, 964-975.
https://doi.org/10.1016/j.neuroimage.2016.09.058
Remiszewski, N., Bryant, J. E., Rutherford, S. E., Marquand, A. F., Nelson, E., Askar, I., ... &
Kraguljac, N. V. (2022). Contrasting Case-Control and Normative Reference Approaches to
Capture Clinically Relevant Structural Brain Abnormalities in Patients With First-Episode
Psychosis Who Are Antipsychotic Naive. *JAMA psychiatry*.
https://doi.org/10.1001/jamapsychiatry.2022.3010
Reychav, I., Beeri, R., Balapour, A., Raban, D. R., Sabherwal, R., & Azuri, J. (2019). How reliable
are self-assessments using mobile technology in healthcare? The effects of technology
identity and self-efficacy. *Computers in Human Behavior*, *91*, 52-61.
https://doi.org/10.1016/j.chb.2018.09.024
Rokicki, J., Wolfers, T., Nordhøy, W., Tesli, N., Quintana, D. S., Alnæs, D., ... & Westlye, L. T.
(2021). Multimodal imaging improves brain age prediction and reveals distinct abnormalities
in patients with psychiatric and neurological disorders. *Human brain mapping*, *42*(6), 1714-
1726. https://doi.org/10.1002/hbm.25323
Salih, A., Boscolo Galazzo, I., Raisi-Estabragh, Z., Rauseo, E., Gkontra, P., Petersen, S. E., ... &
Menegaz, G. (2021). Brain age estimation at tract group level and its association with daily
life measures, cardiac risk factors and genetic variants. *Scientific reports*, *11*(1), 1-14.
https://doi.org/10.1038/s41598-021-99153-8
Shaked, D., Leibel, D. K., Katzel, L. I., Davatzikos, C., Gullapalli, R. P., Seliger, S. L., ... &
Waldstein, S. R. (2019). Disparities in diffuse cortical white matter integrity between
socioeconomic groups. *Frontiers in Human Neuroscience*, *13*, 198.
https://doi.org/10.3389/fnhum.2019.00198
Smith, S. M. (2002). Fast robust automated brain extraction. *Human brain mapping*, *17*(3), 143-155.
https://doi.org/10.1002/hbm.10062

Smith, S. M., Vidaurre, D., Alfaro-Almagro, F., Nichols, T. E., & Miller, K. L. (2019). Estimation of
brain age delta from brain imaging. *Neuroimage*, *200*, 528-539.
https://doi.org/10.1016/j.neuroimage.2019.06.017
Smith, S. M., Jenkinson, M., Johansen-Berg, H., Rueckert, D., Nichols, T. E., Mackay, C. E., ... &
Behrens, T. E. (2006). Tract-based spatial statistics: voxelwise analysis of multi-subject
diffusion data. *Neuroimage*, *31*(4), 1487-1505.
https://doi.org/10.1016/j.neuroimage.2006.02.024
Smith, S. M., Jenkinson, M., Woolrich, M. W., Beckmann, C. F., Behrens, T. E., Johansen-Berg,
H., ... & Matthews, P. M. (2004). Advances in functional and structural MR image analysis
and implementation as FSL. *Neuroimage*, *23*, S208-S219.
https://doi.org/10.1016/j.neuroimage.2004.07.051
Sone, D., Beheshti, I., Shinagawa, S., Niimura, H., Kobayashi, N., Kida, H., ... & Shigeta, M. (2022).
Neuroimaging-derived brain age is associated with life satisfaction in cognitively unimpaired
elderly: A community-based study. *Translational psychiatry*, *12*(1), 1-6.
https://doi.org/10.1038/s41398-022-01793-5
Subramaniapillai, S., Suri, S., Barth, C., Maximov, I. I., Voldsbekk, I., van der Meer, D., ... & de
Lange, A. M. G. (2022). Sex-and age-specific associations between cardiometabolic risk and
white matter brain age in the UK Biobank cohort. *Human brain mapping*, *43*(12), 3759-3774.
https://doi.org/10.1002/hbm.25882
Sudlow, C., Gallacher, J., Allen, N., Beral, V., Burton, P., Danesh, J., ... & Collins, R. (2015). UK
biobank: an open access resource for identifying the causes of a wide range of complex
diseases of middle and old age. *PLoS medicine*, *12*(3), e1001779.
https://doi.org/10.1371/journal.pmed.1001779
van der Harten, B., de Leeuw, F. E., Weinstein, H. C., Scheltens, P., & Biessels, G. J. (2006). Brain
imaging in patients with diabetes: a systematic review. *Diabetes care*, *29*(11), 2539-2548.
https://doi.org/10.2337/dc06-1637
Veraart, J., Fieremans, E., & Novikov, D. S. (2016). Diffusion MRI noise mapping using random
matrix theory. *Magnetic resonance in medicine*, *76*(5), 1582-1593.
https://doi.org/10.1002/mrm.26059
Vidal-Pineiro, D., Wang, Y., Krogsrud, S. K., Amlien, I. K., Baaré, W. F., Bartres-Faz, D., ... & Fjell,
A. (2021). Individual variations in 'brain age'relate to early-life factors more than to
longitudinal brain change. *Elife*, *10*, e69995. https://doi.org/10.7554/eLife.69995
Wade, D. T., & Halligan, P. W. (2017). The biopsychosocial model of illness: a model whose time
has come. *Clinical rehabilitation*, *31*(8), 995-1004.
https://doi.org/10.1177/0269215517709890
Westlye, L. T., Walhovd, K. B., Dale, A. M., Bjørnerud, A., Due-Tønnessen, P., Engvig, A., ... &
Fjell, A. M. (2010). Lifespan changes of the human brain white matter: diffusion tensor
imaging (DTI) and volumetry. *Cerebral cortex*, *20*(9), 2055-2068.
https://doi.org/10.1093/cercor/bhp280
Wrighten, S. A., Piroli, G. G., Grillo, C. A., & Reagan, L. P. (2009). A look inside the diabetic brain:
Contributors to diabetes-induced brain aging. *Biochimica et Biophysica Acta (BBA)-*
Molecular Basis of Disease, *1792*(5), 444-453. https://doi.org/10.1016/j.bbadis.2008.10.013
Wood, D. A., Kafiabadi, S., Al Busaidi, A., Guilhem, E., Montvila, A., Lynch, J., ... & Booth, T. C.
(2022). Accurate brain-age models for routine clinical MRI examinations. *NeuroImage*, *249*,
118871. https://doi.org/10.1016/j.neuroimage.2022.118871
Zhao, B., Zhang, J., Ibrahim, J. G., Luo, T., Santelli, R. C., Li, Y., ... & Zhu, H. (2021). Large-scale
GWAS reveals genetic architecture of brain white matter microstructure and genetic overlap
with cognitive and mental health traits (n= 17,706). *Molecular psychiatry*, *26*(8), 3943-3955.
https://doi.org/10.1038/s41380-019-0569-z

6   Manuscript Contents

6.1   Figures

Figure 1. Overview of the Variables Used

Figure 2. Overview of the Statistical Models Used

Figure 3. Marginal R^2 Values for Statistical Models across Diffusion Approaches

Figure 4. Sociodemographic Model Predictors with 95% Confidence Interval

* indicates Bonferroni-corrected $p < .05$

Figure 5. Health Model Predictors with 95% Confidence Interval

* indicates Bonferroni-corrected $p < .05$

Figure 6. Well-being Model Predictors with 95% Confidence Interval

* indicates Bonferroni-corrected $p < .05$

Figure 7. Cognition Model Predictors with 95% Confidence Interval

* indicates Bonferroni-corrected $p < .05$

6.2 Tables

Table 1: Linear models relating multimodal brain age to bio-psycho-social factors

Variable	Level or Metric	Variable value	BAG[1]	N[2]	Marginal R^2 diff	Log Likelihood$_{diff}$	χ^2 p_{diff}[4]	β[3]	p_{pred}[4]
Brain age	Mean ± SD	-0.003 ± 4.534	-0.003	32174					
Demographics									
Scanner Site	% Cheadle	57.559	-0.011	18519					
	% Newcastle	26.403	0.090	8495					
	% Reading	15.913	-0.129	5120					
	% Bristol	0.124	-0.148	40					
Sex	% Male	47.122	0.259	17013				-1.07	.001
	% Female	52.878	-0.237	15161					
Age	Mean ± SD	64.473 ± 7.614	-0.003	32174				0.62	<.001
Socio-demographics									
Ethnicity	European	96.800	-0.034	31160	-1.8 x 10^{-5}				
	Non-European	2.970	1.010	956	-1.8 x 10^{-5}	4	.007	-0.31	.007
	Prefer not to say	(0.180)	-0.023	58					
Income[5]	% less £18k	10.363	-0.529	3310	1.4 x 10^{-4}				
	% £18k-£30	24.253	-0.874	7747	1.4 x 10^{-4}	11	<.001	-0.28	<.001
	% £30k-£52k	27.713	-0.107	8852	1.4 x 10^{-4}	11	<.001	-0.29	<.001
	% £52k-100k	21.586	1.220	6895	1.4 x 10^{-4}	11	<.001	-0.18	.016
	% > £100k	6.956	2.070	2222	1.4 x 10^{-4}	11	<.001	-0.30	.003
	Don't know	3.040	-1.20	972					
	Prefer not to say	6.086	-1.16	1944					
Higher education	% Yes	49.326	0.215	15870	5.2 x 10^{-5}	1	.177	5.3 x 10^{-2}	.176
	% No	50.674	-0.215	16304	5.2 x 10^{-5}	1	.177		
Cognitive test scores									
Matrix puzzles solved	Mean ± SD	8.011 ± 0.500	0.127	21755	6.8 x 10^{-4}	27	<.001	-0.08	<.001
Tower rearranging correct attempts	Mean ± SD	9.920 ± 2.123	0.139	21587	5.0 x 10^{-4}	17	<.001	-0.04	<.001
Prospective memory	Mean ± SD	1.067 ± 0.397	0.054	30300	-1.7 x 10^{-6}	0	.312	-0.05	.311
Fluid intelligence	Mean ± SD	6.631 ± 0.397	0.075	29786	7.3 x 10^{-4}	23	<.001	-0.07	<.001
Digits remembered	Mean ± SD	6.675 ± 1.540	0.082	23070	1.6 x 10^{-4}	15	<.001	-0.06	<.001
Mean number of incorrect pair matches across trials	Mean ± SD	2.214 ± 1.279	0.097	20770	2.9 x 10^{-4}	3	.014	-0.05	.015
Life satisfaction[6]									
Job satisfaction	Mean ± SD	4.511 ± 0.863	1.069	18399	-5.5 x 10^{-6}	1	.494	0.02	.494
Financial	Mean ± SD	4.714 ± 0.828	-0.0004	31909	2.6 x 10^{-4}	10	<.001	-0.10	<.001

satisfaction Health satisfaction	Mean ± SD	4.470 ± 0.766	-0.0005	31911	0.001	52	<.001	-0.26	<.001
Overall health rating	Mean ± SD	3.030 ± 0.630	0.0007	31934	0.001	66	<.001	-0.36	<.001
Family relation satisfaction	Mean ± SD	4.814 ± 0.846	0.002	31737	2.7×10^{-4}	17	<.001	-0.10	<.001
Friend relationship satisfaction	Mean ± SD	4.784 ± 0.846	0.0003	31640	2.0×10^{-6}	0	.485	-0.02	.485
Happiness	Mean ± SD	4.542 ± 0.686	-0.004	31884	1.2×10^{-4}	3	.011	-0.07	.011
health and lifestyle factors									
BMI	Mean ± SD	26.319 ± 4.269	-0.011	31052	5.9×10^{-4}	41	<.001	0.04	<.001
Pulse pressure	Mean ± SD	60.027 ± 14.540	-0.014	28184	0.002	66	<.001	0.02	<.001
WHR	Mean ± SD	0.872 ± 0.088	-0.015	31138	0.004	129	<.001	4.86	<.001
Smoking	% Yes	2.629	1.570	838	7.5×10^{-5}	4	.005	0.34	.006
	% No	97.374	-0.0368	31033	7.5×10^{-5}	4	.005		
Diabetes	% Yes	1.688	0.494	543	5.1×10^{-4}	22	<.001	0.99	<.001
	% No	98.312	-0.012	31631	5.1×10^{-4}	22	<.001		
Hypertension	% Yes	19.680	-0.215	6332	0.002	151	<.001	0.86	<.001
	% No	80.320	0.049	25842	0.002	151	<.001		
High Cholesterol	% Yes	12.202	-1.07	3926	-7.4×10^{-5}	18	<.001	0.26	<.001
	% No	83.798	0.145	28248	-7.4×10^{-5}	18	<.001		
Vascular Diagnosis	% Yes	22.726	-0.386	7312	0.002	136	<.001	0.78	<.001
	% No	87.274	0.109	24862	0.002	136	<.001		
Birth weight (kg)	Mean ± SD	3.358 ± 0.619	0.386	19409	3.0×10^{-4}	10	<.001	-0.18	<.001
Daily coffee intake (cups)	Mean ± SD	2.065 ± 1.815	0.001	31973	3.0×10^{-4}	20	<.001	0.06	<.001

Note: Linear models were used to observe each of the bio-psycho-social variables of interest in a model of [brain age ~ age + sex + age*sex + bio-psycho-social variable] with scanner site as a random factor. The model was then compared to a baseline model of [brain age ~ age + sex + age*sex] with scanner site as a random factor, and hence, R^2 values refer to variance uniquely contributed by the single bio-psycho-social variable when added to the baseline model, referring to the marginal R^2. For sex and age, fixed effects, and site, the random effect are being presented from the baseline model.

1 Brain age gap (BAG) values refer here to the differences between model predictions, i.e., the brain age and the chronological age. Importantly, we used an uncorrected BAG. These values might differ because of missing data in the associated bio-psycho-social factors and will be influenced by the model's age bias.

2 Sample sizes differ due to missing data in the bio-psycho-social variables.

3 p_{pred} refers to the p-values of the predictors/bio-psycho-social variables' β, p_{diff} refers to the differences between baseline model and the respective model including the bio-psycho-social variable of interest.

4 β values of the bio-psycho-social variables of interest.

5 Likert-type scales were applied for self-rating scales ranging from 1 "extremely unhappy" to 6 "extremely happy" with the exception of overall health ranging from 1 "poor" to 4 "excellent".

962   Table 2: Performance of Brain Age Prediction Models

Approach§	Number of MRI features	R^2 (SD)	RMSE (SD)	MAE (SD)	Prediction-Age Correlation* [95% CI]
BRIA	690	0.550 (0.012)	5.007 (0.057)	4.002 (0.042)	0.742 [0.737, 0.747]
DKI	207	0.576 (0.015)	4.958 (0.077)	3.975 (0.068)	0.754 [0.755, 0.764]
DTI	276	0.571 (0.014)	4.983 (0.072)	3.984 (0.062)	0.756 [0.751, 0.761]
SMT	276	0.531 (0.010)	5.214 (0.053)	4.183 (0.036)	0.729 [0.724, 0.734]
mcSMT	276	0.519 (0.011)	5.175 (0.045)	4.153 (0.036)	0.721 [0.716, 0.726]
WMTI	207	0.585 (0.012)	4.903 (0.065)	3.928 (0.050)	0.765 [0.761, 0.770]
Mean	28	0.393 (0.012)	5.932 (0.051)	4.812 (0.046)	0.627 [0.621, 0.634]
Full	1932	0.645 (0.011)	4.534 (0.041)	3.624 (0.037)	0.804 [0.800, 0.808]

Table logic: R^2, RMSE, MAE are displayed in the format Mean (Standard Deviation), Pearson's correlations are
displayed in the format Correlation Score 95% Confidence Interval [Lower Bound, Upper Bound].

Mean refers to the mean multimodal model calculated on diffusion metrics averaged over the skeleton for all six diffusion
approaches. Full refers to the full multimodal model based on all diffusion data from the six diffusion approaches, i.e.
mean multimodal data in addition to metrics averaged over the JHU atlas regions. R^2 = variance explained, RMSE = root
mean squared error, MAE = mean absolute error.

§ For an overview of the metrics contained in each of the diffusion approaches see Supplementary Table 1.

* All correlation were significant at *p* < .001.

7 Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

8 Author Contributions

M.K.: Study design, Software, Formal analysis, Visualizations, Project administration, Writing – original draft, Writing – review & editing
T.P.G. Writing – review & editing
A.M.d.L.: Software, Writing – review & editing
D.v.d.M.: Software, Writing – review & editing
A.L.: Writing – review & editing, Funding acquisition
E.E.: Writing – review & editing, Funding acquisition
D.B.: Writing – review & editing
O.A.A.: Writing – review & editing, Funding acquisition
L.T.W.: Writing – review & editing, Funding acquisition
I.I.M. supervision, Study design, Data pre-processing and quality control, Writing – review & editing, Funding acquisition

9 Funding

This research was funded by the Research Council of Norway (#223273).

10 Acknowledgments

This study has been conducted using UKB data under Application 27412. UKB has received ethics approval from the National Health Service National Research Ethics Service (ref 11/NW/0382). The work was performed on the Service for Sensitive Data (TSD) platform, owned by the University of Oslo, operated and developed by the TSD service group at the University of Oslo IT-Department (USIT). Computations were performed using resources provided by UNINETT Sigma2 – the National Infrastructure for High Performance Computing and Data Storage in Norway. Finally, we want to thank all UKB participants and facilitators who made this research possible.

11 Data Availability Statement

All raw data are available from the UK Biobank (www.ukbiobank.ac.uk). Synthetic datasets of the diffusion MRI data were created with the synthpop R package (Nowok et al., 2016) based on the original data for all six diffusion approaches (resulting in six datasets) which are openly available at the Open Science Framework: (https://osf.io/nv8ea/). Furthermore, a synthetic version of the data used to analyze group differences and associations of brain age with bio-psycho-social factors can be found here: https://osf.io/nk43a/. Synthetic datasets are simulated datasets closely mimicking the statistical characteristics of the original data while protecting data privacy and anonymity. Analysis code can be found on GitHub (https://github.com/MaxKorbmacher/Brain-Age-Variability).

Bio-Psycho-Social Factors observed in the Context of Brain Age

Brain Age

Biological Dimension

Health and Lifestyle

- BMI
- WHR
- Pulse pressure
- Diabetes
- Smoking
- High cholesterol
- Diagnosed vascular problem
- Birth weight
- Sleeping hours
- Daily coffee intake

Social Dimension

Sociodemographics

- Age
- Sex
- Ethnicity
- Income
- Education
- Number of social visits

Psychological Dimension

Cognitive Scores

- Prospective Memory
- Fluid Intelligence
- Symbol Digit Substitution
- Pair Matches
- Matrix Puzzles
- Tower Arranging
- Digit Memorisation

Life Satisfaction

- Financial & Job
- Health
- Relationships
- Happiness

Modelling Brain Age from Bio-Psycho-Social Factors

Baseline Model
- Age
- Sex
- Scanner Site

Model 1
Sociodemographics

- Age
- Sex
- Scanner Site
- Ethnicity
- Income
- Education
- Number of Social Visits

Model 2
Cognitive Scores

- Age
- Sex
- Scanner Site
- Prospective Memory
- Fluid Intelligence
- Symbol Digit Substitution
- Pair Matches
- Matrix Puzzles
- Tower Arranging
- Digit Memorization

Model 2
Life Satisfaction

- Age
- Sex
- Scanner Site
- Financial & Job
- Health Rating & Satisf.
- Friend & Family Relation Satisf.
- Happiness

Model 4
Health and Lifestyle

- Age
- Sex
- Scanner Site
- BMI
- WHR
- Pulse Pressure
- Diabetes
- Smoking
- High Cholesterol
- Diagnosed Vascular Probl.
- Birth Weight
- Sleeping Hours
- Daily Coffee Intake