# Establishing Causality in the Relationship between Sleep and Migraine in a Global Sample: A Bayesian Approach

Emily C. Stanyer[1], Jack Brookes[2], Jia Rong Pang[3], Alexandre Urani[3], Philip R. Holland[1], Jan Hoffmann[1,4*]

[1]Wolfson Centre for Age-related Diseases, Institute for Psychiatry, Psychology, and Neuroscience, King's College London, London, United Kingdom

[2]Independent Researcher

[3]Healint Pte Ltd.

[4]NIHR-Wellcome Trust King's Clinical Research Facility/SLaM Biomedical Research Centre, King's College Hospital, London, United Kingdom

*Corresponding author: Dr Jan Hoffmann, jan.hoffmann@kcl.ac.uk

# Abstract

**Background:** There is a bidirectional link between sleep and migraine, however causality is difficult to determine. Previous studies rely on retrospective questionnaires, and small samples to support their findings. This study aimed to overcome this by teasing apart this relationship using sleep assessment and concomitant migraine data collected from a smartphone application using Bayesian modelling.

**Methods:** Anonymized self-reported data on sleep and migraine from 11,166 global users (aged 18-81 years, mean: 41.21, standard deviation: 11.49) were collected from the Migraine Buddy application (Healint Pte. Ltd.) between 30th June and 31st December 2021. Measures included: demographics, start and end times of each sleep episode and migraine attack, and the pain intensity for each migraine attack (visual analogue scale 0-10). Bayesian regression models were used to predict occurrence of a migraine attack the next day based on users' deviations from mean monthly sleep, number of sleep interruptions, and hours slept the night before in those reporting ≥4 and <25 migraine attacks on average per month. Conversely, we modelled whether attack occurrence and pain intensity could predict hours slept that same night.

**Results**: Once exclusion criteria were applied, there were 724 users (129 males, 412 females, 183 unknown) with an average age of 41.88 years ($SD = 11.63$), with a mean monthly number of attacks of 9.94. A greater number of sleep interruptions (95% Highest Density Interval (95% HDI [0.112 – 0.205]) and deviation from a user's mean sleep the night before (95% HDI [0.040 – 0.080]) were significant predictors of a next day migraine attack. Total hours slept was not a significant predictor (95% HDI [-0.04

– 0.04]). Pain intensity, but not attack occurrence was a positive predictor of hours

slept.

**Discussion:** Sleep fragmentation and deviation from typical sleep are the main drivers

of the relationship between sleep and migraine, whereas overall sleep duration is not.

Conversely, simply having a migraine attack does not predict sleep duration, it is the

pain associated with an attack which alters sleep. This study has shed light on the

causal mechanisms of sleep and migraine and highlights sleep hygiene as crucial in

migraine management.

## Introduction

Both clinically and preclinically, there is an established relationship between sleep and

migraine[1,2], and sleep and pain more generally[3]. For example, migraine attacks are

often preceded by premonitory symptoms, the most common of which is fatigue[4].

Paradoxically, sleep disruption, both too little or too much is reported to trigger attacks

in approximately 50% of patients[5] and represents a key risk factor for chronification

(progression to >15 attacks per month)[6]. Conversely, patients report sleep as a

strategy for attack resolution[7]. This relationship is further complicated by the fact that

many migraine treatments affect the sleep cycle[8] and affective disorders involving

sleep disturbance are three times more likely in people with migraine than the general

population[9]. In a recent meta-analysis, we reported poorer subjective sleep quality and

reduced rapid-eye-movement (REM) sleep in migraine patients compared to healthy

controls[10]. However, what remains unanswered is whether poor sleep increases the

likelihood of migraine attacks or whether having an attack causes disrupted sleep. This

is important as migraine is a severely disabling disorder which impacts more than one

billion people globally[11] and current treatments are ineffective in many patients[12], thus

understanding the relationship with sleep could lead to more targeted prevention.

Previous studies have small samples[13,14] or focus solely on one specific population at

one time point[15,16] and provide conflicting findings. For example, hours slept on the

previous night is predictive of pain[17] and migraine[14] the next day. Conversely, sleep

disturbance but not reduced hours slept has been shown to increase pain[18], and is

associated with higher odds of migraine the next day[19]. Fewer studies have

investigated the impact of migraine attacks on subsequent sleep. However, numerous

studies have demonstrated that pain generally during both the day and night results in

poor subsequent sleep and sleep quality[20–23].

The current study aimed to reconcile this disparity and investigate the relationship

between sleep and migraine using a longitudinal sample from across 99 different

countries collected from a smartphone application. Moreover, we aimed to tease apart

causality and establish whether sleep disruption predicts occurrence of attacks or

alternatively whether attacks predict disrupted sleep. This analysis differs from

previous literature as it avoids issues with retrospective reporting, as users report

migraine attacks as and when they happen, and sleep variables are automatically

detected by the application based on usage of the mobile phone and confirmed by the

user the next morning.

We hypothesize, in line with previous literature on sleep fragmentation, that more

sleep interruptions the night before will predict the occurrence of a migraine attack the

next day, but the number of hours slept will not. Similarly, we hypothesize that

deviation from average monthly sleep will predict the occurrence of an attack the next

day. On the other hand, we hypothesize that experiencing a migraine attack and the

pain intensity will both predict a reduction in hours slept compared to average sleep

on the evening on which the attack begins.

## Methods

### Design and inclusion criteria

This was a retrospective cross-sectional study on self-reported data gathered from users of the Migraine Buddy mobile application (Healint Pte. Ltd., https://healint.com). Data were collected between 30th June 2021 and 31st December 2021. Only application users which were active for at least 6 months and reported an activity in each month during the collection period were included in the dataset. 6 months' worth of data was chosen as it is unlikely many users would progress to chronic migraine during this short period, and it also avoids issues with potential seasonal migraine triggers and circannual periodicity[24]. Users were required to have signed up to the application before the 7th January 2021 and had to have 25 or more days of user-confirmed sleep records in each month during the period. There were no restrictions on the geographical location, age, or gender of the participants and these were optional for the participant to report within the application. No ethical review board was required due to there being no user-identifiable information collected or stored. When users signed up to the application they agreed to the terms and conditions of the Healint Pte. Ltd. policy of data collection and disclosure.

### Data Transformation and pre-processing

All data pre-processing and statistical analysis was conducted in RStudio (Version 2022.02.3). The following self-reported variables were gathered by the application: demographics (if reported: age, gender, country), start and end times of each sleep episode, and start and end times of each reported migraine attack, whether an attack was thought to be triggered by menstruation by the participant (yes/no), the pain intensity for each migraine attack (visual analogue scale from 0 to 10), and the symptoms experienced with each attack. For the purposes of demographic reporting,

users were categorised by age as follows: young adults <40 years, and older adults

≥40 years. Participants could choose from a range of default symptoms for each attack

or alternatively use a free-text response to enter a non-default symptom.

To account for the different time zones in the dataset, times (sleep and migraine attack

start and end times) were converted from Coordinated Universal Time (UTC) to the

user's local time zone. The mean number of attacks per month over the 6 months for

each participant was computed from the start and end times of attacks. Users were

then categorised into different frequencies of migraine as follows: Very Low Frequency

Episodic Migraine (VLFEM) = 0-3 migraines per month, Low Frequency Episodic

Migraine (LFEM) = 4-7, High Frequency Episodic Migraine (HFEM) = 8-14, Chronic

Migraine (CM) = ≥15 attacks per month. For the purposes of demographic reporting,

the percentage of attacks which were reported to be triggered by menstruation was

calculated for each individual.

**Measurement of Sleep**

The measurement of sleep by the application is based upon when a user picks up their

mobile phone during the night. The user can then confirm their sleep detected by the

application the following morning. The number of hours slept was calculated from the

start and end times and number of unique sleep episode identifiers. Based on the start

time of the sleep, it was assigned to a night, and defined as the same night if a sleep

started before 08:00am local time. This aims to avoid two sleep episodes (e.g. one

beginning at 11:59 and another at 00:01) being classified as a different night's sleep.

Based on this information, if multiple sleep episodes were recorded on one night this

was counted as a sleep interruption. To generate the number of interruptions variable

we used the number of separate sleep episodes per night minus one. The time in bed

variable was calculated as the difference between the start time of the first sleep

episode of the night and the end time of the final sleep event for that night. Wake time

was calculated as the time in bed minus the number of hours slept. Sleep efficiency

was calculated as time asleep divided by time in bed multiplied by 100 based on

convention. To calculate deviations from monthly mean hours slept, for each user we

computed a Z-score of their total hours slept and took the absolute value of the Z-

score of an individual's deviation, and for each night calculated how many standard

deviations away from the mean the sleep duration was that night.

**Measurement of migraine**

A similar approach was taken to classify migraine attacks. Each migraine episode and

duration was classified based on the start and end time. Users who reported on

average more than 25 migraine episodes per month were excluded ($n = 61$), as if users

had daily attacks, it would be difficult to establish attack onset and fit a predictive

model. We removed any individual migraine attacks which were greater than 96 hours,

as typical attacks last between 4 and 72 hours[25] and these were likely to be data entry

errors. As migraine attacks typically span multiple days, we calculated the duration of

migraine experienced on each day as well as the date that the attack started. For

example, if it started at 20:00 on one day but ended at 16:00 the next day attack

duration would be classified as four hours on day one, and 16 hours on day two.

For all variables the mean and standard deviation were computed.

**Bayesian Modelling**

We used Bayesian estimation techniques to infer distributions of possible parameter

values in each model. We chose a Bayesian approach rather than traditional

frequentist statistics as this avoids issues surrounding the interpretation of *p*-values

and statistical significance[26,27]. Bayesian linear regression allows quantification of

evidence in favour of, or against, a substantive model compared with a baseline

model[28].

To establish whether sleep variables could predict occurrence of a migraine attack on

a given day, we created a generalized linear model with a constant plus three predictor

variables: deviation from an individual's mean sleep duration (*Z*-score), number of

sleep interruptions, and total hours slept, and whether an attack occurred or not

(yes/no) as the binary dependent variable.

Conversely, we created a linear model supposing that total hours slept on a given

night could be predicted from a constant plus two predictor variables: occurrence of

an attack (0 or 1) and pain intensity of an attack (0 – 10).

We used Bayesian inference to estimate the posterior distribution of every parameter

value, which quantifies the relationship between that predictor variable and the

outcome variable. In both models, parameter values were estimated per participant,

and assumed to come from a normal distribution, with the mean and standard

deviation of the distribution also being estimated. Only users who reported ≥4 attacks

for every month across the six months were included in the Bayesian modelling due

to difficulties fitting a predictive model to those which do not have frequent migraines.

We implemented these analyses using the *BRMS* package[29] in *R. BRMS* uses Stan[30]

and implements a Hamiltonian Markov Chain Monte Carlo algorithm. We estimated

the posterior distribution *P*(Parameters | Data) using the No-U-Turn algorithm[31]

implemented in *BRMS.* For the first model with attack occurrence as the outcome

variable, Bernouli was used as the family function, and for the second where hours

slept was the outcome variable, the default Gaussian was used. Four separate chains

of 4,000 samples were taken from the posterior distribution, the first 1,000 samples

were discarded as warmup samples. We verified the chains did not diverge using

trace-plots after sampling, and $\hat{R}$ values were close to 1.000 in both models for all

parameters. Parameters' priors were all set to be uninformative normal distributions

with mean = 0 and standard deviation = 100. We computed 95% highest density

intervals (HDIs) on population mean parameters and used the positioning in relation

to 0 to assess significance. As well as the 95% HDIs we reported the means of the β

value estimates for each model.

## Results

### Demographics

There were 11,166 app users in the overall dataset. We removed any users which

reported their age as <18 years as sleep architecture can change with age[32] and we

were interested in studying the relationship between sleep and migraine in adults. This

left 11,086 users, 7239 of which did not report their age, leaving age data for 3847

users. The mean age of those which reported it was 41.21 years (range = 18-81 years,

SD = 11.50) (see Figure 1). There may have been some users under 18 years of age

which did not disclose their age, therefore it is possible that some users under 18 were

included in the analysis.

When categorizing adults as younger (<40 years of age) and older (≥40 years), there

were 1729 younger adults and 2118 older adults. 7239 could not be categorized. Of

the 11,086 users, 5911 self-reported as female (53.32%), 1276 as male (11.51%), and

3899 did not report their gender (35.17%). Users were reportedly based in 99 different

countries (see Figure 1), with most of the app users residing in the United States of

America (39.34%), Japan (10.54%), Germany (6.68%), United Kingdom (6.43%), and

France (6.29%) at the time at which they signed up to the app. The sleep and migraine variables according to geographical region are shown in .

**Figure 2**.

**Sleep and Migraine Attack Characteristics**

The descriptive statistics for the sleep and migraine measures according to age group and gender are shown in Tables 1 and 2, respectively. The proportion of users classified into the different frequencies of migraine based on the number of attacks per month is shown in Table 3.

**Bayesian Modelling**

Some users were excluded based on the migraine frequency requirements ($\geq 4$ and $\leq 25$ attacks per month), thus in the Bayesian modelling there were 724 users (129 males, 412 females, 183 gender unknown) with an average age of 41.88 years (SD = 11.63) and mean monthly attack occurrence of 9.94. Posterior distributions show the population mean parameter fits across the two models (see Figures 3 and 4). These distributions tell us the estimate of the possible values of the parameters given the data.

**Sleep parameters predicting migraine**

Deviation from mean monthly sleep duration ($M = 0.06$, 95% highest density interval (HDI) [min: 0.04 – max: 0.08]) and number of sleep interruptions yesterday ($M = 0.16$, 95% HDI [min: 0.11 – max: 0.21]) were positive predictors of next day migraine attack occurrence. This means that for every deviation from mean sleep, there is a 6.1% increase in odds of having an attack, and for every sleep interruption, there is 17.4% increased odds of an attack occurring. Total hours slept yesterday was not a positive

or negative predictor of attack occurrence ($M$ = -0.00, 95% HDI [min: -0.04 – max:

0.04]).

**Migraine parameters predicting sleep**

Having a migraine attack was not a predictor of hours slept that evening ($M$ = -0.03,

95% HDI [min: -0.07 – max: 0.01]), whereas the pain intensity reported during an

attack was a positive predictor ($M$ = 0.13, 95% HDI [min: 0.06 – max: 0.20]), indicating

that for every increase of 1 on the VAS for pain intensity, hours slept increased by 0.13

hours (~8 minutes).

# Discussion

This analysis aimed to investigate the relationship between sleep and migraine in a

global sample using measures collected from a smartphone application. We

demonstrated that fragmented sleep, and deviation from typical sleep can predict a

migraine attack, but sleep duration does not, thus in line with our hypotheses. On the

other hand, having a more painful migraine attack is predictive of increased sleep

duration that same evening, but simply having a migraine attack is not. This contrasts

with our hypothesis as we hypothesized that both having an attack and pain intensity

would predict reduced sleep duration. Thus, we were able to disentangle this

bidirectional relationship, in that poor sleep leads to migraine, but having a migraine

attack is not sufficient to disrupt sleep – but rather it is the pain intensity associated

with attacks which alters sleep.

This is a novel finding which is supported by previous literature demonstrating that

pain results in altered sleep[20–23], and that sleep fragmentation but not hours slept is

predictive of migraine[18-19]. This also aligns with a prospective questionnaire study in

pediatric migraine which found that headache intensity was a significant predictor of

delayed or disturbed sleep[33]. However, it is surprising that more pain led to increased

sleep duration, rather than reduced sleep duration given previous studies report a

decrease or disruption.

**Strengths and limitations**

The strengths of this study are that it is a large sample size, and the data were

anonymized and retrospective, meaning that there is little potential for self-report bias,

forgetting, or demand characteristics, as patients reported the start and end times of

migraine attacks in real time. Users were from 99 different countries, across a 6-month

period in comparison to previous studies which often focus on one region and time

point suggesting that the results are generalizable to a wide-range of cultures and

geographical locations.

There are some limitations to consider with this study. Most notably, we used

estimates of sleep from a smart phone application which determines whether a

person is asleep based on whether they pick up their mobile phone or not. This

means that we could fail to capture interruptions where the individual did not go on

their mobile phone, and subtler changes to sleep architecture such as microsleeps,

or changes in sleep stages. However, users approve their sleep episode the next

morning suggesting that it provides an accurate subjective reflection of their sleep.

Nonetheless, these results show that despite the cause of the sleep disruption

(intrinsic or extrinsic), it can have a considerable impact on migraine likelihood.

Moreover, even with this coarse-grained measure we were able to see that changes

in sleep are a significant predictor of migraine attacks, suggesting the utility of large

datasets collected from smartphone applications for investigating neurological

conditions. That being said it is unclear whether the changes in sleep are a trigger or

simply a manifestation of the migraine attack premonitory symptoms. For example,

polyuria is a common premonitory symptom[34] and therefore could be contributing to

the interrupted sleep. Nonetheless, changes in sleep could be used as a biomarker

of an ensuing migraine attack, and prophylactic treatment could therefore be

delivered in a timely manner. Moreover, another consideration is that although the

app collects information covering the International Classification of Headache

Disorders (ICHD-3) criteria (symptoms, duration, location of pain etc.), we did not

verify that each individual user had a migraine diagnosis. Future studies could collate

this information and verify users which could be diagnosed with migraine according

to the criteria. This would also allow for more detailed investigations of headache

disorders in relation to sleep variables (e.g. migraine with versus without aura).

**Clinical implications**

The implications of this study for clinicians are that sleep hygiene should be

considered an inherent part of migraine management. This might be aided by the use

of sleep diaries and/or wearable sleep monitors[35]. Rather than focusing on the

absolute duration of sleep, clinical care and subsequent clinical research should

concentrate on ensuring sleep maintenance and reducing awakenings, in order to

reduce the likelihood of attack occurrence. For example, auditory closed-loop

stimulation during non-rapid-eye-movement sleep is a potential target for non-invasive

enhancement of slow wave sleep and ensuring sleep continuity and maintenance[36].

Focusing on reducing pain associated with attacks through treatments which are

unlikely to cause medication overuse headache[37], rather than focusing on total attack

prevention, could lessen the impact on subsequent sleep and the likelihood of further

attack occurrence and chronification.

**Conclusion**

This study aimed to disentangle the relationship between sleep and migraine. We found that sleep interruptions and deviation from average hours slept are associated with greater likelihood of a migraine attack occurring, whereas overall sleep duration is not.

On the other hand, simply experiencing a migraine attack does not predict altered sleep duration but having a more intensely painful attack predicts increased sleep duration.

Clinicians should ensure sleep hygiene is intrinsic to migraine management. Longitudinal studies with polysomnography, and more detailed investigations of headache disorders classified according to ICHD criteria are required to confirm these findings, and methods for improving sleep continuity should be explored as a potential migraine intervention.

## References

1.  Holland PR. Headache and sleep: Shared pathophysiological mechanisms. Cephalalgia. 2014;34:725–744.

2.  Dodick DW, Eross EJ, Parish JM. Clinical, Anatomical, and Physiologic Relationship Between Sleep and Headache. Headache: The Journal of Head and Face Pain. 2003;43:282–292.

3.  Moldofsky H. Sleep and pain. Sleep Medicine Reviews. 2001;5:385–396.

4.  Gago-Veiga AB, Vivancos J, Sobrado M. The premonitory phase: a crucial stage in migraine. Neurología (English Edition) [online serial]. Epub 2019 Jul 23. Accessed at: http://www.sciencedirect.com/science/article/pii/S2173580819301038. Accessed April 8, 2020.

5.  Kelman L. The triggers or precipitants of the acute migraine attack. Cephalalgia. 2007;27:394–402.

6.  Bigal ME, Lipton RB. What predicts the change from episodic to chronic migraine? Current Opinion in Neurology. 2009;22:269–276.

7.   Kelman L, Rains JC. Headache and Sleep: Examination of Sleep Patterns and
Complaints in a Large Clinical Sample of Migraineurs. Headache: The Journal of
Head and Face Pain. 2005;45:904–910.

8.   Nesbitt AD, Leschziner GD, Peatfield RC. Headache, drugs and sleep.
Cephalalgia. SAGE Publications Ltd STM; 2014;34:756–766.

9.   Hamelsky SW, Lipton RB. Psychiatric Comorbidity of Migraine. Headache: The
Journal of Head and Face Pain. 2006;46:1327–1333.

10.  Stanyer EC, Creeney H, Nesbitt AD, Holland PRR, Hoffmann J. Subjective Sleep
Quality and Sleep Architecture in Patients With Migraine: A Meta-analysis.
Neurology [online serial]. Wolters Kluwer Health, Inc. on behalf of the American
Academy of Neurology; Epub 2021 Sep 22. Accessed at:
https://n.neurology.org/content/early/2021/09/22/WNL.0000000000012701.
Accessed September 24, 2021.

11.  Vos T, Abajobir AA, Abate KH, et al. Global, regional, and national incidence,
prevalence, and years lived with disability for 328 diseases and injuries for 195
countries, 1990–2016: a systematic analysis for the Global Burden of Disease
Study 2016. The Lancet. 2017;390:1211–1259.

12.  Zobdeh F, ben Kraiem A, Attwood MM, et al. Pharmacological treatment of
migraine: Drug classes, mechanisms of action, clinical trials and new treatments.
British Journal of Pharmacology. 2021;178:4588–4607.

13.  Vollono Catello, Gnoni Valentina, Testani Elisa, et al. Heart Rate Variability in
Sleep-Related Migraine without Aura. Journal of Clinical Sleep Medicine.
American Academy of Sleep Medicine; 2013;09:707–714.

14.  Houle TT, Butschek RA, Turner DP, Smitherman TA, Rains JC, Penzien DB.
Stress and sleep duration predict headache severity in chronic headache
sufferers. PAIN®. 2012;153:2432–2440.

15.  Engstrøm M, Hagen K, Bjørk MH, et al. Sleep quality, arousal and pain thresholds
in migraineurs: a blinded controlled polysomnographic study. The Journal of
Headache and Pain. 2013;14:12.

16.  Karthik N, Sinha S, Taly AB, Kulkarni GB, Ramachandraiah CT, Rao S. Alteration
in polysomnographic profile in 'migraine without aura' compared to healthy
controls. Sleep Medicine. 2013;14:211–214.

17.  Edwards RR, Almeida DM, Klick B, Haythornthwaite JA, Smith MT. Duration of
sleep contributes to next-day pain report in the general population. PAIN®.
2008;137:202–207.

18.  Smith MT, Edwards RR, McCann UD, Haythornthwaite JA. The Effects of Sleep
Deprivation on Pain Inhibition and Spontaneous Pain in Women. Sleep.
2007;30:494–505.

19.  Bertisch SM, Li W, Buettner C, et al. Nightly sleep duration, fragmentation, and
quality and daily risk of migraine. Neurology. Epub 2019 Dec
16.:10.1212/WNL.0000000000008740.

20.  Mulligan EP, Brunette M, Shirley Z, Khazzam M. Sleep quality and nocturnal pain
in patients with shoulder disorders. Journal of Shoulder and Elbow Surgery.
2015;24:1452–1457.

21.  Marty M, Rozenberg S, Duplan B, et al. Quality of sleep in patients with chronic
low back pain: a case-control study. Eur Spine J. 2008;17:839–844.

22.  Keilani M, Crevenna R, Dorner TE. Sleep quality in subjects suffering from
chronic pain. Wien Klin Wochenschr. 2018;130:31–36.

23.  Wylde V, Rooker J, Halliday L, Blom A. Acute postoperative pain at rest after hip
and knee arthroplasty: Severity, sensory qualities and impact on sleep.
Orthopaedics & Traumatology: Surgery & Research. 2011;97:139–144.

24.  Alstadhaug KB, Bekkelund S, Salvesen R. Circannual periodicity of migraine?
European Journal of Neurology. 2007;14:983–988.

25.  Kelman L. Pain Characteristics of the Acute Migraine Attack. Headache: The
Journal of Head and Face Pain. 2006;46:942–953.

26.  Wasserstein RL, Lazar NA. The ASA Statement on p-Values: Context, Process,
and Purpose. The American Statistician. Taylor & Francis; 2016;70:129–133.

27.  Amrhein V, Greenland S, McShane B. Scientists rise up against statistical
significance. Nature. 2019;567:305–307.

28.  Gadie A, Shafto M, Leng Y, Cam-CAN, Kievit RA. How are age-related
differences in sleep quality associated with health outcomes? An epidemiological
investigation in a UK cohort of 2406 adults. BMJ Open. British Medical Journal
Publishing Group; 2017;7:e014920.

29.  Bürkner P-C. brms: An R Package for Bayesian Multilevel Models Using Stan.
Journal of Statistical Software. 2017;80:1–28.

30.  Stan Development T. RStan: the R interface to Stan. R package version 217 3.
Epub 2018.

31.  Hoffman MD, Gelman A. The No-U-Turn Sampler: Adaptively Setting Path
Lengths in Hamiltonian Monte Carlo [online]. arXiv; 2011. Accessed at:
http://arxiv.org/abs/1111.4246. Accessed July 18, 2022.

32.  Dorffner G, Vitr M, Anderer P. The Effects of Aging on Sleep Architecture in
Healthy Subjects. In: Vlamos P, Alexiou A, editors. GeNeDis 2014. Cham:
Springer International Publishing; 2015. p. 93–100.

33.  Heyer GL, Rose SC, Merison K, Perkins SQ, Lee JEM. Specific Headache
Factors Predict Sleep Disturbances Among Youth With Migraine. Pediatric
Neurology. 2014;51:489–493.

34. Charles A. The pathophysiology of migraine: implications for clinical
management. The Lancet Neurology. 2018;17:174–182.

35. Bianchi MT. Sleep devices: wearables and nearables, informational and
interventional, consumer and clinical. Metabolism. 2018;84:99–108.

36. Stanyer EC, Baniqued PDE, Awais M, et al. The impact of acoustic stimulation
during sleep on memory and sleep architecture: A meta-analysis. Journal of
Sleep Research. 2022;31:e13385.

37. Saengjaroentham C, Strother LC, Dripps I, et al. Differential medication overuse
risk of novel anti-migraine therapeutics. Brain. 2020;143:2681–2688.

## Authors Contributions

439  E.C.S and J.H. conceptualized the study. E.C.S and J.B performed the statistical

440  analysis. E.C.S drafted the manuscript. JR.P. extracted and prepared the data from

441  the Healint platform. J.B., JR.P., A.U., and J.H. revised the manuscript for intellectual

442  content.

## Funding and Disclosures

This work has been supported by a Medical Research Council PhD studentship (ECS;

MR/N013700/1). Healint provided in-kind dataset, prior data-collection, data

preparation, and intellectual support.

J. Hoffmann reports honoraria for consulting activities and/or serving on advisory

boards from Allergan, Autonomic Technologies Inc., Cannovex BV, Chordate Medical

AB, Eli Lilly, Hormosan Pharma, Lundbeck, Novartis, Sanofi and Teva. He received

personal fees for Medico-Legal work as well as from Oxford University Press,

Quintessence Publishing, Sage Publishing and Springer Healthcare. He also reports

a research grant from Bristol Myers Squibb. J. Hoffmann serves as Associate Editor

for Cephalalgia, Cephalalgia Reports, Journal of Oral & Facial Pain and Headache as

well as for Frontiers in Pain Research. He is an elected member of the Board of

Trustees of the International Headache Society (IHS) and serves as a Council Member

and Treasurer of the British Association for the Study of Headache (BASH). All these

activities are unrelated to the submitted work.

P.R.H. reports, unrelated to the current project, honoraria for educational and advisory

purposes from Allergan, Novartis and Teva as well as research funding from Eli Lilly,

Amgen, Kallyope and Cellgene/Bristol Myers Squibb. P.R.H serves as an associate

editor for Cephalalgia and the British Journal of Pharmacology.

JR.P and A.U. are employees of Healint Pte Ltd

**ORCID**

J. Brookes 0000-0002-4600-7561

J. Hoffmann: 0000-0002-2103-9081

P.R. Holland: 0000-0003-3345-4853

E.C. Stanyer: 0000-0002-7672-5652

**Acknowledgements**

The authors would like to thank Dr Alexander Nesbitt for useful discussions about the

results.

## Tables

### Table 1: Descriptive Statistics for the sleep variables according to user demographics

| | Overall sample M (SD) | Older adults M (SD) | Young adults M (SD) | Age (years) unknown M (SD) | Female M (SD) | Male M (SD) | Gender unknown M (SD) |
|---|---|---|---|---|---|---|---|
| Total Sleep Time | 7.45 (1.00) | 7.46 (1.n04e) | 7.53 (0.95) | 7.43 (1.00) | 7.55 (1.00) | 7.36 (0.96) | 7.34 (1.01) |
| Sleep Start Time from Midnight (hours) | 23.55 (1.34) | 23.40 (1.42) | 23.75 (1.41) | 23.55 (1.29) | 23.50 (1.38) | 23.62 (1.35) | 23.60 (1.27) |
| Sleep End Time (hours) | 31.21 (1.24) | 31.10 (1.27) | 31.54 (1.23) | 31.17 (1.22) | 31.28 (1.25) | 31.15 (1.31) | 31.13 (1.19) |
| Time in Bed (hours) | 7.66 (1.22) | 7.71 (1.28) | 7.79 (1.27) | 7.62 (1.18) | 7.78 (1.25) | 7.53 (1.10) | 7.54 (1.18) |
| Time Awake (hours) | 0.21 (0.69) | 0.25 (0.76) | 0.26 (0.78) | 0.19 (0.64) | 0.23 (0.75) | 0.17 (0.51) | 0.19 (0.63) |
| Sleep Efficiency (%) | 0.99 (0.034) | 0.98 (0.038) | 0.98 (0.037) | 0.99 (0.032) | 0.99 (0.036) | 0.99 (0.028) | 0.99 (0.032) |
| Number of Sleep Interruptions | 0.12 (0.18) | 0.13 (0.20) | 0.13 (0.18) | 0.12 (0.18) | 0.12 (0.19) | 0.097 (0.17) | 0.13 (0.17) |

*M* = Mean, *SD* = Standard Deviation

**487** **Table 2: Descriptive statistics for migraine variables according to user**
**488** **demographics**

| | Overall sample M (SD) | Older adults M (SD) | Young adults M (SD) | Age (years) unknown M (SD) | Female M (SD) | Male M (SD) | Gender unknown M (SD) |
|---|---|---|---|---|---|---|---|
| Pain intensity (Scale 1-10) | 5.26 (1.64) | 5.34 (1.63) | 5.50 (1.55) | 5.19 (1.65) | 5.42 (1.59) | 4.90 (1.77) | 5.14 (1.64) |
| Attack duration (hours) | 18.15 (14.86) | 19.46 (15.65) | 17.91 (13.75) | 17.70 (14.71) | 19.16 (15.22) | 14.69 (13.07) | 17.72 (14.65) |
| Number of attacks overall | 19.77 (24.17) | 21.72 (24.94) | 18.14 (21.87) | 19.32 (24.22) | 20.62 (24.44) | 23.41 (27.71) | 17.27 (22.21) |
| Attacks triggered by menstruation (%) | 0.23 (0.29) | 0.18 (0.28) | 0.31 (0.30) | 0.24 (0.29) | 0.27 (0.30) | 0.0 (0.02) | 0.25 (0.29) |
| Attacks per month | 2.82 (3.45) | 3.10 (3.56) | 2.59 (3.12) | 2.76 (3.46) | 2.95 (3.49) | 3.34 (3.96) | 2.47 (3.17) |
| Migraine start from midnight (hours) | 11.95 (3.31) | 11.46 (3.14) | 12.96 (3.18) | 11.96 (3.35) | 5.42 (1.59) | 4.89 (1.77) | 5.14 (1.64) |
| Migraine end time (hours) | 30.12 (15.10) | 30.91 (15.88) | 31.07 (14.04) | 29.66 (14.95) | 19.18 (15.25) | 14.77 (13.10) | 17.73 (14.66) |

**489** *M* = Mean, *SD* = Standard Deviation.

**490** **Table 3: Proportion and percentage of app users categorised in different**
**491** **frequencies of migraine based on number of migraine attacks per month**

| | Proportion of users | | | | Percentage of users (%) | | | |
|---|---|---|---|---|---|---|---|---|
| Gender | CM (≥15) | HFEM (8-14) | LFEM (3-7) | VLFEM (0-3) | CM (≥15) | HFEM (8-14) | LFEM (3-7) | VLFEM (0-3) |
| Female | 81 | 326 | 1053 | 3672 | 1.6 | 6.4 | 20.5 | 71.6 |
| Male | 23 | 91 | 247 | 728 | 2.1 | 8.4 | 22.7 | 66.9 |
| Unknown | 39 | 135 | 568 | 2631 | 1.2 | 4 | 16.8 | 78 |
| Total | 1435 | 552 | 1868 | 7,031 | 1.50 | 5.80 | 19.50 | 73.30 |

**492**

**493**

# Figure legends

**495** **Figure 1: Demographics of the users in the dataset.** A) Choropleth map showing

**496** the distribution and number of users across the world. B) Population pyramid showing

**497** the age (for those which reported it), and gender of the users in the dataset.

**Figure 2: Sleep and migraine parameters according to region.** A) Mean hours slept according to region. B) Mean sleep start time according to region. C) Migraine start and end (D) time according to region. E) Migraine attack duration according to region F) Histogram of the start time of all reported migraine attacks

**Figure 3: Bayesian modelling results. Sleep variables the night before can predict next day occurrence of a migraine attack.** Larger deviations from typical sleep (A) and a greater number of sleep interruptions (B) predict occurrence of a next day migraine attack, but total hours slept does not (C). Density plots show posterior distributions for the population means of the three tested sleep-related predictor variables on whether a migraine attack occurred or not. Error bars indicate the 95% highest density interval.

**Figure 4: Bayesian modelling results. Migraine pain intensity can predict total hours slept on the same night of an attack.** Having an attack (A) did not predict alterations to sleep, but greater pain intensity (B) results in more hours slept the following night. Density plots show posterior distributions for the two tested models of the population means of migraine-related predictor variables on total hours slept. Error bars indicate the 95% highest density interval. VAS = visual analogue scale.

**A.** Deviation from Mean Sleep Yesterday (z)

**B.** Number of Sleep Interruptions Yesterday

**C.** Total Hours Slept Yesterday

**A.** Had Migraine Attack

**B.** Pain Intensity (VAS 0-10)