

An update to the critical appraisal of Milwaukee protocol

Ferhat Arslan¹, Haluk Vahaboglu^{1*}

Istanbul Medeniyet Universitesi

Abstract:

Rabies is a highly fatal encephalitis. Currently, there is no approved treatment. Inducing therapeutic coma during the first week of symptomatic rabies patient, called Milwaukee protocol, had been suggested as promising. However, recent evidence failed to support the use of the Milwaukee protocol. This mini-review analyzed the reports of patients managed with therapeutic coma since 2014 to provide an update for the critical appraisal of this protocol.

Introduction:

Rabies is an almost uniformly fatal encephalitis caused by the members of the genus *Lyssavirus*. During the last thirty years, the global burden of rabies is declined due to increased living conditions and the availability of vaccines in rural areas (1). Bats and infected dogs are the primary vehicles of human rabies. Two clinical forms are described as paralytic and furious, the latter even more dramatic (2). The virus enters the axons from neuromuscular junctions after entering through broken skin or mucosa. Viruses are transferred to the dorsal root ganglia and, with secondary transport, reach the brain (3). Naturally, in a maximum of 14 days from the first symptoms appear, patients die due to respiratory or cardiac arrest (4).

The pathophysiology of neuronal damage is yet to be elucidated. Autopsy findings of patients who died of rabies vary; some show extensive damage to the brain tissue (5), while others do not (6). However, animal studies displayed extensive neuronal damage attributed to mitochondrial dysfunction and apoptosis (7, 8). Despite the progress in understanding the pathophysiology of neuronal damage in rabies, approved treatment has not yet been developed (9). In 2005 a 15-year-old girl developed rabies-like encephalitis after bat exposure and survived with the experimental Milwaukee protocol (10). The Milwaukee protocol recommends inducing therapeutic coma by ketamine and midazolam during rabies patients' first week of ICU admittance (8). Despite the great excitement of this first report, Milwaukee protocol failed to replicate the initial success during the following years. It has been proposed to abandon therapeutic coma induction during ICU care of rabies patients (8, 11). However, since the first warning in 2015 to leave inducing therapeutic coma in rabies management has been ignored, the Milwaukee protocol is still being used (8).

This mini-review aimed to recruit survival evidence on Milwaukee protocol since 2015.

Data collection:

We screened the Web of Science using "rabies (All fields) AND Milwaukee (All fields)," refined dates from 2015 to 2022, carefully inspected abstracts and identified data for 13 cases (Table 1). Next, we screened the Web of Science using "rabies (All fields) AND survived (All fields) AND case OR patient (All fields) to identify survived cases since 2015 who have not managed with induced therapeutic coma.

Results:

Table 1 outlines the characteristics of 12 cases managed with induced therapeutic coma published after 2014 in the medical literature. Briefly, only one patient survived (patient 12). This patient, however, was applied to the hospital with atypical symptoms such as acute abdominal pain, sore throat, and a mild cough. The initial diagnosis of patient 12 was aspiration pneumonia, and the exposure was a cat scratch. The diagnosis was made with anti-rabies antibody titration, and rabies antigen or RNA has never been shown. Patient 12 was managed under induced coma for ten days, and following the cessation of therapeutic coma patient gradually improved and was finally discharged with minimal neurologic deficit. Case reports stated the death reasons for patients 1, 2, 7, 8, and 10 (Table 1). Death among patients with therapeutic coma mostly was due to respiratory and cardiac failure. Interestingly, nosocomial infections were not reported if they ever occurred. Six patients received post-exposure prophylaxis (PEP) 3-5 times, and one patient received both rabies immunoglobulin (Ig) and a full course of rabies vaccine (RABV).

Table 2 displays patients managed with conventional ICU care and survived. Three dead patients were also included in the table because these were presented in the case series with surviving patients (12). A total of eight patients survived; all had 3-5 PEP, and two had post-exposure Ig. The follow-up period of survived patients was a minimum of six months and a maximum of four and a half years.

Discussion:

This mini-review aimed to address whether all rabies survivors had managed through induced therapeutic coma and found that this is not the case. Although we found one patient survived with Milwaukee protocol management, the diagnosis for that case was not robust. The patient was admitted with pneumonia. Moreover, she improved gradually after ending the therapeutic coma. The authors did not discuss what would happen if they had completed the therapeutic coma before ten days or even not inducing it at all.

The induced therapeutic coma may cause severe side effects such as electrolyte disturbances and cardiorespiratory arrest (13). Potential side effects, including facilitating the acquisition of nosocomial infections and causing immune depression, were discussed previously (8). Our mini-review also provided evidence that some patients died of nosocomial infections acquired under therapeutic coma. Uncontrolled hypotension, acidosis, and respiratory failure might be because of sepsis and, eventually, septic shock developed during the therapeutic coma. However, the authors neither gave any information nor discussed nosocomial infections.

The definitive diagnosis of rabies is almost always difficult in the early course of the disease (4). Therefore, applying potentially hazardous and experimental Milwaukee protocol, namely the therapeutic coma, might also harmful to patients with other mimicking encephalitides.

Finally, we agree with the authors who previously warned about abandoning Milwaukee protocol (inducing therapeutic coma) in the management of rabies (8, 11).

Table 1. Characteristics of rabies cases applied Milwaukee protocol since 2015

	Age/gender	Year ¹	Died of	Hosp. stay (days)	PEP	outcome	LAMA	ref
1	33/M	2021	Uncontrolled hypotension	18	none	Died		(14)
2	27/F	2021	Respiratory failure	580	5x RABV	Died		(15)
3	51/M	2021	Not available	7	Ig & 5x RABV	Died		(16)
4	29/M	2020	Not stated	28	5x RABV	Died		(17)
5	58/F	2020	Not stated	17	3x RABV	Died	yes	(17)
6	34/M	2020	Not stated	2	3x RABV	Died		(17)
7	12/M	2019	Autonomic dysregulation	15	none	Died		(18)
8	58/F	2017	Heart block, profound acidosis	8	none	Died		(6)
9	10/M	2016	Not stated	4	none	Died		(19)
10	9/F	2015	Refractory hypotension	75	none	Died		(5)
11 ²	8/F	2015			none	Survived		(20)

¹ Publication year

² Cat scratch

Table 2. Characteristics of rabies cases not applied Milwaukee protocol since 2015 and survived

	Age/gender	Year ³	Died (days)	PEP	outcome	follow-up	ref
1	33/F	2021		4x RABV	Survived	12 months	(21)
2	4/M	2020		3x RABV	Survived	12 months	(22)
3	13/M	2019	150	3x RABV	Died		(12)
4	10/M	2019	180	3x RABV	Died		(12)
5	5/F	2019	240	3x RABV	Died		(12)
6	3/F	2019		5x RABV	Survived	6 months	(12)
7	5/F	2019		Ig & 4x RABV	Survived	7.5 months	(12)
8	4/M	2019		Ig & 3x RABV	Survived	10 months	(12)
9	26/M	2019		4x RABV	Survived	13 months	(12)
10	9/M	2019		4x RABV	Survived	13 months	(12)
11	8/M	2015		5x RABV	Survived	4.5 years	(23)

³ Year, publication year

References:

1. Gan H, Hou X, Wang Y, et al. Global burden of rabies in 204 countries and territories, from 1990 to 2019: results from the Global Burden of Disease Study 2019. *Int J Infect Dis* 2022;
2. Fooks AR, Cliquet F, Finke S, et al. Rabies. *Nat Rev Dis Prim* 2017;3
3. Etesami R, Conzelmann KK, Fadai-Ghotbi B, et al. Spread and pathogenic characteristics of a G-deficient rabies virus recombinant: An in vitro and in vivo study. *J Gen Virol* 2000;81:2147–53
4. Jackson AC. Human Rabies: a 2016 Update. *Curr Infect Dis Rep* 2016;18:1–6
5. Caicedo Y, Paez A, Kuzmin I, et al. Virology, immunology and pathology of human rabies during treatment. *Pediatr Infect Dis J* 2015;34:520–8
6. Pathak S, Horton DL, Lucas S, et al. Diagnosis, management and post-mortem findings of a human case of rabies imported into the United Kingdom from India: A case report. *Virology* 2014;11:1–7
7. Scott CA, Rossiter JP, Andrew RD, et al. Structural abnormalities in neurons are sufficient to explain the clinical disease and fatal outcome of experimental rabies in yellow fluorescent protein-expressing transgenic mice. *J Virol* 2008;82:513–21
8. Zeiler FA, Jackson AC. Critical Appraisal of the Milwaukee Protocol for Rabies: This Failed Approach Should Be Abandoned. *Can J Neurol Sci* 2015;43:44–51
9. Appolinario CM, Jackson AC. Antiviral therapy for human rabies. *Antivir Ther* 2015;20:1–10
10. Willoughby RE, Tieves KS, Hoffman GM, et al. Survival after Treatment of Rabies with Induction of Coma. *N Engl J Med* 2005;352:2508–14
11. Wilde H, Hemachudha T. The "milwaukee Protocol" for Treatment of Human Rabies Is No Longer Valid. *Pediatr Infect Dis J* 2015;34:678–9
12. Mani RS, Damodar T, Divyashree S, et al. Case report: Survival from rabies: Case series from India. *Am J Trop Med Hyg* 2019;100:165–9
13. Ledesma LA, Lemos ERS, Horta MA. Comparing clinical protocols for the treatment of human rabies: the Milwaukee protocol and the Brazilian protocol (Recife). *Rev Soc Bras Med Trop* 2020;53:e20200352
14. Goravey W, Husain A, Ali GA, et al. Antemortem diagnosis of human rabies: A case report. *Clin Case Reports* 2021;9:711–3
15. Zarate AA, Sanchez M, Flores-Lovon K, et al. Pregnant woman from southern Peru with rabies infection who delivered a healthy baby. *Trop Doct* 2021;51:457–8
16. Mohindra R, Suri V, Chatterjee D, et al. Measuring antibody titres following rabies postexposure prophylaxis in immunosuppressed

patients: a norm rather than the exception. *BMJ Case Rep* 2021;14

17. Kumar S, Gupta P, Panda P. Death from rabies: The reason being poor compliance to vaccination or it's failure. *J Fam Med Prim Care* 2020;9:4437
18. Dhayhi NS, Arishi HM, Ibrahim AAYA, et al. First confirmed case of local human rabies in Saudi Arabia. *Int J Infect Dis* 2019;87:117–8
19. Agarwal A. The "Milwaukee protocol" (MP) hope does not succeeds for rabies victim. *Med J Dr DY Patil Univ* 2017;10:184–6
20. Lu A, Shah P, Shen P, et al. Temporal evolution on MRI of successful treatment of rabies. *Clin Imaging* 2015;39:893–6
21. Sumirini AA, Manchanda B V, Mehra S. The Rare Case of a Human Rabies Survivor & the Comparative Study of All Documented Human Rabies Survivors Till 2021. *J Neuroinfect Dis* 2022;0–5
22. John B, Kumar S, Kumar S, et al. Child survivor of rabies in India: a case report. *Paediatr Int Child Health* 2020;40:255–60
23. Netravathia M, Udani V, Manic RS, et al. Unique clinical and imaging findings in a first ever documented PCR positive rabies survival patient: A case report. *J Clin Virol* 2015;70:83–8