USES OF THE SKEW-LOGISTIC FUNCTION FOR MULTI-WAVE FUNCTIONS

Dr. Russell Cheng

University of Southampton
Highfield, SO17 5BJ
United Kingdom
cheng@btinternet.com

Dr. Brian Williams

SACEMA
Stellenbosch University,
South Africa
williamsbg@me.com

ABSTRACT

Skew-Logistic (SL) function has been proposed to model a real-life dynamic process which rises monotonically to peak followed by a monotonic falling back. It was originally introduced to model the first stage of the covid-19 pandemic when it first appeared with the purpose of forecasting the likely behiour of covid. In its subsequent behaviour, as covid-19 rose and fell under the influence of different controls and with the onset variants, the prevalence and incidence of covid rose and fell repeatedly producing what we might call a Multi-Wave (MW) behaviour, the waves not nessarily the same size. The purpose of the paper is to show how the method of using the SL function for one wave can be easily modified to model the MW situation. To illustrate our extended method, we apply it to two examples. One is to covid -19, to model its most recent behavour and examine how things might change. We also apply it to climate change, undoubtedly the most serious issue, as without ensuring the world becomes rapidly carbon will bring to and end the known world

Keywords: covid-19 waves, climate change, carbon negative

1   INTRODUCTION

The Skew-Logistic (SL) function was introduced in Dye et al. (2020) to compare the first wave of the COVID-19 epidemics in different European countries. The mathematical form is described in the Supplemental Materials of the paper, which also describes how the model can be fitted to real data using standard statistical techniques. This paper shows how the method can be used to firstly fit to the individual waves of a sample of Multi-Wave (MW) data and how all these individual waves can be then combined to create an overall simultaneous fit the entire data sample. It will be convenient to summarise the single wave fitting first, which is done in Section 2. Then, in Section 3, we discuss the MW fitting method.

Our first application is described in Section 4. In this example we fit the most recent covid-19 UK Active Cases that occurred from March 2022 till November 2022 covering the most recent 3 waves. We then use this to show predict the likely case level in the next few weaks.

In our second we fit the SL-MW model to the annual atmospheric carbon dioxide level, in parts per million (CO_2 ppm) from 1800 to 2022. We also fit the model to the world average Temperature (^{0}C). We then use the fits to predict the CO_2 and Temperature levels until 2100, showing what might happen under different scenarios.

2   SINGLE WAVES

Fitting the SL function to the observation of one wave of the process of interest forms the basis for fitting several SL functions to MW data. We therefore start by summarising the description of the SL function given in Dye et al. (2020).

The SL function takes the form:θ

$$D(t, \boldsymbol{\theta}) = a \frac{e^{b(t-\tau)}}{[1+e^{\frac{1}{2}(b-d)(t-\tau)}]^2} = a \frac{e^{d(t-\tau)}}{[1+e^{-\frac{1}{2}(b-d)(t-\tau)}]^2} \qquad (1)$$

where $D(t,\boldsymbol{\theta})$ is the prevalence or incidence of the quantity of interest, this being the number of active covid-19 cases in Example 1 and CO_2 in Example 2 All four parameters are readily interpretable. The parameter a, is close to the maximum value of $D(t)$, whilst τ indicates the location of the maximum. The parameters b and d, respectively indicate the rates of rise and decline of $D(t)$.

As shown in Equation 1, b and d are mathematically identical.

We write $\boldsymbol{\theta} = (\sigma, a, b, d, \tau)$ where σ is the standard deviation of an individual observation. As dependence of $D(t)$ on the the parameters is obvious, $\boldsymbol{\theta}$ is usually omitted.

We assume that $b > 0$ and $d < 0$ so that the term $(b - d)/2$ in the denominator is always positive. $D(t) \cong e^{b(t-\tau)}$ if $t \to -\infty$, so that b gives the exponential rate of increase of $D(t)$ as t increases from $-\infty$, and $D(t) \cong e^{d(t-\tau)}$ if $t \to \infty$, so that $d < 0$ gives the exponential rate decrease.

Either form of $D(t)$ in Equation 1 can thus be used when fitting to data, with the signs of b and d showing which role they have.

The explicit maximum value of $D(t)$ is:

$$D_{max} = ar^{(\frac{2r}{1+r})}(1 + r)^{-2} \qquad (2)$$

where $r = -b/d$. The maximum point is at

$$t_{max} = \tau + \frac{2}{b-d}\log(-\frac{b}{d}). \qquad (3)$$

Equations (2) and (3) show that, when b and $-d$ are close in value, a will be close to the maximum of D, and τ wil be close to the maximum point.

As described in Dye et al. (2020), the SL function can be fitted to data using the method of maximum likelihood (ML) using Nelder-Mead search. This latter needs initial parameter values to be provided. This is not always straightforward, especially with multimodal functions, particularly when the number of parameters is large. A very attractive feature with the SL function is that, its readily interpretable parameters enables the user to intervene and interactively select appropriate initial parameters are close the best. This enables Nelder-Mead to reliably obtain an accurate optimum. We can simply visually examine the data to obtain an initial estimate of its position and size. The rates of increase and decrease to be roughly estimated to give starting values for b and d. An initial value for a can then be obtained using Equation (2), whilst the observed maximum and Equation (3) can be used to give the initial value of τ.

In applying ML optimization, we include an extra parameter σ, the standard deviation of the observational error. Estimation of σ is included in the ML method, so that the quality of the fit is also assessed.

We turn now to the MW situation

3   MULTI-WAVE

3.1 Function Formula and Data Examples

We assume a very simple additive functional form for when there are N waves. Namely:

$$D(t) = \sum_{i=1}^{i=N} a_i \, e^{b_i(t-\tau_i)}\{[1 + e^{\frac{1}{2}(b_i-d_i)(t-\tau_i)}]^{-2}\}, \qquad (4)$$

where typically, but not invariably, the τ_i are in increasing order $\tau_1 < \tau_2 \ldots < \tau_N$ indicating the (approximate) location of each of the maxima of the waves. We give two data samples where $D(t)$ might be fitted.

The first, Example 1, comprises the observed Daily New Cases of Covid-19 from 3 March 2022 to 17 November 2022. Figure 1 shows a chart of the sample.

Figure 1. *Covid-19 observations from 17 May to 16 November 2022*

Example 2, comprises the observations of the global CO₂ level from 1800 to 2022. A plot of this sample is shown in Figure 2.

Figure 2. *CO₂ −282 ppm Level from 1800 to 2022*

There is a noticable peak in the sample plot where a wave is particular prominent, but this of course depends on the juxtaposition of the waves, so that where a wave peaks may not be obvious from the full sample plot. In Example 1, one might conjecture that there are three waves. In Example 2, the number of waves is not so clear if the are one or waves.

We consider our proposed fitting method next.

3.2 The Basic Method

The proposed method of fitting $D(t)$ to data takes two stages.

In the first stage the data is grouped into separate sections, with each representing the position where the data is most dependent on a particular wave. The division process could be made automatic, but if speed is not important, and this is the case our two examples, then it is simplest for the user to carry out this process by eye. As will be seen, in Example 1, the number of waves looks likely to be three. As previously intimated, the choice is not so obvious in Example 2. However the initial slope variation suggests an earlier wave that becomes dominated by a second wave. Here more adjustment is needed to fix the position of the first wave and our final choice turns out to be two waves.

With a choice of the number of waves and their likely positions made, a single wave model is separately fitted to each section of the data. There are five parameters, a, b, d, τ and σ for each section so that the number of parameters is 5 in each. In Example 1 with three waves fitted the total number of parameters is 13 as we assume that σ remains the same for both waves. Varying σ is possible but there was no evidence this was needed in this case. For Example 2 the number of parameters in the full fit is 9.

In the second stage the model of Equation 1 is then fitted to the entire data set using, as the initial parameter estimates, the values obtained in the first stage for each of the single waves.

To find the maximum when using search methods like Nelder-Mead, a good choice of initial parameters is extremely important, particularly when the parameter count is high, when the search domain is high dimensional. The problem is particularly demanding when the function is multimodal. This is the case in our problem. In our case, visual evidence of a good fit of a regression line to data, together with confidence level assessment, provides reassurance that our method is satisfactory.

In the next section we study the Examples in more detail

4   FITTING METHOD

4.1 Covid-19 Example

The fitting problem is very amenable to being tackled visually and user interaction. We indicate the general procedure which is very flexible. There are no hard and fast rules and, depending on the data, adjustments are easy to make. How a simple overall approach is to divide the fitting into two steps. We consider Example 1.

As the first step we make a visual inspection of the sample. This suggests choosing 3 waves with waves 1,2 and 3 contributing respectively to data points in the ranges $S_1 = \{1, 86\}$, $S_2 = \{87,172\}$, $S_3 = (173,243\}$, where the sample size is $n = 243$. We then separately fit the single wave SL model of $D(t)$, as given in Equation (1), to each of the data sets S_j, $j = 1,2,3$. The fitting process is described in Cheng et al. (2020), Section 2 and in Dye et al. (2020), Supplementary Materials, and will not be repeated here. Figure 3 gives the results of this step.

Figure 3. *The three fitted waves for the COVID-19 sample*

The fitted parameter values are given in Table 1 below.

Table 1: *Parameter estimates obtained by dividing full sample into 3 subsamples and fitting a single wave separately to each subsample.*

Parameters	1-86: Wave 1	87-172 Wave 2	173-243 Wave 3
σ	4490	1630	748
a	254000	101000	27500
b	0.215	0.103	0.0492
d	-0.0596	-0.0713	-0.174
τ	44600	4470	44900

Once the single waves have been fitted, we proceed to the second step which is to fit the *N*=3, Multi-wave *D*(*t*) of Equation (4) taking $\theta = \{\sigma, a_1, b_1, d_1, \tau_1, \ a_2, b_2, d_2, \tau_2, \ a_3, b_3, d_3, \tau_3\}$ as the vector of initial parameter values. The subscripts correspond to the number of the fitted Wave.

The initial Multi-wave *D*(*t*) obtained, using the initial parameter values of Table1, is shown (in green) in the chart of Figure 4. Note that the last segment of this curve, corresponding to Wave 3, is noticeably different from the fit to Wave 3 shown in the third plot of Figure 3.

Figure 4 also shows the final fit (red line) obtained for the full sample, where it will be seen that the optimization greatly improves the overall fit.

The optimized parameter estimates are shown in Table 2

Figure 4. *The N = 3 Multi-wave fit to the full COVID-19 sample*

Table 2:
*Parameter estimates obtained by an N=3
Multi-wave to the full COVID-19 sample*

Parameters	$i = 1$	$i = 2$	$i = 3$
σ	4610		
a_i	27000	102000	28300
b_i	0.198	0.0817	0.0483
d_i	-0.0637	-0.0875	-0.135
τ_i	44600	4470	44900

4.2 CO_2 Example

In this example, visual inspection of the data sample shown in Figure 2 suggests fitting two waves.

Figure 5. *The two Single Wave fits to the Global CO_2 ppm sample*

Table 3: *Parameter estimates obtained by dividing full GlobalCO₂ sample into 2 subsamples and fitting a single wave separately to each subsample.*

Parameters	1-76: Wave 1	77-145 Wave 2
σ	1.46	675.0
a	111.0	0.0269
b	0.0394	0.0269
d	-0.0196	-0.784
τ	1939.0	208.0

Use of these single wave parameter estimates as initial values produces the *N=2* Multi-Wave fit shown in Figure 6 below.

Figure 6. *The N=2 Multi-wave fit to the full Global CO_2 ppm sample
and the corresponding estimates on the Temperature using the SL method
and the fit using the T-Quarticmethod. Also shown are the Jonas fit (the
most optimistic) and the Denning calculation. The single point is the current
Temperature.*

Also shown in the Figure are the CO_2 fits using the SL method and a fit reflecting the quartic terms of the Jonas calculation. These are discussed in the next Subsection.

4.3 Temperature Calculation

In this subsection we explain the calculation of the change in temperature from 1800 to 2022 that is also plotted in Figure 6. We make the simple first order linear assumption that $T(t)$ the average global temperature in ^0C is

$$T(t) = \alpha + \beta D(t) = \alpha + \beta(C(t) - 280). \qquad (5)$$

From 1800 to 2022, $C(t)$ is global concentration level in ppm at time t has increased by about 48% from 280 ppm to 420 ppm, an increase of (420-280)/280 = 0.5, whilst $T(t)$ has increased by about 1.5 degrees to 3.9 degrees. We have $C(t) = D(t) + 280$ so from Equation 5 we have

$$\alpha = 2.4 \text{ and } \beta = \frac{1.5}{140} = 0.0107$$

Jonas gives a formula

$$Rc_y = 5.35 * ln(C_y/C_0) - j * ((T_0 + Tc_{y-1})^4 - T_0^4)$$

giving the between the yearly change in CO_2 in terms of the corresponding change of year. Simply as reflection of this formula, but without using any of the considered reasoning given in (Jonas, M) we used following quartic polynomial giving the reverse calculation:

$$C(T) = a + b(T - f) + c(T - f)^2 + d(T - f)^3 + e(T - f)^4$$

It will be seen in Figure 6 that the calculation gives comparable results to the other fits. The difference of the curve based on Equation (5) and the other fits arise from different base T values used as the initial year.

5 ERROR ESTIMATION

5.1 Bootstrap Analysis

We use bootstrap analysis to calculate confidence levels of the fitted quantities. The method is well-known (see for example, Cheng 2017; Dye et al. 2020, Cheng et al. 2020), so full details are not repeated here.

In summary, to estimate confidence levels for $D(t)$, we generate a parametric bootstrap sample ,$E(t_i)$ $i= 1,2,…,n$, where

$$E(t_i) = D\left(t_i, \widehat{\boldsymbol{\theta}}\right) + \varepsilon_i, \ i = 1,2, …, n$$

and $\widehat{\boldsymbol{\theta}}$ is the ML estimate of $\boldsymbol{\theta}$, and

$$\varepsilon_i \sim NID(0, \sigma^2), \ i = 1,2, …, n,$$

is a random sample of mutually independent distributed, normal pseudorandom variables with variance σ^2.

This is carried out B times so that we have

$$\{ E^{*(j)}(t_i) = D^{(j)}\left(t_i, \widehat{\boldsymbol{\theta}}\right) + \varepsilon_{(i)}^{(j)}, i = 1,2, …, n \ \}, \ j=1,2,…,B$$

with the asterisk denoting that the observation is bootstrapped.

We can now estimate the parameters from each of the samples giving the bootstrap parameter estimates and bootstraps functions:

$$\widehat{\boldsymbol{\theta}}^{(j)}; \{ D^{(j)}(t) = D(t, \widehat{\boldsymbol{\theta}}^{(j)}) \ i = 1,2,…, n \ \}.$$

From these a confidence interval can be obtained for each parameter using the ranked values of that parameter.

A similar method can be carried out for $D(t)$. Note however that this only gives the confidence interval level at each t_i separately. the confidence level is reduced if several different t_i are considered simultaneously. A more complicated method is needed using maximized likelihood regions. See Cheng (2017) for details. We could have used the simpler method in this paper, but made a modification to model the dependence of an observation on previous observations. Specifically the observations assumed to be a first order autoregressive process : Thus we set

$$E(t_i) = \alpha E(t_{i-1}) + \beta \varepsilon_{(i-1)}, i = 1,2,…, n.$$

The standard deviation of an observation is therefore not σ, but is s, which is approximately

$$s = \frac{\beta}{1-\alpha}\sigma$$

Thus values of $\alpha = 0.75, \beta = 0.25$, would make $s = \sigma$.

In the experiments reported here we actually varied the value of β slightly to remove possible bias in the estimate of σ. This is not of great practical consequence as our results are only for discussion and are not used in practice. In any case our interest is centred on the SL parameters, and a well known property is that the estimate of σ is asymptotically independent of the other parameters.

5.1 Covid 19 Bootstrap Results

Figure 7 illustrates the results of a representative set of three bootstrap scatterplots of pairs of parameters, obtained from $B = 50$ bootstraps. These indicate the typical scatter.

Figure 7. *Some Bootstrap Scatterplots*

Figure 8 gives shows the resulting $D(t)$ plots of the MLE and the Upper and Lower CI curves with confidence level 90%.

Figure 8. *The N=3 Multi-wave fit to the COVID-19 sample*

5.1 CO₂ Bootstrap Results

Figure 9 illustrates the results of a representative set of eight bootstrap scatterplot pairs of parameters obtained from $B = 50$ bootstraps. These indicate the typical scatter.

Figure 9. *Some Scatterplots for CO₂ Example*

Figure 10. *The N=2 Multi-wave fit to the CO₂ sample*

===

6   IMPLICATONS OF CO₂ SCENARIO

To understand the impact that *Homo sapiens* has had on the earth's climate it is necessary to deal with very large numbers. The weight of the earth is 5.97×10^{24} kg of which 9×10^{19} kg is carbon in the earth's crust. The carbon in plants weighs 4.50×10^{14} kg very much less than in the earth's crust. The carbon in 1 ppm CO_2 in the atmosphere weighs 2.12×10^{12} kg. Before the industrial revolution, the concentration of CO_2 in the atmosphere fluctuated, between about 190 ppm and 270 ppm for 800k years, with a period of about 100k years, while the rises tended to be faster than the declines. The weight of carbon in the atmosphere was therefore 4.87×10^{14} kg, about the same as the weight of carbon in plants, fluctuating from about 4.03×10^{14} to 5.72×10^{14} kg.

Since the industrial revolution the concentration of CO_2 in the atmosphere is now about 415 ppm (Figure 6) as the result of burning fossil fuels, so that the weight of carbon has increased to 8.80×10^{14} kg or about twice weight of carbon in plants. The consequence of this is that the earth's atmosphere has already experience an increase in temperature of about 2°C (Figure 6) with the prospect that, if the concentration of CO_2 continues to increase, following the current trend, the temperature could increase by 6°C above preindustrial levels which would be catastrophic for our survival. We therefore need to sequester CO_2 on a massive scale.

In order to return to pre-industrial levels of atmospheric carbon and therefore temperatures, we need to remove about 4×10^{14} kg of carbon from the atmosphere or 1.5×10^{10} kg every day from now until the year 2100. The biomass of rain forests is 500 metric tons per hectare or 50 kilograms/m^2 so that one would have to expand the worlds rain forest, which currently occupies a 7 million square metres, by about 12k square meters every day until the end of the century. Efficient ways to sequester carbon from the atmosphere are desperately needed.

A recent discussion suggests that systematically scattering iron-rich dust onto target areas in oceans around the world could sequester perhaps 3×10^{13} kg carbon per year, or about 10^{11} kg per day, if the world's deep oceans were to be treated annually. Using a new method of carbon capture it seems that one could remove carbon at a cost of US\$0.5/kg of carbon which means that it would cost about 10^9 US\$ per day, every day from now until 2100 while the worlds GDP is 2.7×10^{10} per day. These studies suggest that it may be technically feasible to sequester carbon on the necessary scale but this would have to be done on an extraordinary scale and much more efficiently than is now possible. The long term survival of our species depends on it.

==

REFERENCES

Cheng R C H (2017). *Non-Standard Parametric Statistical Inference*. Oxford University Press, Oxford.

Dye C, Cheng R C H, Dagpunar J and Williams B G (2020). The scale and dynamics of COVID-19 epidemics across Europe. *R. Soc. Open Sci.* . **7**: 201726 https://doi.org/10.1098/rsos.201726

Jonas, M. The Mathematics of Carbon Dioxide, Part 1. https://archive.ph/FC7zE

AUTHOR BIOGRAPHIES

RUSSELL CHENG retired from the University of Southampton in 2007 where he had been Head of the Operational Research Group, having held previous positions at Cardiff University and the University of Kent at Canterbury. https://www.southampton.ac.uk/maths/about/staff/rchc.page

BRIAN WILLIAMS is Senior Research Fellow at the South African Centre for Epidemiological Modelling and Analysis (SACEMA) having held the position of Epidemiologist at the World Health Organization from which he retired in 2008.