Novel Community-Based Rabies Surveillance (CBRS) In Urban Lilongwe, Malawi

Precious Innocent Mastala[¶1,2*], Melaku Tefera[¶1†], Edson Chiweta[¶2], Madeline Nyamwanza[¶2], Thoko Kapalamula[¶1]

[1]Faculty of Veterinary Medicine, Lilongwe University of Agriculture and Natural Resources, Lilongwe, Malawi
[2]Lilongwe Society for the Protection and Care of Animals, Lilongwe, Malawi

*Corresponding Author

Precious Innocent Mastala

Email: preciousmastala@gmail.com

[¶]These authors contributed equally to this work.

Abstract

Introduction: Rabies is a neglected tropical disease that is endemic to Malawi, killing almost 500 people every year. Domestic dogs are implicated as the main reservoir for rabies. However, there is limited capacity to detect canine rabies in the country, particularly in urban areas where there are high dog densities. The Lilongwe Society for the Protection and Care of Animals designed and implemented a canine community-based rabies surveillance (CBRS) in Lilongwe city between January and June 2020.

Objectives: This study assessed the effectiveness of the CBRS program in canine rabies detection compared to the traditional passive routine surveillance data of the years between 2015 and 2019. Furthermore, the study determined rabies Post-Exposure Prophylaxis (PEP) issuance trends in emergent cases.

Methods: Suspected rabies cases were recorded and followed. The dogs that were found alive were quarantined for 10 days for observation. Brain samples were collected post-mortem for direct-fluorescent antibody testing for antigen detection.

Results: There were 610 suspected cases, and only 217 (39.2%) were followable. Out of these, 150 (62.8%) were followed. A total of 9 dogs (6%) tested positive, and there were 11 (7.3%) probable cases. This represents a 4.97-fold increase in rabies cases detected compared to the pre-surveillance period, with a resultant of 3.33 cases/month from 0.67 cases/month. The program also helped to initiate PEP treatment in exposed people. There were 141 bite victims and 136 (96.5%) reported for PEP. Out of these, 89 (65.4%) were from healthy bites and 33 (24.4%) from suspect cases indicating a **63.97% to 88.24%** rate of irrelevant PEP issuance.

Conclusion: The CBRS program established relevance in increasing canine rabies detection and subsequent prevention of related human infections. It also established a high rate of PEP issuance following healthy bites. Therefore, we recommend enhanced multi-sectoral collaborative efforts aimed at increasing rabies detection and optimizing PEP usage.

Keywords: Community-based rabies surveillance, Lilongwe, Malawi, One Health, Post-Exposure Prophylaxis, Rabies

One Health Impact Statement

This novel implementation research features a One Health approach in which several arms of the public and private sectors including the community collaborate towards ending rabies. This collaboration provides synergy towards accelerating the identification of rabid animals in the community and effectively guides people to take the necessary precautions including prompt Post-Exposure Prophylaxis in cases of exposure. This ensures ultimate health promotion at the human-animal interface.

Introduction

Rabies is a viral zoonotic encephalitis infection of the central nervous system of homeothermic animals, including humans, that usually ends in the death of the host once clinical symptoms appear (Quinn *et al.*, 2016). The virus is usually transmitted to humans through contact with infected animals, primarily by bites, scratches, or other contact with saliva (Health Protection Agency, 2013).

In Malawi, rabies is endemic and often referred to as "a notifiable disease" (Mallewa *et al.*, 2007). Despite that, there are about 59,000 and 500 (WHO, 2015) human deaths globally and in Malawi per annum, respectively. About 99% of these human rabies cases can be mostly attributed to dogs. Rabies is neglected in most developing countries, (Hampson *et al*, 2015).

Rabies causes huge losses, which can be attributed mainly to premature death, but also direct costs of Rabies post-exposure prophylaxis (PEP), additional costs to communities from livestock losses, lost income whilst seeking Rabies PEP, and very little in dog vaccination. Despite the losses, rabies prevention and control can be easily addressed by increasing awareness, vaccinating 70–80% of the dog population and prompt rabies PEP (WHO *et al*, 2018). Thus, controlling dog rabies substantially reduces human exposure to the virus.

Despite the important role of dogs in the transmission dynamics of the virus, issues surrounding the propagation and maintenance of dog rabies are still not clear. Achieving the 'Zero canine rabies-related human deaths by 2030' goal requires effective and well-coordinated surveillance of the disease (Bourhy et al, 2016). In Malawi, there is no nationwide instituted rabies surveillance program, but specific district non-governmental organizations run surveillance programmes in southern and central Malawi. Samples from suspected cases are mostly delivered for rabies testing through well-informed owners or consulting veterinary or paraveterinary professionals. This leaves room for the cycle of neglect and, as such, undermines the true burden of the disease on the population. Several elements can be attributed to the failure in control and elimination of rabies, these include: failure to employ effective tools; organizational failure in the planning, implementation, or evaluation of the rabies program specifically, a failure to gather and properly present information about disease risk and vulnerabilities, and a failure to generate systematic synergies with other programs and stakeholders to ensure efficient capacity deployment, and conduct thorough risk assessment on the sustainability of the rabies program (Hampson *et al*, 2015). The reporting of the disease is hindered by difficulties with the system and the lack of diagnostic facilities, (Mallewa *et al*, 2007).

Malawi has a One Health approach to addressing rabies which is played through a coordination between state animal health and public health departments. This coordination ensures that when a person is potentially exposed to the rabies virus through an animal, they should report to the veterinary department for advice and follow up of the suspected case where possible. Depending on the case, the veterinarian recommends PEP treatment for the victim. The global call for the elimination of dog-mediated human rabies by 2030 has prompted many countries to invest in dog vaccinations (Sambo *et al.*, 2013), including the Malawi government, with alliances with two non-governmental organisations; Lilongwe Society for Prevention and Care of Animals and Mission Rabies, where Memorandums of understanding (MOU) are in place for rabies surveillance and

monitoring (WHO, 2019). This will revamp the One Health platform, which has long existed and
encourage surveillance. The platform ensures that there is Integrated Bite Case Management
(IBCM), where every dog bite case is systemically followed by both medical and veterinary
support, and necessary management is done following each dog's confirmation of rabies infection.

Employing the Community Based Rabies Surveillance (CBRS), where the community is at the
core, could be important in detecting possible rabies cases and reporting them for confirmation
and intervention systems. This paper details the process of developing the program, which
comprised of both active and passive surveillance components. The active surveillance was done
involved actively looking for rabies suspicion cases in the communities, while the passive
surveillance involved cases reported to the surveillance team on bite and PEP cases reported by
the DALHLD and the referral hospital for PEP respectively. The study involved the assessment
of data on animal rabies cases between January and June 2020, and a comparison to the pre-
surveillance period of 2015–2019. It also assesses the rabies PEP issuance trends for the study
period.

Materials and Methods: Study Design, study area, and sample population

The study was conducted in urban Lilongwe, a district in the central region of Malawi (Figure 1).
According to the National Statistical Office (NSO), (2018), the government of Malawi has
230,265 households with a total population of 989,318.

Figure 1: The study area: Lilongwe City with its designated areas of residence

CBRS Design
The surveillance program was piloted from January to June 2020. The stakeholders of the
program were the Lilongwe Society for the Protection and Care of Animals (LSPCA), the District

Animal Health and Livestock Department (DAHLD), the Lilongwe District Health Office (DHO), and Lilongwe City Council (LCC). The LSPCA was the lead of the program. The DAHLD reported cases of suspect human-animal exposures, the DHO helped to report cases of human exposures presenting for rabies PEP at the hospital, and the LCC helped in reporting roadside killed dogs. The data captured and reported for follow-up included each person's name, phone number, and physical address. The LSPCA conducted the follow-ups of all reported cases (Figure 2).

Taskforce

The program taskforce had four rabies project officers (RPOS) with a para-veterinary profession, a veterinarian, the DHO, the DAHLD (The bite reporting section and the Central Veterinary Laboratory), and Community Rabies Action Groups (RAGs). The members of the RAGs included community leaders, community police, human and animal health surveillance officers, church leaders, school managers and market vendors association chairpersons.

Procedures

In December 2019 the RPOs were trained on the principles of rabies and its prevention in both animals and people. After the training, they received rabies pre-exposure vaccinations, and additional training in animal rabies surveillance and bite investigation. Since suspected cases were to be reported and handled through the LSPCA, a toll-free number (172) was established under one of the Network Proving Companies called TNM (Telecommunications Malawi), followed by publicity of the program to the communities through radios, televisions, newspapers, flyers, and other document materials and roving.

The RPOs were stationed in nine areas that were distinguished as hotspots of human exposures using the 2018 data obtained from DAHLD on dog bites, and DHO on rabies PEP treatment. The RPOs roved around the areas to identify any rabies suspected animals. Further to this the RPOSs followed up on reported exposure cases from the DAHLD and DHO first via telephone followed by physical visit to the reported site regardless of being a hotspot or not. The follow-up efforts were done to locate the infected animal. All animals cases were diagnosed and defined of their rabies status based on the case definition by Wallace *et al*, (2015) in the following box (

Table *1*).

Table 1: Animal Rabies Surveillance Program Case Definitions

Animal Rabies Surveillance Program Case Definitions
Confirmed Rabies Cases
▪ Diagnostic confirmation of rabies virus by Direct Fluorescent Antibody Test (dFAT)
Probable Rabies Cases
▪ Animals that were not tested for rabies and
▪ Animals that died during the observation

- Animals that did not pass observation (escaped animals, not found by further investigation)
- Animals that developed one or more clinical signs and died after being bitten by a suspect/probable/confirmed rabid animal

Suspect Rabies Cases
- Animals reported to animal rabies Project Officers that could not be assessed or
- Animals that had less than 2 signs of rabies and test results were inconclusive
- Animals that died after being hit on the road

Non-cases
- Animals that are healthy after the 14-day observation period or
- Animals that test negative by dFAT

Depending on the initial case definition, the animal would be kept under home quarantine for a
minimum of 10 days if it did not show obvious clinical signs of rabies. Within the quarantine
period, the animal would be followed to note any changes towards rabies definition. If after the
period or during an investigation the animal showed clinical signs, the animal was humanely
captured and transported to LSPCA Clinic for euthanasia.

The euthanasia of the animal involved a series of steps as recommended by the World Society
for the Protection of Animals (WSPA) (2010). Following euthanasia, the animal was tagged with
an ID, properly wrapped and placed in a sample bag, and transported to CVL where dFAT was
performed on a brain sample.

Test results from the laboratory were reported through a WhatsApp forum to the stakeholders
and by telephone call to the bite victim and reporting health facility or in person where telephone
communication was not possible. If a case tested positive, the bite victim would be encouraged
to continue with PEP treatment or start the treatment if not started yet.

Figure 2: The CBRS Case Reporting and Investigation structure infographic

Results

Active and Passive Surveillance

During the surveillance period, a total of 610 cases were reported during the surveillance period, of which 239 (39.2%) were followable, and only 150 (24.6% of the total and 62.8% of the followable cases) cases were followed (

Figure *4*). Of the followed cases, 46 were under active surveillance and 104 through passive

surveillance. 13 (8.67%) were tested cases and 9 (69.23%) were confirmed (Figure 4). All cases

sent for testing were from Active surveillance.

Overall, there were 150 cases, 9 (6%) positive cases, 11 (7.3%) probable cases, 33 (22%) suspect

cases and 97 (64.6%) non-cases. This entails that the program contributed 20 (13.6%) positive

and probable cases. The active surveillance contributed 75% (28.6% of total active surveillance),

whilst, the passive surveillance contributed 35% (6.7% of total passive surveillance cases).

According to Table 2, the animals involved in bites were dogs, and only 2 cats were reported.

Notice that there was 1 confirmed donkey case detected through active surveillance but was not

included in the study because it was not detected in the area of study.

Figure 3: Figure shows the number of cases that were reported during the study period versus the cases that were followable and actually followed. It also splits the followed cases into the actual case definitions

179
180
181

183   Figure 4: Reported Cases

185   Table 2: Animals followed during the investigations

Figure 5: Investigated cases during the study period and the outcome diagnosis based on case definitions

Species	Case Review			
	Non-Cases	Confirmed	Probable	Suspect
Dog	96(64.9%)	9 (6.1%)	10(6.8%)	33(22.3%)
Cat	1 (50%)	0(0%)	1(50%)	0(0%)
Total (n (%))	97 (64.7)	9(6%)	11(7.3%)	33(22%)

The most common signs of illness among all defined cases as shown in Table 3, were, unusual
aggression and biting, with exception of Paresis/paralysis (Odds Ratio (OR)=.039-.148 and .045-
.159), hypersalivation (OR=2.994-71.4 and 3.46-66.667), restlessness (OR=2.188-29.411 and
1.656-45.454) and anorexia (OR=2.105-62.5 and 236-22.727) which were common among

confirmed and probable cases respectively. Lethargy (OR=6.758-29.4) and hydrophobia
(OR=6.757-25.64) were among the confirmed cases only.

Table 3: Clinical signs observed during the investigation of rabies suspect animals by case status

Observed Signs	Case Review						
	Non-Cases (n (%)) #	Confirmed (n (%))	ODDS RATIOS (OR)	Probable (n (%))	ODDS RATIOS (OR)	Suspect (n (%))	ODDS RATIOS (OR)
Paresis/ paralysis	0 (0%)	1(11.1%)	.076(.039-.148)	2(18.2%)	.085(.045-.159)	0 (0%)	§
aggression	11(11.6%)	4(44.4%)	6.25(1.457-27.027)	5(45.5%)	6.536(1.701-25)	1(3%)	.244(.03-1.969)
Abnormal vocalisation	5(5.3%)	2(22.2%)	5.263(.86-32.258)	2(18.2%)	4.082(0.691-24.39)	0 (0%)	1.359(1.224-1.508)
Biting	94(98.9%)	6(66.7%)	.064(.011-.386)	9(81.8%)	0.134(.021-.975)	33(100%)	.74(.668-.82)
Hypersalivation	5(5.3%)	4(44.4%)	14.706(2.994-71.4)	5(45.5%)	15.385(3.46-66.667)	0 (0%)	1.358(1.224-1.508)
Lethargy	0 (0%)	1(11.1%)	13.158(6.758-29.4)	0 (0%)	§	0 (0%)	§
Restlessness	4(4.2%)	1(11.1%)	2.9(2.188-29.411)	3(27.3%)	8.696(1.656- 45.454)	0 (0%)	1.355(1.221-1.504)
Anorexia	4(4.2%)	3(33.3%)	11.628(2.105-62.5)	1(9.1%)	2.326(.236-22.727)	0 (0%)	1.355(1.221-1.504)
Hydrophobia	0 (0%)	1(11.1%)	13.158(6.757-25.64)	0 (0%)	§	0 (0%)	§

#Non-cases category was used as a reference for comparison

§ No statistics were computed because the designated sign was a constant.

There were 141 bite victims. Out of these, 91 (64.54%) were from healthy bites; 8 (5.67%) from confirmed cases; 9 (6.38%) from probable cases; and 33 (23.4%) from suspect cases. Out of these in the different exposure categories, 136 (96.45%) of the bite victims reported rabies PEP. Out of the 136, 87 (63.97%) from the non-cases, 7 (5.15%) from confirmed cases (1 person was reluctant to go for PEP after communication of lab results and advice due to religious convictions), 9 (6.62%) from probable cases, and 44 (32.35%) from suspect cases. This indicates between 63.97% to 88.24% of PEP issuance following healthy bites.

Rabies Testing at CVL During the CBRS Study Period and Pre-Surveillance Period

A total of 22 suspected case samples from Lilongwe urban were tested at CVL of these 13 cases were from active surveillance. 10 (45.5%) of the tested samples were positive and 12(54.5%) tested negative.

Prior to the surveillance period (2015-2019), the LSPCA and government carried out mass rabies vaccinations with a passive surveillance system. During this period there were a total of 40 confirmed cases, averaging 8 cases per year and 0.67 cases per month (Figure 6).

Figure 6: Camparison of rabies dFAT tests for cases between the pre-surveillance and surveillance period at Central Veterinary Laboratory

Spatial Distribution of Rabies

In Figure 7, confirmed and probable cases were detected from Area 25, Area 18, Area 22, Area 9, Area 22, Area 23, Area 24, Area 36, Area 39, Area 49, Area 50, Area 56, and Area 57.

Figure 7: The spatial distribution of rabies cases during the study period

Discussion

In low-resource settings like in Malawi, the absence of reliable surveillance data makes it difficult to monitor and understand the disease burden, and implementing appropriate prevention and control measures. The CBRS under the case definitions justified by the Odds Ratios as by Wallace *et al*, (2015), identified 20 confirmed and probable cases within the study period. This indicates a 3.3 cases per month compared to the pre-surveillance period which had 0.67 cases per month and translates into a 5-fold increase in cases. The surveillance program also helped to initiate PEP treatment in exposed individuals. This indicates the importance of surveillance in the efficient detection of rabies cases which without the CBRS could not be detected as was also found by Wallace *et al*, (2015) who with surveillance there was a 10-fold increase in the detection of rabies in animals within the first 2 years of the surveillance. Herein, we demonstrate that data from

CBRS can help to reduce state losses from human death in exposed individuals, as there is a
follow-up to encourage the exposed individuals to go for PEP treatment.

When compared to the reported cases in this study, we saw a significant decrease in the number
of cases that were followed. Only 150 (24.6%) of the 610 reported cases were followed. This
was due to the fact that cases from DHO and DALHD were frequently reported many days after
they had been recorded in the respective departments. Additionally, it was challenging to follow
these cases because the majority of the reported data from DALHD lacked the contact
information for victims and dog owners, as appropriate. Additionally, some residents of the
community were unaware of the surveillance program, which led them to dismiss any phone-
based investigation into the cases because they believed it might be a common study period
fraudster scam. There are frequently few perceived incentives for carrying out rabies surveillance,
as mentioned by Wallace et al. (2015), Lembo et al. (2010), and Kitala et al. (2000), in the absence
of systems capable of prompt investigation and reporting for clinical and public health purposes.
Rabies and animal bites have a significant impact on public health. Therefore, accurate and timely
data are essential to understanding the epidemiology and burden of rabies (Adomako et al., 2018).
This means that poor and discrepant data underestimate the true burden of rabies and negate
the advocacy efforts needed to achieve control and elimination (Scott, et al 2017). Subsequently,
this can result in a lack of awareness of the case burden, reduced funding for control, and poor
community engagement around prevention.

247
The study found that the majority of people who sought rabies PEP had not been exposed. It was
discovered that only 14.18% of the population had been exposed to positive and probable cases.
Additionally, we discovered that some religious beliefs have an impact on prompt PEP treatment.
These religious convictions put the believers at risk to dying from the disease. The established
One Health platform for rabies in Malawi assures that the state veterinarians and para-
veterinarians make the decision regarding whether or not PEP should be administered to the
bitten victim(s) in addition to wound care. The observed irregularity in PEP issuance to mostly
healthy bite victims may indicate a serious issue with the veterinary practitioners' decision-making
and necessitates an investigation of what influences their decisions. But it should be noted that in
regions like Malawi where rabies is enzootic in dogs and wildlife, the WHO advises that PEP be
implemented right away unless sufficient laboratory surveillance and data show that the species
in question is not a rabies vector (WHO (2014), Ngugi et al, (2016) and Obonyo et al, (2016)).
Likewise, even if the suspect animal is not available for testing or observation, rabies PEP
administration should still begin. However, the same may be discontinued if the animal involved
is vaccinated and, following observation for 10 days, the animal remains healthy or is humanely
killed and declared negative for rabies by a WHO-prescribed laboratory test.

Given the scarcity of resources, PEP must be used more effectively for those who have been
exposed. To reduce inappropriate PEP administration, Moore (2000) recommends increasing
vaccination rates for pet (owned) animals. Furthermore, with advanced training of the staff and
adhering to WHO recommendations, this could be accomplished. There is a need to teach the

communities about responsible dog ownership including the need for dog population management and vaccination and animal behaviour to reduce and prevent animal-bite cases.
To ensure that the exposed individuals and the anti-rabies dispensary are promptly informed to stop the PEP if the involved animal is deemed healthy, it is important to establish good coordination between the involved parties, including the diagnosing laboratory, the veterinary team, the human health department, and the community/patient. It is also important to educate communities on responsible dog ownership including the need for dog population management and vaccination and animal behaviour to reduce and prevent animal-bite cases.

The highest rabies cases were recorded during the study period in Areas 25, Area 18, and Area 22, according to the spatial distribution. Adjustments to comprehend the true significance of the observations are however prevented by the lack of statistics or studies on dog populations. The majority of nations around the world lack such data on pet populations (Carvelli et al. 2020). It is crucial to have thorough knowledge of zoonoses, infectious diseases, and risk factors associated with their spatial and temporal distribution in the pet population (Day et al. (2012) and Carvelli et al. 2020). As seen here, one of the most important tools for comprehending the population's burden is knowledge of animal populations' sizes, demographics, and ownership patterns.

Limitation
Due to COVID-19-related public health restrictions, the surveillance team's activities were heavily affected and as such some cases could have gone unreported.

Conclusion and Recommendations

This is the first time to employ a community based canine rabies surveillance in Malawi. The study has illustrated how a multisectoral, One Health approach can enhance canine rabies surveillance. The CBRS program has demonstrated its value in enhancing the detection of canine rabies cases and assisting the exposed victims in initiating PEP for rabies in a timely manner. Additionally, a high rate of PEP issuance following healthy bites has been established. In order to increase rabies detection and maximize PEP use, we advise increased multi-sectoral collaborative efforts. Every major stakeholder must work together to strengthen surveillance, make the best use of data, and maximize the use of available resources if the goal of ending rabies-related human deaths is to be accomplished.

Funding statement
This research was funded by the Lilongwe University of Agriculture and Natural Resources (LUANAR), the Lilongwe Society for the Protection and Care of Animals and the Martyn Edelsten (of The University of Edinburgh) Research and Internship Grant Fund.

Conflict of interest declaration
The authors declare that there is no conflict of interest

Ethics statement confirming that all relevant guidelines were followed

Ethical Approval for research was provided by Animal Health Committee reference number DAHLD/AHC/12/2019/1.

Acknowledgements

We would like to thank Kondwani Chiumya and the entire Lilongwe Society for the Protection and Care of Animals (LSPCA) team for their expertise and assistance throughout all aspects of this study.

References

Adomako B.Y., Baiden F.,Sackey S.,Ameme D.K.,Wurapa F.,Nyarko K.M.,Kenu E., and Afari E. (2018) *Dog Bites and Rabies in the Eastern Region of Ghana in 2013–2015: A Call for a One-Health Approach*. Hindawi, Journal of Tropical Medicine, Volume 2018, Article ID 6139013, https://doi.org/10.1155/2018/6139013.

Hampson K., Coudeville L., Lembo T., Sambo M., Kieffer A., Attlan M., Barrat J., Blanton J.D., Briggs J.D., Cleaveland S., Costa P., Freuling C.M., Hiby E., Knopf L., Leanes F., Meslin F., Metlin A., Miranda M.E., Muller T., Nel L.H., Recuenco S., Rupprecht6 C.E., Schumacher C., Taylor L., Vigilato M.A.N., Zinsstag J., Dushoff J. (2015). *Estimating the Global Burden of Endemic Canine Rabies*. PLOS Neglected Tropical Diseases, DOI: 10.1371/journal.pntd.0003709

Health Protection Agency (HPA) (2013). *HPA guidelines on managing rabies post-exposure prophylaxis: January 2013*. VW-1197.01.

Kitala P.M., McDermott J.J., Kyule M.N., Gathuma J.M. (2000). *Community-Based Active Surveillance for Rabies in Machakos District, Kenya.* Prev Vet Med, 2000. 44(1–2): p. 73–85. PMID: 10727745

Lembo T., Hampson K., Kaare M.T., Ernest E., Knobel D., Kazwala R.R.,Haydon D.T., Cleaveland S. (2010). *The feasibility of canine rabies elimination in Africa: dispelling doubts with data*. PLoS Negl Trop Dis, 2010. 4(2): p. e626. doi: 10.1371/journal.pntd.0000626 PMID: 20186330

Mallewa M., Fooks A.R., Banda D., Chikungwa P., Mankhambo L., Molyneux E., Molyneux M.E., and Solomon T. (2007). *Rabies Encephalitis in Malaria-Endemic Area, Malawi, Africa.* Emerging Infectious Diseases. 2007; 13(1):136–9. PMID:17370529.

Moore D.A. (2000). *Animal bite epidemiology and surveillance for rabies postexposure prophylaxis*. J

Am Vet Med Assoc, Volume 217.

National Statistical Office (NSO) (2019). *Malawi Population and Housing Census: Main Report.*

Government of Malawi.

Ngugi J.N., Maza A.K., Omolo O.J. and Mark Obonyo M. (2018). *Epidemiology and surveillance of*

human animal-bite injuries and rabies postexposure prophylaxis, in selected counties in Kenya, 2011–

2016. BMC Public Health, 18:996 https://doi.org/10.1186/s12889-018-5888-5.

Obonyo M, Akoko JM, Orinde AB, Osoro E, Boru WG, Njeru I, Fèvre E.M. (2016). *Suspected*

rabies in humans and animals, Laikipia County, Kenya. Emerg Infect Dis. 22(3):551-553

Quinn P.J., Markey B.K., Leonard F.C., FitzPatrick E.S., Fanning S, and Hartigan P.J. (2016).

Concise Review of Veterinary Microbiology. 2nd edition. John Wiley & Sons Ltd, The Atrium,

Southern Gate, Chichester, West Sussex, PO19 8SQ, UK.

Sambo M, Cleaveland S, Ferguson H, Lembo T, Simon C, Urassa H. and Hampson K. (2013).

The burden of rabies in Tanzania and its impact on local communities. PLoS Negl Trop Dis 7: e2510.

doi: 10.1371/journal.pntd.

Scott T.P., Coetzer A, Fahrion A.S., and Nel L.H. (2017). *Addressing the disconnect between the*

estimated, reported, and true rabies data: the development of a regional African Rabies Bulletin,"

Frontiers in Veterinary Science, vol. 4.

Wallace R.M., Reses H., Frank R., Dilius P., Fenelon F., Orciari L., Etheart M., Destine A.,

Crowdis K., Blanton J.D., Francisco C., Ludder F., Vilas V.D.R., Haim J, and Millien M. (2015).

Establishment of a Canine Rabies Burden in Haiti through the Implementation of a Novel Surveillance

Program. PLoS Negl Trop Dis 9(11): e0004245. doi: 10.1371/journal.pntd.0004245.

World Health Organization (WHO) (2014) *WHO Guide for Rabies Pre and Post-Exposure*

Prophylaxis in Humans. https://www.who.int/rabies/PEP_Prophylaxis_guideline_15_2014.pdf

Accessed on 12/09/2020.

World Health Organization (WHO) (2015). *Rabies in Malawi: A mother's first-hand account.*

Accessed from https://www.who.int/news-room/feature-stories/detail/rabies-in-malawi-a-

mother-s-first-hand-account on 16th January, 2020.

World Health Organization (WHO) (2019). *Joint External Evaluation of IHR Core Capacities of the*

Republic of Malawi. World Health Organization.

World Health Organization (WHO), Food and Agriculture Organization of the United Nations

(FAO) and World Organisation for Animal Health (OIE) (2018). *Zero by 30: The Global Strategic*

Plan to End Human Deaths from Dog-Mediated Rabies by 2030.

World Society for the Protection of Animals (WSPA). (2010). *Surveying roaming dog populations:*

guidelines on methodology 2010. World Society for the Protection of Animals

Carvelli A, Scaramozzino P, Iacoponi F, Condoleo R, Della Marta U (2020) Size, demography,

ownership profiles, and identification rate of the owned dog population in central Italy. PLoS

ONE 15(10): e0240551. https://doi.org/10.1371/journal.pone.0240551

Day MJ, Breitschwerdt E, Cleaveland S, Karkare U, Khanna C, Kirpensteijn J, et al. Surveillance

of zoonotic infectious diseases transmitted by small companion animals. Emerg Infect Dis. 2012;

2022. http://dx.doi.org/10.3201/eid1812.120664