1   **COVID-19 in Cambodia: Epidemiology, Response, and Lessons Learned, 27 January**

2   **2020–30 June 2022**

3
4

Authors: Srean Chhim[1], Grace Marie Ku[2], Sovathiro Mao[1], Willem van de Put[2], Wim Van

Damme[2, 3], Por Ir[1], Chhea Chhorvann[1], Vandine Or[4]

Affiliations:

[1] National Institute of Public Health, Phnom Penh, Cambodia

[2] Institute of Tropical Medicine, Antwerp, Belgium

[3] Vrije Universiteit Brussel, Belgium

[4] Ministry of Health, Phnom Penh, Cambodia

13
14
15   *** Corresponding author:** Srean Chhim (Lot # 80, Samdach Penn Nouth Blvd. (St. 289), Phnom
16   Penh, postal 120804), srean168@gmail.com
17
18   *Keywords*: COVID-19, SARS-CoV-2, Cambodia
19
20   *Word count 4063/5000*
21
22

ABSTRACT

Objective: This review describes the COVID-19 epidemiology and response between 27 January 2020 and 30 June 2022, and analyses the lessons learned for future public health responses in Cambodia.

Methods: We analysed the national COVID-19 database and reviewed literature.

Results: We classified Cambodia's COVID-19 epidemiology into three phases. Phase 1 (27 January 2020–19 February 2021) was when COVID-19 was well contained with only 578 cases, mainly imported (n = 320, 55.4%), and no deaths. Phase 2 (20 February 2021–31 October 2021) was when a large community outbreak occurred, with 255,792 cases and 2,805 deaths recorded. Phase 3 (1 November 2021–30 June 2022) was after the country's full re-opening, with 47,314 cases and 251 deaths. In phase 1, Cambodia responded using six non-pharmaceutical measures using the "5 Es" principle—*Early detection, Early isolation, Early tracing, Early treatment, and Early education*. Boosting the vaccination awareness campaign and coverage, and imposing a lockdown were the essential measures added to the above in phase 2. In phase 3, all non-pharmaceutical measures were lifted after high vaccination coverage of 93.1% of the population.

Conclusion: Cambodia successfully contained the spread of SARS-CoV-2 in the first year and quickly attained high vaccine coverage by the second year of the response. The core of this success was the strong political will and high cooperation from the public. However, Cambodia needs to improve its medical infrastructure, human resources, and laboratory capacity for future health emergencies.

Keywords: COVID-19, SARS-CoV-2, Response, Cambodia, Lessons learned

SUMMARY BOX

WHAT IS ALREADY KNOWN ON THIS TOPIC

- COVID-19 spread globally, but how the pandemic played out in each country depended on various factors, including respective government responses and the general public's adherence to COVID-19 measures.

WHAT DOES THIS STUDY ADD

- Early response—*Early detection, Early isolation, Early tracing, Early treatment, and Early education*—is the core of successful SARS-CoV-2 containment.
- Achieving high vaccination coverage quickly leads to a decline in the number of deaths and to eventual full re-opening of the country.
- Responding to the pandemic requires decisive leadership, that refers to decisions being made quickly, in a timely manner, and without delay.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- This study highlights lessons learned and success stories in Cambodia for organising and investing in responses for any future outbtreaks.

INTRODUCTION

Starting in early 2020, COVID-19 spread globally. However, how the epidemic played out in each country depended on various factors, including respective government responses and the general public's adherence to COVID-19 measures.[1]

The early COVID-19 response in Cambodia was not, as yet, been well-documented in scientific publications. While several articles described the early phases of the COVID-19 situation and the results of the COVID-19 responses, including non-pharmaceutical interventions (NPIs) to contain the spread of SARS-CoV-2, many important aspects, such as laboratory utilisation and vaccination, were overlooked.[2, 3]

Our review aims to describe the COVID-19 epidemiology and responses and their effects in preventing and reducing transmission. The review covers the period between 27 January 2020 (when the first case, a 60-year-old Chinese man traveling with his family from Wuhan, China, to Cambodia for a tour, was detected) and 30 June 2022, several months after Cambodia's full re-opening. It also expands the lessons learned from implementing these responses through a policymakers' perspective. This can plausibly help to better prepare for future health shocks.

Setting and context

Located in Southeast Asia, Cambodia borders Thailand, Vietnam, and Laos. Cambodia is a low-income country of 15.55 million people (as of 2019).[4] The overall population density is 87 inhabitants per square kilometre (km^2), with averages of 874 inhabitants per km^2 in urban and 55 inhabitants per km^2 in rural settings. Approximately 60% of the population lives in rural areas.[4] Phnom Penh is the capital city, with a population of 2.3 million in 2019 or 3,361 inhabitants per

km^2.[4] Cambodia's population is relatively young, with people aged 50 years or older

accounting for only 18.4%.[4]

The Cambodian health system was already weak even before the pandemic. In 2014, the

World Health Organization (WHO) estimated that Cambodia had 0.2 physicians per 1,000

population.[5, 6] The ratio of hospital beds was also low at 0.9 per 1,000 population.[6, 7] There

was no infrastructure for isolation of patients with highly infectious diseases.

Prior to the pandemic, it was found that Cambodia was not ready to respond to infectious

disease outbreaks. In 2016, the Joint External Evaluation of the International Health Regulation

(IHR) assessed Cambodia's capacity to prevent, detect, and rapidly respond to public health

threats. The total score was 12.5% out of 100, indicating limited capacity.[8]

Throughout the pandemic, the COVID-19 responses in Cambodia have been constantly

evolving, following the dynamics of the disease and availability of resources and tools (e.g.,

vaccine). In Cambodia, the responses can be classified into three phases: containment phase,

mitigation phase, and post country full re-opening phase.

METHODOLOGY

We analysed secondary data and reviewed relevant literature and documents to describe the

epidemiology, responses, and strategies between 27 January 2020 and 30 June 2022 in

Cambodia.

Data for descriptive epidemiology

Tests, cases, and deaths

The Inter-ministerial COVID-19 Response Committee provided the COVID-19 data for tests,

cases, and deaths recorded between 27 January 2020 and 30 June 2022. The COVID-19 data is

under the governance and stewardship of the Inter-ministerial COVID-19 Response Committee,

with Cambodia's Communicable Disease Control Department (CCDC) being the central body

responsible for collection and storage. The data were stored at the individual level, allowing

them to be stratified by age group, gender, and geographical areas at the provincial and district

levels. These individual data were collected from all laboratories in Cambodia, using a

laboratory form (see supplementary file 1). The completed forms were entered into each

laboratory database and exported to a CSV or Excel file and submitted to the CCDC. At the

CCDC, these files were checked for standardisation, and imported into the National COVID-19

Database.

Data for response and themes

Relevant events

We documented relevant events to draw a more comprehensive picture of the COVID-19

response in Cambodia. Our data team recorded all the policies and guidelines issued by the

government, and triangulated these documents with other documents (e.g., WHO website

updates, situation reports). The information we collected included dates of border closures,

school closures, cancellation/prohibition of public events and gatherings, enforcement of

facemask usage in public areas, and lockdowns.

Vaccination data

The National COVID-19 Vaccination Committee provided vaccination data. These data included

the number of vaccine doses that the government obtained through purchasing and donations,

and can be stratified by vaccine brand and donor country. It also included vaccination coverage

data: the number of vaccinated individuals, and the number of doses administered (e.g., one

dose/incomplete immunisation, two doses/completed immunisation, three doses, four doses, and

five doses).

RESULTS

COVID-19 epidemiology

Figure 1 presents the daily number of COVID-19 cases, tests done, and deaths in Cambodia

between the first documented case (27 January 2020) and 30 June 2022. We identified three

significant timepoints of the pandemic in Cambodia.

Phase 1 (27 January to 19 February 2021): containment phase

Phase 1 lasted for 389 days. COVID-19 was well-contained in this phase. Cambodia had only

578 cases, mainly imported (n = 320, 55.4%), and no deaths (Figure 1 and Table 1).

Phase 2 (20 February to 31 October 2021): mitigation phase

Phase 2 was known as the large community outbreak phase and lasted for 252 days with 255,792

cases (Figure 1 and Table 1). Of these, only a small proportion were imported cases (n = 19,480,

8.2%). A total of 2,805 COVID-19 deaths were reported, giving a case-fatality rate of 1.1%

(Figure 2 and Table 1). Deaths occurred disproportionately in the population aged 50 years and

over, with case-fatality of 4.9% (Figure 3). The most affected period was between March and

October 2021 (Figure 3).

Phase 3 (1 November 2021 to 30 June 2022): full re-opening

Phase 3 had 47,314 cases, in which 3,595 cases (7.6%) were imported (Figure 1 and Table 1). In

this phase, 251 deaths were reported, resulting in a case-fatality rate of 0.5%.

Figure 1. COVID-19 daily cases, tests, and deaths in Cambodia, 2020–2022 *[Insert*

Figure 1]

Table 1. Reported COVID-19 cases and deaths by phase in Cambodia, 27 January 2020 to 30
June 2022

Phase of COVID-19 outbreak	Theme	All cases	Imported cases (%)	Death (%)
Phase 1. 27 January 2020 to 19 February 2021:	Containment with NPIs	578	320 (55.4)	0 (0.0)
27 January–30 April 2020	Initial response	144	88 (61.1)	0 (0.0)
1 May–31 October 2020	Relaxing after long period of low cases	167	85 (50.9)	0 (0.0)
1 November 2020–19 February 2021	Intensifying response for larger scale outbreak	267	147 (55.1)	0 (0.0)
Phase 2. 20 February 2021 to 31 October 2021	Large community outbreak and vaccination	255,792	19,480 (7.6)	2,805 (1.1)
Phase 3. 1 November 2021 to 30 June 2022	Full re-opening	47,314	3,595 (7.6)	251 (0.5)

158   Abbreviations: NPIs, Non-pharmaceutical interventions

160   **Figure 2. The death trends in Cambodia, 2021–2022** *[Insert Figure 2]*

161   **Figure 3. The COVID-19 death rate in percent by age groups in Cambodia, 2021–**

162   **2022** *[Insert Figure 3]*

Response

The government implemented eight significant measures during phases 1 and 2: *(1) detect, isolate/quarantine; (2) face coverings, hand hygiene, and physical distancing measures; (3) risk communication and community engagement; (4) school closures; (5) border closures; and (6) public event and gathering cancellation; (7) vaccination; and (8) lockdown* (Table 2). Each of these measures was implemented when the need for such intervention was seen to be warranted. We note that the first six measures were implemented during phase 1, and measures 7 and 8 were added in phase 2.

Table 2: Description of the eight measures used in Cambodia in response to the COVID-19 outbreak

No	Measure	Description	Timeframe
1	Detect, isolate/quarantine	This measure was used to detect cases for isolation and find contacts for quarantine. Contact tracing was the primary measure to trace the contacts, test them, and quarantine them. Contact tracing commenced within 24 hours after a confirmed case was detected. The contacts were classified into high, medium, and low risk. High and medium-risk contacts were tested at least two times during the quarantine. Low-risk contacts were tested if they had any signs or symptoms during the quarantine period.[9]	▪ Phase 1 and early phase 2
2	Face coverings, hand hygiene, and physical distancing	This measure was known as "3 DOs and 3 DON'Ts." The "3 DOs" refer to (1) wearing a face mask properly, (2) washing hands frequently with alcohol, gel, or soap, and (3) maintaining a physical distance of at least 1.5 meters from others. The "3 DON'Ts" refers to (1) not going to places that are not well-ventilated, (2) not going to crowded places, and (3) not touching others, shaking hands, or hugging.	▪ All phases, to 30 June 2022
3	Risk communication and community engagement	It refers to the regular sharing of official information through various MoH's communication channels to timely raise awareness and dispel circulating myths and rumours, especially on social media.	▪ All phases, to 30 June 2022
4	School closure	School closure refers to the halt of in-person	▪ Phase 1

		teaching in schools. The nationwide school closure commenced on 16 March when Cambodia had detected 24 cases. On 6 September 2020, the government re-opened the schools countrywide after a low possibility of a community outbreak. On 1 December 2020, the government closed all schools countrywide again after a large cluster was detected. School closures were continued until the country's full re-opening on 1 November 2021.	o Closure 1: 16 March to 6 September 2020 o Closure 2: 1 December 2020 to 1 November 2021
5	Border restriction/closure	On 16 March 2020, Cambodia closed all border gates with Vietnam. On 16 April 2020, it implemented travel restrictions for passengers from France, Germany, Iran, Italy, Spain, and the United States of America. On 20 May 2020, the government required all passengers to have a COVID-19 negative certificate within 72 hours before flying (effective until 31 October 2021); it also required health insurance with a minimum amount of USD 50,000, and another test on arrival and quarantine for 14 days at a government-designated place (effective until 1 November 2021).	▪ Phase 1 o Closure 1: 16 March to 19 May 2020, unless otherwise indicated
6	Public event and gathering cancellation	This measure was used to cancel all public events (e.g., Khmer New Year) and gatherings (e.g., wedding celebrations, usual funerals, memorial services, and all festivals).	▪ All phases, to 30 June 2022
7	Vaccination	The goal of COVID-19 vaccination is to reduce transmission, severe cases, hospitalisations and deaths from COVID-19. On 10 February 2021, 10 days before Cambodia declared the large community outbreak, Cambodia started vaccine rollout with the Sinopharm vaccine. The vaccination campaign accelerated in April 2021 as cases increased exponentially. By mid-September 2021, 93.1% of the 15.55 million population had received at least two doses/completed immunisation.	▪ Late Phase 1, Phase 2, and Phase 3
8	Lockdown	Lockdown is a large-scale local travel restriction. Cambodia started a countrywide lockdown between 15 April and 15 May 2021. By "lockdown", we refer to restrictions on movement *between* villages or communes. Individuals could travel *within* the village or communes; however, they were advised to stay at home.	▪ Phase 2 o Lockdown: 15 April to 15 May 2021

177   We describe the responses in each phase as follows:

Phase 1. 27 January 2020 to 19 February 2021: containment with NPIs

After detecting its first case on 30 January 2020 Cambodia started implementing mandatory

temperature screenings for all travellers entering Cambodia at all points of entry (airports and

international borders). In March and April 2020, Cambodia detected several clusters of imported

cases that were not identified through the temperature screening system. These cases were

detected through notifications from the WHO's IHR channel and event-based surveillance

(screening local and international news).

　　To respond to these clusters, intensive contact tracing was undertaken to trace, test and

quarantine the contacts. This strategy was an essential component of "*Measure 1: detect,*

isolate/quarantine". Contact tracing commenced within 24 hours after a confirmed case was

detected. All contacts were quarantined at quarantine centres or homes for 14 days from the last

date of exposure. In 2020, Cambodia missed several opportunities to detect imported COVID-19

cases at the international border gates. From these primary cases, the virus spread locally.

Contact tracing was used to find cases for isolation, and contacts for quarantine, thereby

containing the spread.

　　Alongside measure 1, Cambodia also implemented *Measure 2: face covering, hand*

hygiene, and physical distancing since it detected its first case. The government strongly advised

people to take measures for self-protection such as wearing face masks, hand-washing, and

physical distancing, which were complied to by the population.

　　The government, through the Ministry of Health (MoH) also implemented *Measure 3: risk*

communication and community engagement since it detected its first case. Regular sharing of

official information through various MoH communication channels helped to dispel circulating

myths, false information and rumours, especially on social media.[10] The MoH reiterated

simple messages to the public, known as *"3 DOs and 3 DON'Ts."* The "3 DOs" refer to (1)

wearing a face mask properly, (2) frequent hand disinfection using alcohol, gel-based

formulations, or soap, and (3) maintaining a physical distance of at least 1.5 meters from others.

The "3 DON'Ts" refer to (1) not going to places that are not well-ventilated, (2) not going to

crowded places, and (3) not touching others, shaking hands, or hugging. These nationwide

communication campaigns stimulated cooperation from the public. They were launched by the

MoH through the "Sub-Committee on Education, Training, and Public Communication of the

Inter-ministerial Committee to Fight COVID-19" with the theme *"Acting together to stop*

COVID-19 transmission".

On 16 March 2020, when the country's total number of detected cases reached 24,

Measure 4: school closures were effected nationwide. In as much as 23 of the cases (95.8%)

were imported, Cambodia also implemented *Measure 5: border closures.* Cambodia closed all

border gates with Vietnam before full closures with other countries. On 26 March 2020, when

Cambodia reached 90 cases, Measure *6: cancellation of all public events* (e.g., Khmer New

Year) *and gatherings* (e.g., wedding celebrations, usual funeral memorial services, all festivals)

were imposed. Travel restrictions for passengers from France, Germany, Iran, Italy, Spain, and

the United States of America were implemented on 16 April 2020.

By 20 May 2020, following the increasing trends of cases, travel restrictions were

modified and the government required all passengers to have a COVID-19 negative certificate

within 72 hours before flying, health insurance with a minimum amount of USD 50,000, another

test on arrival, and quarantine for 14 days at a government-designated place.

After a few months with low cases, the government considered Cambodia had a low-risk

of transmission outbreak and decided to re-open all schools nationwide on 6 September 2020.

On 3 November 2020, the Hungarian Foreign Minister had a two-day visit to Cambodia

and met several hundred government officials including the Prime Minister, students, and

teachers, on different occasions. On 6 November 2020, he tested positive for SARS-CoV-2 when

he arrived in Bangkok, Thailand. It was locally known as the "3 November Event". Everyone

who attended the gatherings and their contacts (over 2,000 people) were tested and quarantined

at home for 14 days since the last date of exposure. Four cases were detected as part of this

cluster. It was unclear who the source of infection was, as the Hungarian Foreign Minister tested

negative within 72 hours before flying to Cambodia. The "3 November Event" prompted the

government to tighten all measures due to its high potential for community transmission. On 23

November 2020, the government announced the closure of the "3 November Event" after no

more cases were detected.

On 28 November 2020, Cambodia announced its first community transmission after a

woman was hospitalised with respiratory symptoms and later tested positive for SARS-CoV-2,

locally known as the "28 November community outbreak". Six of her family members also

tested positive for SARS-CoV-2. Her family members who tested positive had no history of

travel abroad. However, she and her family had trips to several provinces in Cambodia and

shopping centres for "Black Friday". As a response, massive testing and quarantine were

implemented, and where those affected were noted to have cooperated. All individuals who were

at sites where one or more of the confirmed cases had visited were tested. Certain commercial

businesses (e.g., AEON supermarket) were closed for 14 days. On 1 December 2020, the

government re-instated nationwide school closures for in-person classes and ordered online

classes to be administered instead. It remained in effect until the country's full re-opening.

Within the span of one month, over 20,000 people were tested and home-quarantined as part of

this cluster. Forty-one cases (no deaths) were detected from this event. On 29 December 2020,

the government announced the closure of the "28 November community outbreak".

Phase 2. 20 February 2021 to 31 October 2021: mitigation with boosting vaccination

coverage

Ten days before this phase, on 10 February 2020, Cambodia started *Measure 7: rolling out*

COVID-19 vaccination with the Sinopharm vaccine after China donated one million doses of the

said vaccine.

On 19 February 2021, two foreign women were known to have escaped the quarantine

hotels to visit a night club while waiting for testing results before they could leave the quarantine

site. Their results were positive for SARS-CoV-2. It triggered the government to start a large-

scale testing of the night club's visitors. Over 20 cases were identified. On the same date, the

laboratory of the National Institute of Public Health also detected a few cases among those who

had requested testing to secure a travel certificate.

On 20 February 2021, Cambodia announced a second community outbreak event, locally

known as the "20 February community outbreak".

To respond to this new event, two new measures—*Measure 7*: *vaccination and Measure*

8: lockdown—were added to the six measures implemented in Phase 1.

Measure 1: trace, isolate/quarantine was intensified. All individuals in the buildings or

sites where one or more confirmed cases had visited were tested. Around 8,000 tests were

performed per day which was the full capacity of Cambodia's laboratories at that time. However,

the cases increased exponentially (Figure 1). The other five measures were also intensified.

Measure 8: lockdown was implemented covering the period between 15 April and 15

May 2021. Despite these more stringent measures, cases continued to rise (Figure 1).

During the same period, from April 2021, the vaccination campaign was *sped up* and ran

without interruption. On average, 50,000 doses were provided daily and quickly achieved

vaccination coverage in more than 93.1% of the total population within five months (n =

14,485,634 out of 15,552,211) by 15 September 2021.

After achieving high vaccination coverage, the government stopped active surveillance, but

monitored two indicators—severe cases and deaths—to ensure that the COVID-19 situation

remains under control.

Phase 3. 1 November 2021 to 30 June 2022: full re-opening

Without any unusual increase in severe cases and deaths for 30 days in a row, the government

decided to lift all compulsory containment measures from 1 November 2021.[11] Soon after, all

non-pharmaceutical measures were lifted. However, a significant proportion of people in urban

cities still choose to wear masks until the end of analysis period.

Strategy

We retrospectively identified six themes from the COVID-19 responses in Cambodia.

Theme 1: Setting up and managing a new response system

Cambodia has learned quickly and adapted to the dynamics of the disease. When the number of

COVID-19 cases started to increase from March 2020, the Prime Minister led the national

response. He created and chaired the National COVID-19 Response Committee. This committee

was composed of high-level committees to give advice on the various aspects of the country's

COVID-19 response. The high-level committees included 25 Provincial COVID-19 Committees,

each chaired by the provincial governors to monitor the responses in each province, the High-

level Ministry of Health Task Force to mobilise resources and the Technical Working Group led

by Cambodia's CCDC with technical support from governmental and non-governmental

stakeholders to provide technical advice on the COVID-19 emergency response.

Theme 2: Containing the spread with early response

As described previously, six measures were implemented to contain the spread of the virus.

These measures had a background-strategy, the so-called "5 Es"—E1 for Early Detection, E2 for

Early Isolation, E3 for Early Tracing, E4 for Early Treatment, and E5 for Early Education.

Theme 3: Strengthening the identification of cases and contacts

Cambodia trained new rapid response teams (RRTs), increasing from around 2,000 to over 5,000

by the end of phase 1. The capacities of existing laboratories were increased and 18 new

laboratories in Phnom Penh and 11 provinces were set up. Testing capacity increased from 500

PCR tests per day in April 2020, to around 15,000 PCR tests per day in July 2021.

Theme 4: Strengthening care for COVID-19 patients

To reduce the case-fatality rate, Cambodia had to strengthen the capacity of its health care

system.[12] Cambodia addressed this issue by increasing the number of doctors and nurses who

had the capability to manage the disease.

During phase 1, Cambodia trained public healthcare providers from over 1,300 public

healthcare facilities, including health centres, to treat and care for COVID-19 patients. In

addition, it trained doctors- and nurses-in-training to treat and care for COVID-19 patients.

School buildings, wedding buildings, and abandoned hotels were temporarily repurposed

as isolation and treatment centres.

Theme 5: Boosting vaccination coverage

While the cases seemed unstoppable, in April 2021, Cambodia tried to boost the vaccination rate

to build herd immunity.

Cambodia supplied basic doses with the Sinovac, Sinopharm, Moderna, or Janssen

vaccines. The AstraZeneca and Pfizer vaccines were provided as adult booster doses. Children

received the Sinovac and/or Pfizer (paediatric) vaccines and no booster doses were given.

A total of 53.47 million doses of the COVID-19 vaccine were obtained (Table 3). Over

half (53%) were purchased using the state budget and the remainder were provided through

foreign aid. The majority of the vaccines was the Sinovac vaccine (64.4%), followed by the

Sinopharm vaccine (14.6%), the Pfizer vaccine (9.6%), the AstraZeneca vaccine (9.1%), and the

remainder were the Janssen and Moderna vaccines (2.4%).

Table 3. Vaccine doses obtained by the Cambodian government in 2021

Resource	Sinovac	Sinopharm	AstraZeneca	Pfizer	Janssen	Pfizer (Paediatric)	Moderna	Total	%
National budget	24,500,000	4,000,000						28,500,000	53.3%
China's aid	9,500,000	3,800,000						13,300,000	24.9%
COVAX	424,800		2,253,900	1,999,530	1,060,100		188,160	5,926,490	11.1%
Australia's aid				2,350,530		186,800		2,537,330	4.7%
Italy's aid			1,056,000					1,056,000	2.0%
Hungary's aid			523,100					523,100	1.0%
Poland's aid			300,000			143,990		443,990	0.8%
UK's aid			415,040					415,040	0.8%
India's aid			325,000					325,000	0.6%
Spain's aid						300,000		300,000	0.6%
ASEAN				139,230				139,230	0.3%
Total	34,424,800	7,800,000	4,873,040	4,489,290	1,060,100	630,790	188,160	53,466,180	100%
%	64.4%	14.6%	9.1%	8.4%	2.0%	1.2%	0.4%	100%	

Theme 6: Supporting the disadvantaged groups

Lockdown negatively impacts those in disadvantaged groups, particularly, daily-wage earners

(e.g., garment workers, Tuk-tuk (motorcycle), taxi drivers, construction workers, and street

vendors).[13] The government and local and international development partners strived to

respond in a timely manner to help the disadvantaged groups.[13] Initially, delivering food

supplies in Cambodia had been too slow and challenging due to the absence of a food

distribution infrastructure specifically for the poor.[3, 13, 23]. Nevertheless, between May 2020

and May 2021, the government provided social assistance through the COVID-19 Cash Transfer

Programme to 700,000 low-income households, equivalent to more than 2.8 million individuals.

By June 2021, each household had received an average of USD 366 in nine instalments.[14]

DISCUSSION

This analysis describes how the COVID-19 epidemiology evolved in Cambodia and how the

country responded between 27 January 2020 (when the first case was detected) and 30 June 2022

(several months after the country announced full re-opening). It also highlights the strategies

behind the responses.

The COVID-19 epidemiology in Cambodia is split into three phases. The first phase from

27 January 2020 to 19 February 2021 when the COVID-19 outbreak was well contained without

a single death. According to WHO data, by 7 February 2021, Cambodia had the 7[th] lowest

number of cases per 100,000 population, among 214 countries and territories.[15] However, we

acknowledge that it is difficult to compare number of cases with other countries because they are

severely impacted by different surveillance efforts and testing capacities. The second phase from

20 February to 31 October 2021 was the large community outbreak stage. During this phase,

Cambodia recorded 2,805 deaths out of 15,552,211 population, giving a mortality rate of 180 per

million. This mortality rate ranked 74[th] out of 214 countries and territories reported to the

WHO.[16] In the third phase, after the country fully re-opened on 1 November 2021, 47,314

cases and 251 deaths were recorded, but the cases were also largely under-reported due to the

relaxation of all interventions, including testing. Nevertheless, we note that the country still

monitored severe cases and deaths from COVID-19, but we could not access the number of

severe cases.

Cambodia responded with NPIs during the first phase, and added lockdown and vaccines

in the second phase. The six measures in the first phase included *(1) detect, isolate/quarantine;*

(2) face coverings, hand hygiene, and physical distancing measures; (3) risk communication and

community engagement; (4) school closures; (5) border closures; and (6) public event and

gathering cancellation. These measures worked well in the Cambodian context with high

cooperation from the public. It also may have benefited from early action when the community

transmission has not yet existed. In the second phase, the implementation of the lockdown had

not achieved the expected results. The number of cases continued to increase. This may be due to

authorities locking down the areas instead of houses, allowing people to move within the area. In

this second phase, high coverage of the COVID-19 vaccine might have been the only measure

that helped reduce infections and deaths and help Cambodia re-open the country.

Cambodia responded with several strategies: *(1) setting up and managing a new response*

system, (2) containing the spread with early response, (3) strengthening the identification of

cases and contacts, (4) strengthening care for COVID-19 patients, (5) boosting vaccination

coverage, and (6) supporting the disadvantaged groups. The core of these strategies relied on

strong governance and decisive leadership (fast, at the right time, without delay) to respond early

to prevent infections and deaths, and to support those who required livelihood support.

Lessons learned

From Cambodia's COVID-19 response, we can learn several lessons. First, the COVID-19

response requires decisive leadership. Second, the response must be early and include contact

tracing, risk communication, community engagement, and expansion of the laboratory capacity.

Third, the lockdown did not produce the expected results. Fourth, rapid nationwide vaccination

achieving a high coverage rate was crucial and provided Cambodia a ticket to re-open the

country. Fifth, cooperation from public was a relevant and necessary factor.

1. COVID-19 response requires decisive leadership

The COVID-19 pandemic is a "wicked problem" [17] and many decisions were made amidst

uncertainties. Leadership is a core part of governance, referring to the government's ability to

provide guidance, plan, and make appropriate modifications to the laws or regulations at all

levels and sectors to enable an effective response.

We learned that responding to the pandemic requires decisive leadership, referring to

decisions being taken fast, timely, and without delay. With this characteristic, Cambodia

responded promptly, despite low pre-pandemic capacity. Literature suggests that decisive

leadership led to success in responding to COVID-19.[18-22] Similar leadership characteristics

and outcome at the early phase were seen in China, New Zealand, Taiwan, and South Korea.[18-

22]

2. The response must be early and include contact tracing, risk communication, and

strengthening laboratory capacity, which were the core element of the early

response

Cambodia could contain the spread of the virus for more than a year with early

responses/actions. Contact tracing was not only to find the cases, isolate them, and find their

contacts, but it also provided real-time information, which was necessary for the response teams

to decide where to impose additional restrictions. Effective risk communication and community

engagement was critical to educate people to receive reliable information and gain trust, and

strengthened cooperation from the public. However, the laboratory capacity determines the speed

of the response. The response could be slower or faster based on the number of tests that can be

performed within a time period.

3. Lockdown must accompany support to disadvantaged groups

Lockdown is a controversial measure for many settings as it affects the livelihoods of many

individuals. Supporting the needs of low-income families was essential to reducing the burden of

COVID-19.

4. Vaccination gave Cambodia a ticket to re-open the country

As a low-income country, Cambodia was unique in achieving high vaccination coverage quickly.

Upon achieving a 93.5% vaccination coverage in mid-September 2021, a marked decline in the

number of deaths was noted by October and a massive decrease in the number of cases

approximately one month later. Cambodia was then able to fully re-open the country.

5. Cooperation from the public was the necessary factor

From the measures described above, adherence and compliance from the general public is

required.[2] The government called it solidarity. This compliance is a fundamental factor for

success in containing the spread in the early phase, and the very successful vaccination campaign

in the latter stage.

Strengths

This review is among the few that presents the COVID-19 epidemiology and response, and

highlights lessons learned in Cambodia for organising and investing in future responses.

Limitations

Our review has a few limitations. First, the relationship between epidemiology and response may

or may not be causal. Second, the COVID-19 data may be highly under-reported and may not

reflect the actual trends due to limited testing capacity and changing surveillance strategies.

However, we are confident that what may have been missed were cases not requiring

hospitalization and/or medical attention, or the asymptomatic and mild COVID-19 infections.

CONCLUSION

Despite low levels of preparedness for the pandemic, Cambodia successfully contained the

spread of SARS-CoV-2 in the first year and quickly attained high vaccine coverage by the

second year of the response. The core of this success was the strong political will and high

cooperation from the public. However, Cambodia needs to improve its preparedness, especially

on medical infrastructure to isolate cases, control infectious disease outbreaks, and further

develop its human resources and laboratory capacity.

ACKNOWLEDGMENTS

We would like to thank all the Inter-ministerial COVID-19 Response Committee and

Cambodia's Communicable Disease Control Department (CCDC) for providing data.

AUTHOR CONTRIBUTIONS

SC, GK, SM, WP, WD, PI, CC, and VO, designed the study; SC analyzed data; SC, GK, SM,

WP, WD, PI, CC, and VO wrote the manuscript.

FUNDING

This work received no external funding.

CONFLICT OF INTERESTS

All authors have declared no competing interest.

ETHICS APPROVAL

Not applicable.

DATA AVAILABILITY STATEMENT

Data are available on reasonable request. Data are available on reasonable request to Dr. Vandine

Or (vandine@online.com.kh).

REFERENCES

1. Van Damme W, Dahake R, Delamou A, *et al.* The COVID-19 pandemic: diverse contexts; different epidemics—how and why? *BMJ Glob Health* 2020;5(7):e003098. doi: 10.1136/bmjgh-2020-003098

2. Nit B, Samy AL, Tan SL, *et al.* Understanding the Slow COVID-19 Trajectory of Cambodia. *Public Health Pract (Oxf)* 2021;2:100073. doi: 10.1016/j.puhip.2020.100073 [published Online First: 2021/02/02]

3. Tatum M. Cambodia ends controversial COVID-19 restrictions. *Lancet* 2021;397(10289):2035. doi: 10.1016/S0140-6736(21)01196-X

4. National Institute of Statistic (NIS). General Population Census of the Kingdom of Cambodia 2019, Phnom Penh, Cambodia: NIS, 2019. Available: https://nis.gov.kh/nis/Census2019/Final%20General%20Population%20Census%202019-English.pdf [Accessed 16 December 2022].

5. The World Bank. Physicians (per 1,000 per people) - Cambodia, Washington DC: The World Bank, 2014. Available: https://data.worldbank.org/indicator/SH.MED.PHYS.ZS?locations=KH [Accessed 16 December 2022].

6. World Health Organization. The Global Health Observatory, Geneva, Switzerland: WHO, 2020. Available: https://www.who.int/data/gho/data/indicators/indicator-details/GHO/hospital-beds-(per-10-000-population) [Accessed 16 December 2022].

7. The World Bank. Beds (per 1,000 per people) - Cambodia, Washington DC: The World Bank, 2016. Available: https://data.worldbank.org/indicator/SH.MED.BEDS.ZS?locations=KH [Accessed 16 December 2022].

8. World Health Organization. Joint External Evaluation of IHR Core Capacities, Geneva, Switzerland: WHO, 2017. Available: https://apps.who.int/iris/handle/10665/254705 [Accessed 16 December 2022].

9. Chhim S. Applied Epidemiology in Cambodia. The Australian National University (Australia), 2021.

10. World Health Organization. 2019 Novel Coronavirus (2019-nCoV): Strategic Preparedness and Response Plan, Geneva, Switzerland: WHO, 2020. Available: https://www.who.int/publications/i/item/strategic-preparedness-and-response-plan-for-the-new-coronavirus [Accessed 16 December 2022].

11. Kunthear M. PM declares Cambodia fully reopen from Nov 1, Phnom Penh: The Phnom Penh Post, 2021. Available: https://www.phnompenhpost.com/national/pm-declares-cambodia-fully-reopen-nov-1 [Accessed 16 December 2022].

12. Sheikhbardsiri H, Raeisi AR, Nekoei-Moghadam M, *et al.* Surge Capacity of Hospitals in Emergencies and Disasters With a Preparedness Approach: A Systematic Review. *Disaster Med Public Health Prep* 2017;11(5):612-20. doi: 10.1017/dmp.2016.178 [published Online First: 2017/03/08]

13. Macdonald S. Learning from Cambodia's COVID-19 Response: Foreign Policy in Focus (FPIF), 2021. Available: https://fpif.org/learning-from-cambodias-covid-19-response/ [Accessed 19 Oct 2022 2022].

14. Wendy Karamba, Isabelle Salcher, Tong K. Results from the High-Frequency Phone Survey of Households Round 5 (1–21 March 2021), Phnom Penh, Cambodia: World Bank Group, 2021. Available:

https://documents1.worldbank.org/curated/en/833701624563697124/pdf/Main-
Report.pdf [Accessed 16 December 2022].

15. World Health Organization. Weekly epidemiological update - 9 February 2021, Geneva,
Switzerland: WHO, 2021. Available: https://www.who.int/publications/m/item/weekly-
epidemiological-update---9-february-2021 [Accessed 16 December 2022].

16. Our World in Data. Coronavirus (COVID-19) Cases, 2022. Available:
https://ourworldindata.org/covid-cases [Accessed 16 December 2022].

17. Schiefloe PM. The Corona crisis: a wicked problem. *Scand J Public Health* 2020;49(1):5-8.
doi: 10.1177/1403494820970767

18. Nicola M, Sohrabi C, Mathew G*, et al.* Health policy and leadership models during the
COVID-19 pandemic: A review. *Int J Surg* 2020;81:122-29. doi:
10.1016/j.ijsu.2020.07.026

19. Al Saidi AMO, Nur FA, Al-Mandhari AS*, et al.* Decisive leadership is a necessity in the
COVID-19 response. *Lancet* 2020;396(10247):295-98. doi: 10.1016/S0140-
6736(20)31493-8

20. Stoller JK. Reflections on leadership in the time of COVID-19. *BMJ Leader* 2020;4:77-79.
doi: 10.1136/leader-2020-000244

21. Kaul V, Shah VH, El-Serag H. Leadership During Crisis: Lessons and Applications from the
COVID-19 Pandemic. *Gastroenterology* 2020;159(3):809-12. doi:
10.1053/j.gastro.2020.04.076

22. Phillips G, Kendino M, Brolan CE*, et al.* Lessons from the frontline: Leadership and
governance experiences in the COVID-19 pandemic response across the Pacific region.
Lancet Reg Health West Pac 2022;25 doi: 10.1016/j.lanwpc.2022.100518

23. Johnson S, Srey V. Thousands of Cambodians go hungry in strict lockdown zones, Phnom
Penh, Cambodia: The Guardian, 2021. Available: https://www.theguardian.com/global-
development/2021/may/13/thousands-of-cambodians-go-hungry-in-strict-lockdown-
zones [Accessed 16 December 2022].