1   **COVID-19 in Cambodia, January 2020–June 2022: a success story**

2
3

Authors: Srean Chhim[1], Grace Marie Ku[2], Sovathiro Mao[1], Willem van de Put[2], Wim Van

Damme[2, 3], Por Ir[1], Chhea Chhorvann[1], Vandine Or[4]

Affiliations:

[1] National Institute of Public Health, Phnom Penh, Cambodia

[2] Institute of Tropical Medicine, Antwerp, Belgium

[3] Vrije Universiteit Brussel, Belgium

[4] Ministry of Health, Phnom Penh, Cambodia

12
13
14   ** Corresponding author:* Srean Chhim (Lot # 80, Samdach Penn Nouth Blvd. (St. 289), Phnom
15   Penh, postal 120804), srean168@gmail.com

16
17   *Keywords***:** COVID-19, SARS-CoV-2, Cambodia
18
19

ABSTRACT

As a member state of the International Health Regulation 2005, Cambodia has been continuously strengthening its capacity to respond to health emergencies and prevent the international spread of diseases. Despite this, Cambodia's capacity to prevent, detect, and rapidly respond to public health threats remained limited at the onset of the pandemic. This paper describes epidemiological phases, response phases, strategy, and lessons learned in Cambodia between 27 January 2020 and 30 June 2022. We classified epidemiological phases in Cambodia into three phases, in which Cambodia responded using eight measures: *(1) detect, isolate/quarantine; (2) face coverings, hand hygiene, and physical distancing measures; (3) risk communication and community engagement; (4) school closures; (5) border closures; (6) public event and gathering cancellation; (7) vaccination;* and *(8) lockdown.* The measures corresponded to six strategies: *(1) setting up and managing a new response system, (2) containing the spread with early response, (3) strengthening the identification of cases and contacts, (4) strengthening care for COVID-19 patients, (5) boosting vaccination coverage,* and *(6) supporting disadvantaged group*s. Finally, ten lessons were learned for future health emergency responses. Findings suggest that Cambodia successfully contained the spread of SARS-CoV-2 in the first year and quickly attained high vaccine coverage by the second year of the response. The core of this success was the strong political will and high level of cooperation from the public. However, Cambodia needs to further improve its infrastructure for quarantining and isolating cases and close contacts and laboratory capacity for future health emergencies.

Keywords: COVID-19, SARS-CoV-2, Response, Cambodia, Lessons learned

SUMMARY BOX

- COVID-19 spread globally, but how the pandemic played out in each country depended on various factors, including government responses and the general public's adherence to COVID-19 measures.

- Early response—*Early detection, Early isolation, Early tracing, Early treatment, and Early education*—is the core of successful SARS-CoV-2 containment.

- Achieving high vaccination coverage quickly leads to a decline in the number of deaths and to eventual full re-opening of the country.

- Responding to the pandemic requires decisive leadership and good governance, that refers to decisions being made quickly, in a timely manner, and without delay.

- High level of cooperation from the public is a fundamental factor for success in containing the spread in the early phase, and the massively successful vaccination campaign in the later stage.

INTRODUCTION

Starting in early 2020, COVID-19 spread globally. However, how the epidemic played out in each country depended on various factors, including the government responses and the general public's adherence to COVID-19 measures.[1]

Cambodia's health system has significantly improved in the last three decades contributing to a rise in life expectancy.[2] However, its readiness in responding to a health emergency was still limited at the onset of the pandemic. In 2014, the World Health Organization (WHO) estimated that Cambodia had only 0.2 physicians per 1,000 population.[3, 4] The ratio of hospital beds was also low at 0.9 per 1,000 population.[4, 5] There was limited infrastructure for isolation of patients with highly infectious diseases.

As a member state of the International Health Regulation (IHR) 2005, Cambodia committed to implementing the IHR's framework and continuously strengthening its capacity to respond to health emergencies, including outbreaks. In spite of this, the Joint External Evaluation by the IHR—a process of evaluating a country's capacity to prevent, detect, and rapidly respond to public health threats—indicated that Cambodia had limited capacity in 2016.[6]

The COVID-19 response in Cambodia has not been well-documented in scientific publications. While several papers described the early phases of COVID-19 responses including non-pharmaceutical interventions (NPIs), many other important aspects, such as rapid scale up of laboratory capacity and vaccination, were overlooked.[7, 8]

Our paper describes the epidemiological phases, response phases, strategy, and lessons learned in Cambodia between 27 January 2020 (when the first case was detected) and 30 June 2022, several months after Cambodia's full re-opening.

METHODOLOGY

We analysed secondary data and reviewed relevant literature and documents to describe the COVID-19 epidemiology, responses, and strategies between 27 January 2020 and 30 June 2022 in Cambodia.

Data for descriptive epidemiology

Tests, cases, and deaths

The Inter-ministerial COVID-19 Response Committee provided the data of COVID-19 tests, and we used publicly available data from WHO's COVID-19 dashboard for cases and deaths.[9] The tests, cases, and deaths recorded between 27 January 2020 and 30 June 2022. The COVID-19 data is under the governance and stewardship of the Inter-ministerial COVID-19 Response Committee, with Cambodia's Communicable Disease Control Department (CCDC) being the central body responsible for collection and storage. The data were aggregated and reported to WHO as part of the IHR 2005's commitment. The data were collected from all laboratories in Cambodia, using a laboratory form (see supplementary file 1). The completed forms were entered into each laboratory database and exported to a CSV or Excel file and submitted to the CCDC. At the CCDC, these files were checked for standardisation, and imported into the National COVID-19 Database.

Data for response and themes

Relevant events

We documented relevant events to draw a more comprehensive picture of the COVID-19 response in Cambodia. Our data team recorded all the policies and guidelines issued by the government, and triangulated these documents with other documents (e.g., WHO website

updates, situation reports). The information we collected included dates of border closures,

school closures, cancellation/prohibition of public events and gatherings, enforcement of

facemask usage in public areas, and lockdowns.

Vaccination data

The National COVID-19 Vaccination Committee provided vaccination data. These data included

the number of vaccine doses that can be stratified by vaccine brand. It also included vaccination

coverage data: the number of vaccinated individuals, and the number of doses administered (e.g.,

one dose/incomplete immunisation, two doses/completed immunisation, first booster doses,

second booster doses, and third booster doses).

RESULTS AND DISCUSSION

EPIDEMIOLOGICAL PHASES

Figure 1 presents the daily number of COVID-19 cases, and PCR tests in Cambodia. In

Cambodia, we identified three significant phases of the pandemic: containment phase, mitigation

phase, and full re-opening phase.

Phase 1 (27 January to 19 February 2021): containment phase

In Phase 1, COVID-19 was well-contained. Cambodia had only 484 cases, mainly imported and

no deaths (Figure 1 and Table 1).

Phase 2 (20 February to 31 October 2021): mitigation phase

In Phase 2, a large community outbreak occurred that lasted for 252 days with 118,129 cases

(Figure 1 and Table 1). A total of 2,781 COVID-19 deaths were reported (Figure 2 and Table 1).

Phase 3 (1 November 2021 to 30 June 2022): full re-opening

Phase 3 had 17,669 cases (Figure 1 and Table 1). In this phase, 275 deaths were reported.

Figure 1. COVID-19 daily cases, tests, and deaths in Cambodia, 2020–2022 *[Insert*

Figure 1]

Note: Data is based on the official data from the MOH.

Table 1. Reported COVID-19 cases and deaths by phase in Cambodia, 27 January 2020 to 30
June 2022

Phase of COVID-19 outbreak	Theme	All cases	Death
Phase 1. 27 January 2020 to 19 February 2021:	Containment with NPIs	484	0
27 January–30 April 2020	Initial response • Intensive contact tracing • Quarantine/isolation • Border restriction	122	0
1 May–31 October 2020	Relaxing after long period of low cases	169	0
1 November 2020–19 February 2021	Intensifying response for larger scale outbreak	193	0
Phase 2. 20 February 2021 to 31 October 2021	Large community outbreak and vaccination • Vaccine rolling out since 10 February 2021 • City lockdown • Travel restrictions • School closures • Home treatment/isolation • Government cash transfers to disadvantaged groups	118,129	2,781
Phase 3. 1 November 2021 to 30 June 2022	Full re-opening • Achieved goal: 10 million population vaccinated • Continue with 3 DOs and 3 DON'Ts • Continue to provide booster dose and make it routine • Continue to support the disadvantaged groups	17,669	275

Abbreviations: NPIs, Non-pharmaceutical interventions

Figure 2. COVID-19 death trends in Cambodia, 2021–2022 *[Insert Figure 2]*

Note: Data is based on the official data from the MOH.

RESPONSE PHASES

The government implemented eight significant measures during phases 1 and 2: *(1) detect,*

isolate/quarantine; (2) face coverings, hand hygiene, and physical distancing measures; (3) risk

communication and community engagement; (4) school closures; (5) border closures; (6) public

event and gathering cancellation; (7) vaccination; and (8) lockdown (Table 2). Each of these

measures was implemented quickly when the need for such intervention was seen to be

warranted. The first six measures were implemented in phase 1, and measures 7 and 8 were

added in phase 2.

Table 2: Description of the eight measures used in Cambodia in response to the COVID-19

outbreak

No	Measure	Description	Timeframe
1	Detect, isolate/quarantine	This measure was used to detect cases for isolation and find contacts for quarantine. Contact tracing was the primary measure to trace the contacts, test them, and quarantine them. Contact tracing commenced within 24 hours after a confirmed case was detected. The contacts were classified into high, medium, and low risk. High and medium-risk contacts were tested at least two times during the quarantine. Low-risk contacts were tested if they had any signs or symptoms during the quarantine period.[10]	▪ Phase 1 and early phase 2
2	Face coverings, hand hygiene, and physical distancing	This measure was known as "3 DOs and 3 DON'Ts." The "3 DOs" refer to (1) wearing a face mask properly, (2) washing hands frequently with alcohol, gel, or soap, and (3) maintaining a physical distance of at least 1.5 meters from others. The "3 DON'Ts" refers to (1) not going to places that are not well-ventilated, (2) not going to crowded places, and (3) not touching others, shaking hands, or hugging.	▪ All phases, to 30 June 2022
3	Risk communication and community engagement	It refers to the regular sharing of official information (e.g., intensified press release, press conference, media talks, etc) through various MoH's communication channels to timely raise awareness and dispel circulating myths and rumours, especially on social media.	▪ All phases, to 30 June 2022
4	School closure	School closure refers to the halt of in-person teaching in schools. The nationwide school closure commenced on 16 March when Cambodia had detected 24 cases. On 6 September 2020, the government re-opened the schools countrywide after a low possibility of a community outbreak. On 1 December 2020, the government closed all schools countrywide again	▪ Phase 1 o Closure 1: 16 March to 6 September 2020 o Closure 2: 1 December

		after a large cluster was detected. School closures were continued until the country's full re-opening on 1 November 2021.	2020 to 1 November 2021
5	Border restriction/closure	On 16 March 2020, Cambodia closed all border gates with Vietnam. On 16 April 2020, it implemented travel restrictions for passengers from France, Germany, Iran, Italy, Spain, and the United States of America. On 20 May 2020, the government required all passengers to have a COVID-19 negative certificate within 72 hours before flying (effective until 31 October 2021); it also required health insurance with a minimum amount of USD 50,000, and another test on arrival and quarantine for 14 days at a government-designated place (effective until 1 November 2021).	▪ Phase 1 ○ Closure 1: 16 March to 19 May 2020, unless otherwise indicated
6	Public event and gathering cancellation	This measure was used to cancel all public events (e.g., Khmer New Year) and gatherings (e.g., wedding celebrations, usual funerals, memorial services, and all festivals).	▪ All phases, to 30 June 2022
7	Vaccination	The goal of COVID-19 vaccination is to reduce transmission, severe cases, hospitalisations and deaths from COVID-19. On 10 February 2021, 10 days before Cambodia declared the large community outbreak, Cambodia started vaccine rollout with the Sinopharm vaccine. The vaccination campaign accelerated in April 2021 as cases increased exponentially. By mid-September 2021, 93% of the 15.55 million population had received at least two doses/completed immunisation.	▪ Late Phase 1, Phase 2, and Phase 3
8	Lockdown	Lockdown is a large-scale local travel restriction. Cambodia started a countrywide lockdown between 15 April and 15 May 2021. By "lockdown", we refer to restrictions on movement *between* villages or communes. Individuals could travel *within* the village or communes; however, they were advised to stay at home.	▪ Phase 2 ○ Lockdown: 15 April to 15 May 2021

Phase 1. 27 January 2020 to 19 February 2021: containment with NPIs

After detecting the first case, on 30 January 2020, Cambodia started implementing mandatory

temperature screenings for all travellers entering Cambodia at all points of entry (airports and

international borders). In March and April 2020, Cambodia detected several clusters of imported

cases that were not identified through the temperature screening system, but through

notifications from the WHO's IHR channel and event-based surveillance (screening local and

international news). From these primary cases, the virus spread locally.

To respond to these clusters, intensive contact tracing was undertaken to trace, test and

quarantine the contacts. This strategy was an essential component of "*Measure 1: detect,*

isolate/quarantine". Contact tracing commenced within 24 hours after a case was detected. All

contacts were quarantined at quarantine centres or homes for 14 days from the last date of

exposure.

Cambodia also implemented *Measure 2: face covering, hand hygiene, and physical*

distancing since it detected its first case. The government strongly advised people to take such

measures for self-protection and compliance by the population was very high.

The government, through the Ministry of Health (MoH) also implemented *Measure 3: risk*

communication and community engagement from the onset. Regular sharing of official

information through various MoH communication channels helped to dispel circulating myths,

false information, and rumours, especially on social media.[11] The MoH reiterated simple

messages to the public, known as *"3 DOs and 3 DON'Ts"*. The "3 DOs" refer to (1) wearing a

face mask properly, (2) frequent hand disinfection using alcohol, gel-based formulations, or

soap, and (3) maintaining a physical distance of at least 1.5 meters from others. The "3 DON'Ts"

refer to (1) not going to places that are not well-ventilated, (2) not going to crowded places, and

(3) not touching others, shaking hands, or hugging. These nationwide communication campaigns

stimulated cooperation from the public. They were launched by the MoH through the "Sub-

Committee on Education, Training, and Public Communication of the Inter-ministerial

Committee to Fight COVID-19" with the theme *"Acting together to stop COVID-19*

transmission".

On 16 March 2020, when the country's total number of detected cases reached 24,

Measure 4: school closures were imposed nationwide. As 23 of the cases (95.8%) were

imported, Cambodia also implemented *Measure 5: border closures.* Cambodia first closed all

border gates with Vietnam before full closure with other countries. On 26 March 2020, when

Cambodia reached 90 cases, Measure *6: cancellation of all public events* (e.g., Khmer New

Year) *and gatherings* (e.g., wedding celebrations, usual funeral memorial services, all festivals)

were imposed. Travel restrictions for passengers from France, Germany, Iran, Italy, Spain, and

the United States of America were implemented on 16 April 2020.

By 20 May 2020, following the increasing trend of cases, travel restrictions were

modified and the government required all passengers to have a COVID-19 negative certificate

within 72 hours before flying, health insurance with a minimum cover of USD 50,000, another

test on arrival, and quarantine for 14 days at a government-designated place.

After a few months with low cases, the government considered Cambodia had a low-risk

of transmission outbreak and decided to re-open all schools nationwide on 6 September 2020.

On 3 November 2020, the Hungarian Foreign Minister came on a two-day visit to

Cambodia and met several hundred government officials including the Prime Minister, as well as

students, and teachers, on different occasions. On 6 November 2020, he tested positive for

SARS-CoV-2 on his return in Bangkok, Thailand. This became locally known as the "3

November Event". Everyone who attended these gatherings and their contacts (over 2,000

people) were tested and quarantined at home for 14 days since the last date of exposure. Four

cases were detected as part of this cluster. It was unclear who the source of infection was, as the

Hungarian Foreign Minister tested negative within 72 hours before flying to Cambodia. The "3

November Event" prompted the government to tighten all measures due to its high potential for

community transmission. On 23 November 2020, the government announced the closure of the

"3 November Event" after no more cases were detected.

On 28 November 2020, Cambodia announced its first community outbreak after a woman

hospitalised with respiratory symptoms and later tested positive for SARS-CoV-2, locally known

as the "28 November community outbreak". Six of her family members also tested positive.

They had no history of travel abroad. However, she and her family visited several provinces in

Cambodia and shopping centres for "Black Friday". As a response, massive testing and

quarantine were implemented. All individuals who were at sites where one or more of the

confirmed cases had visited were tested. Certain commercial businesses (e.g., AEON

supermarket) were closed for 14 days. On 1 December 2020, the government re-instated

nationwide school closures for in-person classes and ordered online classes to be administered

instead. This remained in effect until the country's full re-opening. Within the span of one

month, over 20,000 people were tested and home-quarantined as part of this cluster. Forty-one

cases (no deaths) were detected from this event. On 29 December 2020, the government

announced the closure of the "28 November community outbreak".

Phase 2. 20 February 2021 to 31 October 2021: mitigation with vaccination campaign

Ten days before this phase, on 10 February 2021, Cambodia started *Measure 7: rolling out*

COVID-19 vaccination with the Sinopharm vaccine after China donated one million doses of the

said vaccine.

On 19 February 2021, two foreign women escaped the quarantine hotels to visit a night

club while waiting for test results. Their results were positive for SARS-CoV-2. It triggered a

large-scale testing of the night club's visitors. Over 20 cases were identified. On the same date,

the laboratory of the National Institute of Public Health also detected several cases among those

who had requested pre-travel testing.

On 20 February 2021, Cambodia announced a second community outbreak, locally

known as the "20 February community outbreak".

Measure 1: trace, isolate/quarantine was intensified. All individuals in the buildings or

sites where one or more confirmed cases had visited were tested. Around 8,000 tests were

performed per day, which was the full capacity of Cambodia's laboratories at that time.

However, as the cases increased exponentially (Figure 1), the other five measures were also

intensified.

Measure 8: lockdown was implemented covering the period between 15 April and 15

May 2021. Despite these more stringent measures, cases continued to rise (Figure 1).

During the same period, from April 2021, the vaccination campaign was *sped up* and ran

without interruption. On average, 50,000 doses were provided daily and quickly achieved

vaccination coverage of 93% of the total population within five months (n = 14,485,634 out of

15,552,211) by 15 September 2021.

After achieving such high vaccination coverage, the government stopped active surveillance,

but monitored two indicators—severe cases and deaths—to ensure that the COVID-19 situation

remained under control.

Phase 3. 1 November 2021 to 30 June 2022: full re-opening

Without any unusual increase in severe cases and deaths for 30 days in a row, the government

decided to lift all compulsory containment measures from 1 November 2021. However, the

government still advised people to continue with the "3 DOs and 3 DON'Ts", provided booster

doses routinely, and continued to support the disadvantaged groups.

STRATEGY

We identified six themes from the COVID-19 responses in Cambodia.

Theme 1: Setting up and managing a new response system

Cambodia learned quickly and adapted to the dynamics of the disease. When the number of

COVID-19 cases started to increase from March 2020, the Prime Minister led the national

response. He created and chaired the National COVID-19 Response Committee. This committee

was composed of high-level committees to give advice on the various aspects of the country's

COVID-19 response. The high-level committees included 25 Provincial COVID-19 Committees,

each chaired by the provincial governors, a High-level Ministry of Health Task Force to mobilise

resources and the Technical Working Group led by Cambodia's CCDC with technical support

from governmental and non-governmental stakeholders to provide technical advice on the

COVID-19 emergency response.

Theme 2: Containing the spread with early response

As described previously, six measures were implemented to contain the spread of the virus.

These measures had a background-strategy, the so-called "5 Es"— Early Detection, Early

Isolation, Early Tracing, Early Treatment, and Early Education.

Theme 3: Strengthening the identification of cases and contacts

Cambodia trained new rapid response teams (RRTs), increasing from around 2,000 to over 5,000

by the end of phase 1. The capacities of existing laboratories were increased and 18 new

laboratories in Phnom Penh and 11 provinces were set up. Testing capacity increased from 500

PCR tests per day in April 2020, to around 15,000 PCR tests per day in July 2021.

Theme 4: Strengthening care for COVID-19 patients

To reduce the case-fatality rate, Cambodia had to strengthen the capacity of its health care

system.[12] Cambodia addressed this issue by training many doctors and nurses to manage

COVID-19 cases.

During phase 1, Cambodia trained public healthcare providers from over 1,300 public

healthcare facilities, including health centres, to treat and care for COVID-19 patients. In

addition, it trained doctors- and nurses-in-training to treat and care for COVID-19 patients.

School buildings, wedding buildings, and abandoned hotels were temporarily repurposed

as isolation and treatment centres.

Theme 5: Boosting vaccination coverage

In April 2021, when the cases seemed unstoppable, Cambodia boosted the vaccination rate to

build herd immunity.

After basic vaccination with the Sinovac, Sinopharm, Moderna, AstraZeneca, or Janssen

vaccines, adult booster doses were given with AstraZeneca and Pfizer vaccines. Children

received the Sinovac and/or Pfizer (paediatric) vaccines and booster doses.

A total of 54.06 million doses of COVID-19 vaccine were obtained (Table 3). The

majority of the vaccines was Sinovac (63.7%), followed by Sinopharm (15.0%), Pfizer (adults

and paediatric; 10.0%), AstraZeneca (7.8%), Covishield (1.2%), and the remainder were Janssen

and Moderna (2.3%).

290   Table 3. Vaccine doses obtained by the Cambodian government till 05 February 2023

Type of vaccine	Doses received	Percent
Sinovac	34,424,800	63.7
Sinopharm	8,100,000	15.0
Pfizer	4,489,290	8.3
Pfizer Paediatric	923,990	1.7
AstraZeneca	4,224,040	7.8
Janssen	1,060,100	2.0
Covishield	649,000	1.2
Moderna	188,160	0.3
Total	54,059,380	100.0

Abbreviation: ASEAN, Association of Southeast Asian Nations

Theme 6: Supporting disadvantaged groups

Lockdown negatively impacts those in disadvantaged groups, particularly, daily-wage earners

(e.g., garment workers, Tuk-tuk (motorcycle) and taxi drivers, construction workers, and street

vendors).[13] The government and local and international development partners strived to

respond in a timely manner to help the disadvantaged groups.[13] Initially, delivering food

supplies was too slow and challenging due to the absence of a food distribution infrastructure

specifically for the poor.[8, 13, 14] Nevertheless, between May 2020 and May 2021, the

government provided social assistance through the COVID-19 Cash Transfer Programme to

700,000 low-income households, equivalent to more than 2.8 million individuals. By June 2021,

each household had received an average of USD 366 in nine instalments.[15]

LESSONS LEARNT

From Cambodia's COVID-19 response, we highlight ten lessons learnt.

1. COVID-19 response requires decisive leadership and good governance

The COVID-19 pandemic is a "wicked problem" [16] and many decisions were made

amidst uncertainties. Leadership is a core part of governance, referring to the government's

ability to provide guidance, plan, and make appropriate modifications to the laws or regulations

at all levels and sectors to enable an effective response. We learned that responding to the

pandemic requires decisive leadership, referring to decisions being taken fast, timely, and

without delay. The decisive leadership must be supported by the good governance to effectively

implement the newly introduced policies or guideline. With this characteristic, Cambodia

responded promptly, despite limited pre-pandemic capacity. Literature suggests that decisive

leadership led to success in responding to COVID-19 at the early phase in China, New Zealand,

Taiwan, and South Korea.[17-21]

2. Laboratory capacity determine the response speed

The country should be able to conduct tests quickly without having to ship samples overseas.

Laboratory capacity limit the speed of the response, meaning delay in releasing of results also

delays response, impacting community transmission and deaths due to COVID-19. A rapid scale-

up of laboratory capacity was essential. An example of high mortality and excess deaths due to

delays in providing COVID-19 results can be seen in Ecuador.[22] Taiwan and Singapore

learned from the SARS response in 2000s and understood well the need to extend laboratory

capacity and took early actions that may have significantly contributed to its success in

containing the virus spread in the early stage and overall mortality rate over time.[23] In

Cambodia, the Rapid Respond Team (RRT) demands an immediate laboratory result (within 24

hours), so that they can decide in the field. Cambodia managed to scale up its laboratory rapidly

to response to the need of response team and dynamic of the disease.

3. Risk communication and community engagement could fight the infodemics, but must be

consistent

Effective risk communication and community engagement have been critical to educating people

to receive reliable information, gain trust and strengthen engagement with the public.[24]

Fighting the infodemic must be intense. It is important to agree that only one person or group

shares the COVID-19 information to make it consistent and avoid public confusion.

4. Contact tracing worked well in a small-scale response and stopped working when the

community outbreak started

Contact tracing was not only to find the cases, isolate them, and find their contacts, but it also

provided real-time information, which was necessary for the response teams to decide where to

impose additional restrictions. However, it only works in settings where cooperation from the

public is high. Moreover, it only works before community outbreak occurs. Contact tracing

should be stopped when the evidence of community outbreak is confirmed to save resources for

a mitigation plan.

5. School closures caused class disruption, but it saved elderly lives

Schools are a perfect environment for transmission, and a risk for elderlies (aged 50 and over)

who usually live with their children and grandchildren. However, school closures were highly

controversial due to their effects on children's learning. Cambodia implemented early school

closures and empty schools were used as quarantine centers. This measure may have been

necessary to help Cambodia successfully contain the virus for more than a year and prevent

deaths from COVID-19.

6. Face coverings, hand hygiene, physical distancing, border restriction/closures, and

public event and gathering cancellation were not controversial in Cambodia

People in Cambodia tended to easily accept face coverings, hand hygiene, and physical

distancing, border restriction/closure, and public event and gathering cancellation. The

cooperation was high.

7. Lockdown must accompany support to disadvantaged groups

Lockdown is a controversial measure for many settings as it affects the livelihoods of many

individuals.[25] Supporting the needs of low-income families was essential to reducing the social

burden of COVID-19. The slow support was due to the limited existing registration system of

people who need support. However, the government managed to provide food and cash support

to around 700,000 households or 2.8 million individuals.[15] From this experience, it is

important to register and regularly update those who need support or so-called IDPoor in

Cambodia.[26]

8. Vaccination gave Cambodia a ticket to re-open the country

As a low-income country, Cambodia was unique in achieving high vaccination coverage quickly.

Upon achieving a 93% vaccination coverage in mid-September 2021, a marked decline in the

number of deaths was noted by October 2021, approximately one month later. Cambodia was

then able to fully re-open the country.

9. Cooperation from the public was the necessary factor

From the measures described above, adherence and compliance from the general public is required.[7] The government called it solidarity. This compliance is a fundamental factor for success in containing the spread in the early phase, and the very successful vaccination campaign in the later stage.

10. The background-strategy of 5Es was central for success

The background strategy of the "5 Es" was behind the intensified implementation of each measure. In responding to the pandemic, every action must be timely to avoid running behind the transmission.

CONCLUSION

Despite limited levels of preparedness prior to the pandemic, Cambodia successfully contained the spread of SARS-CoV-2 in the first year and quickly attained high vaccine coverage by the second year of the response. The core of this success was the strong political will and high cooperation from the public. However, Cambodia needs to improve its infrastructure in order to quarantine contacts isolate cases and increase its laboratory capacity to better prepare for future health emergencies.

This review is among the few that presents the COVID-19 epidemiology and response, and highlights lessons learned in Cambodia for organising and investing in future responses.

This paper has not explicitly covered a few important aspects such as good governance, resource mobilization, and multilateral cooperation (e.g., role of authorities and other ministries).

398   Additionally, the COVID-19 data presented in this paper may be highly under-reported and may

399   not reflect the actual trends due to limited testing capacity and changing surveillance strategies.

ACKNOWLEDGMENTS

We would like to thank all the Inter-ministerial COVID-19 Response Committee and Cambodia's Communicable Disease Control Department (CCDC) for providing data.

AUTHOR CONTRIBUTIONS

SC, GK, SM, WP, WD, PI, CC, and VO designed the study; SC analyzed data; SC, GK, SM, WP, WD, PI, CC, and VO wrote the manuscript.

FUNDING

This work received no external funding.

CONFLICT OF INTERESTS

All authors have declared no competing interest.

ETHICS APPROVAL

Not applicable.

DATA AVAILABILITY STATEMENT

Data is publicly available. It can be accessed via the WHO online database (Cambodia: WHO Coronavirus Disease (COVID-19) Dashboard With Vaccination Data | WHO Coronavirus (COVID-19) Dashboard With Vaccination Data).

REFERENCES

1. Van Damme W, Dahake R, Delamou A, *et al.* The COVID-19 pandemic: diverse contexts; different epidemics—how and why? *BMJ Glob Health* 2020;5(7):e003098. doi: 10.1136/bmjgh-2020-003098

2. World Health Organization. Regional Office for the Western P. The Kingdom of Cambodia health system review, Manila: WHO Regional Office for the Western Pacific, 2015. Available: https://apps.who.int/iris/bitstream/handle/10665/208213/9789290616917_eng.pdf [Accessed 16 December 2022 2022].

3. The World Bank. Physicians (per 1,000 per people) - Cambodia, Washington DC: The World Bank, 2014. Available: https://data.worldbank.org/indicator/SH.MED.PHYS.ZS?locations=KH [Accessed 16 December 2022].

4. World Health Organization. The Global Health Observatory, Geneva, Switzerland: WHO, 2020. Available: https://www.who.int/data/gho/data/indicators/indicator-details/GHO/hospital-beds-(per-10-000-population) [Accessed 16 December 2022].

5. The World Bank. Beds (per 1,000 per people) - Cambodia, Washington DC: The World Bank, 2016. Available: https://data.worldbank.org/indicator/SH.MED.BEDS.ZS?locations=KH [Accessed 16 December 2022].

6. World Health Organization. Joint External Evaluation of IHR Core Capacities, Geneva, Switzerland: WHO, 2017. Available: https://apps.who.int/iris/handle/10665/254705 [Accessed 16 December 2022].

7. Nit B, Samy AL, Tan SL, *et al.* Understanding the Slow COVID-19 Trajectory of Cambodia. *Public Health Pract (Oxf)* 2021;2:100073. doi: 10.1016/j.puhip.2020.100073 [published Online First: 2021/02/02]

8. Tatum M. Cambodia ends controversial COVID-19 restrictions. *Lancet* 2021;397(10289):2035. doi: 10.1016/S0140-6736(21)01196-X

9. World Health Organization (WHO). Cambodia: WHO Coronavirus Disease (COVID-19) Dashboard, 2023. Available: https://covid19.who.int/region/wpro/country/kh/ [Accessed 07February 2023 2023].

10. Chhim S. Applied Epidemiology in Cambodia. The Australian National University (Australia), 2021.

11. World Health Organization. 2019 Novel Coronavirus (2019-nCoV): Strategic Preparedness and Response Plan, Geneva, Switzerland: WHO, 2020. Available: https://www.who.int/publications/i/item/strategic-preparedness-and-response-plan-for-the-new-coronavirus [Accessed 16 December 2022].

12. Sheikhbardsiri H, Raeisi AR, Nekoei-Moghadam M, *et al.* Surge Capacity of Hospitals in Emergencies and Disasters With a Preparedness Approach: A Systematic Review. *Disaster Med Public Health Prep* 2017;11(5):612-20. doi: 10.1017/dmp.2016.178 [published Online First: 2017/03/08]

13. Macdonald S. Learning from Cambodia's COVID-19 Response: Foreign Policy in Focus (FPIF), 2021. Available: https://fpif.org/learning-from-cambodias-covid-19-response/ [Accessed 19 October 2022].

14. Johnson S, Srey V. Thousands of Cambodians go hungry in strict lockdown zones, Phnom Penh, Cambodia: The Guardian, 2021. Available: https://www.theguardian.com/global-

development/2021/may/13/thousands-of-cambodians-go-hungry-in-strict-lockdown-
zones [Accessed 16 December 2022].

15. Wendy Karamba, Isabelle Salcher, Tong K. Results from the High-Frequency Phone Survey
of Households Round 5 (1–21 March 2021), Phnom Penh, Cambodia: World Bank
Group, 2021. Available:
https://documents1.worldbank.org/curated/en/833701624563697124/pdf/Main-
Report.pdf [Accessed 16 December 2022].

16. Schiefloe PM. The Corona crisis: a wicked problem. *Scand J Public Health* 2020;49(1):5-8.
doi: 10.1177/1403494820970767

17. Nicola M, Sohrabi C, Mathew G*, et al.* Health policy and leadership models during the
COVID-19 pandemic: A review. *Int J Surg* 2020;81:122-29. doi:
10.1016/j.ijsu.2020.07.026

18. Al Saidi AMO, Nur FA, Al-Mandhari AS*, et al.* Decisive leadership is a necessity in the
COVID-19 response. *Lancet* 2020;396(10247):295-98. doi: 10.1016/S0140-
6736(20)31493-8

19. Stoller JK. Reflections on leadership in the time of COVID-19. *BMJ Leader* 2020;4:77-79.
doi: 10.1136/leader-2020-000244

20. Kaul V, Shah VH, El-Serag H. Leadership During Crisis: Lessons and Applications from the
COVID-19 Pandemic. *Gastroenterology* 2020;159(3):809-12. doi:
10.1053/j.gastro.2020.04.076

21. Phillips G, Kendino M, Brolan CE*, et al.* Lessons from the frontline: Leadership and
governance experiences in the COVID-19 pandemic response across the Pacific region.
Lancet Reg Health West Pac 2022;25 doi: 10.1016/j.lanwpc.2022.100518

22. Torres I, Sippy R, Sacoto F. Assessing critical gaps in COVID-19 testing capacity: the case
of delayed results in Ecuador. *BMC Public Health* 2021;21(1):637. doi: 10.1186/s12889-
021-10715-x

23. Cheng H-Y, Li S-Y, Yang C-HJJotFMA. Initial rapid and proactive response for the COVID-
19 outbreak—Taiwan's experience. 2020;119(4):771.

24. Tambo E, Djuikoue IC, Tazemda GK*, et al.* Early stage risk communication and community
engagement (RCCE) strategies and measures against the coronavirus disease 2019
(COVID-19) pandemic crisis. *Global Health Journal* 2021;5(1):44-50. doi:
https://doi.org/10.1016/j.glohj.2021.02.009

25. World Health Organization. Coronavirus disease (COVID-19): Herd immunity, lockdowns
and COVID-19: WHO, 2020. Available: https://www.who.int/news-room/questions-and-
answers/item/herd-immunity-lockdowns-and-covid-19 [Accessed 12 October 2022].

26. Kaba MW, Baesel K, Poch B*, et al.* IDPoor: a poverty identification programme that enables
collaboration across sectors for maternal and child health in Cambodia. *BMJ*
2018;363:k4698. doi: 10.1136/bmj.k4698