Type of article: Systematic review article

Title of the article: Clinical Virology and Effect of Vaccination and Monoclonal Antibodies
against SARS-CoV-2 Omicron Sub Variant BF.7 (BA.5.2.1.7): A systematic review.

Name of the Authors: Santenna Chenchula*[1], Krishna Chaitanya Amerneni, Mohan Krishna
Ghanta, Padmavathi R[2], Madhu Bhargavi Chandra, Madhu Babu Adusumilli,Madhavrao
Chavan[1], Sofia Mudda, Rupesh Gupta, Bhawna Lakhawat.

Running Title: The BF.7 (BA.5.2.1.7) variant Virology and Preventing Strategies.

The affiliation(s) of the author(s):

1. *Santenna Chenchula, PhD, Department of Pharmacology, All India Institute of
Medical Sciences (AIIMS)Bhopal, Madhya Pradesh, India. Email: csanten7@gmail.com

2. Krishna Chaitanya Amarneni, Faculty, Western Michigan University, Kalamazoo,
Michigan. Email: krishnachaitanya.amerneni@med.wmich.edu

3. Mohan Krishna Ghanta, Assistant Professor, Department of Pharmacology, MVJ
Medical College and Research Hospital, Bangalore, Karnataka, Email:
mohanakrishna.ghanta@gmail.com

4. Padmavathi R, Resident, SVS Medical College and Hospital, Telangana, India. Email:
pad.mythili@gmail.com

5. Madhu Bhargavi Chandra, Intern, All India Institute of Medical sciences, Bhopal.
email: madhubhargavi196@gmail.com

6. Madhu Babu Adusumilli, Intern, All India Institute of Medical sciences, Bhopal. email:
madhu.adusumilli3@gmail.com

7. Sofia Mudda, Medical Officer, All India Institute of Medical sciences, Bhopal. Email:
sophia.dr@gmail.com

8. Madhavrao Chavan, Department of Pharmacology, All India Institute of Medical Sciences (AIIMS) Mangalagiri, Andhra Pradesh, India. Email: madhavrao.pharm@aiimsmangalagiri.edu.in

9. Rupesh Gupta, Assistant Professor, Department of Internal Medicine, Government Medical College, Shahdol, Madhyapradesh, India. Email: neolog.raj@gmail.com

10. Bhawna Lakhawat , Tutor, department of Pharmacology, All India Institute of Medical sciences, Bhopal, Email: bhawnaajmer001@gmail.com

*Corresponding Author: Santenna Chenchula, Department of Pharmacology, 3rd floor, All India Institute of Medical Sciences, Bhopal,462020, Madhya Pradesh, India.

Phone no: 7987220348

Email Id: csanten7@gmail.com.

The total number of pages:11

Total number of Tables/Graphs:0

Total number of Images: 0

Word counts: For abstract: 187

For the text: 2133

Prior publication: Nil

Sources of Support: Nil

Conflicts of interest: Nil

Read and approved by all authors: Yes.

Abstract:

Since its identification, the novel coronavirus "severe acute respiratory syndrome coronavirus 2 "(SARS-CoV-2) In in late 2019 AT Wuhan, China, by the World Health Organization (WHO), which cause the coronavirus disease 2019, is rapidly spreading, resulting in the global pandemic. As of 19 December 2022, more than 64 million confirmed cases and 6,645,812 deaths have been reported across the world. Over time, the SARS-CoV-2 acquired genetic mutations resulting in multiple types of SARS-CoV-2 variants and subvariants that have been confirmed. The Omicron (B.1.1.529) variant was identified later in November 2021, with enhanced immune escape and was followed with various sublineages due to mutations in the spike protein of the SARS-CoV-2. However, rapid resurge in COVID-19 reports by Omicron subvariant BF.7(BA.2.75.2) in China and other countries, alarming global threat. The present systematic review was conducted using the MeSH terms and keywords "Omicron" AND "BA.5.2.1.7" OR "BF.7" in Pub Med, Google Scholar and MedRXiv database and grey literature from the authentic database and websites. We identified a total of 14 eligible studies. We have reviewed all the eligible available studies to understand the viral mutations, and factors associated with the increase in the reports of COVID-19 cases in China and across the world and to evaluate the effectiveness of vaccination and monoclonal antibodies against the BF.7 variant.

Introduction:

Since its identification in late 2019 the novel coronavirus, severe acute respiratory syndrome

coronavirus 2 (SARS-CoV-2) In Wuhan, China, by the World Health Organization (WHO),

which cause the coronavirus disease 2019, it is rapidly spreading, resulting in the global

pandemic [1]. As of 19 December 2022, more than 64 million confirmed cases and

6,645,812 deaths have been reported across the world [2]. Over time, the SARS-CoV-2 acquired

genetic mutations resulting in multiple types of SARS-CoV-2 variants and subvariants that have

been confirmed [3]. Certain variants have gained keen attention because of their characteristics

of rapid transmissibility, enhanced immune escape and severity of the infection and these are

considered variants of concern (VOC) that continue to threaten public health [3].

Later in November 2021, the Omicron (B.1.1.529) variant with enhanced immune escape was

first reported from Botswana and thereafter from South Africa with an increased infection [3-4].

Very soon it has been spread swiftly to several other countries, across the world with subtle raise

in the number of COVID-19 infections [4-5]. Subsequently, Omicron sublineages with

increasingly greater replication advantages emerged, replacing the previous predominant sub

lineage [3]. There are about 200 sublineages of Omicron [3]. The original Omicron variant was

sublineage BA.1, BA.2, BA.3, BA.4 and BA.5 [5-6]. Other Omicron sublineages, such as BQ.1,

BQ.11, BF.7, BA.2.75, and XBB, which evolved from various previously circulating

sublineages, have been increasing in prevalence worldwide [7]. Each sublineage differs from the

others by several mutations in the spike protein except for BA.4 and BA.5, which have identical

spike proteins [8]. Evidence has shown that all omicron sub-variants are distinct from pre-

omicron variants, including BA.1, BA.2 and BA.5 omicron sub-variants are also antigenically

distinct from each other [9]. The BF.7 also known as BA.2.75.2 Variant was first identified on

May 13, 2022, in Belgium. In response to the current surge in the COVID-19 reports by Omicron

subvariant BF.7 also known as BA.2.75.2 in China and other countries, triggering global alarm

the present review was conducted to understand the virology, factors associated with increased

reports of COVID-19 infections with BF.7 variant in China and possible urgent preventing

strategies to be taken to curtail the novel omicron variants outbreak across the world.

Methodology:

We have performed a comprehensive literature search from inception to till December 2022,

using the MeSH terms and keywords "Omicron" AND "BA.5.2.1.7" OR "BF.7" in Pub Med,

Google Scholar and MedRXiv database and grey literature from the authentic database and

websites to find relative information about Omicron subvariant BF.7 (BA.5.2.1.7) and data

assessed to understand viral mutations, factors associated with increased reports of COVID-19

infections with BF.7 variant in China and on literature evidence on vaccine effectiveness and

monoclonal antibodies(mAbs) against BF.7 to understand possible urgent preventing strategies

to be taken to curtail the novel omicron variants outbreak in China and across the world.

Results:

In the present review, after removing duplications and literature not mentioned regarding BF.7 or

BA.5.2.1.7, a total of 14 studies were found eligible among the 79 studies **[Figure1].** The

SARS-CoV-2 Omicron BF.7 (BA.5.2.1.7) a subvariant of BA.5, is responsible for the ongoing

resurgence of cases since late September 2022 in China [10].To date, BF.7 (BA.5.2.1.7)

subvariant has been reported in Belgium, China, Denmark, Norway, France, Germany, India,

Mongolia, the United Kingdom and the United States[11-12]. However, Before China, the BF.7

variant has been circulating at an incredibly intense level in USA and Europe since August 2022[8-10]. The current reports from China indicate BF.7 has the strongest infection ability out of the Omicron subvariants, being quicker to transmit than other variants, having a shorter incubation period, and with a greater capacity to infect people who have had a previous COVID infection, or been vaccinated, or both [11]. The most common symptoms of an infection with BF.7 were similar to those associated with other Omicron subvariants, primarily upper respiratory symptoms [12]. Patients may have a fever, cough, sore throat, running nose and fatigue, and gastrointestinal symptoms like vomiting and diarrhoea among other symptoms [11-12]. The basic reproduction number Rt of omicron BF.7(BA.5.2.1.7) variant was 10 to 18.6. which is significantly high compared to BA.1 Omicron variant which is only 5.08[11]. A study by Kathy Leung et al., to track the effective reproduction number Rt of Omicron BF.7 in Beijing from November to December 2022 found that the Rt increased to 3.42 (95% CrI: 2.79 – 4.17) on November 18, the infection incidence peaked on December 10, and the cumulative infection attack rate was 42.5% (95% CrI: 20.3 – 63.9) on December 14[13]. The high transmission rate of BF.7(BA.5.2.1.7), is due to the novel mutations in the spike protein [11-15]. Evidence from the molecular modelling studies revealed the mechanisms of antibody-mediated immune evasion by R346T, K444T, F486S, and D1199N mutations on the receptor-binding domain (RBD) [15]. However, the omicron subvariant BF.7 carries an additional specific mutation, R346T in the spike protein of SARS-CoV-2 which is derived from the BA.4/5 subvariant, which is why the BF.7 variant has a 4.4-fold higher neutralization resistance than the original D614G variant [15]. The Arg346 mutations R346T in the BF.7 variant spike glycoprotein, particularly those on the RBD has been associated with an increased immune escape capability to neutralize antibodies generated by vaccines or previous infection [11-15]. A study by Wang, Qian et al., investigated

the viral receptor affinities to dimeric human ACE2 by surface plasmon resonance and antibody

evasion properties of the novel omicron subvariants BA.2, BA.4/5, BA.4.6, BA.4.7, BA.5.9, and

BF.7. Findings of the study have shown that all the spike proteins from BA.4/5 sublineages,

including BF.7 have shown a higher affinity for human ACE2[16].

Effect of monoclonal antibodies and Vaccines against BF.7:

The sensitivity profile of novel omicron subvariants to neutralisation by a panel of 23 mAbs

found that the neutralisation profiles of BF.7, BA.4.6, BA.4.7 and BA.5.9 was similar to BA.4/5

except for mAbs in RBD class 3 (Bebtelovimab, Sotrovimab, romlusevimab, XGv289, XGv282,

2-7,ciglavimab, BD55-5840, BD55-3152,BD-804) which showed a substantial reduction in their

neutralisation potency due to R346T mutation, which is associated with eliminated or weakened

hydrogen bonds or salt bridges, or both, between R346 and some RBD class 3 mAbs [16]. A

monoclonal antibody cocktail combination of cilgavimab and tixagevimab, has received an

emergency use authorisation for the prevention of COVID-19, could also not neutralise BF.7 and

other novel variants BA.4.6, BA.4.7, and BA.5.9[16].

A study by Sullivan, David J et al., to evaluate polyclonal antibodies from individuals both with

at least 3 vaccine doses and also recently recovered from Omicron COVID-19(VaxCCP)

neutralizing activity against new Omicron lineages in 740 individual patient samples from 37

separate cohorts defined by boosted vaccinations with or without recent Omicron COVID-19, as

well as infection without vaccination found that more than 96% of the plasma samples from

individuals in the recently i.e., within 6 months boosted VaxCCP study cohorts neutralized

BQ.1.1 with 79%, XBB.1 with 22% and BF.7 with 94% variants respectively [17].

Another study by Sullivan DJ et al. found that the BA.5-bivalent booster dose vaccines elicited

better neutralization against the newly emerged Omicron sublineages including BF.7 variant than

the parental mRNA vaccine and those individuals with SARS-CoV-2 infection history develop

higher and broader neutralization antibodies against the ongoing novel Omicron sublineages

after the BA.5-bivalent booster [18].

Another study by Zhu, Ka-Li Jiang et al., found that two-dose CoronaVac or a third-dose

ZF2001 booster elicits limited neutralization against Omicron subvariants 6 months after

vaccination but hybrid immunity as well as Delta, BA.1, and BA.2 breakthrough infections

induced long-term persistence of the antibody response, and over 70% of sera neutralized BF.7.

as well as BA.1, BA.2, BA.4/BA.5[19].

In an in-vitro study by Wang, Qian et al., to investigate the antibody evasion properties of

BA.4.6, BA.4.7, BA.5.9, and BF.7, we assessed the sensitivity of their corresponding

pseudoviruses to neutralisation with serum samples from healthy individuals who had received

three doses of a COVID-19 mRNA vaccine BNT162b2 or mRNA-1273 and patients with either

BA.1, BA.2, or BA.4/5 breakthrough infection after vaccination found that the 50% inhibitory

dose (ID_{50}) titres of the booster dose vaccinated serum samples against BA.4.6, BA.4.7, BA.5.9,

and BF.7 were similar to that against BA.4/5, with no more than 1·5-fold deviation in the

geometric mean values and neutralizing antibody titres against new emerging omicron

subvariants were higher than those of the serum samples of previous omicron variants[16].

Another systematic review of 27 included studies across the world on the effect of booster dose

vaccination against Omicron and its subvariants by Chenchula S et al., found that booster-dose

vaccines have produced significant efficacy against the SARS-CoV-2 and its subvariants [20].

Discussion:

The present review focuses on the variant of concern Omicron BF.7 with, increased binding to angiotensin-converting enzyme-2 (ACE2) receptor, increased expression of the RBD, enhanced infectivity and transmissibility potential with an increased immune evasion and resistance to the majority of available monoclonal antibodies is due to the R346T mutation in spike glycoprotein, particularly those on the RBD[15-16]. Being quicker to transmit than other Omicron variants, having an increased binding to angiotensin-converting enzyme-2 (ACE2) receptor, increased expression of the RBD, shorter incubation period, increased evasion of antibodies and a greater capacity to infect people who have had a previous COVID-19 infection, and/or been vaccinated, or both, the BF.7 variant is causing a significant increase in number COVI-19 cases in China and across the world [16]. A SARS-CoV-2 transmission model study by Kathy Leung et al., from China, using the data from recent outbreaks in Hong Kong and Shanghai to compare different scenarios in China concluded that the hospitals will be overwhelmed if infections rise as rapidly because of the easing of Zero COVID policy restrictions and will probably result in about one million deaths over the next few months, the study forecasts [21]. The study also suggests that, if 85% of the population gets a booster fourth dose of the COVID -19 vaccine other than the inactivated-virus vaccines, it could slow the rise in infections and reduce the number of severe infections and deaths by up to 35% [22]. According to this study, if China lifts the zero-COVID policy the Omicron could infect between 160 million and 280 million people and 1.55 million deaths largely among unvaccinated older adults [22]. The study also suggests that, if 85% of the population gets the fourth dose of a vaccine other than the inactivated-virus vaccine could slow the increase in infections, the number of severe infections and deaths [22]. According to reports from China, more than 90% of the population has been fully vaccinated [23]. However, less than

half of people aged 80 and over have received three doses of the vaccine [23]. Evidence also

shows that the inactivated vaccine CoronaVac from China produced lower levels of neutralizing

antibodies and these antibody levels also dropped quickly over time [23]. In addition, studies

also have shown that protection from severe disease was also very less in preventing severe

disease and death among 80 years and older [22]. Evidence from the literature suggests that the

majority of monoclonal antibodies have not shown reduced neutralization potential against BF.7

variant. However, booster dose vaccination was significantly associated with increased efficacy

against BF.7 and other emerging Omicron variants [16-20]. Therefore, the R346T mutation in

the BF.7 variant was associated with reduced sensitivity to monoclonal antibodies, increased

binding to angiotensin-converting enzyme-2 (ACE2) receptor, and increased expression of the

RBD. Based on the evidence, COVID-19 infection with SARS-CoV-2 variant BF.7 may

associate with a serious illness among the non-vaccinated population, with weaker immune

systems such as the elderly and those with concomitant comorbidities [13-15].Limitations of the

present review includes, there are only few number of studies available on BF.7 variant. Meta

analysis was not done because of majority of studies are in vitro and there was no comparator.

Conclusion:

Findings of the present review suggest that infection with SARS-CoV-2 subvariant BF.7 with

R346T mutation on the spike glycoprotein RBD, may be associated with increased

transmissibility, and serious illness among the non-vaccinated and immune-compromised

individuals such as the elderly, children and those with concomitant comorbidities. Therefore we

recommend that, in addition to regular surveillance activities, the protection strategies need to

rapidly achieve by global vaccination to nonvaccinated populations and booster dose vaccination

with other than inactivated vaccines, especially among immunocompromised and vulnerable

populations with comorbidities, those aged ≥60 years more and in children, to prevent possible

outbreaks of novel SARS-CoV-2 Omicron subvariants and to decrease severity, mortality and to

mitigate the healthcare and economic impacts due to the emerging Omicron sublineages and

future VOCs.

Acknowledgement:

None to declare.

Conflict of Interests: The authors declare that there are no conflicts of interest.

Grants and funding: None.

Author Contributions: Santenna Chenchula conducted the literature search and data extraction

and drafted the manuscript. Krishna Chaitanya Amarneni, Mohan Krishna Ghanta, Padmavathi

R, Madhu Bhargavi Chandra, Madhu Babu Adusumilli, Bhawna Lakhawat, Madhavrao Chavan,

Sofia Mudda, Rupesh Gupta revised the final manuscript. All authors reviewed and approved the

final version of the manuscript.

Data Availability:

The data used in this systematic review is available from the corresponding author with a

reasonable request.

References:

1. WHO Coronavirus (COVID-19) dashboard. https://covid19.who.int/. Accessed: 20

December 2022.

2. WHO Classification of omicron (B.1.1.529): SARS-CoV-2 variant of concern. Statement 26

November 2021.Accessed: 20 December, 2022: https://www.who.int/news/item/26-11-2021-

classification-of-omicron-(b.1.1.529)-sars-cov-2-variant-of-concern.

3. Classification of Omicron (B.1.1.529): SARS-CoV-2 Variant of Concern. Available from:

https://www.who.int/news/item/26-11-2021-classification-of-omicron-(b.1.1.529)-sars-cov-

2-variant-of-concern. Accessed on: 20 December 2022.

4. Viana R, Moyo S, Amoako DG, et al. Rapid epidemic expansion of the SARS-CoV-2

Omicron variant in southern Africa. Nature. 2022;603(7902):679-686.

5. Arora P, Zhang L, Rocha C, et al. Comparable neutralisation evasion of SARS-CoV-2

omicron subvariants BA.1, BA.2, and BA.3. Lancet Infect Dis. 2022;22(6):766-767.

6. Rössler A, Netzl A, Knabl L, Schäfer H, Wilks SH, Bante D, et al . BA.2 and BA.5 omicron

differ immunologically from both BA.1 omicron and pre-omicron variants. Nat Commun.

2022 Dec 13;13(1):7701. doi: 10.1038/s41467-022-35312-3.

7. Implications of the emergence and spread of the SARS-CoV-2 variants of concern BA.4 and

BA.5 for the EU/EEA. European Centre for Disease Prevention and Control. Available at:

https://www.ecdc.europa.eu/en/news-events/implications-emergence-spread-sars-cov-2-

variants-concern-ba4-and-ba5 . Accessed on 20 December, 2022.

8. Tegally H, Moir M, Everatt J, Giovanetti M, Scheepers C, Wilkinson E et al., Emergence of

SARS-CoV-2 Omicron lineages BA.4 and BA.5 in South Africa. Nat Med. 2022;28(9):1785.

9. Jian F, Yu Y, Song W, Yisimayi A, Yu L, Gao Y, et al. Further humoral immunity evasion of

emerging SARS-CoV-2 BA.4 and BA.5 subvariants. Lancet Infect Dis. 2022

Nov;22(11):1535-1537. doi: 10.1016/S1473-3099(22)00642-9.

10. Mallapaty S. Can China avoid a wave of deaths if it lifts strict zero COVID policy? Nature.

2022 ;612(7939):203. doi: 10.1038/d41586-022-04235-w.

11. BF.7: What to know about the Omicron COVID variant. Available at :

https://www.cbsnews.com/news/bf7-new-omicron-coronavirus-variant-covid/. Accesed on

20 December 20222.

12. Aleem A, Akbar Samad AB, Slenker AK. Emerging Variants of SARS-CoV-2 And Novel

Therapeutics Against Coronavirus (COVID-19). 2022 Oct 10. In: StatPearls [Internet].

Treasure Island (FL): StatPearls Publishing; 2022 Jan–. PMID: 34033342.

13. Kathy Leung, Eric H. Y. Lau, Carlos K. H. Wong, Gabriel M. Leung, Joseph T. Wu.

Estimating the transmission dynamics of Omicron in Beijing, November to December

2022.medRxiv 2022.12.15.22283522; doi: https://doi.org/10.1101/2022.12.15.22283522.

14. SARS-CoV-2 Omicron BF.7 variant. Available at : https://en.wikipedia.org/wiki/SARS-

CoV-2_Omicron_BF.7_variant. Accessed on 20 December 2022.

15. Qu P, Evans JP, Faraone JN, Zheng YM, Carlin C, Anghelina M, et al. Enhanced

neutralization resistance of SARS-CoV-2 Omicron subvariants BQ.1, BQ.1.1, BA.4.6, BF.7,

and BA.2.75.2. Cell Host Microbe. 2022 Nov 22:S1931-3128(22)00568-6. doi:

10.1016/j.chom.2022.11.012.

16. Wang Q, Li Z, Ho J, Guo Y, Yeh AY, Mohri H, et al. Resistance of SARS-CoV-2 omicron

subvariant BA.4.6 to antibody neutralisation. Lancet Infect Dis. 2022 Dec;22(12):1666-1668.

doi: 10.1016/S1473-3099(22)00694-6.

17. Cai J, Deng X, Yang J, Sun K, Liu H, Chen Z, et al . Modeling transmission of SARS-CoV-2

Omicron in China. Nat Med. 2022 Jul;28(7):1468-1475. doi: 10.1038/s41591-022-01855-7.

18. China Covid: What is China's policy and how many cases are there?.BBC: Reality check.

Available from: https://www.bbc.com/news/59882774. Accessed on 20 December 2022.

19. Mallapaty S. China's COVID vaccines have been crucial - now immunity is waning. Nature.

2021 Oct;598(7881):398-399. doi: 10.1038/d41586-021-02796-w.

20. Sullivan DJ, Franchini M, Senefeld JW, Joyner MJ, Casadevall A, Focosi D. Plasma after

both SARS-CoV-2 boosted vaccination and COVID-19 potently neutralizes BQ.1.1 and

XBB.1.   bioRxiv   [Preprint].   2022   Nov   30:2022.11.25.517977.   doi:

10.1101/2022.11.25.517977.

21. Zhu, Ka-Li Jiang, Xiao-Lin Wang, Xue-Jun Xia, Xian Zhan, Bing-Dong Cao, Guo-Ping

Wang, et al.,.Durability of Neutralization Against Omicron Subvariants after Vaccination

and   Breakthrough   Infection.   Available   at

SSRN: https://ssrn.com/abstract=4281072 or http://dx.doi.org/10.2139/ssrn.4281072

22. Chenchula S, Karunakaran P, Sharma S, Chavan M. Current evidence on efficacy of

COVID-19 booster dose vaccination against the Omicron variant: A systematic review. J

Med Virol. 2022 ;94(7):2969-2976. doi: 10.1002/jmv.27697.

Figure 1: PRISMA Flow Diagram for the study selection.

Identification of studies via in Pub Med, Google Scholar and MedRXiv by Using Following MESH or Key words

"Omicron" AND "BA.5.2.1.7" OR "BF.7"

Identification

Records identified from databases: (n=80) → Records excluded based on title and abstract review (n=35)

Screening

Records Screened (n = 45)

Eligibility

Full-text articles assessed for eligibility (n=22) → Records excluded with reasons (n= 23)
Duplicate data (n=10)
BF.7 variant not mentioned (n=13)

Included

Studies included in qualitative analysis (n=14)