**Research article: Ageing Increases the Incidence Rate of Post Coronary Artery Shunt Complications**

Running tittle: Current surgical management advances

**Authors' list: Basheer Abdullah Marzoog[1,*], Averina Daria Romanovna[2]**

[1] National Research Mordovia State University. Address: Bolshevitskaya Street, 68, Saransk, Rep. Mordovia, Russia. Postal address:  Russia, Mordovia republic, Saransk, Bolshevitskaya Street, 31, 430005.

[2] Sechenov First Moscow State Medical University. Address: Trubetskaya St., bld. 2, Moscow, 119991. ORCID: 0009-0004-9377-2241

[*]Corresponding author: Basheer Abdullah Marzoog, undergraduate medical school student at National Research Mordovia State University (marzug@mail.ru, +79969602820). Address: Bolshevitskaya Street, 68, Saransk, Rep. Mordovia, 430005. Postal address:  Mordovia republic, Saransk, Bolshevitskaya Street, 31. ORCID: 0000-0001-5507-2413, Scopus ID: 57486338800

Competing interests: No competing interests regarding the publication.

All authors read and approved the final version of the manuscript

**Abstract:**

**Background**: Several complications reported after coronary artery bypass graft surgery (CABG), such as postoperative arrythmia and postoperative stroke.

**Aims:** The study aimed to assess the role of age advancement in the incidence rate of post coronary artery shunt complications.

**Objectives:** Aging is a pathophysiological process that experience every living cell and starts from the early period of development, before even born.

**Materials and methods:** A retrospective analysis of 290 patients who underwent a coronary artery bypass graft at Mordovia republic hospital for the period 2017-2021. The sample was subdivided into two groups, the first group 126 patients (mean age: min; max, 55.21:41.00;60.00) and the second group 163 (M age: min; max, 66.11: 61.00; 80.00). For statistical analysis, we used the T test, the one-way ANOVA test, ROC analysis, and the Pearson correlation test using the Statistica 12 programme.

**Results**: Post-CABG arrythmia developed in elderly patients, $p<0.012528$. Subsequently, after CABG arrythmia, the ICU and total hospitalisation days, $p< 0.000000$; $p< 0.000072$, respectively.  Elderly people are at higher risk of psychosis after CABG surgery, $p <0.007379$. psychosis significantly increases hospitalisation days in the ICU, $p < 0.000140$. Postoperative stroke occurs more frequently in elderly people, $p < 0.037736$. Subsequently, postoperative stroke increases the ICU hospitalization days, $p <0.000747$. The number of ITAs used is less in the elderly, $p < 0.016145$. In terms of correlation, exist a direct association between age and ICU / total hospitalization days/number of complications, $r= 0.189046, 0.141415, 0.138565$; respectively.

**Conclusions:** The number of complications is determined by age, CPB time, aortic cross-clamp time, the days of hospitalisation in the ICU and the total days of hospitalisation. Elderly people undergoing CABG are at higher risk of psychosis, arrythmia, longer total and ICU hospitalization days, and stroke.

**Others:** Total hospitalization days depend on the presence of arrythmia, which is commonly seen in elderly patients > 63 years old.

**Keywords**: Surgery, Myocardial Infarction, Angioplasty, CABG, CVD & CVS, progressive Angina

**Background**

Coronary artery graft surgery (CABG) performed on patients with ischemic heart disease to relieve the discrepancy between demand and supply of heart tissue. CABG surgery is performed in two versions, with stopped heart (on-pump) or working heart (off-pump) with cardiopulmonary bypass (CPB). CABG surgery involves revascularization of the coronary arteries through grafting of vessels from the saphenous vein, brachial vein, internal thoracic arteries, internal mammary arteries, and intercostal thoracic arteries [1]. The surgical approach to CABG is performed endoscopically or by thoracotomy. Recent surgeries are performed using the Da-Venice robot to minimize the risk of complications and improve prognosis by reducing the recovery period [2,3].

Post CABG complications are frequent and seen in all types of CABG surgery, off-pump and on-pump. The remarkable advancement in the current CABG techniques reduced the complications frequency, but the underlying risk factors and pathophysiological mechanisms of development of each complication remain unclear. Depending on the type of surgery, the risk of development of complication is different. Several complications are reported after the postoperative period, such as postoperative arrythmia, post-operative psychosis, postoperative stroke, postoperative myocardial infarction, postoperative hydrothorax, re-sternotomy, postoperative dyspnea, and sternal wound infection [3,4]. The risk factors for development of post CABG include diabetes mellites, arterial hypertension, dyspnea, progressive angina, and post myocardial infarction (MI) sclerosis, advanced age, long CPB time, prolonged Aortic cross-clamp time , long ICU hospitalization days, and long total hospitalization days [5,6].

**Methods**

The sample was divided by age into two groups A (60≥age in years ≥41) and B (80≥age in years ≥61). Accordingly, we used the stepwise model builder-logistic regression. The used model variables in the test are ejection fraction, age in numbers, BMI, CPB, XCL time, days reanimation, total days after surgery, and number of complications. We found that age plays important role. (*Figure 1*) (*Table 1*)

Table 1: Predicted and real values of the sample divided by age.

| | Classification of cases (stat in copy of post coronary artery shunt complications)Odds ratio: infinityLog odds ratio: infinity | | | | | |
|---|---|---|---|---|---|---|
| | Predicted: A | Predicted: B | Percent correct | Predicted: A - Testing | Predicted: B - Testing | Percent correct - Testing |
| Observed: A | 50 | 0 | 100 | 69 | 5 | 93,2432432 |

| | Classification of cases (stat in copy of post coronary artery shunt complications)Odds ratio: infinityLog odds ratio: infinity | | | | | |
|---|---|---|---|---|---|---|
| | Predicted: A | Predicted: B | Percent correct | Predicted: A - Testing | Predicted: B - Testing | Percent correct - Testing |
| Observed: B | 0 | 61 | 100 | 7 | 91 | 92,8571429 |

Therefore, the sample has been divided by age into two groups, the first group included 126 patients (mean age: min; max, 55.21:41.00;60.00) and the second group 163 (mean age: min; max, 66.11: 61.00; 80.00). The data were collected from the archive of the hospital for the period 2017-2021. Retrospective analysis has been carried out to assess the role of age advancement in postoperative complications. The mean age of the sample by year is presented in Figure 2.

For statistical analysis, we used the T test, the one-way ANOVA test, and the Pearson correlation test using the Statistica 12 programme. (StatSoft, Inc. (2014). STATISTICA (data analysis software system), version 12. www.statsoft.com.)

**Results**

In 2017, 33 (11.37931 %) patients performed coronary bypass surgery. In 2018, the number increased to 35 (12.06897 %) patients. In 2019, the number significantly increased to 61 (21.03448%) patients. In 2020, the highest number of bypass surgeries performed, 84 (28.96552 %) patients. In 2021, 77 (26.55172%) patients underwent coronary bypass surgery. Body mass index (BMI) range from 16.360-45.350 $Kg/m^2$ (mean; standard error: 45.350; 0.241918). Ejection fraction ranged from 29.000-77.000 ml/min (mean; standard error: 55.772; 0.438403). 278 (95.86207%) patients underwent on-pump CABG surgery and 12 (4.13793 %) patients underwent off-pump CABG surgery. The period of artificial circulation in CPB surgeries ranged from 0.000-431.000 min (M; ±m: 431.000; 2.661081). The ischemia time ranged from 0.000-147.000 min (M; ±m: 55.466; 1.341570). Postoperative hospitalisation (hereafter, postoperative and post-CABG will be used interchangeably) in the intensive care unit ranged from 1.000 to 27.000 days (M; ±m: 2.821; 0.140122). The total hospitalization days (ICU hospitalization days plus recovery hospitalization days) after surgery ranged 7.000-50.000 days (M; ±m: 11.722; 0.292762).

In the first group, the mean age is 55,21 years (min; max, 41,00; 60,00) and the mean BMI is 28,8208 kg/m2 (min; max, 16,36; 45,35). Moreover, in the first group, mean ejection fraction is 54,85% (min; max, 29,00; 76,00) and the mean CPB 101,19 mints (min; max, 0,00; 270,00). Furthermore, the mean XCL time is 56,29 min (min; max, 0,00; 105,00) and the mean hospitalisation days in the intensive care unit is 2,4758 days (min; max, 1,00; 6,00).

Furthermore, the mean total hospitalization days after surgery is 10,96 days (min; max, 7,00000; 26,0000) and the mean number of complications is 0,2857 (min; max, 0,00; 3,00). Non-parametric parameters have been evaluated in table 2. With age, the number of shunts increases in the first group. However, a reduction in the incidence rate of complications has been observed in the second group (*Figure 3,4*).

Table 2: Age in the determination of preoperative conditions and post coronary artery shunt surgery complications. *The missed data are treated as part of the of the relative value but not the absolute value. ** values in bold are statically significant.

| Factor | Index | Group A (40-60 years old); 126 patients | Group B (61-80 years old); 164 patients | $X^2$ | P value |
|---|---|---|---|---|---|
| Sex | M | 121 (96.03 %) | 123 (75.00%) | 23.616 | **<0.001**\*\* |
| | F | 5 (3.96 %) | 41 (25.00 %) | | |
| Progress. Angina | Yes | 50 (39.68 %) | 62 (37.80 %) | 0.081 | 0.776 |
| | No | 76 (60.31 %) | 101 (61.58 %) | | |
| DM | Yes | 21 (16.66 %) | 45 (27.43 %) | 4.827 | **0.029**\*\* |
| | No | 105 (83.33 %) | 118 (71.95 %) | | |
| DM complication | Yes | 9 (7.14 %) | 19 (11.58 %) | 1.655 | 0.199 |
| | No | 117 (92.85 %) | 144 (87.80 %) | | |
| Dyspnea | Yes | 25 (19.84 %) | 30 (18.29 %) | 0.095 | 0.758 |
| | No | 101 (80.15 %) | 133 (81.09 %) | | |
| Arrythmia before surgery | Yes | 20 (15.87 %) | 49 (29.87 %) | 7.871 | **0.006**\*\* |
| | No | 106 (84.12 %) | 114 (69.51 %) | | |
| Post mi sclerosis | Yes | 84 (66.66 %) | 94 (57.31 %) | 2.432 | 0.119 |
| | No | 42 (33.33 %) | 69 (42.07 %) | | |
| Aneurysm LV | Yes | 7 (5.55 %) | 2 (1.219 %) | 4.413 | **0.036**\*\* |
| | No | 119 (94.44 %) | 161 (98.17 %) | | |
| Type of surgery | On-pump | 121 (96.03 %) | 157 (95.73 %) | 0.016 | 0.899 |
| | Off-pump | 5 (3.96 %) | 7 (4.26 %) | | |
| Number of used internal thoracic arteries | One | 68 (53.96 %) | 123 (75.00 %) | 14.018 | **<0.001**\*\* |
| | Two | 58 (46.03 %) | 41 (25.00 %) | | |
| Coronary artery shunts and plastic surgery of the LV aneurism | Yes | 6 (4.76 %) | 1 (0.60 %) | 5.215 | **0.023**\*\* |
| | No | 120 (95.23 %) | 163 (99.39 %) | | |
| Post-op. aneurism LV | Yes | 1 (0.79 %) | 2 (1.22%) | 0.126 | 0.723 |
| | No | 125 (99.20 %) | 162 (98.78 %) | | |
| Post-op. arrythmia | Yes | 8 (6.35%) | 26 (15.85 %) | 6.597 | **0.011**\*\* |
| | No | 118 (93.65 %) | 134 (81.70 %) | | |
| Post-op. psychosis * | Yes | 3 (2.38 %) | 11 (6.70 %) | 2.999 | 0.084 |
| | No | 121 (96.03%) | 148 (90.24 %) | | |
| post-op. Stroke * | Yes | 1 (0.79 %) | 3 (1.82 %) | 0.584 | 0.445 |
| | No | 123 (97.61 %) | 156 (95.12 %) | | |
| Post-operative MI * | Yes | 1 (0.79 %) | 1 (0.60 %) | 0.031 | 0.860 |
| | No | 123 (97.61 %) | 158 (96.34 %) | | |
| post-op. hydrothorax * | Yes | 13 (10.31 %) | 21 (12.80 %) | 0.489 | 0.485 |

| | | | | | | |
|---|---|---|---|---|---|---|
| | No | 111 (88.09 %) | 138 (84.14 %) | | | |
| Re-sternotomy * | Yes | 5 (3.96 %) | 4 (2.43 %) | 0.520 | 0.471 | |
| | No | 119 (94.44 %) | 155 (94.51 %) | | | |
| Post operative dyspnea * | Yes | 4 (3.17 %) | 7 (4.26 %) | 0.258 | 0.612 | |
| | No | 120 (95.23 %) | 152 (92.68 %) | | | |
| Postoperative superficial sternum infection * | Yes | 0 (0.00 %) | 1 (0.60 %) | 0.715 | 0.398 | |
| | No | 104 (82.54%) | 145 (88.41%) | | | |
| Postoperative deep sternum infection * | Yes | 0 (0.00 %) | 5 (3.04 %) | 3.634 | 0.057 | |
| | No | 104 (82.54%) | 141 (85.97 %) | | | |
| Status after surgery | Alive | 125 (99.20 %) | 160 (97.56 %) | 1.138 | 0.286 | |
| | Dead | 1 (0.79 %) | 4 (2.43 %) | | | |

In table 3, a comparative demonstration of various factors in the development of different complications in pre- and post- CABG survivals.

Table 3: Comparation demonstration of various factors in the development of different complications in post CABG survivals.

| Variable | T-tests; Grouping: Aneurysm LV (stat in Post cor shunt comp) Group 1: No Group 2: Yes | | | | | | |
|---|---|---|---|---|---|---|---|
| | Mean No | Mean Yes | t-value | df | p | Valid N No | Valid N Yes |
| Ejection fraction | 56,139 | 44,333 | 4,85773 | 287 | 0,000002 | 280 | 9 |
| Variable | T-tests; Grouping: Arrythmia Before surgery (stat in Post cor shunt comp) Group 1: No Group 2: Yes | | | | | | |
| | Mean No | Mean Yes | t-value | df | p | Valid N No | Valid N Yes |
| ECMO time | 106,507 | 0,000 | 9,009503 | 288 | 0,000000 | 278 | 12 |
| MI | 57,860 | 0,000 | 9,936552 | 288 | 0,000000 | 278 | 12 |
| Variable | T-tests; Grouping: Coronary Shunt +plastic LV aneurysm repair (stat in Post cor shunt comp) Group 1: No Group 2: Yes | | | | | | |
| | Mean No | Mean Yes | t-value | df | p | Valid N No | Valid N Yes |
| Ejection fraction | 56,110 | 42,143 | 5,10667 | 287 | 0,000001 | 282 | 7 |
| MI | 55,004 | 74,143 | -2,20413 | 288 | 0,028306 | 283 | 7 |
| Variable | T-tests; Grouping: Post-operative aneurysm LV (stat in Post cor shunt comp) Group 1: No Group 2: Yes | | | | | | |
| | Mean No | Mean Yes | t-value | df | p | Valid N No | Valid N Yes |
| Ejection fraction | 55,682 | 64,333 | -2,01070 | 287 | 0,045292 | 286 | 3 |
| ECMO time | 100,655 | 240,333 | -5,58113 | 288 | 0,000000 | 287 | 3 |
| MI | 54,840 | 115,333 | -4,72802 | 288 | 0,000004 | 287 | 3 |
| Variable | T-tests; Grouping: Post-operative arrythmia. (stat in Post cor shunt comp) Group 1: No Group 2: Yes | | | | | | |
| | Mean No | Mean Yes | t-value | df | p | Valid N No | Valid N Yes |
| Age | 60,976 | 64,161 | -2,51307 | 281 | 0,012528 | 252 | 31 |
| Days reanimation | 2,548 | 5,032 | -5,83306 | 281 | 0,000000 | 252 | 31 |
| Total hospitalization days after surgery | 11,310 | 15,000 | -4,02907 | 281 | 0,000072 | 252 | 31 |
| Variable | T-tests; Grouping: Post-operative psychosis (stat in Post cor shunt comp) Group 1: Yes Group 2: No | | | | | | |

| Variable | Mean Yes | Mean No | t-value | df | p | Valid N Yes | Valid N No |
|---|---|---|---|---|---|---|---|
| Age | 66,000 | 61,082 | 2,69885 | 281 | 0,007379 | 14 | 269 |
| Days reanimation | 5,143 | 2,699 | 3,86094 | 281 | 0,000140 | 14 | 269 |

| Variable | T-tests; Grouping: Post-operative Stroke (stat in Post cor shunt comp) Group 1: Yes Group 2: No | | | | | | |
|---|---|---|---|---|---|---|---|
| | Mean Yes | Mean No | t-value | df | p | Valid N Yes | Valid N No |
| Age | 68,250 | 61,226 | 2,087585 | 281 | 0,037736 | 4 | 279 |
| Days reanimation | 6,750 | 2,763 | 3,409293 | 281 | 0,000747 | 4 | 279 |

| Variable | T-tests; Grouping: Post-operative hydrothorax (stat in Post cor shunt comp) Group 1: Yes Group 2: No | | | | | | |
|---|---|---|---|---|---|---|---|
| | Mean Yes | Mean No | t-value | df | p | Valid N Yes | Valid N No |
| BMI | 27,346 | 29,085 | -2,33929 | 281 | 0,020021 | 34 | 249 |
| ECMO time | 123,265 | 97,044 | 3,83233 | 281 | 0,000157 | 34 | 249 |
| MI | 64,618 | 53,827 | 2,71109 | 281 | 0,007119 | 34 | 249 |
| Days reanimation | 4,912 | 2,534 | 5,80811 | 281 | 0,000000 | 34 | 249 |
| total days after surgery | 17,088 | 10,980 | 7,37431 | 281 | 0,000000 | 34 | 249 |

| Variable | T-tests; Grouping: Post-operative dyspnea (stat in Post cor shunt comp) Group 1: Yes Group 2: No | | | | | | |
|---|---|---|---|---|---|---|---|
| | Mean Yes | Mean No | t-value | df | p | Valid N Yes | Valid N No |
| Days reanimation | 6,909 | 2,654 | 6,22816 | 281 | 0,000000 | 11 | 272 |
| Total hospitalization days after surgery | 21,364 | 11,324 | 7,17446 | 281 | 0,000000 | 11 | 272 |

| Variable | T-tests; Grouping: Post-operative Subendocardial MI (stat in Post cor shunt comp) Group 1: No Group 2: Yes | | | | | | |
|---|---|---|---|---|---|---|---|
| | Mean No | Mean Yes | t-value | df | p | Valid N No | Valid N Yes |
| Total hospitalization days after surgery | 11,586 | 30,000 | -3,66815 | 248 | 0,000299 | 249 | 1 |

| Variable | T-tests; Grouping: Post-operative transmural. MI (stat in Post cor shunt comp) Group 1: No Group 2: Yes | | | | | | |
|---|---|---|---|---|---|---|---|
| | Mean No | Mean Yes | t-value | df | p | Valid N No | Valid N Yes |
| Days reanimation | 2,747 | 5,600 | -2,61051 | 248 | 0,009591 | 245 | 5 |
| Total hospitalization days after surgery | 11,404 | 24,200 | -5,87749 | 248 | 0,000000 | 245 | 5 |

| Variable | T-tests; Grouping: status (stat in Post cor shunt comp) Group 1: live Group 2: died | | | | | | |
|---|---|---|---|---|---|---|---|
| | Mean live | Mean died | t-value | df | p | Valid N live | Valid N died |
| ECMO time | 99,937 | 256,750 | -7,50135 | 288 | 0,000000 | 286 | 4 |
| MI | 55,000 | 88,750 | -2,97365 | 288 | 0,003192 | 286 | 4 |

| Variable | T-tests; Grouping: sex (stat in Post cor shunt comp) Group 1: M Group 2: F | | | | | | |
|---|---|---|---|---|---|---|---|
| | Mean M | Mean F | t-value | df | p | Valid N M | Valid N F |
| Age | 60,630 | 65,239 | -4,37030 | 287 | 0,000017 | 243 | 46 |
| BMI | 28,444 | 30,695 | -3,46865 | 287 | 0,000603 | 243 | 46 |
| Ejection fraction | 55,354 | 57,978 | -2,20457 | 287 | 0,028278 | 243 | 46 |

| Variable | T-tests; Grouping: Progressive Angina (stat in Post cor shunt comp) Group 1: No Group 2: Yes | | | | | | |
|---|---|---|---|---|---|---|---|
| | Mean No | Mean Yes | t-value | df | p | Valid N No | Valid N Yes |

| Variable | Mean No | Mean Yes | t-value | df | p | Valid N No | Valid N Yes |
|---|---|---|---|---|---|---|---|
| BMI | 28,421 | 29,406 | -1,99512 | 287 | 0,046975 | 177 | 112 |

| Variable | T-tests; Grouping: DM (stat in Post cor shunt comp) Group 1: No Group 2: Yes | | | | | | |
|---|---|---|---|---|---|---|---|
| | Mean No | Mean Yes | t-value | df | p | Valid N No | Valid N Yes |
| Age | 60,794 | 63,288 | -2,65962 | 287 | 0,008262 | 223 | 66 |
| BMI | 28,390 | 30,196 | -3,18160 | 287 | 0,001625 | 223 | 66 |
| Days reanimation | 2,592 | 3,580 | -3,02226 | 282 | 0,002739 | 218 | 66 |
| Total hospitalization days after surgery | 11,229 | 13,347 | -3,10156 | 282 | 0,002120 | 218 | 66 |

| Variable | T-tests; Grouping: DM complication (stat in Post cor shunt comp) Group 1: No Group 2: Yes | | | | | | |
|---|---|---|---|---|---|---|---|
| | Mean No | Mean Yes | t-value | df | p | Valid N No | Valid N Yes |
| BMI | 28,610 | 30,600 | -2,45502 | 287 | 0,014681 | 261 | 28 |
| Days reanimation | 2,582 | 5,011 | -5,41994 | 282 | 0,000000 | 256 | 28 |
| Total hospitalization days after surgery | 11,160 | 16,854 | -6,16561 | 282 | 0,000000 | 256 | 28 |

| Variable | T-tests; Grouping: dyspnea (stat in Post cor shunt comp) Group 1: Yes Group 2: No | | | | | | |
|---|---|---|---|---|---|---|---|
| | Mean Yes | Mean No | t-value | df | p | Valid N Yes | Valid N No |
| Total hospitalization days after surgery | 13,080 | 11,375 | 2,25940 | 280 | 0,024627 | 50 | 232 |

| Variable | T-tests; Grouping: arrythmia. Before surgery (stat in Post cor shunt comp) Group 1: No Group 2: Yes | | | | | | |
|---|---|---|---|---|---|---|---|
| | Mean No | Mean Yes | t-value | df | p | Valid N No | Valid N Yes |
| Age | 60,776 | 63,319 | -2,75395 | 286 | 0,006265 | 219 | 69 |

| Variable | T-tests; Grouping: post MI cardio-sclerosis (stat in Post cor shunt comp) Group 1: Yes Group 2: No | | | | | | |
|---|---|---|---|---|---|---|---|
| | Mean Yes | Mean No | t-value | df | p | Valid N Yes | Valid N No |
| Ejection fraction | 53,579 | 59,288 | -6,81607 | 287 | 0,000000 | 178 | 111 |
| Days reanimation | 3,053 | 2,450 | 2,10757 | 282 | 0,035950 | 175 | 109 |

In terms of post-operative complications, patients with coronary artery bypass graft (CABG) and plastic surgery repair of the left ventricle aneurysm had longer Aortic cross-clamp time, t-value -2.20413, p <0.028306. Additionally, patients with CABG have less ejection fraction, t-value 5.10667, p < 0.000001. Patients with post-CABG left ventricle (LV) aneurysm had less ejection fraction, t-value -2.01070, p <0.045292. Furthermore, patients with post-CABG LV aneurysm had a longer CPB time, t-value -5.58113, p < 0.000000.  Furthermore, patients with LV aneurysm had longer Aortic cross-clamp time, t-value -4.72802, p < 0.000004.

Post-CABG arrythmia developed in elderly patients, t-value -2.51307, p<0.012528. Subsequently, after CABG arrythmia, the ICU and total hospitalisation days, t value -5.83306, p< 0.000000; t-value -4.02907, p< 0.000072, respectively.  Elderly people are at increased risk of psychosis after CABG surgery, t value 2.69885, p <0.007379. Postoperative psychosis significantly increases ICU hospitalization days, t-value 3.86094, p < 0.000140.

Postoperative stroke occurs more frequently in elderly people, t-value 2.087585, p < 0.037736. Subsequently, postoperative stroke increases the ICU hospitalization days, t-value 3.409293, p <0.000747. The incidence rate of complications in the first and second groups is shown in Figure 5.

The number of ITAs used is less in the elderly, t-value 2.41992, p < 0.016145. In terms of correlation, we found a direct association between age and ICU / total hospitalisation days / number of complications, r= 0.189046, 0.141415, 0.138565; respectively. No statistical significance in the correlation between number of the complications and the number of the shunts as well as between the number of used internal thoracic arteries and number of complications.

**Discussion**

In light of results, prolonged Aortic cross-clamp time is observed in coronary artery bypass graft (CABG) and plastic surgery repair of the left ventricle aneurysm and patients with preoperative LV aneurysm. The prolonged Aortic cross-clamp time results in the use of two internal thoracic arteries. The prolonged aortic cross-clamp time increases the risk of hydrothorax and the number of shunts, which subsequently increases the CPB time. Patients with low BMI are at high risk of hydrothorax. Furthermore, hydrothorax and the use of two internal thoracic arteries increase the time of the CPB. The coexistence of CABG and left ventricle aneurysm increases the time required for CPB.

The post CABG ICU hospitalization days is related with the presence of post-operative arrythmia, post-operative psychosis, post-operative stroke, post-operative hydrothorax, post-operative dyspnea, and superficial sternal wound infection. Total hospitalization days depend on the presence of arrythmia, which is commonly seen in elderly patients > 63 years old. Additionally, deep sternal wound infection increases total hospitalisation days and hospitalisation days. The ejection fraction and the days of hospitalisation in the ICU are closely associated with the total days of hospitalisation. Ejection fraction is age, sex, and presence of CABG and or CABG with LV aneurysm plastic repair dependent.

Age is a crucial factor in terms of postoperative complications. Where advancement in age increases the days of hospitalisation in the ICU, the total days of hospitalisation, and the number of complications. Elderly people are at high risk for stroke, psychosis, and arrythmia [4,7–14]. However, elderly people passed one internal thoracic artery CABG.

The number of complications is not associated with the death and alive status of patients or with the number of shunts. Furthermore, there were no statistical differences in the number of complications and the number of shunts and the number of internal thoracic arteries used.

However, the number of complications depends on the existence of preoperative progressive angina and PMIMS. In addition, the number of complications is determined by age increase, CPB time, the Aortic cross-clamp time, the days of hospitalization in the ICU, and total hospitalization days.

The results of our study are consistent with those of other studies. Several studies reported that post-CABG complications include arrythmia, particularly atrial fibrillation [4,15–21]. However, the potential underlying pathophysiological mechanisms remain unclear and further elaboration is required. However, the potential pathophysiological pathway is multifactorial and involves autophagy dysfunction, upregulated sympathetic tone, mitochondrial dysfunction, inflammation, abnormal atrial conduction, and systemic inflammatory response indicated by elevated C-reactive protein and leukocytes [22–31]. Some studies showed that on-pump CABG surgery has higher risk of atrial fibrillation than off-pump [32–35].

**Conclusions**

Hydrothorax is seen in a low BMI and a long Aortic cross-clamp time which leads to longer CPB time. Progressive angina and PMIMS increase the risk of postoperative complications, particularly hydrothorax. Elderly people undergoing CABG are at higher risk of psychosis, arrythmia, longer total and ICU hospitalization days, and stroke.

**List of abbreviations**

CVD; Cardiovascular disease, COVID-19; Corona virus infectious disease-19, CABG; coronary artery bypass graft, CPB; cardiopulmonary bypass, BMI; body mass index, MI; myocardial infarction, DM; Diabetes mellites, ICU; intensive care unit, PMIMS; post-myocardial infarction myocardial sclerosis, LV; left ventricle, ITA; internal thoracic artery

# Declarations

1. Ethics approval and consent to participate: The study approved by the National Research Mordovia State University, Russia, from "Ethics Committee Requirement N8/2 from 30.06.2022".

2. Consent for publication: Written informed consent was obtained from the participants for publication of study results and any accompanying images.

3. Availability of data and materials: applicable on reasonable request.

4. Competing interests: The authors declare that they have no competing interests regarding publication.

5. Funding's: Not applicable (This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors)

6. Authors' contributions: MB is the writer, researcher, collected and analyzed data, and revised the manuscript. **Daria Romanovna  Averina** inserted the data from the patients history into excel table. All authors have read and approved the manuscript.

7. Acknowledgments: not applicable

8. Authors' information: Basheer Abdullah Marzoog, National Research Mordovia State University (marzug@mail.ru, +79969602820). Address: Bolshevitskaya St., 68, Saransk, Rep. Mordovia, 430005. ORCID: 0000-0001-5507-2413. **Daria Romanovna Averina**, Sechenov First Moscow State Medical University. Address: Trubetskaya St., bld. 2, Moscow, 119991. ORCID: 0009-0004-9377-2241

9. The paper has not been submitted elsewhere

**Reference**

1. Gaudino M, Benedetto U, Fremes S, et al. Radial-Artery or Saphenous-Vein Grafts in Coronary-Artery Bypass Surgery. *N Engl J Med*. 2018;378(22):2069-2077. doi:10.1056/NEJMoa1716026

2. Leyvi G, Forest SJ, Srinivas VS, et al. Robotic Coronary Artery Bypass Grafting Decreases 30-Day Complication Rate, Length of Stay, and Acute Care Facility Discharge Rate Compared with Conventional Surgery. *Innov Technol Tech Cardiothorac Vasc Surg*. 2014;9(5):361-367. doi:10.1097/imi.0000000000000095

3. Pooria A, Pourya A, Gheini A. Postoperative complications associated with coronary artery bypass graft surgery and their therapeutic interventions. *Future Cardiol*. 2020;16(5):481-496. doi:10.2217/fca-2019-0049

4. Palmerini T, Savini C. Risks of Stroke After Coronary Artery Bypass Graft – Recent Insights and Perspectives. Di Eusanio M, ed. *Interv Cardiol Rev*. 2011;9(2):77. doi:10.15420/icr.2011.9.2.77

5. Sabik JF, Blackstone EH, Gillinov AM, Smedira NG, Lytle BW. Occurrence and Risk Factors for Reintervention After Coronary Artery Bypass Grafting. *Circulation*. 2006;114(1_supplement). doi:10.1161/CIRCULATIONAHA.105.001149

6. Montrief T, Koyfman A, Long B. Coronary artery bypass graft surgery complications: A review for emergency clinicians. *Am J Emerg Med*. 2018;36(12):2289-2297. doi:10.1016/j.ajem.2018.09.014

7. Brown PP, Kugelmass AD, Cohen DJ, et al. The Frequency and Cost of Complications

Associated With Coronary Artery Bypass Grafting Surgery: Results from the United States Medicare Program. *Ann Thorac Surg*. 2008;85(6):1980-1986. doi:10.1016/J.ATHORACSUR.2008.01.053

8. Roach GW, Kanchuger M, Mangano CM, et al. Adverse Cerebral Outcomes after Coronary Bypass Surgery. *N Engl J Med*. 1996;335(25):1857-1864. doi:10.1056/NEJM199612193352501

9. Palmerini T, Biondi-Zoccai G, Reggiani LB, et al. Risk of stroke with coronary artery bypass graft surgery compared with percutaneous coronary intervention. *J Am Coll Cardiol*. 2012;60(9):798-805. doi:10.1016/J.JACC.2011.10.912

10. Van Linden J Der, Hadjinikolaou L, Bergman P, Lindblom D. Postoperative stroke in cardiac surgery is related to the location and extent of atherosclerotic disease in the ascending aorta. *J Am Coll Cardiol*. 2001;38(1):131-135. doi:10.1016/S0735-1097(01)01328-6

11. Naylor AR, Mehta Z, Rothwell PM, Bell PRF. Carotid artery disease and stroke during coronary artery bypass: A critical review of the literature. *Eur J Vasc Endovasc Surg*. 2002;23(4):283-294. doi:10.1053/EJVS.2002.1609

12. Charlesworth DC, Likosky DS, Marrin CAS, et al. Development and validation of a prediction model for strokes after coronary artery bypass grafting. *Ann Thorac Surg*. 2003;76(2):436-443. doi:10.1016/S0003-4975(03)00528-9

13. Hornero F, Martín E, Rodríguez R, et al. A multicentre Spanish study for multivariate prediction of perioperative in-hospital cerebrovascular accident after coronary bypass surgery: The PACK2. *Interact Cardiovasc Thorac Surg*. 2013;17(2):353-358. doi:10.1093/ICVTS/IVT102

14. Mathisen L, Andersen MH, Hol PK, et al. Preoperative cerebral ischemic lesions predict physical health status after on-pump coronary artery bypass surgery. *J Thorac Cardiovasc Surg*. 2005;130(6):1691-1697. doi:10.1016/J.JTCVS.2005.08.008

15. Khan MF, Herle A, Movahed MR. Risk Factors for Post-Coronary Artery Bypass Grafting (CABG) Atrial Fibrillation and the Role of Aspirin and Beta Blockers in its Prevention. *J Atr Fibrillation*. 2013;5(5):800. doi:10.4022/JAFIB.800

16. Karkouti K, Cohen MM, McCluskey SA, Sher GD. A multivariable model for predicting the need for blood transfusion in patients undergoing first-time elective coronary bypass graft surgery. *Transfusion*. 2001;41(10):1193-1203. doi:10.1046/j.1537-2995.2001.41101193.x

17. Sellman M. Reoperations for bleeding after coronary artery bypass procedures during

25 years. *Eur J Cardio-Thoracic Surg*. 1997;11(3):521-527. doi:10.1016/S1010-7940(96)01111-6

18.     Horwich P, Buth KJ, Légaré JF. New onset postoperative atrial fibrillation is associated with a long-term risk for stroke and death following cardiac surgery. *J Card Surg*. 2013;28(1):8-13. doi:10.1111/JOCS.12033

19.     Amar D, Shi W, Hogue CW. Clinical prediction rule for atrial fibrillation after coronary artery bypass grafting. *ACC Curr J Rev*. 2004;13(12):50. doi:10.1016/J.ACCREVIEW.2004.11.027

20.     Villareal RP, Hariharan R, Liu BC, et al. Postoperative atrial fibrillation and mortality after coronary artery bypass surgery. *J Am Coll Cardiol*. 2004;43(5):742-748. doi:10.1016/J.JACC.2003.11.023

21.     McKhann GM, Grega MA, Borowicz LM, Baumgartner WA, Selnes OA. Stroke and encephalopathy after cardiac surgery: An update. *Stroke*. 2006;37(2):562-571. doi:10.1161/01.STR.0000199032.78782.6C

22.     Marzoog BA, Vlasova TI. Myocardiocyte autophagy in the context of myocardiocytes regeneration: a potential novel therapeutic strategy. *Egypt J Med Hum Genet*. 2022;23(1):41. doi:10.1186/s43042-022-00250-8

23.     Carnes CA, Chung MK, Nakayama T, et al. Ascorbate Attenuates Atrial Pacing-Induced Peroxynitrite Formation and Electrical Remodeling and Decreases the Incidence of Postoperative Atrial Fibrillation. *Circ Res*. 2001;89(6). doi:10.1161/hh1801.097644

24.     De Vecchi E, Pala MG, Di Credico G, et al. Relation between left ventricular function and oxidative stress in patients undergoing bypass surgery. *Heart*. 1998;79(3):242-247. doi:10.1136/hrt.79.3.242

25.     Dimmer C, Tavernier R, Gjorgov N, Van Nooten G, Clement DL, Jordaens L. Variations of autonomic tone preceding onset of atrial fibrillation after coronary artery bypass grafting. *Am J Cardiol*. 1998;82(1):22-25. doi:10.1016/S0002-9149(98)00231-8

26.     Kalman JM, Munawar M, Howes LG, et al. Atrial fibrillation after coronary artery bypass grafting is associated with sympathetic activation. *Ann Thorac Surg*. 1995;60(6):1709-1715. doi:10.1016/0003-4975(95)00718-0

27.     Budeus M, Hennersdorf M, Röhlen S, et al. Prediction of atrial fibrillation after coronary artery bypass grafting: The role of chemoreflexsensitivity and P wave signal averaged ECG. *Int J Cardiol*. 2006;106(1):67-74. doi:10.1016/j.ijcard.2004.12.062

28. Tsikouris JP, Kluger J, Song J, White CM. Changes in P-wave dispersion and P-wave duration after open heart surgery are associated with the peak incidence of atrial fibrillation. *Hear Lung*. 2001;30(6):466-471. doi:10.1067/mhl.2001.118363

29. Engelmann MDM, Svendsen JH. Inflammation in the genesis and perpetuation of atrial fibrillation. *Eur Heart J*. 2005;26(20):2083-2092. doi:10.1093/eurheartj/ehi350

30. Bruins P, Velthuis H te, Yazdanbakhsh AP, et al. Activation of the Complement System During and After Cardiopulmonary Bypass Surgery. *Circulation*. 1997;96(10):3542-3548. doi:10.1161/01.CIR.96.10.3542

31. Abdelhadi RH, Gurm HS, Van Wagoner DR, Chung MK. Relation of an exaggerated rise in white blood cells after coronary bypass or cardiac valve surgery to development of atrial fibrillation postoperatively. *Am J Cardiol*. 2004;93(9):1176-1178. doi:10.1016/j.amjcard.2004.01.053

32. Tomic V, Russwurm S, Mo□ller E, et al. Transcriptomic and Proteomic Patterns of Systemic Inflammation in On-Pump and Off-Pump Coronary Artery Bypass Grafting. *Circulation*. 2005;112(19):2912-2920. doi:10.1161/CIRCULATIONAHA.104.531152

33. Puskas JD, Edwards FH, Pappas PA, et al. Off-Pump Techniques Benefit Men and Women and Narrow the Disparity in Mortality After Coronary Bypass Grafting. *Ann Thorac Surg*. 2007;84(5):1447-1456. doi:10.1016/J.ATHORACSUR.2007.06.104

34. Afilalo J, Rasti M, Ohayon SM, Shimony A, Eisenberg MJ. Off-pump vs. on-pump coronary artery bypass surgery: An updated meta-analysis and meta-regression of randomized trials. *Eur Heart J*. 2012;33(10):1257-1267. doi:10.1093/EURHEARTJ/EHR307

35. Møller CH, Penninga L, Wetterslev J, Steinbrüchel DA, Gluud C. Off-pump versus on-pump coronary artery bypass grafting for ischaemic heart disease. *Cochrane Database Syst Rev*. Published online March 14, 2012. doi:10.1002/14651858.CD007224.PUB2

**Figure legend**

Figure 1: Receiver Operating Characteristic (ROC) Curve Analysis and Area Under the curve (AUC) analysis of the Sample.

Figure 2: Mean age of the patients who passed coronary artery shunt surgery (whole sample). At the same time this mean age shows average age of occurrence of indications for CABG, which is usually associated with the acute coronary syndrome development.

Figure 3: Reduction in the incidence rate of complications in the second group > 60 years.

Figure 4: With age, the number of shunts increases in the first group < 60 years.

Figure 5: The incidence rate of complications in the first (<60 years) and second group (> 60 years), respectively. (H; hydrothorax, R; resternotomy, A; arrythmia, P; psychosis, MI; myocardial infarction, D; dyspnea, DSI; deep sternum wound infection, SSI; superficial sternum wound infection)