Research article: Risk Factors for Traumatic Brain Injury in Men and Women

Running title: Recent Epidemiological Research Advances

Authors' list: **Basheer Abdullah Marzoog[1,*]**

[1] National Research Mordovia State University. Address: Bolshevitskaya Street, 68, Saransk,

Rep. Mordovia, 430005. Postal address: Mordovia republic, Saransk, Bolshevitskaya Street, 31.

[*]Corresponding author: Basheer Abdullah Marzoog, undergraduate medical school student at

National Research Mordovia State University (marzug@mail.ru, +79969602820). Address:

Bolshevitskaya Street, 68, Saransk, Rep. Mordovia, 430005. Postal address: Mordovia republic,

Saransk, Bolshevitskaya Street, 31. ORCID: 0000-0001-5507-2413, Scopus ID: 57486338800

Competing interests: No competing interests regarding the publication.

Abstract

Objectives: The study covers the current status of TBI and provides a statistical recommendation

to health organizations in the world.

Design: A retrospective analytical study. The descriptive results showed that a difference exists

between males and females and for these reasons the sample was divided into two groups

according to the primary descriptive statistics. The first group included 212 (31.59 %) females,

while the second group included 459 (68.40%) males.

Setting: Assess the risk factors for TBI in both sexes and compare them with the results of

international studies.

Participants: The study involved 671 patients for the period from 07/01/2017- 17/12/2018. The

primary data were collected from the republic hospital.

Interventions: Retrospectively analysed these patients using T test, one- and two-way ANOVA

test, and the Pearson correlation test using the Statistica 12 program.

Main Outcome Measures: The most frequently reported cause of TBI in men and women is

domestic accidents. Men are affected by TBI in early life compared to women.

Results: The mean age of the men is 44.41 years (Std. error 0.75). The mean age of the women

is 49.50 years (Std. error 1.38). In the male group, 354 (77.12 %) patients live in the city and 105

(22.87%) live in the village. However, in the female group, 170 (80.18%) patients live in the city

and 42 (19.81%) live in the village. In the men group, 172 (37.47%) patients had TBI due to

domestic accidents. The most frequently reported etiology of TBI in women is domestic

accident, reported in 122 (57.54 %). In female group, the mean age of patients with domestic

accident associated TBI is 53.76 years (Std. error 1.85). In male group, the mean age of patients

with domestic accident associated TBI is 50.74 years (Std. error 1.09). Total hospitalization days

of the patients were associated with the age of the patients, r= 0.12. Where men are hospitalized

longer than women, t value -2.261, p < 0.024. In the male and female groups, there is a direct

correlation between age and the total hospitalization days in the male group, r=0.173; r=0.148,

respectively.

Conclusion: The most frequently reported cause of TBI in men and women is domestic

accidents. Men are affected by TBI in early life compared to women.

Keywords: Traumatic brain Injury; Surgery; Risk factor; Gender; Epidemiology; Glasgow coma

scale

Background

Traumatic brain injury is reported in multiple studies, and it is suggested that gender plays an important role in post-TBI outcomes [1]. A recent multicenter study concluded that gender plays an important role in the severity of TBI using the Glasgow coma scale (GCS) score [2]. Traumatic brain injury can be contusion and concussion, and penetrating wound injury as well as anoxic brain injuries.

Several factors play a role in the determination of the risk of TBI in men and women, such as age, hormonal status, and the presence of poor health care practice [3,4]. The classical causes of TBI include car accidents, falls from hills, sport associated injuries, and domestic violence [5]. The domestic violence is the primary cause of TBI and post TBI outcomes and complications [6].

Traumatic brain injury clinical manifestations are severity-dependent and age-dependent. Various sensory symptoms, physical symptoms, and cognitive, behavioral, or mental symptoms [7].

The dramatic increase in the number of traumatic brain injuries exposes the community to a high prevalence of disability associated with TBI associated disability[8]. Therefore, TBI is an emergency and requires urgent action to reduce the incidence rate of TBI occurrence [9]. The study sought to determine the risk factors most frequently reported for TBI occurrences.

Methods

A retrospective analytical study involved 671 patients for the period of 07/01/2017- 17/12/2018. The primary data were collected from the republic hospital for the past 2 year and retrospectively analyzed. The descriptive results showed that a difference exists between males and females and for these reasons the sample was divided into two groups according to the primary descriptive statistics. The first group included 212 (31.59 %) females, while the second group included 459 (68.40%) males. The consent of the patients has been taken for scientific purposes to analyze and publish the results of the study.

For statistical analysis, the T test, one- and two-way ANOVA test, and the Pearson correlation test were used using the Statistica program. (StatSoft, Inc. (2011). STATISTICA (data analysis software system), version 10. www.statsoft.com.).

Results

The descriptive statistical analysis showed that the mean age of the participants is 46.016 years old (min; max: 15.00; 92.00). The mean hospitalization days ranged from 0.00 - 49.00 days (mean; Std. Error: 12.75; 0.26). The study involved 212 (31.59 %) women and 459 (68.40%) men. The demographic distribution of the sample included 524 (78.09 %) who live in the city and 147 (21.90%) who live in the village. The frequency of traumatic brain injury (TBI) due to alcohol-related brain injury is observed in 139 (20.71%) patients, due to criminal injury 75

(11.17%) patients, due to domestic accident 294 (43.81%) patients, due to work injury 16

(2.38%) patients, due to motor vehicle injury 111 (16.54 %) patients, due to postoperative

haematoma 2 (0.29%) patients, due to convulsive syndrome 7 (1.043%) patients, due to sports

injury 2 (0.29%) patients, due to unknown reason 25 (3.72%) patients. Number of the patients

required blood transfusion is 3 (0.44%) and 668 (99.55%) required only conservative therapy.

Of 671, the need for surgical treatment was observed in 198 (39.91%) patients, missing data in

181 (26.97%), and 292 (43,51%) did not require surgical intervention. Out of 198, 125 (63.13%)

patients required surgical debridement of wounds. 42 (21.32%) patients required scull

trepanation with removal of acute subdural hematoma. 6 (3.03 %) patients required scull

trepanation and removal of acute epidural haematoma. 8 (4.12%) patients required scull

trepanation and removal of chronic subdural hematoma. 4 (2.03 %) patients required scull

trepanation and removal of acute intracerebral haematoma. 3 (1.54 %) patients required scull

trepanation and osteoplasty of fracture. 3 (1.52 %) patients required osteoplasty of fracture. 4

(2.03%) patients required Bulau pleural drainage. 1 (0.50 %) patient required surgical

debridement of wounds and pleural drainage by Bulau. 2 (1.01%) patients required scull

trepanation and removal of acute subdural haematoma with removal of acute intracerebral

hematoma.

The mean age of patients affected by TBI who live in the city is 47.101 years. The mean age of

the patients affected by TBI who live in the village is 42.14 years. The people who live in the

village are at increased risk of TBI, t value 3.02971, p< 0.002.

Total hospitalization days of the patients were associated with the age of the patients, r= 0.12.

Where men are hospitalized longer than women, t value -2.26, p < 0.024.

Mean age of patients with alcohol-related brain injury is 46.079 years (Std. error 1.17). Mean age

of patients with domestic accident is 50.74 years (Std. error 1.098). Mean age of patients with

motor vehicle-related injury is 38.072 years (Std. error 1.68). The mean age of patients with

criminal injury is 37.94 years (Std. error 1.59). The mean age of patients with work injury is

41.75 years (Std. error 3.081). The mean age of patients with convulsive syndrome is 50.142

years (Std. error 5.65). The mean age of patients with postoperative haematoma is 55.00 years

(Std. error 2.00). Mean age of patients with sports injury is 26 years (Std. error 0.00). The mean

age for patients with TBI of unknown reasons is 52.04 years (Std. error 2.77).

In terms of gender-dependent risk factors, men are frequently exposed to domestic accidents as a

risk factor for the development of TBI. In the men group, 172 (37.47 %) patients had TBI due to

domestic accidents. However, the mean hospitalization days of these patients was 13.30 (Std.

error 0.56).

The mean age of the men is 44.41 years (Std. error 0.75). The mean age of the women is 49.50

years (Std. error 1.38). Indicates that men are affected by TBI at an early age compared to women, t value 3.50, p <0.0004. (Figure 2)

In the male group, 354 (77.12 %) patients live in the city and 105 (22.87%) live in the village. However, in the female group, 170 (80.18%) patients live in the city and 42 (19.81%) live in the village.

The most frequently reported etiology of TBI in women is domestic accident, reported in 122 (57.54 %). However, the most frequently reported cause of TBI in males is domestic accident too. (Figure 3)

In the male group, the frequency of domestic accident-associated TBI seen in 172(37.47 %) patients, alcohol-related brain injury-associated TBI in 122 (26.57 %) patients, motor vehicle-related injury associated TBI in 68 (14.81 %) patients, criminal injury associated TBI in 56 (12.20 %) patients, work injury associated TBI in 13 (2.83 %) patients, convulsive syndrome-associated TBI in 5 (1.089 %) patients, sports injury associated TBI in 2 (0.43%) patients, postoperative haematoma-associated TBI in 1 (0.21 %), and unknown reason in 20 (4.35 %) patients. In the female group, domestic accidents were reported in 122 (57.54 %) patients, motor vehicle injury reported in 43 (20.28%) patients, criminal injury reported in 19 (8.96%) patients, alcohol-related brain injury reported in 17 (8.018%) patients, work injury reported in 3 (1.41 %) patients, convulsive syndrome reported in 2 (0.94340 %) patients, postoperative hematoma reported in 1 (0.47%) patient and unknown reason seen in 5 (2.35 %) patients.

In the male group, 458 (99.78 %) patients required conservative therapy and 1 (0.21%) required blood transfusion. However, in the female group, 210 (99.05 %) required conservative surgery and 2 (0.94 %) were treated without conservative therapy.

In male group, the mean age of patients with alcohol-related brain injury is 46.079 years (Std. error 1.17). The mean age of patients with TBI associated with domestic accidents is 50.74 years (Std. error 1.098). The mean age of patients with motor vehicle injury-associated TBI is 38.072 years (Std. error 1.68). Mean age of patients with criminal injury associated TBI is 37.94 years (Std. error 1.59). Mean age of patients with work injury associated TBI is 41.75 years (Std. error 3.081). The mean age of patients with convulsive syndrome associated TBI is 50.14 years (Std. error 5.65). The mean age of patients with sports injury-associated TBI is 26 years (Std. error 0.00). The mean age of patients with postoperative haematoma-associated TBI is 57 years (Std. error 0.00). The mean age of patients with unknown reason associated TBI is 52.04 years (Std. error 2.77). In the female group, the mean age of patients with alcohol-related brain injury is 43.52 years (Std. error 3.63). The mean age of patients with domestic accident-associated TBI is 53.76 years (Std. error 1.85). Mean age of patients with motor vehicle-related injury associated TBI is 40.30 years (Std. error 3.069). Mean age of patients with criminal injury associated TBI is

37.94 years (Std. error 1.59). Mean age of patients with work injury associated TBI is 45.33 years (Std. error 4.17). The mean age of patients with convulsive syndrome associated TBI is 57.5 years (Std. error 5.50). The mean age of patients with sports injury-associated TBI is 26 years (Std. error 0.00). The mean age of patients with postoperative haematoma-associated TBI is 53 years (Std. error 0.00). The mean age of patients with unknown reason is 61.20 years (Std. error 6.39).

In male group, out of 459 patients with TBI, 158 (46.19%) required surgical treatment and 184 (53.80%) resolved without surgical treatment, missing data in 117 (25.54%) patients. Of 158 patients, 3 (1.89%) required fracture osteoplasty. 94 (59.49%) patients required surgical debridement of the wounds. 11 (6.96%) required scull trepanation and removal of chronic subdural haematoma. 1 (0.63%) required scull trepanation with removal of acute subdural hematoma. 38 (24.05%) required scull trepanation and removal of acute subdural hematoma. 2 (1.26%) required scull trepanation with fracture osteoplasty. 1 (0.63%) required surgical debridement of wounds with pleural drainage by Bulau. 5 (3.16%) required scull trepanation with removal of acute epidural haematoma. 3 (1.89%) required scull trepanation with removal of acute intracerebral haematoma. 2 (1.26%) required scull trepanation and removal of acute subdural haematoma, as well as removal of acute intracerebral hematoma. In the female group, out of 212 patients, 40 (18.87%) patients required surgical treatment, missing data in 64 (30.18 %) patients. Of 40, 31 (77.50%) patients required surgical debridement of wounds, 4 (10.00%) required scull trepanation with removal of acute subdural haematoma, 1 (2.50%) required scull trepanation with removal of acute epidural haematoma, 2 (5.00%) required scull trepanation with removal of chronic subdural haematoma, 1 (2.50%) required scull trepanation with removal of acute intracerebral haematoma, and 1 (2.50%) required scull trepanation with fracture osteoplasty. (*Table 1*)

Table 1: Non parametric characteristics of the sample.

Factor	Index	Group M ()	Group F ()	X^2	P value
Sex	M	459 (100.00 %)	0 (0.00 %)	671.000	<0,001
	F	0 (0.00 %)	212 (100.00%)		
Living place	City	354 (77,12 %)	170 (80,18 %)	0.796	0.373
	Village	105 (22,87 %)	42 (19,81 %)		
Conservative therapy	Yes	458 (99,78 %)	210 (99,05 %)	0.373	0.191
	Blood transfusion	1 (0,21 %)	2 (0,94 %)		

In the male group, the mean hospitalization days for patients with TBI who live in the city are

12.83 days and 14.29 days for patients who live in the village. In the female group, the mean

hospitalization day is 11.87 days (Std. error 0.40). However, in the female group, the mean days

for TBI patients living in the city are 11.79 days and 12.19 days for patients living in the village.

The Mean total hospitalization days of patients with alcohol-related brain injury is 12.00 days

(Std. error 1.85). The mean total hospitalization of patients with domestic accidents is 12.64 days

(Std. error 0.40). The mean total hospitalization of patients with motor vehicle-related injury is

13.53 days (Std. error 0.66). The mean total hospitalization of patients with criminal injury is

9.026 days (Std. error 0.61). The mean total hospitalization of patients with work injury is 12.81

days (Std. error 2.029). The mean total hospitalization of patients with convulsive syndrome is

15.71 days (Std. error 3.037). The mean total hospitalization of patients with sports injuries is 8.5

days (Std. error 3.50). The mean total hospitalization of patients with postoperative hematoma is

16.50 days (Std. error 3.50). The mean total hospitalization of patients for unknown reason is

15.00 days (Std. error 1.34).

In the male group, the mean total hospitalization days of patients with alcohol-related brain

injury is 14.09 days (Std. error 0.62). The mean total hospitalization of patients with domestic

accidents is 13.30 days (Std. error 0.56). The mean total hospitalization of patients with motor

vehicle-related injury is 13.92 days (Std. error 0.94). The mean total hospitalization of patients

with criminal injury is 9.17 days (Std. error 0.76). The mean total hospitalization of patients

with work injury is 12.30 days (Std. error 2.47). The mean total hospitalization of patients with

convulsive syndrome is 15.4 days (Std. error 4.38). The mean total hospitalization for sports

injuries is 8.5 days (standard error 3.5). The mean total hospitalization of patients with

postoperative haematoma is 8.5 days (Std. error 3.50). The mean total hospitalization of patients

for unknown reason is 15.4 days (Std. error 1.62). In the female group, the mean total

hospitalization days of patients with alcohol-related brain injury is 12.00 days (Std. error 1.85).

The mean total hospitalization of patients with domestic accident is 11.71 days (Std. error 0.55).

The mean total hospitalization of patients with motor vehicle-related injury is 12.90 days (Std.

error 0.85). The mean total hospitalization of patients with criminal injury is 8.57 days (Std.

0.85). The mean total hospitalization of patients with work injury is 15.00 days (Std. error 1.73).

The mean total hospitalization of patients with convulsive syndrome is 15.4 days (Std. error

4.38). The mean total hospitalization days of patients with sports injury is 8.5 days (Std. error

3.5). The mean total hospitalization of patients with postoperative haematoma is 8.5 days (Std.

error 3.50). The mean total hospitalization of patients for unknown reason is 13.40 days (Std.

error 1.80).

In the male group, the mean age of patients with TBI who live in the city is 44.85 years and

42.92 years for patients who live in the village. Whereas, in the female group, the mean age of

patients with TBI who lives in the city is 51.79 years old and 40.21 years old for patients living

in the village, statically significant difference in the age of the TBI female patients, t value 3.42,

p< 0.0007.

In the male and female groups, there is a direct correlation between age and the total

hospitalization days, r=0.17; r=0.14, respectively.

Discussion

The city population affected by TBI in older age in compare to village citizens. The total

hospitalization days are longer in men than in women. Longer hospitalization days are observed

in older people. Living in the village posses' higher risk of TBI, probably due to the lack of

primary health care facilities. Men are affected by TBI in early life compared to women. The

frequency of TBI causes is associated with the age of the patients. Consuming alcohol is a huge

risk factor for TBI. Poor hygienic practice and bad habits such as heavy alcohol drinking are

avoidable risk factors.

The limitation of a study is the absence of the control group to compare the incidence rate

between men and women. The outcomes of post TBI depend on the gender [2,10–18]. Different

studies suggested that age plays a central role in increasing the risk of TBI [19]. In particular,

women under 45 years of age and older than 65 years of age are at increased risk than men of the

same age [10]. Furthermore, being a male, the risk factor for TBI is two times higher as being a

woman. However, a noticeable increase in the incidence rate of TBI in women is observed

compared to previous studies [20,21]. Probably associated with women's increase in the

participation in different kinds of contact sports (such as rugby), military services, and home

violence [5,22–25].

Conclusions

The most frequently reported cause of TBI in men and women is domestic accidents. Men are

affected by TBI in early life compared to women. The people who live in the village are at

increased risk of TBI. Gender plays a crucial role in the severity and incidence rate of post-TBI

outcomes and complications. The results of our study expand the knowledge in the field of

traumatic brain injury risk factors in a gender-dependent manner. Also, the results of our study

are in constant with the results of the previous studies [15,17,26,27].

List of abbreviations

TBI; traumatic brain injury

1.   Ethics approval and consent to participate: applicable.

2.   Consent for publication: applicable on reasonable request.

3.   Availability of data and materials: The data are applicable.

4. Competing interests: The authors declare that they have no competing interests regarding the
publication.
5. Funding: Not applicable (This research did not receive any specific grant from funding
agencies in the public, commercial, or not-for-profit sectors).
6. Authors' contributions: MB collected data, analyzed the statistical data, wrote the draft, and
revised the final version of the paper. All authors have read and approved the manuscript.
7. Acknowledgments: none.
8. Authors' information: Basheer Abdullah Marzoog, medical school student at National
Research Mordovia State University (marzug@mail.ru, +79969602820). Address:
Bolshevitskaya st., 68, Saransk, Rep. Mordovia, 430005. ORCID: 0000-0001-5507-2413.
Scopus ID: 57486338800.
9. The paper has not been submitted elsewhere.
10. The study approved by the National Research Mordovia State University, Russia, from
"Ethics Committee Requirement N8/2 from 30.06.2021". Written informed consent was
obtained from the patients for publication of study results and any accompanying images.

References

1. Mac Donald CL, Barber J, Jordan M, Johnson AM, Dikmen S, Fann JR, et al. Early
clinical predictors of 5-year outcome after concussive blast traumatic brain injury. JAMA
Neurol. 2017;74:821–9.
2. Eom KS, Kim JH, Yoon SH, Lee S jong, Park KJ, Ha SK, et al. Gender differences in
adult traumatic brain injury according to the Glasgow coma scale: A multicenter
descriptive study. Chinese J. Traumatol. [Internet]. 2021 [cited 2022 Aug 10];24:333.
Available from: /pmc/articles/PMC8606602/
3. Ma C, Wu X, Shen X, Yang Y, Chen Z, Sun X, et al. Sex differences in traumatic brain
injury: a multi-dimensional exploration in genes, hormones, cells, individuals, and society.
Chinese Neurosurg. J. [Internet]. 2019;5:24. Available from:
https://cnjournal.biomedcentral.com/articles/10.1186/s41016-019-0173-8
4. Biegon A. Considering Biological Sex in Traumatic Brain Injury. Front. Neurol.
2021;12:39.
5. Monahan K, Goldfine A, Biegon A. Traumatic brain injuries in victims of intimate partner
violence: an underappreciated source of neurological morbidity. Future Neurol. [Internet].
2017;12:189–91. Available from: https://www.futuremedicine.com/doi/10.2217/fnl-2017-
0026
6. James SL, Theadom A, Ellenbogen RG, Bannick MS, Montjoy-Venning W, Lucchesi LR,
et al. Global, regional, and national burden of traumatic brain injury and spinal cord

304   injury, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016.

305   Lancet Neurol. [Internet]. 2019;18:56–87. Available from:

306   https://linkinghub.elsevier.com/retrieve/pii/S1474442218304150

307 7. McMahon P, Hricik A, Yue JK, Puccio AM, Inoue T, Lingsma HF, et al.

308   Symptomatology and functional outcome in mild traumatic brain injury: Results from the

309   prospective TRACK-TBI study. J. Neurotrauma. 2014;31:26–33.

310 8. Lasry O, Liu EY, Powell GA, Ruel-Laliberté J, Marcoux J, Buckeridge DL. Epidemiology

311   of recurrent traumatic brain injury in the general population. Neurology [Internet].

312   2017;89:2198–209. Available from:

313   https://www.neurology.org/lookup/doi/10.1212/WNL.0000000000004671

314 9. Albrecht JS, Hirshon JM, McCunn M, Bechtold KT, Rao V, Simoni-Wastila L, et al.

315   Increased Rates of Mild Traumatic Brain Injury Among Older Adults in US Emergency

316   Departments, 2009-2010. J. Head Trauma Rehabil. [Internet]. 2016;31:E1–7. Available

317   from: https://journals.lww.com/00001199-201609000-00008

318 10. Mikolić A, van Klaveren D, Oude Groeniger J, Wiegers EJA, Lingsma HF, Zeldovich M,

319   et al. Differences between Men and Women in Treatment and Outcome after Traumatic

320   Brain Injury. J. Neurotrauma [Internet]. 2020;neu.2020.7228. Available from:

321   https://www.liebertpub.com/doi/10.1089/neu.2020.7228

322 11. Gupte RP, Brooks WM, Vukas RR, Pierce JD, Harris JL. Sex Differences in Traumatic

323   Brain Injury: What We Know and What We Should Know. J. Neurotrauma [Internet].

324   2019;36:3063–91. Available from:

325   https://www.liebertpub.com/doi/10.1089/neu.2018.6171

326 12. Rosene JM, Raksnis B, Silva B, Woefel T, Visich PS, Dompier TP, et al. Comparison of

327   Concussion Rates Between NCAA Division I and Division III Men's and Women's Ice

328   Hockey Players. Am. J. Sports Med. [Internet]. 2017;45:2622–9. Available from:

329   http://journals.sagepub.com/doi/10.1177/0363546517710005

330 13. Colvin AC, Mullen J, Lovell MR, West RV, Collins MW, Groh M. The role of concussion

331   history and gender in recovery from soccer-related concussion. Am. J. Sports Med.

332   2009;37:1699–704.

333 14. Covassin T, Elbin RJ, Harris W, Parker T, Kontos A. The role of age and sex in

334   symptoms, neurocognitive performance, and postural stability in athletes after concussion.

335   Am. J. Sports Med. 2012;40:1303–12.

336 15. Styrke J, Sojka P, Björnstig U, Bylund PO, Stålnacke BM. Sex differences in symptoms,

337   disability and life satisfaction three years after mild traumatic brain injury: A

338   populationbased cohort study. J. Rehabil. Med. 2013;45:749–57.

16. Lippa SM, Brickell TA, Bailie JM, French LM, Kennedy JE, Lange RT. Postconcussion

Symptom Reporting after Mild Traumatic Brain Injury in Female Service Members:

Impact of Gender, Posttraumatic Stress Disorder, Severity of Injury, and Associated

Bodily Injuries. J. Head Trauma Rehabil. 2018;33:101–12.

17. Yue JK, Levin HS, Suen CG, Morrissey MR, Runyon SJ, Winkler EA, et al. Age and sex-

mediated differences in six-month outcomes after mild traumatic brain injury in young

adults: a TRACK-TBI study. Neurol. Res. 2019;41:609–23.

18. Yu W-Y, Hwang H-F, Lin M-R. Gender Differences in Personal and Situational Risk

Factors for Traumatic Brain Injury Among Older Adults. J. Head Trauma Rehabil.

[Internet]. 2022;37:220–9. Available from:

https://journals.lww.com/10.1097/HTR.0000000000000708

19. Varriano B, Tomlinson G, Tarazi A, Wennberg R, Tator C, Tartaglia MC. Age, Gender

and Mechanism of Injury Interactions in Post-Concussion Syndrome. Can. J. Neurol. Sci.

2018;45:636–42.

20. Roos KG, Wasserman EB, Dalton SL, Gray A, Djoko A, Dompier TP, et al.

Epidemiology of 3825 injuries sustained in six seasons of National Collegiate Athletic

Association men's and women's soccer (2009/2010–2014/2015). Br. J. Sports Med.

[Internet]. 2017;51:1029–34. Available from:

https://bjsm.bmj.com/lookup/doi/10.1136/bjsports-2015-095718

21. Gerritsen H, Samim M, Peters H, Schers H, Laar F van de. Incidence, course and risk

factors of head injury: a retrospective cohort study. BMJ Open [Internet].

2018;8:e020364. Available from: https://bmjopen.bmj.com/lookup/doi/10.1136/bmjopen-

2017-020364

22. Koreny M, Sterz F, Uray T, Schreiber W, Holzer M, Laggner A, et al. Effect of cooling

after human cardiac arrest on myocardial infarct size. Resuscitation. 2009;80:56–60.

23. Ma R, Lopez V, Weinstein MG, Chen JL, Black CM, Gupta AT, et al. Injury Profile of

American Women's Rugby-7s. Med. Sci. Sports Exerc. 2016;48:1957–66.

24. Spano M, Risucci DA, Etienne M, Petersen KH. Epidemiology of sports related

concussion in brazilian jiu-jitsu: A cross-sectional study. Sports. 2019;7.

25. Wu V, Huff H, Bhandari M. Pattern of physical injury associated with intimate partner

violence in women presenting to the emergency department: A systematic review and

meta-analysis. Trauma, Violence, Abus. 2010;11:71–82.

26. Preiss-Farzanegan SJ, Chapman B, Wong TM, Wu J, Bazarian JJ. The Relationship

Between Gender and Postconcussion Symptoms After Sport-Related Mild Traumatic

Brain Injury. PM R. 2009;1:245–53.

27. Bazarian JJ, Blyth B, Mookerjee S, He H, McDermott MP. Sex differences in outcome

after mild traumatic brain injury. J. Neurotrauma. 2010;27:527–39.

Figure legend

Figure 1: Total hospitalization days and etiology in men and women, respectively (in women

statically, no significant different).

Figure 2: Graphical presentation of the direct association between the total hospitalization days

and the age and sex of the patients.

Figure 3: Graphical presentation of the risk markers in males and females in the republic. (The number 1 indicates males and number 2 indicates females; the nearest values to 1 demonstrates higher incidence in males and oppositely, the value nearest to 2, indicates higher incidence in female).